

# Kainic Acid-Induced Neurotoxicity: Targeting Glial Responses and Glia-Derived Cytokines

Xing-Mei Zhang<sup>1</sup> and Jie Zhu<sup>1,2,\*</sup>

<sup>1</sup>Department of Neurobiology, Care Sciences and Society, Karolinska Institute, Stockholm, Sweden; <sup>2</sup>Department of Neurology, The First Hospital of Jilin University, Changchun, China

**Abstract:** Glutamate excitotoxicity contributes to a variety of disorders in the central nervous system, which is triggered primarily by excessive Ca<sup>2+</sup> influx arising from overstimulation of glutamate receptors, followed by disintegration of the endoplasmic reticulum (ER) membrane and ER stress, the generation and detoxification of reactive oxygen species as well as mitochondrial dysfunction, leading to neuronal apoptosis and necrosis. Kainic acid (KA), a potent agonist to the  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA)/kainate class of glutamate receptors, is 30-fold more potent in neurotoxicity than glutamate. In rodents, KA injection resulted in recurrent seizures, behavioral changes and subsequent degeneration of selective populations of neurons in the brain, which has been widely used as a model to study the mechanisms of neurodegenerative pathways induced by excitatory neurotransmitter. Microglial activation and astrocytes proliferation are the other characteristics of KA-induced neurodegeneration. The cytokines and other inflammatory molecules secreted by activated glia cells can modify the outcome of disease progression. Thus, anti-oxidant and anti-inflammatory treatment could attenuate or prevent KA-induced neurodegeneration. In this review, we summarized updated experimental data with regard to the KA-induced neurotoxicity in the brain and emphasized glial responses and glia-oriented cytokines, tumor necrosis factor- $\alpha$ , interleukin (IL)-1, IL-12 and IL-18.

**Keywords:** Kainic acid, excitotoxicity, microglia, astrocytes, cytokines.

## INTRODUCTION

Excitotoxicity mediated by glutamate receptors may underlay the pathology of a number of neurological abnormalities, including Huntington's chorea, Alzheimer's disease (AD), and Parkinson's disease (PD). Excitotoxic cell death is commonly induced experimentally by the administration of kainic acid (KA), a potent agonist to the  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA)/kainate class of glutamate receptors [1-3]. In rodents, injections of KA resulted in recurrent seizures, behavioral changes and subsequent degeneration of selective populations of neurons in the brain [4, 5]. Thus, administration of KA in rodents has been widely used as a model to study the mechanisms of neurodegenerative pathways induced by excitatory neurotransmitters.

## 1. OVERVIEW OF EXCITOTOXICITY INDUCED BY GLUTAMATE

L-glutamate, the major excitatory transmitter in the brain and spinal cord, is associated with learning, cognition, memory and neuro-endocrine functions [6, 7]. The glutamate receptors can be divided into two broad categories: the ionotropic receptors that mediate fast postsynaptic potentials by activating ion channels directly, and the metabotropic receptors that results in the expression of slow postsynaptic potentials through second messengers [8, 9]. The action of glutamate on the ionotropic receptors is always excitatory [10, 11]. There are three major subtypes of ionotropic

glutamate receptors: AMPA, kainate, and N-methyl-D-aspartate (NMDA), named according to the types of synthetic agonists that activate them, respectively [12, 13]. The NMDA glutamate receptor is blocked by specific antagonists such as D(-)-2-amino-5-phosphonovalerate (APV) [14, 15]. Both AMPA and kainate receptors are blocked by 6-cyano-7-nitroquinoxalin-2,3-dione (CNQX) [16, 17]. Thus the AMPA and kainate receptors are sometimes referred to together as non-NMDA receptors. The ion channel of NMDA receptor is a tetrameric structure that results from up to seven genes coding for seven subunits termed GluN1, GluN2A, GluN2B, GluN2C, GluN2D, GluN3A and GluN3B [18, 19]. The AMPA receptor family is composed of four subunits, GluA1-4 [20, 21]. The kainate receptor family comprises five genes, divided into two subfamilies, including GluK4-5 and GluK1-3. GluK4 and GluK5 exhibit higher affinity for kainate than GluK1-3 [22, 23]. Herein, we used the new nomenclature for glutamate receptors recommended by the International Union of Pharmacology Committee on Receptor Nomenclature and Drug Classification (NC-IUPHAR) [24]. NC-IUPHAR recommended and previous nomenclatures of ionotropic glutamate receptor subunits are listed in Table 1.

Excessive amounts of glutamate are highly toxic to neurons, an action termed glutamate excitotoxicity [25, 26]. Glutamate excitotoxicity is triggered primarily by excessive Ca<sup>2+</sup> influx arising from overstimulation of the NMDA subtype of glutamate receptors, followed by disintegration of the endoplasmic reticulum (ER) membrane and ER stress, the generation of reactive oxygen species (ROS) as well as mitochondrial dysfunction, leading to neuronal apoptosis and necrosis [27, 28]. There is increasing realization that the mitochon-

\*Address correspondence to this author at the Division of Neurodegeneration (NOVUM, plan 5), Department of NVS, Karolinska Institute, Karolinska University Hospital Huddinge, Stockholm, Sweden, SE-141 86; Tel: +46 8 58585494; Fax: +46 8 58585470; E-mail: Jie.Zhu@ki.se

**Table 1. NC-IUPHAR Recommended and Previous Nomenclatures of Ionotropic Glutamate Receptor Subunits**

<b>Ionotropic Glutamate Family</b>	<b>NC-IUPHAR Subunit Nomenclature</b>	<b>Previous Nomenclatures</b>	<b>Human Gene Name</b>	<b>Human Chromosomal Location</b>
NMDA	GluN1	GLU <sub>N1</sub> , NMDA-R1, NR1, GluR $\xi$ 1	GRIN1	9q34.3
	GluN2A	GLU <sub>N2A</sub> , NMDA-R2A, NR2A, GluR $\epsilon$ 1	GRIN2A	16p13.2
	GluN2B	GLU <sub>N2B</sub> , NMDA-R2B, NR2B, hNR3, GluR $\epsilon$ 2	GRIN2B	12p12
	GluN2C	GLU <sub>N2C</sub> , NMDA-R2C, NR2C, GluR $\epsilon$ 3	GRIN2C	17q25
	GluN2D	GLU <sub>N2D</sub> , NMDA-R2D, NR2D, GluR $\epsilon$ 4	GRIN2D	19q13.1
	GluN3A	GLU <sub>N3A</sub> , NMDA-R3A, NMDAR-L, chi-1	GRIN3A	9q31.1
	GluN3B	GLU <sub>N3B</sub> , NMDA-R3B,	GRIN3B	19p13.3
AMPA	GluA1	GLU <sub>A1</sub> , GluR1, GluRA, GluR-A, GluR-K1, HBGR1	GRIA1	5q31.1
	GluA2	GLU <sub>A2</sub> , GluR2, GluRB, GluR-B, GluR-K2, HBGR2	GRIA2	4q32-q33
	GluA3	GLU <sub>A3</sub> , GluR3, GluRC, GluR-C, GluR-K3	GRIA3	Xq25-q26
	GluA4	GLU <sub>A4</sub> , GluR4, GluRD, GluR-D	GRIA4	11q22
Kainate	GluK1	GLU <sub>K5</sub> , GluR5, GluR-5, EAA3	GRIK1	21q22.11
	GluK2	GLU <sub>K6</sub> , GluR6, GluR-6, EAA4	GRIK2	6q16.3-q21
	GluK3	GLU <sub>K7</sub> , GluR7, GluR-7, EAA5	GRIK3	1p34-p33
	GluK4	GLU <sub>K1</sub> , KA1, KA-1, EAA1	GRIK4	11q22.3
	GluK5	GLU <sub>K2</sub> , KA2, KA-2, EAA2	GRIK5	19q13.2

drial dysfunction occupies the center stage in these processes [28, 29]. In many cell types, glutamate neurotoxicity is induced by NMDA as well as non-NMDA receptors [25, 26, 30].

## 2. KA ADMINISTRATION TO RODENTS INDUCES SEIZURES, SELECTIVE NEURODEGENERATION AND BEHAVIORAL CHANGES

Kainic acid (KA) is a non-degradable analog of glutamate and 30-fold more potent in neurotoxicity than glutamate [31-33]. Administration of KA to rodents caused a well characterized seizure syndrome, as described by Ben-Ari and other research groups [34, 35]. The seizure activity caused by intravenous, intraperitoneal, intranasal injections or microinjection into amygdala or hippocampus is divided in several distinct phases. During the first 20-30 min, the animals have "staring" spells, followed by head nodding and numerous wet-dog shakes for another 30 min. One hour after KA administration, the animal starts recurrent limbic motor seizures, including masticatory and facial movements, forepaws tremor, rearing and loss of postural control. The seizures then become progressively severer, with a reduction in the intermission. In the following 1-2 h, the animal displays a full status epilepticus [34-36].

KA-induced damage seriously impact the hippocampus. The hippocampus is particularly vulnerable to KA-induced neurotoxicity due to the high density of kainate receptors [37]. The hilar neurons are sensitive to KA-induced neurotoxicity, but neuron loss in the other areas of the hippocampus differs between animal species and strains [38-40]. In

rats, the systemic injections of KA produced widespread neuronal death, primarily in the hippocampus hilus, CA1 and CA3 areas [30, 41]. Mouse strains vary significantly in their sensitivity to KA-induced neurodegeneration [39, 40, 42, 43]. In general, the C57BL/6, C57BL/10, and (C57BL/6 x CBA/J) F1 strains are resistant to KA-induced neurodegeneration, while the FVB/N, ICR and DBA/2 J strains are vulnerable [40]. C57BL/6, the "relatively" resistant mouse strain, reveals significant neuronal damage in CA1 and CA3, and to a lesser extent, in the polymorphic layer of the dentate gyrus 12 h post-treatment of KA systemically detected by cupric-silver and Fluoro-Jade B staining [44, 45]. CA3 region has the highest abundance of kainate receptors, the activation of which can elevate the concentration of ROS and impair the normal function of mitochondria [46-48]. CA3 neurons are directly excited by stimulation of their KA receptors and indirectly, by increased glutamate efflux secondary to KA stimulation of mossy fibers [49, 50]. CA3 synchroization produces spreading epileptiform activity that extends to CA1 and other limbic structures [51, 52] (Fig. 1).


CA1 pyramidal neurons receive two distinct excitatory inputs that are capable of influencing hippocampal output and involving in spatial memory and memory consolidation [53, 54]. Damage in CA1/CA3 regions of hippocampus induced by KA mainly results in the spatial learning deficits [55, 56]. KA-treated Wistar rats are impaired in the water maze and object exploration tasks, and hyperactive in the open field test, which can be improved by physical exercise and the selective cyclooxygenase (COX)-2 inhibitor [57, 58]. Intraperitoneal injections of KA into the developing rat

brains induce the impaired short-term spatial memory in the radial-arm maze, deficient long-term spatial learning and retrieval in the water maze, and a greater degree of anxiety in the elevated plus maze [59, 60]. Mice with a single unilateral injection of KA into the dorsal hippocampus exhibit a decrease in depression-like behavior in the forced swimming test and retarded acquisition as well as impaired retention of visual-spatial information in the Morris water maze test [61].


### 3. KA MEDIATES GENERATION OF OXIDATIVE STRESS

KA receptors have both presynaptic modulatory and direct postsynaptic excitatory actions [62, 63]. The activation of KA receptors produces membrane depolarization and results in alteration in intracellular calcium concentrations, which is required to trigger the neuronal death cascade (Fig. 2)

[64]. KA can also induce the release of lactate dehydrogenase (LDH), and a decrease in 3-(4, 5-dimethylthiazole-2-yl)-2, 5-diphenyl tetrazolium bromide (MTT), which result in damage of mitochondrial function [2]. KA administration increases the generation of ROS and reactive nitrogen species (RNS). There is growing evidence that free radical generation plays a key role in the neuronal damage [65]. KA has been shown to immediately induce COX-2 expression that might be involved in hippocampal neuronal death [66]. Early induced COX-2 facilitates the recurrence of hippocampal seizures, and late synthesized COX-2 stimulates hippocampal neuron loss after KA administration [67]. COX catalyzes the first step in the synthesis of prostanoids, including prostaglandins (PGs), prostacyclin, and thromboxanes. PGE(2) is pathologically increased in the brain after KA treatment, and has been proven to be closely associated with neuronal death [68]. In addition, lipid peroxides play critical



**Fig. (1). The Input and Output Pathways of Hippocampal Formation.** Entorhinal cortex (EC) is the main input to the hippocampus. EC projects to the dentate gyrus (DG) via perforant fiber pathway and provides the critical input to CA3 via mossy fiber pathway, then to CA1 by means of the Schaffer collateral pathway. Additionally, EC can also project directly to CA3, CA1 and subiculum (Sub). Meantime, EC is the major output of the hippocampus. Arrows denote the direction of impulse flow.




**Fig. (2). Schematic Overview of KA-Mediated Neuronal Death.** (1) By stimulating glutamate receptors (GluR), kainic acid (KA) elicits the increase of intracellular  $Ca^{2+}$ , activation of  $Ca^{2+}$ -dependent enzyme and production of free radicals; (2) Excessive  $Ca^{2+}$  and free radicals cause mitochondrial dysfunction, release of mitochondrial factors, activation of caspase-3, leading to neuronal apoptosis; (3) KA causes the disintegration of the endoplasmic reticulum (ER) and ER stress with the activation of the ER proteins Bip, Chop, and caspase-12, involved in neuronal apoptosis; (4)  $Ca^{2+}$  overload and excessive free radicals cause directly mitochondrial swelling, leading to neuronal necrosis. COX: cyclooxygenase; ROS: reactive oxygen species; RNS: reactive nitrogen species.

roles in the initiation and modulation of inflammation and oxidative stress upon KA insult. Seizures can induce the products of lipid peroxidation, such as F(2)-isoprostanes and Isofurans, which have been thought to be the reliable indices of oxidative stress *in vivo* [69]. Moreover, KA causes the disintegration of the ER membrane in hippocampal neurons and ER stress with the activation of the ER proteins Bip, Chop, and caspase-12 [70]. ER stress appears to act at an early stage of the cell death process prior to disruption of calcium homeostasis, excessive accumulation of ROS, and mitochondrial dysfunction [71]. Old astrocyte specifically induced substance (OASIS) is involved in the endoplasmic reticulum stress response [72]. A recent study showed that OASIS expressed in astrocytes plays an important role in protection against neuronal damage induced by KA [1].

#### 4. GLIAL CELLS ARE ACTIVATED UPON KA ADMINISTRATION

KA-induced neuronal death is accompanied by increased activation of microglia and astrocytes [73-75]. Additionally, the activated glial cells cluster at the hippocampal lesions and the immunostaining reactivity is particularly strong around areas of debris (Fig. 3).



**Fig. (3).** Glial cells activation accompanied the neuronal death 7 days after KA (45 mg/kg body weight) treatment to C57BL/6 mice. (A) Obvious neuronal loss was showed in CA3 area of hippocampus by Nissl's staining. (B) CD11b positive cells (microglia) accumulated in the lesioned CA3 area. (C) GFAP positive cells (astrocytes) spread the whole hippocampus, especially in CA3 area. Arrows in A indicate the areas of neuronal loss.

##### 4.1. Microglia

Microglia account for approximately 20% of the total glial population in the central nervous system (CNS). Microglia are the main effector cell type of the immune and inflammatory responses in the CNS, as earlier reviewed by

Streit and his colleagues [76]. The normal role of microglia could be partly connected to neuroprotection, whereas in pathological conditions microglia may become disease-promoting cells. Upon neuronal injury, microglia rapidly acquire changes in morphology and secrete a variety of soluble mediators [77, 78] (Fig. 4). Some studies suggested that the activated microglia might exert a neuroprotective function, especially in multiple sclerosis (MS) and its animal model, experimental autoimmune encephalomyelitis (EAE) by creating a microenvironment for reparative and regenerative processes [79]. Evidence is also accumulating that activated microglia induce and/or exacerbate neuropathological changes in several CNS diseases such as AD and PD through secreting proinflammatory and neurotoxic factors [80, 81]. In KA-induced hippocampal injury, microglial activation is generally believed to contribute to neuroinflammation and neurodegeneration [74, 82, 83]. A recent study showed that I $\kappa$ B kinase/nuclear factor kappa B (NF- $\kappa$ B) dependent microglial activation participated in KA-mediated injury *in vivo* through induction of inflammatory mediators [82]. However, whether microglial activation initiates the disease progression or merely responds to neuronal death is still unclear.


##### 4.2. Astrocytes

Astrocytes, the most numerous glia cells, have been regarded as passive supporters of neurons in CNS for decades. Studies of the last 20 years, however, challenged this assumption by demonstrating that astrocytes possess functional neurotransmitter receptors [84, 85]. These findings have led to a new concept of neuron-glia intercommunication where astrocytes play an undoubted dynamic role by integrating neuronal inputs and modulating synaptic activity, and so contribute to disease development [86]. Astrocytes have functional receptors for the excitatory neurotransmitter glutamate and respond to physiological concentrations of this substance with oscillations in intracellular Ca<sup>2+</sup> concentrations and spatially propagating Ca<sup>2+</sup> signals [87-89]. *In vitro* studies provided evidence that astrocytes can take up glutamate at synapses and release glutamate in a calcium-dependent manner [90]. A proliferative response of astrocytes at two days after KA treatment has been reported already in 1981 [91]. The expression of glial fibrillary acidic protein (GFAP) has been shown to steadily increase from one/three days up to one month after intra-hippocampal or intraperitoneal injection of KA [92, 93]. Astrogliosis induced by excitotoxicity has been considered as a marker for neurotoxicity [73, 94]. Activated astrocytes produce pro- and anti-inflammatory cytokines, chemokines, neurotrophic factors and other modulators to be involved in neuron-glia communication (Fig. 5). It is believed that astrocytes produce growth factors to prevent neurons from death and to promote proliferation and differentiation of precursor cells [95-97]. Activation of transcription factors, including nuclear factor erythroid-2-related factor 2 (Nrf2) and NF- $\kappa$ B, in astrocytes induces the neuroprotective molecule expression and confers protection to neighboring neurons [98-100].


#### 5. ALTERED CYTOKINE EXPRESSION AFFECTS KA-INDUCED INJURY

Altered expression of cytokines in response to brain injury has diverse actions that can exacerbate, mediate, reduce or inhibit neuronal damage and influence the disease devel-





**Fig. (4). The Categories of Molecules Produced by Activated Microglia.** IL: interleukin; TNF: tumor necrosis factor; TGF: transforming growth factor; IL-1ra: IL-1 receptor antagonist; MIP: macrophage inflammatory protein; MDC: macrophage-derived chemokine; MCP: monocyte-chemoattractant protein; RANTES: regulated on activation normal T cell expressed and secreted; IFN: interferon; IP: IFN-inducible protein; R: receptor; NGF: nerve growth factor; BDNF: brain-derived neurotrophic factor; NT: neurotrophin; MHC: major histocompatibility complex; CR: complement receptor; C1qRp: C1q receptor for phagocytosis enhancement; FasL: Fas ligand; PG: prostaglandin; COX: cyclooxygenase; NO: nitric oxide; ROS: reactive oxygen species; RNS: reactive nitrogen species; MMP: matrix metalloproteinase; NSAIDs: nonsteroidal antiinflammatory drugs.




**Fig. (5). The Categories of Molecules Produced by Activated Astrocytes.** ROS: reactive oxygen species; IL: interleukin; TNF: tumor necrosis factor; IFN: interferon; CSF: colony-stimulating factor; GM-CSF: granulocyte-macrophage CSF; TGF: transforming growth factor; MCP: monocyte-chemoattractant protein; RANTES: regulated on activation normal T cell expressed and secreted; IP: IFN-inducible protein; NGF: nerve growth factor; BDNF: brain-derived neurotrophic factor; bFGF: basic fibroblast growth factor; MMP: matrix metalloproteinase.

opment in a variety of CNS disorders, such as AD, MS, viral or bacterial infections, ischemia, stroke, and various forms of encephalopathies [101-105]. Cytokines can be divided into pro-inflammatory and anti-inflammatory cytokines, which play the neurodestructive and neuroprotective roles, respectively. It is the balance between destructive and protective factors that ultimately determines the net result of the neuro-immune and neuro-inflammation interaction, as reviewed by Kerschensteiner *et al.* [106]. Results from studies using KA model also indicated that cytokines are involved in neuron-glia intercommunication and manipulation of pro- and anti-inflammatory cytokines can modify the outcome with regard to the seizure activity, behavioral changes as well as the neuropathological consequences [75, 107-109] (Fig. 6). The roles of several important cytokines in the CNS disorders are reviewed here, including tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-1 (IL-1), IL-12 and IL-18.

### 5.1. TNF- $\alpha$

Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) is mainly produced by microglia and astrocytes in the CNS. Its functions are medi-

ated through two receptors, TNF receptor (TNFR) 1 (p55) and TNFR2 (p75), both of which are expressed on various cells types [110]. TNF- $\alpha$  over-expression participates in the pathogenesis of several CNS disorders, such as AD [111], bacterial meningitis [112], MS [113] and cerebral malaria [114]. TNF- $\alpha$  potentiates excitotoxic injury to human fetal brain cells [115]. In contrast to its well known deleterious roles, multiple lines of evidence suggested that TNF- $\alpha$  also exhibit neuroprotective properties. This implies an intricate biological function of TNF- $\alpha$  in modulating immune and inflammatory responses. TNF- $\alpha$  knockout worsens *Listeria* infection in the CNS [116] and TNF- $\alpha$  receptor knockout enhances the neuronal damage after excitotoxic [108, 117], ischemic [118] or traumatic injury [119]. Our study showed that mice lacking TNFR1 exhibited a more severe seizure activity, hippocampal neurodegeneration and increased microglial activation, suggesting that TNF- $\alpha$  plays its protective role through TNFR1 signaling [108], which is in agreement with a previous report [120]. Another study also proved that the protective roles of TNF- $\alpha$  in KA-induced neurodegeneration are *via* TNFR2 signaling [117]. Several neuroprotective



**Fig. (6). Cytokines Involved in Neuron-glia Intercommunication.** KA administration enhances further release of endogenous excitatory amino acids, activates microglia and astrocytes. Activated glial cells secrete inflammatory molecules, e.g. cytokines, chemokines, and neurotrophins to influence the outcome of neuronal damage. Glu/KA: glutamate/kainic acid; GluR: glutamate receptors; TGF: transforming growth factor; IL: interleukin; TNF: tumor necrosis factor.

molecules were identified as TNFR1 targets, including members of the Bcl-2 family, DNA repair machinery and cell cycle, developmental, and differentiation factors, neurotransmitters and growth factors, as well as their receptors [121]. The mechanisms by which TNF reduced neuron loss after brain injury may involve the up-regulation of proteins, such as neuronal apoptosis inhibitor protein (NAIP), which maintain calcium homeostasis and reduce free radical generation [122].

### 5.2. IL-1

IL-1 plays a pivotal role in the neuroinflammation that has been well addressed in KA induced neurodegenerative model [109, 123]. Systemic KA administration induced the expression of IL-1 $\beta$ , IL-1 receptor antagonist and IL-1 $\beta$  converting enzyme as early as 3 h, 12 h, and 24 h post-treatment, respectively, which is localized in the areas known to display neuronal and tissue damage upon excitotoxic lesions [124, 125]. Recombinant IL-1 receptor antagonist has dose- and region-dependent effects on neuronal survival after KA treatment and prevents damage-induced changes in amyloid precursor protein and GFAP mRNAs [126]. It has also been shown that IL-1 $\beta$  is activated in the cerebellum with systemic administration of KA, and its type I receptor (IL-1R) is expressed at a soma of cerebellar Purkinje cells [127]. The proconvulsive actions of IL-1 $\beta$  in the hippocampus may depend on the activation of a sphingomyelinase- and Src-family of kinases-dependent pathway which leads to the phosphorylation of the GluN2B subunit [128].


### 5.3. IL-12

IL-12 is a heterodimeric cytokine that consists of a heavy chain, p40, and a light chain, p35 [129]. In the CNS, microglia and astrocytes are the main source of producing IL-12 [130]. Human CNS-derived microglia produce IL-12 *in vitro* after activation with LPS and IFN- $\gamma$  [131]. Murine microglia can be induced to express mRNA encoding the IL-12 receptor (IL-12R) [132], indicating an autocrine regulation path-

way of IL-12 in these cells. IL-12 plays a critical role in several CNS diseases, such as MS and Borna disease. IL-12p40 is increased in cerebrospinal fluid and serum of MS patients [133]. Mice lacking IL-12p40 or administered with anti-IL-12p40 monoclonal antibodies are resistant to the development of EAE [134, 135]. In contrast, IL-12p35 deficient mice are susceptible to EAE [114]. Borna disease virus, which does not trigger disease in most strains, is harmful after infecting the CNS of mice that overexpress IL-12, suggesting an important role of IL-12 in viral infection of the CNS [136]. IL-12 is an active participant in excitotoxic brain injury as suggested by our previous observation that IL-12p35 deficiency alleviates KA-induced hippocampal neurodegeneration [107] and by another finding that IL-12 expression is reduced in hippocampi of transgenic mice protected against KA-induced excitotoxicity by metallothionein overexpression [137].

### 5.4. IL-18

Interleukin (IL)-18 is most closely related to IL-1 $\beta$ . The similarities between both cytokines comprise structure, receptor complex, and pro-inflammatory properties [138]. IL-18 serves as a link between innate and adaptive immune responses, such as stimulating the expression of adhesion molecules, inducing the production of chemokines (IL-8) and cytokines (TNF- $\alpha$  and IL-6), stimulating the activity of NK cells, and promoting T helper 1 (Th1) cells responses in combination with IL-12, Th2 responses in combination with IL-4, and Th17 responses in combination with IL-23 [139]. IL-18 and IL-18 receptor (IL-18R) mRNA have been found in brain tissue and in cultured astrocytes and microglia [140]. IL-18 enhanced postsynaptic AMPA receptor responses in CA1 pyramidal neurons *via* the release of glutamate, thereby facilitating basal hippocampal synaptic transmission [141]. IL-18 deficient mice showed a diminished microglial activation and reduced dopaminergic neuron loss after acute 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine treatment [142]. The roles of IL-18 in KA-induced model are controversial.



**Fig. (7). Schematic Illustration of Anti-Oxidant and Anti-inflammatory Treatments.** By inhibiting one of the major components of the neuroinflammatory response after KA treatment, there could be less inflammation and neuronal loss. The potential treatments include (1) cyclooxygenase (COX)-2 inhibitors and other antioxidants; (2) Phospholipase A2 inhibitors; (3) Free radical scavengers; (4) endoplasmic reticulum (ER) stress inhibitors; and (5) Glia-derived cytokines. Glu/KA: glutamate/kainic acid; GluR: glutamate receptors; ROS: reactive oxygen species; RNS: reactive nitrogen species.

Levels of IL-18 and IL-18R in hippocampus increase progressively from day 1 and peaked at day 3 post-KA treatment [143]. Interestingly, intracerebellar coinjection of IL-18 counteracts the effect of IL-1 $\beta$  in KA-induced ataxia in mice [144]. We showed that exogenous IL-18 administration aggravated the KA-induced injury in normal C57BL/6 mice, while in the condition of IL-18 deficiency, IL-12 could overcompensate the function of IL-18 and worsen the seizure activity as well as hippocampal neurodegeneration [75].

## 6. THERAPEUTIC STRATEGIES

Considering that oxidative stress is central to KA-induced excitotoxic damage, anti-oxidant and anti-inflammatory treatments may attenuate or prevent the KA-mediated neurodegeneration (Fig. 7). The potential role of COX-2 inhibitors as a new therapeutic drug for the neuron loss after KA treatment has been studied. The selective COX-2 inhibitors, celecoxib, NS398, rofecoxib and SC58125, can suppress an elevation of PGE (2) and block hippocampal cell death [57, 145]. The other drugs tested experimentally include curcumin, fluoxetine, ethyl pyruvate and statins, whose neuroprotective effects are associated with their anti-inflammatory and anti-oxidant effects [146]. Free radical scavengers are well known to prevent neuron loss induced by exposure to excitotoxins. Edaravone (Ed), a newly developed free radical scavenger, could inhibit lipid peroxidation and prevent neuron loss when administered after the onset of seizures in a KA-induced neurodegenerative animal model [147]. The pineal secretory product, melatonin, has free-radical-scavenger and antioxidant properties, which attenuates KA-induced neuronal death, lipid peroxidation, and microglial activation [148]. Several phospholipase A (2) inhibitors, quinacrine and chloroquine, arachidonyl trifluoromethyl ketone, bromoenol lactone, cytidine 5-diphosphoamines, and vitamin E, have been shown to prevent the neurodegeneration in KA-mediated neurotoxicity [149]. Moreover, inhibition of ER stress by small molecular

compounds, such as salubrinal, may benefit for treatment of KA-mediated neurotoxicity [70]. Furthermore, the microtubule interacting drug candidate NAP has been shown to protect against KA toxicity *via* regulating the expression of key genes involved in the epileptogenic pathway [150]. A novel approach to engineer patient derived adult stem cells for therapeutic adenosine delivery in autologous cell transplantation has been proved to suppress seizure activity and protect hippocampal neurons from KA-induced damage [151]. Additionally, targeting the pro-inflammatory cytokines, by blocking the unique signal transduction of the specific cytokine is another potential therapeutic strategy.

## 7. CONCLUDING REMARKS

Glutamate excitotoxicity contributes to a variety of CNS diseases, which is involved in overstimulation of glutamate receptors, excessive Ca<sup>2+</sup> influx, ER stress, generation of ROS and mitochondrial dysfunction, leading to neuronal damage. KA is a potent agonist to the AMPA/kainate class of glutamate receptors, which can result in seizures, behavioral changes and neurodegeneration in susceptible brain regions like the hippocampus in rodents.

The activated glia cells and subsequent secretion of inflammatory molecules can modify the outcome of disease progression. By inhibiting one of the major components of the neuroinflammatory response after KA treatment, there could be less inflammation and neuronal loss, therefore an improvement of cognitive function.

## ACKNOWLEDGEMENTS

The work was supported by grants from SADF (Insamlingsstiftelsen för Alzheimer- och Demensforskning) foundation, Swedish Medicine Association, Gamla Tjänarinnor foundation, Gun och Bertil Stohnes foundation and Swedish National Board of Health and Welfare.

## REFERENCES

- [1] Chihara, K.; Saito, A.; Murakami, T.; Hino, S.; Aoki, Y.; Sekiya, H.; Aikawa, Y.; Wanaka, A.; Imaizumi, K. Increased vulnerability of hippocampal pyramidal neurons to the toxicity of kainic acid in OASIS-deficient mice. *J. Neurochem.*, **2009**, *110*(3), 956-965.
- [2] Wang, Q.; Yu, S.; Simonyi, A.; Sun, G.Y.; Sun, A.Y. Kainic acid-mediated excitotoxicity as a model for neurodegeneration. *Mol. Neurobiol.*, **2005**, *31*(1-3), 3-16.
- [3] Yang, D.D.; Kuan, C.Y.; Whitmarsh, A.J.; Rincon, M.; Zheng, T.S.; Davis, R.J.; Rakic, P.; Flavell, R.A. Absence of excitotoxicity-induced apoptosis in the hippocampus of mice lacking the Jnk3 gene. *Nature*, **1997**, *389*(6653), 865-870.
- [4] McKhann, G.M., 2nd; Wenzel, H.J.; Robbins, C.A.; Sosunov, A.A.; Schwartzkroin, P.A. Mouse strain differences in kainic acid sensitivity, seizure behavior, mortality, and hippocampal pathology. *Neuroscience*, **2003**, *122*(2), 551-561.
- [5] Tripathi, P.P.; Sgado, P.; Scali, M.; Viaggi, C.; Casarosa, S.; Simon, H.H.; Vaglini, F.; Corsini, G.U.; Bozzi, Y. Increased susceptibility to kainic acid-induced seizures in Engrailed-2 knockout mice. *Neuroscience*, **2009**, *159*(2), 842-849.
- [6] Herron, C.E.; Lester, R.A.; Coan, E.J.; Collingridge, G.L. Frequency-dependent involvement of NMDA receptors in the hippocampus: a novel synaptic mechanism. *Nature*, **1986**, *322*(6076), 265-268.
- [7] Zahr, N.M.; Mayer, D.; Pfefferbaum, A.; Sullivan, E.V. Low striatal glutamate levels underlie cognitive decline in the elderly: evidence from *in vivo* molecular spectroscopy. *Cereb. Cortex*, **2008**, *18*(10), 2241-2250.
- [8] Bashir, Z.I.; Bortolotto, Z.A.; Davies, C.H.; Berretta, N.; Irving, A.J.; Seal, A.J.; Henley, J.M.; Jane, D.E.; Watkins, J.C.; Collingridge, G.L. Induction of LTP in the hippocampus needs synaptic activation of glutamate metabotropic receptors. *Nature*, **1993**, *363*(6427), 347-350.
- [9] Rainnie, D.G.; Holmes, K.H.; Shinnick-Gallagher, P. Activation of postsynaptic metabotropic glutamate receptors by trans-ACPD hyperpolarizes neurons of the basolateral amygdala. *J. Neurosci.*, **1994**, *14*(11 Pt 2), 7208-7220.
- [10] Stephens, G.J.; Djamgoz, M.B.; Wilkin, G.P. A patch clamp study of excitatory amino acid effects on cortical astrocyte subtypes in culture. *Receptors Channels*, **1993**, *1*(1), 39-52.
- [11] Sugiyama, H.; Ito, I.; Hirono, C. A new type of glutamate receptor linked to inositol phospholipid metabolism. *Nature*, **1987**, *325*(6104), 531-533.
- [12] Loscher, W.; Lehmann, H.; Behl, B.; Seemann, D.; Teschendorf, H.J.; Hofmann, H.P.; Lubisch, W.; Hoyer, T.; Lemaire, H.G.; Gross, G. A new pyrrolyl-quinolinedione series of non-NMDA glutamate receptor antagonists: pharmacological characterization and comparison with NBQX and valproate in the kindling model of epilepsy. *Eur. J. Neurosci.*, **1999**, *11*(1), 250-262.
- [13] Trist, D.G. Excitatory amino acid agonists and antagonists: pharmacology and therapeutic applications. *Pharm. Acta. Helv.*, **2000**, *74*(2-3), 221-229.
- [14] Choi, D.W.; Koh, J.Y.; Peters, S. Pharmacology of glutamate neurotoxicity in cortical cell culture: attenuation by NMDA antagonists. *J. Neurosci.*, **1988**, *8*(1), 185-196.
- [15] Davies, J.; Evans, R.H.; Jones, A.W.; Smith, D.A.; Watkins, J.C. Differential activation and blockade of excitatory amino acid receptors in the mammalian and amphibian central nervous systems. *Comp. Biochem. Physiol. C.*, **1982**, *72*(2), 211-224.
- [16] Evstratova, A.A.; Mironova, E.V.; Dvoretzkova, E.A.; Antonov, S.M. Apoptosis and the receptor specificity of its mechanisms during the neurotoxic action of glutamate. *Neurosci. Behav. Physiol.*, **2009**, *39*(4), 353-362.
- [17] Koh, J.Y.; Goldberg, M.P.; Hartley, D.M.; Choi, D.W. Non-NMDA receptor-mediated neurotoxicity in cortical culture. *J. Neurosci.*, **1990**, *10*(2), 693-705.
- [18] Laube, B.; Kuhse, J.; Betz, H. Evidence for a tetrameric structure of recombinant NMDA receptors. *J. Neurosci.*, **1998**, *18*(8), 2954-2961.
- [19] Salter, M.G.; Fern, R. NMDA receptors are expressed in developing oligodendrocyte processes and mediate injury. *Nature*, **2005**, *438*(7071), 1167-1171.
- [20] Bochet, P.; Audinat, E.; Lambollez, B.; Crepel, F.; Rossier, J. Analysis of AMPA receptor subunits expressed by single Purkinje cells using RNA polymerase chain reaction. *Biochem. Soc. Trans.*, **1993**, *21*(1), 93-97.
- [21] Kwok, K.H.; Tse, Y.C.; Wong, R.N.; Yung, K.K. Cellular localization of GluR1, GluR2/3 and GluR4 glutamate receptor subunits in neurons of the rat neostriatum. *Brain Res.*, **1997**, *778*(1), 43-55.
- [22] Niedzielski, A.S.; Wenthold, R.J. Expression of AMPA, kainate, and NMDA receptor subunits in cochlear and vestibular ganglia. *J. Neurosci.*, **1995**, *15*(3 Pt 2), 2338-2353.
- [23] Raymond, L.A.; Blackstone, C.D.; Haganir, R.L. Phosphorylation and modulation of recombinant GluR6 glutamate receptors by cAMP-dependent protein kinase. *Nature*, **1993**, *361*(6413), 637-641.
- [24] Collingridge, G.L.; Olsen, R.W.; Peters, J.; Spedding, M. A nomenclature for ligand-gated ion channels. *Neuropharmacology*, **2009**, *56*(1), 2-5.
- [25] Choi, D.W. Ionic dependence of glutamate neurotoxicity. *J. Neurosci.*, **1987**, *7*(2), 369-379.
- [26] Frandsen, A.; Drejer, J.; Schousboe, A. Direct evidence that excitotoxicity in cultured neurons is mediated via N-methyl-D-aspartate (NMDA) as well as non-NMDA receptors. *J. Neurochem.*, **1989**, *53*(1), 297-299.
- [27] Nicholls, D.G. Mitochondrial dysfunction and glutamate excitotoxicity studied in primary neuronal cultures. *Curr. Mol. Med.*, **2004**, *4*(2), 149-177.
- [28] Schinder, A.F.; Olson, E.C.; Spitzer, N.C.; Montal, M. Mitochondrial dysfunction is a primary event in glutamate neurotoxicity. *J. Neurosci.*, **1996**, *16*(19), 6125-6133.
- [29] Brunet, N.; Tarabal, O.; Esquerda, J.E.; Caldero, J. Excitotoxic motoneuron degeneration induced by glutamate receptor agonists and mitochondrial toxins in organotypic cultures of chick embryo spinal cord. *J. Comp. Neurol.*, **2009**, *516*(4), 277-290.
- [30] Morales-Garcia, J.A.; Luna-Medina, R.; Martinez, A.; Santos, A.; Perez-Castillo, A. Anticonvulsant and neuroprotective effects of the novel calcium antagonist NP04634 on kainic acid-induced seizures in rats. *J. Neurosci. Res.*, **2009**, *87*, 3687-3697.
- [31] Lee, J.K.; Won, J.S.; Singh, A.K.; Singh, I. Statin inhibits kainic acid-induced seizure and associated inflammation and hippocampal cell death. *Neurosci. Lett.*, **2008**, *440*(3), 260-264.
- [32] McGeer, E.G.; McGeer, P.L. Some factors influencing the neurotoxicity of intrastriatal injections of kainic acid. *Neurochem. Res.*, **1978**, *3*(4), 501-517.
- [33] Olney, J.W.; Rhee, V.; Ho, O.L. Kainic acid: a powerful neurotoxic analogue of glutamate. *Brain Res.*, **1974**, *77*(3), 507-512.
- [34] Ben-Ari, Y. Limbic seizure and brain damage produced by kainic acid: mechanisms and relevance to human temporal lobe epilepsy. *Neuroscience*, **1985**, *14*(2), 375-403.
- [35] Mulle, C.; Sailer, A.; Perez-Otano, I.; Dickinson-Anson, H.; Castillo, P.E.; Bureau, I.; Maron, C.; Gage, F.H.; Mann, J.R.; Bettler, B.; Heinemann, S.F. Altered synaptic physiology and reduced susceptibility to kainate-induced seizures in GluR6-deficient mice. *Nature*, **1998**, *392*(6676), 601-605.
- [36] Chuang, Y.C.; Chang, A.Y.; Lin, J.W.; Hsu, S.P.; Chan, S.H. Mitochondrial dysfunction and ultrastructural damage in the hippocampus during kainic acid-induced status epilepticus in the rat. *Epilepsia*, **2004**, *45*(10), 1202-1209.
- [37] Darstein, M.; Petralia, R.S.; Swanson, G.T.; Wenthold, R.J.; Heinemann, S.F. Distribution of kainate receptor subunits at hippocampal mossy fiber synapses. *J. Neurosci.*, **2003**, *23*(22), 8013-8019.
- [38] Cantalops, I.; Routtenberg, A. Kainic acid induction of mossy fiber sprouting: dependence on mouse strain. *Hippocampus*, **2000**, *10*(3), 269-273.
- [39] Kasugai, M.; Akaike, K.; Imamura, S.; Matsukubo, H.; Tojo, H.; Nakamura, M.; Tanaka, S.; Sano, A. Differences in two mice strains on kainic acid-induced amygdalar seizures. *Biochem. Biophys. Res. Commun.*, **2007**, *357*(4), 1078-1083.
- [40] McLin, J.P.; Steward, O. Comparison of seizure phenotype and neurodegeneration induced by systemic kainic acid in inbred, outbred, and hybrid mouse strains. *Eur. J. Neurosci.*, **2006**, *24*(8), 2191-2202.
- [41] Lee, H.N.; Jeon, G.S.; Kim, D.W.; Cho, I.H.; Cho, S.S. Expression of adenomatous polyposis coli protein in reactive astrocytes in hippocampus of kainic acid-induced rat. *Neurochem. Res.*, **2010**, *35*(1), 114-121.
- [42] Schauwecker, P.E. Genetic basis of kainate-induced excitotoxicity in mice: phenotypic modulation of seizure-induced cell death. *Epilepsy Res.*, **2003**, *55*(3), 201-210.
- [43] Yang, J.; Houk, B.; Shah, J.; Hauser, K.F.; Luo, Y.; Smith, G.; Schauwecker, E.; Barnes, G.N. Genetic background regulates



- semaphorin gene expression and epileptogenesis in mouse brain after kainic acid status epilepticus. *Neuroscience*, **2005**, *131*(4), 853-869.
- [44] Benkovic, S.A.; O'Callaghan, J.P.; Miller, D.B. Sensitive indicators of injury reveal hippocampal damage in C57BL/6J mice treated with kainic acid in the absence of tonic-clonic seizures. *Brain Res.*, **2004**, *1024*(1-2), 59-76.
- [45] Benkovic, S.A.; O'Callaghan, J.P.; Miller, D.B. Regional neuropathology following kainic acid intoxication in adult and aged C57BL/6J mice. *Brain Res.*, **2006**, *1070*(1), 215-231.
- [46] Carriedo, S.G.; Sensi, S.L.; Yin, H.Z.; Weiss, J.H. AMPA exposures induce mitochondrial Ca(2+) overload and ROS generation in spinal motor neurons *in vitro*. *J. Neurosci.*, **2000**, *20*(1), 240-250.
- [47] Lauri, S.E.; Bortolotto, Z.A.; Bleakman, D.; Ornstein, P.L.; Lodge, D.; Isaac, J.T.; Collingridge, G.L. A critical role of a facilitatory presynaptic kainate receptor in mossy fiber LTP. *Neuron*, **2001**, *32*(4), 697-709.
- [48] Reynolds, I.J.; Hastings, T.G. Glutamate induces the production of reactive oxygen species in cultured forebrain neurons following NMDA receptor activation. *J. Neurosci.*, **1995**, *15*(5 Pt 1), 3318-3327.
- [49] Contractor, A.; Sailer, A.W.; Darstein, M.; Maron, C.; Xu, J.; Swanson, G.T.; Heinemann, S.F. Loss of kainate receptor-mediated heterosynaptic facilitation of mossy-fiber synapses in KA2-/- mice. *J. Neurosci.*, **2003**, *23*(2), 422-429.
- [50] Rodriguez-Moreno, A.; Sihra, T.S. Presynaptic kainate receptor facilitation of glutamate release involves protein kinase A in the rat hippocampus. *J. Physiol.*, **2004**, *557*(Pt 3), 733-745.
- [51] Bausch, S.B.; McNamara, J.O. Contributions of mossy fiber and CA1 pyramidal cell sprouting to dentate granule cell hyperexcitability in kainic acid-treated hippocampal slice cultures. *J. Neurophysiol.*, **2004**, *92*(6), 3582-3595.
- [52] Ding, R.; Asada, H.; Obata, K. Changes in extracellular glutamate and GABA levels in the hippocampal CA3 and CA1 areas and the induction of glutamic acid decarboxylase-67 in dentate granule cells of rats treated with kainic acid. *Brain Res.*, **1998**, *800*(1), 105-113.
- [53] Iijima, T.; Witter, M.P.; Ichikawa, M.; Tominaga, T.; Kajiwara, R.; Matsumoto, G. Entorhinal-hippocampal interactions revealed by real-time imaging. *Science*, **1996**, *272*(5265), 1176-1179.
- [54] Speed, H.E.; Dobrunz, L.E. Developmental changes in short-term facilitation are opposite at temporoammonic synapses compared to Schaffer collateral synapses onto CA1 pyramidal cells. *Hippocampus*, **2009**, *19*(2), 187-204.
- [55] Gayoso, M.J.; Primo, C.; al-Majdalawi, A.; Fernandez, J.M.; Garrosa, M.; Iniguez, C. Brain lesions and water-maze learning deficits after systemic administration of kainic acid to adult rats. *Brain Res.*, **1994**, *653*(1-2), 92-100.
- [56] Milgram, N.W.; Isen, D.A.; Mandel, D.; Palantzas, H.; Pepkowski, M.J. Deficits in spontaneous behavior and cognitive function following systemic administration of kainic acid. *Neurotoxicology*, **1988**, *9*(4), 611-624.
- [57] Gobbo, O.L.; O'Mara, S.M. Post-treatment, but not pre-treatment, with the selective cyclooxygenase-2 inhibitor celecoxib markedly enhances functional recovery from kainic acid-induced neurodegeneration. *Neuroscience*, **2004**, *125*(2), 317-327.
- [58] Gobbo, O.L.; O'Mara, S.M. Exercise, but not environmental enrichment, improves learning after kainic acid-induced hippocampal neurodegeneration in association with an increase in brain-derived neurotrophic factor. *Behav. Brain Res.*, **2005**, *159*(1), 21-26.
- [59] Sarkisian, M.R.; Tandon, P.; Liu, Z.; Yang, Y.; Hori, A.; Holmes, G.L.; Stafstrom, C.E. Multiple kainic acid seizures in the immature and adult brain: ictal manifestations and long-term effects on learning and memory. *Epilepsia*, **1997**, *38*(11), 1157-1166.
- [60] Sayin, U.; Sutula, T.P.; Stafstrom, C.E. Seizures in the developing brain cause adverse long-term effects on spatial learning and anxiety. *Epilepsia*, **2004**, *45*(12), 1539-1548.
- [61] Groticke, I.; Hoffmann, K.; Loscher, W. Behavioral alterations in a mouse model of temporal lobe epilepsy induced by intrahippocampal injection of kainate. *Exp. Neurol.*, **2008**, *213*(1), 71-83.
- [62] Campbell, S.L.; Mathew, S.S.; Hablitz, J.J. Pre- and postsynaptic effects of kainate on layer II/III pyramidal cells in rat neocortex. *Neuropharmacology*, **2007**, *53*(1), 37-47.
- [63] Youn, D.H.; Randic, M. Modulation of excitatory synaptic transmission in the spinal substantia gelatinosa of mice deficient in the kainate receptor GluR5 and/or GluR6 subunit. *J. Physiol.*, **2004**, *555*(Pt 3), 683-698.
- [64] Brorson, J.R.; Manzolillo, P.A.; Miller, R.J. Ca<sup>2+</sup> entry via AMPA/KA receptors and excitotoxicity in cultured cerebellar Purkinje cells. *J. Neurosci.*, **1994**, *14*(1), 187-197.
- [65] Ueda, Y.; Yokoyama, H.; Nakajima, A.; Tokumaru, J.; Doi, T.; Mitsuyama, Y. Glutamate excess and free radical formation during and following kainic acid-induced status epilepticus. *Exp. Brain Res.*, **2002**, *147*(2), 219-226.
- [66] Kim, E.J.; Lee, J.E.; Kwon, K.J.; Lee, S.H.; Moon, C.H.; Baik, E.J. Differential roles of cyclooxygenase isoforms after kainic acid-induced prostaglandin E(2) production and neurodegeneration in cortical and hippocampal cell cultures. *Brain Res.*, **2001**, *908*(1), 1-9.
- [67] Takemiya, T.; Matsumura, K.; Yamagata, K. Roles of prostaglandin synthesis in excitotoxic brain diseases. *Neurochem. Int.*, **2007**, *51*(2-4), 112-120.
- [68] Kawaguchi, K.; Hickey, R.W.; Rose, M.E.; Zhu, L.; Chen, J.; Graham, S.H. Cyclooxygenase-2 expression is induced in rat brain after kainate-induced seizures and promotes neuronal death in CA3 hippocampus. *Brain Res.*, **2005**, *1050*(1-2), 130-137.
- [69] Patel, M.; Liang, L.P.; Hou, H.; Williams, B.B.; Kmiec, M.; Swartz, H.M.; Fessel, J.P.; Roberts, L.J., 2nd. Seizure-induced formation of isofurans: novel products of lipid peroxidation whose formation is positively modulated by oxygen tension. *J. Neurochem.*, **2008**, *104*(1), 264-270.
- [70] Sokka, A.L.; Putkonen, N.; Mudo, G.; Pryazhnikov, E.; Reijonen, S.; Khiroug, L.; Belluardo, N.; Lindholm, D.; Korhonen, L. Endoplasmic reticulum stress inhibition protects against excitotoxic neuronal injury in the rat brain. *J. Neurosci.*, **2007**, *27*(4), 901-908.
- [71] Yu, Z.; Luo, H.; Fu, W.; Mattson, M.P. The endoplasmic reticulum stress-responsive protein GRP78 protects neurons against excitotoxicity and apoptosis: suppression of oxidative stress and stabilization of calcium homeostasis. *Exp. Neurol.*, **1999**, *155*(2), 302-314.
- [72] Saito, A.; Hino, S.; Murakami, T.; Kondo, S.; Imaizumi, K. A novel ER stress transducer, OASIS, expressed in astrocytes. *Antioxid. Redox Signal.*, **2007**, *9*(5), 563-571.
- [73] Chen, Z.; Duan, R.S.; Quezada, H.C.; Mix, E.; Nennesmo, I.; Adem, A.; Winblad, B.; Zhu, J. Increased microglial activation and astrogliosis after intranasal administration of kainic acid in C57BL/6 mice. *J. Neurobiol.*, **2005**, *62*(2), 207-218.
- [74] Ravizza, T.; Rizzi, M.; Perego, C.; Richichi, C.; Veliskova, J.; Moshe, S.L.; De Simoni, M.G.; Vezzani, A. Inflammatory response and glia activation in developing rat hippocampus after status epilepticus. *Epilepsia*, **2005**, *46*(Suppl 5), 113-117.
- [75] Zhang, X.M.; Duan, R.S.; Chen, Z.; Quezada, H.C.; Mix, E.; Winblad, B.; Zhu, J. IL-18 deficiency aggravates kainic acid-induced hippocampal neurodegeneration in C57BL/6 mice due to an overcompensation by IL-12. *Exp. Neurol.*, **2007**, *205*(1), 64-73.
- [76] Streit, W.J.; Graeber, M.B.; Kreutzberg, G.W. Functional plasticity of microglia: a review. *Glia*, **1988**, *1*(5), 301-307.
- [77] Kato, H.; Walz, W. The initiation of the microglial response. *Brain Pathol.*, **2000**, *10*(1), 137-143.
- [78] Kreutzberg, G.W. Microglia: a sensor for pathological events in the CNS. *Trends Neurosci.*, **1996**, *19*(8), 312-318.
- [79] Napoli, I.; Neumann, H. Protective effects of microglia in multiple sclerosis. *Exp. Neurol.*, **2010**, *225*(1), 24-28.
- [80] Marinova-Mutafchieva, L.; Sadeghian, M.; Broom, L.; Davis, J.B.; Medhurst, A.D.; Dexter, D.T. Relationship between microglial activation and dopaminergic neuronal loss in the substantia nigra: a time course study in a 6-hydroxydopamine model of Parkinson's disease. *J. Neurochem.*, **2009**, *110*(3), 966-975.
- [81] Venneti, S.; Wiley, C.A.; Kofler, J. Imaging microglial activation during neuroinflammation and Alzheimer's disease. *J. Neuroimmune Pharmacol.*, **2009**, *4*(2), 227-243.
- [82] Cho, I.H.; Hong, J.; Suh, E.C.; Kim, J.H.; Lee, H.; Lee, J.E.; Lee, S.; Kim, C.H.; Kim, D.W.; Jo, E.K.; Lee, K.E.; Karin, M.; Lee, S.J. Role of microglial IKKbeta in kainic acid-induced hippocampal neuronal cell death. *Brain*, **2008**, *131*(Pt 11), 3019-3033.
- [83] Penkowa, M.; Molinero, A.; Carrasco, J.; Hidalgo, J. Interleukin-6 deficiency reduces the brain inflammatory response and increases oxidative stress and neurodegeneration after kainic acid-induced seizures. *Neuroscience*, **2001**, *102*(4), 805-818.
- [84] Hosli, L.; Hosli, E. Receptors for dopamine and serotonin on astrocytes of cultured rat central nervous system. *J. Physiol. (Paris)*, **1987**, *82*(4), 191-195.

- [85] Nedergaard, M. Direct signaling from astrocytes to neurons in cultures of mammalian brain cells. *Science*, **1994**, 263(5154), 1768-1771.
- [86] Vesce, S.; Rossi, D.; Brambilla, L.; Volterra, A. Glutamate release from astrocytes in physiological conditions and in neurodegenerative disorders characterized by neuroinflammation. *Int. Rev. Neurobiol.*, **2007**, 82, 57-71.
- [87] Dani, J.W.; Smith, S.J. The triggering of astrocytic calcium waves by NMDA-induced neuronal activation. *Ciba Found Symp.*, **1995**, 188, 195-205; discussion 205-199.
- [88] van den Pol, A.N.; Finkbeiner, S.M.; Cornell-Bell, A.H. Calcium excitability and oscillations in suprachiasmatic nucleus neurons and glia *in vitro*. *J. Neurosci.*, **1992**, 12(7), 2648-2664.
- [89] Yagodin, S.; Holtzclaw, L.A.; Russell, J.T. Subcellular calcium oscillators and calcium influx support agonist-induced calcium waves in cultured astrocytes. *Mol. Cell Biochem.*, **1995**, 149-150, 137-144.
- [90] Simard, M.; Nedergaard, M. The neurobiology of glia in the context of water and ion homeostasis. *Neuroscience*, **2004**, 129(4), 877-896.
- [91] Murabe, Y.; Ibata, Y.; Sano, Y. Morphological studies on neuroglia. IV. Proliferative response of non-neuronal elements in the hippocampus of the rat to kainic acid-induced lesions. *Cell Tissue Res.*, **1982**, 222(1), 223-226.
- [92] Bendotti, C.; Guglielmetti, F.; Tortarolo, M.; Samanin, R.; Hirst, W.D. Differential expression of S100beta and glial fibrillary acidic protein in the hippocampus after kainic acid-induced lesions and mossy fiber sprouting in adult rat. *Exp. Neurol.*, **2000**, 161(1), 317-329.
- [93] Ding, M.; Haglid, K.G.; Hamberger, A. Quantitative immunohistochemistry on neuronal loss, reactive gliosis and BBB damage in cortex/striatum and hippocampus/amygdala after systemic kainic acid administration. *Neurochem. Int.*, **2000**, 36(4-5), 313-318.
- [94] Torre, E.R.; Lothman, E.; Steward, O. Glial response to neuronal activity: GFAP-mRNA and protein levels are transiently increased in the hippocampus after seizures. *Brain Res.*, **1993**, 631(2), 256-264.
- [95] Braun, A.; Dang, J.; Johann, S.; Beyer, C.; Kipp, M. Selective regulation of growth factor expression in cultured cortical astrocytes by neuro-pathological toxins. *Neurochem. Int.*, **2009**, 55(7), 610-618.
- [96] Dakubo, G.D.; Beug, S.T.; Mazerolle, C.J.; Thurig, S.; Wang, Y.; Wallace, V.A. Control of glial precursor cell development in the mouse optic nerve by sonic hedgehog from retinal ganglion cells. *Brain Res.*, **2008**, 1228, 27-42.
- [97] Sandhu, J.K.; Gardaneh, M.; Iwasiew, R.; Lanthier, P.; Gangaraju, S.; Ribocco-Lutkiewicz, M.; Tremblay, R.; Kiuchi, K.; Sikorska, M. Astrocyte-secreted GDNF and glutathione antioxidant system protect neurons against 6OHDA cytotoxicity. *Neurobiol. Dis.*, **2009**, 33(3), 405-414.
- [98] Lerner-Natoli, M.; Montpied, P.; Rousset, M.C.; Bockaert, J.; Rondouin, G. Sequential expression of surface antigens and transcription factor NF-kappaB by hippocampal cells in excitotoxicity and experimental epilepsy. *Epilepsy Res.*, **2000**, 41(2), 141-154.
- [99] Vargas, M.R.; Johnson, D.A.; Sirkis, D.W.; Messing, A.; Johnson, J.A. Nrf2 activation in astrocytes protects against neurodegeneration in mouse models of familial amyotrophic lateral sclerosis. *J. Neurosci.*, **2008**, 28(50), 13574-13581.
- [100] Vargas, M.R.; Johnson, J.A. The Nrf2-ARE cytoprotective pathway in astrocytes. *Expert Rev. Mol. Med.*, **2009**, 11, e17.
- [101] Allan, S.M.; Rothwell, N.J. Cytokines and acute neurodegeneration. *Nat. Rev. Neurosci.*, **2001**, 2(10), 734-744.
- [102] Basic Kes, V.; Simundic, A.M.; Nikolac, N.; Topic, E.; Demarin, V. Pro-inflammatory and anti-inflammatory cytokines in acute ischemic stroke and their relation to early neurological deficit and stroke outcome. *Clin. Biochem.*, **2008**, 41(16-17), 1330-1334.
- [103] Bennett, J.L.; Stuve, O. Update on inflammation, neurodegeneration, and immunoregulation in multiple sclerosis: therapeutic implications. *Clin. Neuropharmacol.*, **2009**, 32(3), 121-132.
- [104] Rojo, L.E.; Fernandez, J.A.; Maccioni, A.A.; Jimenez, J.M.; Maccioni, R.B. Neuroinflammation: implications for the pathogenesis and molecular diagnosis of Alzheimer's disease. *Arch. Med. Res.*, **2008**, 39(1), 1-16.
- [105] Shiraiishi, M.; Ichiyama, T.; Matsushige, T.; Iwaki, T.; Iyoda, K.; Fukuda, K.; Makata, H.; Matsubara, T.; Furukawa, S. Soluble tumor necrosis factor receptor 1 and tissue inhibitor of metalloproteinase-1 in hemolytic uremic syndrome with encephalopathy. *J. Neuroimmunol.*, **2008**, 196(1-2), 147-152.
- [106] Kerschensteiner, M.; Meinl, E.; Hohlfeld, R. Neuro-immune crosstalk in CNS diseases. *Results Probl. Cell Differ.*, **2010**, 51, 197-216.
- [107] Chen, Z.; Duan, R.S.; Q, H.C.; Wu, Q.; Mix, E.; Winblad, B.; Ljunggren, H.G.; Zhu, J. IL-12p35 deficiency alleviates kainic acid-induced hippocampal neurodegeneration in C57BL/6 mice. *Neurobiol. Dis.*, **2004**, 17(2), 171-178.
- [108] Lu, M.O.; Zhang, X.M.; Mix, E.; Quezada, H.C.; Jin, T.; Zhu, J.; Adem, A. TNF-alpha receptor 1 deficiency enhances kainic acid-induced hippocampal injury in mice. *J. Neurosci. Res.*, **2008**, 86(7), 1608-1614.
- [109] Oprica, M.; Eriksson, C.; Schultzberg, M. Inflammatory mechanisms associated with brain damage induced by kainic acid with special reference to the interleukin-1 system. *J. Cell Mol. Med.*, **2003**, 7(2), 127-140.
- [110] Ware, C.F.; Crowe, P.D.; Vanarsdale, T.L.; Andrews, J.L.; Grayson, M.H.; Jerzy, R.; Smith, C.A.; Goodwin, R.G. Tumor necrosis factor (TNF) receptor expression in T lymphocytes. Differential regulation of the type I TNF receptor during activation of resting and effector T cells. *J. Immunol.*, **1991**, 147(12), 4229-4238.
- [111] Zhao, M.; Cribbs, D.H.; Anderson, A.J.; Cummings, B.J.; Su, J.H.; Wasserman, A.J.; Cotman, C.W. The induction of the TNFalpha death domain signaling pathway in Alzheimer's disease brain. *Neurochem. Res.*, **2003**, 28(2), 307-318.
- [112] Leist, T.P.; Frei, K.; Kam-Hansen, S.; Zinkernagel, R.M.; Fontana, A. Tumor necrosis factor alpha in cerebrospinal fluid during bacterial, but not viral, meningitis. Evaluation in murine model infections and in patients. *J. Exp. Med.*, **1988**, 167(5), 1743-1748.
- [113] Raine, C.S. Multiple sclerosis: TNF revisited, with promise. *Nat. Med.*, **1995**, 1(3), 211-214.
- [114] Grau, G.E.; Pigué, P.F.; Vassalli, P.; Lambert, P.H. Tumor necrosis factor and other cytokines in cerebral malaria: experimental and clinical data. *Immunol. Rev.*, **1989**, 112, 49-70.
- [115] Chao, C.C.; Hu, S. Tumor necrosis factor-alpha potentiates glutamate neurotoxicity in human fetal brain cell cultures. *Dev. Neurosci.*, **1994**, 16(3-4), 172-179.
- [116] Rothe, J.; Lesslauer, W.; Lotscher, H.; Lang, Y.; Koebel, P.; Kontgen, F.; Althage, A.; Zinkernagel, R.; Steinmetz, M.; Bluethmann, H. Mice lacking the tumour necrosis factor receptor 1 are resistant to TNF-mediated toxicity but highly susceptible to infection by *Listeria monocytogenes*. *Nature*, **1993**, 364(6440), 798-802.
- [117] Balosso, S.; Ravizza, T.; Perego, C.; Peschon, J.; Campbell, I.L.; De Simoni, M.G.; Vezzani, A. Tumor necrosis factor-alpha inhibits seizures in mice via p75 receptors. *Ann. Neurol.*, **2005**, 57(6), 804-812.
- [118] Bruce, A.J.; Boling, W.; Kindy, M.S.; Peschon, J.; Kraemer, P.J.; Carpenter, M.K.; Holtsberg, F.W.; Mattson, M.P. Altered neuronal and microglial responses to excitotoxic and ischemic brain injury in mice lacking TNF receptors. *Nat. Med.*, **1996**, 2(7), 788-794.
- [119] Sullivan, P.G.; Bruce-Keller, A.J.; Rabchevsky, A.G.; Christakos, S.; Clair, D.K.; Mattson, M.P.; Scheff, S.W. Exacerbation of damage and altered NF-kappaB activation in mice lacking tumor necrosis factor receptors after traumatic brain injury. *J. Neurosci.*, **1999**, 19(15), 6248-6256.
- [120] Gary, D.S.; Bruce-Keller, A.J.; Kindy, M.S.; Mattson, M.P. Ischemic and excitotoxic brain injury is enhanced in mice lacking the p55 tumor necrosis factor receptor. *J. Cereb. Blood Flow Metab.*, **1998**, 18(12), 1283-1287.
- [121] Taoufik, E.; Petit, E.; Divoux, D.; Tseveleki, V.; Mengozzi, M.; Roberts, M.L.; Valable, S.; Ghezzi, P.; Quackenbush, J.; Brines, M.; Cerami, A.; Probert, L. TNF receptor 1 sensitizes neurons to erythropoietin- and VEGF-mediated neuroprotection after ischemic and excitotoxic injury. *Proc. Natl. Acad. Sci. U S A*, **2008**, 105(16), 6185-6190.
- [122] Thompson, C.; Gary, D.; Mattson, M.; Mackenzie, A.; Robertson, G.S. Kainic acid-induced naip expression in the hippocampus is blocked in mice lacking TNF receptors. *Brain Res. Mol. Brain Res.*, **2004**, 123(1-2), 126-131.
- [123] Simi, A.; Tsakiri, N.; Wang, P.; Rothwell, N.J. Interleukin-1 and inflammatory neurodegeneration. *Biochem. Soc. Trans.*, **2007**, 35(Pt 5), 1122-1126.
- [124] Eriksson, C.; Van Dam, A.M.; Lucassen, P.J.; Bol, J.G.; Winblad, B.; Schultzberg, M. Immunohistochemical localization of interleukin-1beta, interleukin-1 receptor antagonist and interleukin-1beta converting enzyme/caspase-1 in the rat brain after peripheral administration of kainic acid. *Neuroscience*, **1999**, 93(3), 915-930.

- [125] Vezzani, A.; Conti, M.; De Luigi, A.; Ravizza, T.; Moneta, D.; Marchesi, F.; De Simoni, M.G. Interleukin-1beta immunoreactivity and microglia are enhanced in the rat hippocampus by focal kainate application: functional evidence for enhancement of electrographic seizures. *J. Neurosci.*, **1999**, *19*(12), 5054-5065.
- [126] Panegyres, P.K.; Hughes, J. The neuroprotective effects of the recombinant interleukin-1 receptor antagonist rhIL-1ra after excitotoxic stimulation with kainic acid and its relationship to the amyloid precursor protein gene. *J. Neurol. Sci.*, **1998**, *154*(2), 123-132.
- [127] Motoki, K.; Kishi, H.; Hori, E.; Tajiri, K.; Nishijo, H.; Muraguchi, A. The direct excitatory effect of IL-1beta on cerebellar Purkinje cell. *Biochem. Biophys. Res. Commun.*, **2009**, *379*(3), 665-668.
- [128] Balosso, S.; Maroso, M.; Sanchez-Alavez, M.; Ravizza, T.; Frasca, A.; Bartfai, T.; Vezzani, A. A novel non-transcriptional pathway mediates the proconvulsive effects of interleukin-1beta. *Brain*, **2008**, *131*(Pt 12), 3256-3265.
- [129] Kobayashi, M.; Fitz, L.; Ryan, M.; Hewick, R.M.; Clark, S.C.; Chan, S.; Loudon, R.; Sherman, F.; Perussia, B.; Trinchieri, G. Identification and purification of natural killer cell stimulatory factor (NKSF), a cytokine with multiple biologic effects on human lymphocytes. *J. Exp. Med.*, **1989**, *170*(3), 827-845.
- [130] Stalder, A.K.; Pagenstecher, A.; Yu, N.C.; Kincaid, C.; Chiang, C.S.; Hobbs, M.V.; Bloom, F.E.; Campbell, I.L. Lipopolysaccharide-induced IL-12 expression in the central nervous system and cultured astrocytes and microglia. *J. Immunol.*, **1997**, *159*(3), 1344-1351.
- [131] Becher, B.; Dodelet, V.; Fedorowicz, V.; Antel, J.P. Soluble tumor necrosis factor receptor inhibits interleukin 12 production by stimulated human adult microglial cells *in vitro*. *J. Clin. Invest.*, **1996**, *98*(7), 1539-1543.
- [132] Suzumura, A.; Sawada, M.; Takayanagi, T. Production of interleukin-12 and expression of its receptors by murine microglia. *Brain Res.*, **1998**, *787*(1), 139-142.
- [133] Fassbender, K.; Ragoschke, A.; Rossol, S.; Schwartz, A.; Mielke, O.; Paulig, A.; Hennerici, M. Increased release of interleukin-12p40 in MS: association with intracerebral inflammation. *Neurology*, **1998**, *51*(3), 753-758.
- [134] Brok, H.P.; van Meurs, M.; Blezer, E.; Schantz, A.; Peritt, D.; Treacy, G.; Laman, J.D.; Bauer, J.; Hart, B.A. Prevention of experimental autoimmune encephalomyelitis in common marmosets using an anti-IL-12p40 monoclonal antibody. *J. Immunol.*, **2002**, *169*(11), 6554-6563.
- [135] Gran, B.; Zhang, G.X.; Yu, S.; Li, J.; Chen, X.H.; Ventura, E.S.; Kamoun, M.; Rostami, A. IL-12p35-deficient mice are susceptible to experimental autoimmune encephalomyelitis: evidence for redundancy in the IL-12 system in the induction of central nervous system autoimmune demyelination. *J. Immunol.*, **2002**, *169*(12), 7104-7110.
- [136] Freude, S.; Hausmann, J.; Hofer, M.; Pham-Mitchell, N.; Campbell, I.L.; Staeheli, P.; Pagenstecher, A. Borna disease virus accelerates inflammation and disease associated with transgenic expression of interleukin-12 in the central nervous system. *J. Virol.*, **2002**, *76*(23), 12223-12232.
- [137] Penkowa, M.; Florit, S.; Giralt, M.; Quintana, A.; Molinero, A.; Carrasco, J.; Hidalgo, J. Metallothionein reduces central nervous system inflammation, neurodegeneration, and cell death following kainic acid-induced epileptic seizures. *J. Neurosci. Res.*, **2005**, *79*(4), 522-534.
- [138] Dinarello, C.A. IL-18: A TH1-inducing, proinflammatory cytokine and new member of the IL-1 family. *J. Allergy Clin. Immunol.*, **1999**, *103*(1 Pt 1), 11-24.
- [139] Arend, W.P.; Palmer, G.; Gabay, C. IL-1, IL-18, and IL-33 families of cytokines. *Immunol. Rev.*, **2008**, *223*, 20-38.
- [140] Conti, B.; Park, L.C.; Calingasan, N.Y.; Kim, Y.; Kim, H.; Bae, Y.; Gibson, G.E.; Joh, T.H. Cultures of astrocytes and microglia express interleukin 18. *Brain Res. Mol. Brain Res.*, **1999**, *67*(1), 46-52.
- [141] Kanno, T.; Nagata, T.; Yamamoto, S.; Okamura, H.; Nishizaki, T. Interleukin-18 stimulates synaptically released glutamate and enhances postsynaptic AMPA receptor responses in the CA1 region of mouse hippocampal slices. *Brain Res.*, **2004**, *1012*(1-2), 190-193.
- [142] Sugama, S.; Wirz, S.A.; Barr, A.M.; Conti, B.; Bartfai, T.; Shibasaki, T. Interleukin-18 null mice show diminished microglial activation and reduced dopaminergic neuron loss following acute 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine treatment. *Neuroscience*, **2004**, *128*(2), 451-458.
- [143] Jeon, G.S.; Park, S.K.; Park, S.W.; Kim, D.W.; Chung, C.K.; Cho, S.S. Glial expression of interleukin-18 and its receptor after excitotoxic damage in the mouse hippocampus. *Neurochem. Res.*, **2008**, *33*(1), 179-184.
- [144] Andoh, T.; Kishi, H.; Motoki, K.; Nakanishi, K.; Kuraishi, Y.; Muraguchi, A. Protective effect of IL-18 on kainate- and IL-1 beta-induced cerebellar ataxia in mice. *J. Immunol.*, **2008**, *180*(4), 2322-2328.
- [145] Takemiya, T.; Maehara, M.; Matsumura, K.; Yasuda, S.; Sugiura, H.; Yamagata, K. Prostaglandin E2 produced by late induced COX-2 stimulates hippocampal neuron loss after seizure in the CA3 region. *Neurosci. Res.*, **2006**, *56*(1), 103-110.
- [146] Gupta, Y.K.; Briyal, S.; Sharma, M. Protective effect of curcumin against kainic acid induced seizures and oxidative stress in rats. *Indian J. Physiol. Pharmacol.*, **2009**, *53*(1), 39-46.
- [147] Miyamoto, R.; Shimakawa, S.; Suzuki, S.; Ogihara, T.; Tamai, H. Edaravone prevents kainic acid-induced neuronal death. *Brain Res.*, **2008**, *1209*, 85-91.
- [148] Chung, S.Y.; Han, S.H. Melatonin attenuates kainic acid-induced hippocampal neurodegeneration and oxidative stress through microglial inhibition. *J. Pineal Res.*, **2003**, *34*(2), 95-102.
- [149] Farooqui, A.A.; Ong, W.Y.; Horrocks, L.A. Inhibitors of brain phospholipase A2 activity: their neuropharmacological effects and therapeutic importance for the treatment of neurologic disorders. *Pharmacol. Rev.*, **2006**, *58*(3), 591-620.
- [150] Zemlyak, I.; Manley, N.; Vulih-Shultzman, I.; Cutler, A.B.; Graber, K.; Sapolsky, R.M.; Gozes, I. The microtubule interacting drug candidate NAP protects against kainic acid toxicity in a rat model of epilepsy. *J. Neurochem.*, **2009**, *111*(5), 1252-1263.
- [151] Ren, G.; Li, T.; Lan, J.Q.; Wilz, A.; Simon, R.P.; Boison, D. Lenti-viral RNAi-induced downregulation of adenosine kinase in human mesenchymal stem cell grafts: a novel perspective for seizure control. *Exp. Neurol.*, **2007**, *208*(1), 26-37.