

Published in final edited form as:

Carbohydr Res. 2011 September 6; 346(12): 1551–1563. doi:10.1016/j.carres.2011.04.045.

Synthesis and antigenicity of BBGL-2 glycolipids of *Borrelia burgdorferi*, the causative agent of Lyme disease

Vince Pozsgay^{a,*}, Joanna Kubler-Kielb^a, Bruce Coxon^a, Adriana Marques^b, John B. Robbins^a, and Rachel Schneerson^a

^aProgram on Developmental and Molecular Immunity, Eunice Kennedy Shriver National Institute of Child Health and Human Development, Bethesda, MD 20892-2423, United States

^bLaboratory of Clinical Infectious Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892-2423, United States

Abstract

Borrelia burgdorferi is the etiological agent for Lyme disease (LD), the most common vector borne disease in the United States. There is no human vaccine against LD currently available. Our approach to a vaccine is based on its surface-exposed glycolipids. One group of these glycolipids termed BBGL-2 consists of 1,2-di-*O*-acyl-3-*O*-(α -D-galactopyranosyl)-*sn*-glycerol congeners having palmitic, oleic, stearic, linoleic, and myristic acids. In order to delineate the immunodominant region(s) of the BBGL-2 components, we embarked on a synthetic project to provide available structurally defined, homogeneous analogs of BBGL-2 that might help identify the best vaccine candidate. The antigenicity of the synthetic glycolipids was examined by dot-blot analysis using mice sera obtained by immunization with killed *B. burgdorferi* cells, with native BBGL-2 in complete Freund's adjuvant, as well as sera obtained from patients with Lyme disease. We found that the presence of two acyl groups in the glycerol moiety was essential for antigenicity. At least one of these groups must be an oleoyl moiety. Neither the anomeric configuration of the galactose nor the configuration of the glycerol at C-2 was a decisive factor. Based on these findings we designed an 'unnatural' BBGL-2 analog having the structure 3-*O*-(β -D-galactopyranosyl)-1,2-di-*O*-oleoyl-DL-glycerol which is easier and less expensive to synthesize than the other BBGL-2 congeners prepared in this study. This substance proved to be antigenic and is considered a candidate vaccine for Lyme disease.

Keywords

Antigen; *Borrelia burgdorferi*; Vaccine; Diacylglycerol; Dot-blot; Glycolipid

1. Introduction

Lyme disease is a multisystem infection caused by the spirochete *Borrelia burgdorferi* that can involve the skin, nervous system, heart, and joints.¹ It is transmitted by some of the *Ixodes* tick species, and it is the most common vector borne disease in the United States with more than 304,000 LD cases have been reported to the Centers for Disease Control and Prevention from 1995 to 2009. The majority of cases occur in the Middle Atlantic, Northeastern, and North Central states, reaching 30,000 confirmed cases in 2009 together

© 2011 Published by Elsevier Ltd.

*Corresponding author. Tel.: +1 301 295 2266; fax: +1 301 295 1435. pozsgayv@mail.nih.gov, vipo@helix.nih.gov (V. Pozsgay)..

Dedicated to Professor Dr. András Lipták on the occasion of his 75th birthday

with more than 8000 probable cases in the same year.^{2,3} A licensed vaccine (LYMERix™) containing a lipidated recombinant surface protein of *B. burgdorferi* designated as L-OspA, although effective above the age of 12 years when administered with aluminum hydroxide as the adjuvant⁴ was withdrawn from the market in early 2002 after less than 5 years of use, because of inadequate market results. Moreover, there was also the contentious issue of the vaccine's hypothetical potential to induce autoimmunity because of OspA's partial homology to the human leukocyte function-associated antigen 1 in persons with certain HLA-DR alleles^{5,6} but studies have shown no increase in the development of arthritis or other adverse effects.⁷

Currently, there is no vaccine for human use against LD, and prevention of the disease is limited to protective measures to avoid tick bites. An effective vaccine to prevent human Lyme disease would be of great benefit for populations with high risk of acquiring the infection.^{8,9}

B. burgdorferi produces neither a lipopolysaccharide nor a capsular polysaccharide.^{10,11} On the other hand, immunoreactive glycolipids were isolated from *B. burgdorferi* that were shown to be α -galactosyl diacylglycerols.¹² However, neither the location of the acyl groups nor the stereochemistry of the glycerol residue was defined in the early studies. Our laboratory reported the isolation and structural characterization of two groups of surface-exposed glycolipids termed BBGL-1 and BBGL-2.¹¹ Using a variety of chemical and spectroscopic methods, BBGL-1 was identified as 6-*O*-acyl- β -D-galactopyranosyl-cholesterol and BBGL-2 as 1,2-di-*O*-acyl-3-*O*- α -D-galactopyranosyl-*sn*-glycerol.¹¹ These findings were confirmed by other workers who also reported the isolation of cholesteryl β -D-galactopyranoside without an acyl group at the galactose unit as well as 6-*O*-acyl- β -D-glucopyranosyl-cholesterol.^{13,14} In BBGL-2 the most common fatty acid is palmitic acid (1.00 mol), followed by oleic and stearic acids (approx. 0.65 and 0.25 mol, respectively). Myristic and linoleic acids were also detected in 0.15 and 0.12 mol amounts. Based on the relative proportions of the palmitic and oleic acids it was proposed that the major components of BBGL-2 are 3-*O*- α -D-galactopyranosyl-1-*O*-oleoyl-2-*O*-palmitoyl-*sn*-glycerol (**4**) and/or 3-*O*- α -D-galactopyranosyl-2-*O*-oleoyl-1-*O*-palmitoyl-*sn*-glycerol (**3**) whereas the other fatty acids detected in the BBGL-2 fraction remain unaccounted for.¹¹ Because the individual components of the BBGL-2 complex could not be separated, no single structure could be identified. Therefore, the presence of galactosyl-homodiacyl-glycerol derivatives and of those incorporating the minor fatty acids cannot be excluded. Because of the lack of genes in *B. burgdorferi* for the synthesis or elongation of fatty acids,¹⁵ its fatty acids are incorporated from the host or from the environment. This may explain the differences in fatty acid composition reported by different laboratories for in vitro cultivated *B. burgdorferi* cells.^{11,12} In mice and rabbits, BBGL-2 elicited antibodies that reacted with both BBGL-1 and -2, and the sera of LD patients had a strong IgG reaction with BBGL-2.^{11,16} These propensities make the BBGL-2 glycolipids candidates for developing diagnostics and vaccines against *B. burgdorferi* devoid of any immunogenic proteins such as L-OspA that might have the potential to elicit autoantibodies.⁵

Chemical syntheses of BBGL-2 glycolipids having one oleoyl and one palmitoyl group on their glycerol moieties have been reported, but the published synthetic protocols lack rigorous proof of their homogeneity.¹⁷⁻²⁰ An approach by α -galactosylation of the commercially available diglyceride 2-*O*-oleoyl-1-*O*-palmitoyl-*sn*-glycerol (Avanti Polar Lipids) has recently been described.^{19,20} Examination of the ¹³C NMR spectrum of the commercial material, obtained from the same source revealed that it contains up to 25% of an accompanying compound that is most likely 1-*O*-oleoyl-2-*O*-palmitoyl-*sn*-glycerol. The basis of this assumption is that while the ¹H NMR spectrum of the commercial material is fully consistent with the proposed structure, the ¹³C NMR spectrum exhibits, in addition to

two major carbonyl carbon signals, two additional ones that are approximately 1/4th of the major ones, together with several similarly low-intensity peaks. Because in our experience separation of **3** and **4** is not possible, we are tempted to assume that the material reported in Refs. 19 and 20 as compound **3** is a ca. 4:1 mixture of **3** and **4**. An alternative approach^{17,19} starts with the commercially available 3-*O*-benzyl-*sn*-glycerol which is regio-selectively acylated at O-1 followed by a second acylation at O-2 and removal of the *O*-benzyl group with trichloroborane to afford the targeted 1-*O*-acyl(1)-2-*O*-acyl(2)-*sn*-glycerols^{17,19,21} which are then galactosylated. In applying this approach to 1,2-diacyl-*sn*-glycerol precursors, care has to be taken to avoid the well-documented acyl migration that can take place under both acidic and basic conditions.^{22–27} We did not observe acyl migration from the O-1 to the O-2 position of the glycerol moieties, but we did observe migration from O-2 to O-3 over extended periods at room temperature, or even during silica gel column chromatography. In order to prevent undesirable acyl migration, short reaction times and quick chromatographic procedures have been suggested.²⁵ It has been proposed that the regioisomeric purity of acyl glycerols can best be determined by integration of the carbonyl signals of the acyl moieties in high field ¹³C NMR spectra.^{22,23} Because of the low natural abundance (1.1%) of ¹³C combined with the low NMR sensitivity of carbonyl carbons, this method requires long acquisition times to obtain good-quality spectra. Unfortunately, the signal-to-noise ratio of the carbonyl region is insufficient in most reports to allow rigorous assessment of homogeneity.

We are examining approaches to producing BBGL components for use in a vaccine against *B. burgdorferi*. Growing *B. burgdorferi* to produce BBGL's in sufficient quantities for immunization experiments is difficult. In addition, isolation of the glycolipids in a homogeneous form has not been possible, raising reproducibility concerns. To circumvent these difficulties, we are preparing BBGL components by using synthetic chemical methods. So far, we have reported the synthesis of the major BBGL-1 components in their native and bioconjugatable forms²⁸ and prepared a semisynthetic experimental vaccine against *B. burgdorferi* consisting of the BBGL-1 glycolipids covalently linked to bovine serum albumin through an oxime linkage.²⁹ The aim of the present work is to delineate the immunodominant region of BBGL-2 components by assessing their antigenicity. It is expected that such recognition will facilitate the design of a vaccine against LD.

Here we describe experiments aimed at synthesizing the putative BBGL-2 components **1–4**. In order to evaluate the biological importance of various structural features we also describe the synthesis of the diastereoisomers **5–8** that differ from the native ones only in the stereochemistry at C-2 of the glycerol moiety, the saturated analogs **9** and **10**, the mono-*O*-acyl derivatives **11** and **12**, as well as two additional BBGL-2 analogs (**13** and **14**) that contain the galactose moiety in the unnatural, beta glycosidic linkage. (Table 1).

Following the synthetic studies, we disclose salient NMR features of several of the synthetic glycolipids, and discuss the antigenicity of the synthesized glycolipids.

2. Results and discussion

2.1. Synthetic studies

We explored an approach in which the introduction of the α -D-galactopyranosyl moiety to *sn*-3-*O* of the glycerol unit precedes the attachment of the fatty acyl groups, thereby preventing acyl migration to that site.¹⁸ This concept is shown in Scheme 1. Precursor to the target compounds was phenyl 1-thio- β -D-galactopyranoside^{30,31} **15**. In preliminary trials, compound **15** was converted into its tetra-*O*-benzyl derivative, which was reacted with the commercially available glycerol-*sn*-1,2-acetonide **19** under a variety of published thioglycoside activation protocols (not described in the experimental part). Unexpectedly,

the products contained mostly β -interglycosidic linkages, despite the presence of the non-participating benzyl protecting group at the O-2 position of the galactose moiety. We hypothesized that tuning down the reactivity of the galactosyl donor would improve this situation. After attempts with various tri-*O*-acyl derivatives of the thiogalactoside triol **16**, including levulinoyl,³² pentafluoropropionyl,³³ and trifluoroacetyl³³ groups we eventually selected the benzoylated galactosyl chloride **18** obtained by chlorinolysis of thiogalactoside **17**, which in turn was prepared by conventional benzylation of **16**.³⁴ Compound **18** was reacted with acetone **19** under AgOTf activation to afford the desired α -linked intermediate **20** in 61% yield. The 1,2-*cis* (α) interglycosidic linkage was proven by the value of the $J_{1,2}$ coupling constant being 3.6 Hz (see Section 4). The corresponding β -linked glycoside **21** was also isolated, in 19% yield. Its anomeric configuration was indicated by the doublet in its ¹H NMR spectrum at 4.71 ppm, with the $J_{1,2}$ coupling constant being 7.7 Hz. Next, the benzoyl groups were cleaved from **20** by sodium methoxide uneventfully, to afford the triol **22** in excellent yield, followed by careful hydrogenolytic removal of the benzyl group, using ethyl acetate as solvent and a commercial palladium-on-charcoal catalyst, in admixture with 2,4,6-tri-*tert*-butylpyrimidine, thus affording **23** in 87% yield. In our experience, extended hydrogenolysis often leads to a significant drop in pH. A likely reason for this is the presence of residual palladium chloride in the catalyst, from which hydrogen chloride may be generated upon hydrogenation. Treatment of the tetraol **23** with levulinic acid and DCC in the presence of DMAP afforded a nearly stoichiometric yield of the fully protected galactoside **24**.¹⁸ Subsequently, the isopropylidene group was removed from the glycerol moiety by hydrolysis in acetic acid to yield diol **25** in 83% yield, for incorporation of the lipid chains. Employing identical reaction conditions, diastereomer **27** having the galactose moiety at the *sn*-1 position was also synthesized, using the commercially available glycerol derivative **26**, the enantiomer of **19**. (Chart 1).

The synthetic sequence toward BBGL-2 analogs **13** and **14**, that contain the galactose moiety in the unnatural, beta glycosidic linkage started from imidate **28**³⁵ that was reacted with alcohol **19** under activation by TMSOTf, to afford disaccharide **29** in which the β interglycosidic linkage was proven by the value $J_{1,2}$ 7.9 Hz. A two-step replacement of the acyl groups in **29** by levulinoyl groups afforded the fully protected intermediate **30**, from which acetic acid hydrolysis yielded diol **31** in high yield. (Scheme 2).

With the availability of compounds **25**, **27**, and **31** incorporating both the galactose and the glycerol moieties in the required stereo-chemistries, the stage was set for the introduction of the oleoyl, palmitoyl, and stearoyl groups.

For the synthesis of the *sn*-1-*O* and *sn*-2-*O* hetero-di-substituted Gro-moieties, we opted initially to adapt a published protocol¹⁸ that, as we eventually found out by ¹³C NMR spectroscopy, afforded regioisomeric mixtures. In Ref. 18, diol **25** was treated with one equivalent of palmitic or oleic acid in the presence of DCC, apparently without the use of the nucleophilic catalyst DMAP that in our experience is essential to induce O-acylation. Hexane-EtOAc mixtures were used to follow the course of the reaction by TLC, and to isolate the required product 'singly acylated' at *sn*-1-*O*.¹⁸ Using 95:5 and 99:1 v/v mixtures of CH₂Cl₂ and MeOH as the TLC developing solvent allowed us to observe that the 'singly acylated' product is, in fact, a mixture of *sn*-1-*O* and *sn*-2-*O* mono-acylated isomers, with the former predominating in a range of approximately 4:1–9:1. The formation of this mixture remained unnoticed in an earlier protocol,¹⁸ and no physical data have been reported for the 'singly acylated' derivatives either.¹⁸ All of our efforts to separate the 'singly acylated' *sn*-1-*O* and 2-*O* products by TLC or column chromatography using a range of hexane-EtOAc mixtures, reported in the published protocol,¹⁸ proved to be futile.

We also noted the formation of a di-*O*-Gro acylated product before the disappearance of the starting diol, an observation that required the termination of the reaction when some of the diols **25**, **27**, or **31** were still present.

The faster-moving mono-*O*-acylated component was invariably the targeted *sn*-1 acyl product identified by the presence of a characteristic doublet of doublets in its ¹H NMR spectrum at ca. 4.37 ppm, indicating that acylation took place at the primary hydroxyl group of the glycerol moiety, leaving HO-2 unsubstituted. The slightly slower-migrating product was the undesired *sn*-2 acyl isomer showing a diagnostic one-proton multiplet at about 5.1 ppm.

With the homogeneous *O*-1 (*sn*) acylated derivatives in hand, the second acyl group was introduced at *O*-2 of the glycerol moiety, in uneventful reactions using the DCC/DMAP method.

Treatment of the diols **25**, **27**, and **31** with various FAs used in excess under the agency of DCC/DMAP proceeded as expected, and produced the fully substituted intermediates featuring two identical fatty acids on the glycerol moiety.

The singly and the doubly glycerol-acylated protected BBGL-2 congeners prepared in this study are listed in Tables 2 and 3, respectively. In the final stage of the syntheses, the levulinoyl protecting groups were conventionally removed from the galactose moiety by hydrazine acetate in pyridine³² to afford **1–14**.

2.2. Nuclear magnetic resonance studies

A detailed NMR investigation was performed on compounds **1–4**. ¹H NMR spectra were assigned using 2D COSY and high-resolution, 2D TOCSY, the latter technique being used to generate separate 1D sub-spectra for the Gal, Gro, Ole, and Pal residues. Selected ¹H NMR chemical shifts are reported in Table 4 and coupling constants in Table 6. The small $J_{4,5}$ value of the Gal residue resulted in inefficient magnetization transfer from H-4 to H-5 and beyond, so that the assignments for H-5, H-6, and H-6' were obtained from the one-bond, ¹H–¹³C correlations established by 2D HSQC. Selected ¹³C NMR chemical shifts are listed in Table 5. Substituent positions in compounds **1–4** were verified by the inter-residue connectivities observed by high-resolution, 2D HMBC. Anomeric ¹J_{C-1,H-1} coupling constants of **1–4** were measured by ¹H-coupled, 2D HSQC, and were found to lie in the range of 170.1–170.4 Hz, which together with the small values of $J_{1,2}$ 3.7–3.8 Hz confirms the configuration of the glycosidic linkages, as designed.

The structures of regioisomers **3** and **4** are not easily differentiated by ¹H NMR at 500 MHz, but we have found that 1D ¹³C NMR provides an excellent method for distinguishing the structures of the compounds, and assessing their purity. The key parameter is the ¹³C=O chemical shift (see Table 5), which appears to be quite sensitive to the type and chemical environment of the acyl groups. In this regard we note that HMBC was less suitable to detect connectivities in minor isomers. This is because of the limited spectral resolution imposed by small numbers of data points in 2D HMBC, but also because of the restricted signal:noise ratio and dynamic range caused by the small number of scans per FID in 2D NMR. By contrast, 1D ¹³C NMR has better spectral resolution because of the far greater number of points sampled in the FID, as compared with those in the indirectly detected, ¹³C dimension of 2D HMBC.

Analysis of the ¹³C NMR data of pure oleoyl and palmitoyl diglycerides and their mixtures suggests the following rules:

- (a) The $^{13}\text{C}=\text{O}$ chemical shift is primarily influenced by the substituent's position on the glycerol: the 1-*sn*-substituent $^{13}\text{C}=\text{O}$'s resonate at lower field than the 2-*sn*-substituent $^{13}\text{C}=\text{O}$'s, for example, 173.65 versus 173.34 ppm for the Pal substituent at O-1 versus O-2.
- (b) The origin of the minor influence derives from the type of the substituent: the $^{13}\text{C}=\text{O}$'s of the palmitoyl groups resonate at lower field than those of the oleoyl groups, for example, 173.34 versus 173.29 for the Pal versus the Ole group at *sn* O-2.

These differences have proved to be extremely useful in detecting the presence of glyceride mixtures produced by acylation reactions of imperfect regioselectivity, a facet that has not previously been recognized.

2.3. Immunochemistry

That sera from LD patients react with carbohydrate and lipid-containing substances extracted from *B. burgdorferi* was demonstrated many years ago,^{36–38} but not until 2003 was it shown that the surface of this pathogen contains two groups of glycolipids, which were termed BBGL-1 and 2.¹¹ BBGL-2 was reported to contain several FA's of which palmitate and oleate were shown to be the major components, but their exact location on the glycerol moiety is unknown.¹¹ This knowledge may be crucial for the preparation of a chemically well-characterized, synthetic vaccine. In order to determine the structural features necessary for recognition by antibodies, 14 glycolipids have been synthesized, of which 12 contain the Gal residue in the native, a anomeric configuration, and 2 in the unnatural, β one. The synthetic glycolipids were reacted with mouse sera induced by (1) formaldehyde killed *B. burgdorferi* cells and (2) by native purified BBGL-2, and also with serum of patients with late Lyme neuroborreliosis (Fig. 1). There was no reaction of any glycolipid with control sera (not shown). All of the glycolipids exhibited similar antigenic activities to all three sera. No binding occurred with the glycolipids containing only a single FA moiety (compounds **11** and **12**). The lipids containing two saturated FA's were also unreactive: no binding occurred with the compounds containing two palmitates (**1**, **5**, and **13**), two stearates (compound **9**), or one palmitate and one stearate (compound **10**). The observed non-reactivity might be related to the low solubility of these compounds as compared to the rest of the glycolipids in Table 1. All of the compounds having two FA's of which at least one is an oleoyl residue reacted with each of the three sera (**3**, **4**, **7**, and **8**). The glycolipids having two oleates (**2**, **6**, and **14**) were also reactive. Unexpectedly, compounds with the Gal moiety in the unnatural β configuration exhibited the same reactivities as did their natural counterparts: the di-palmitoyl derivative **13** was unreactive, whereas the di-oleoyl congener was seemingly as reactive as compound **2**.

These observations led us to the following conclusions:

- (a) The presence in the glycerol part of two fatty acid moieties is required for antigenicity, leading to the hypothesis that the binding to antibodies involves two fatty acids and two closely located binding sites. Alternatively, one FA may position the other in order to achieve the proper conformation for binding with antibodies.
- (b) Of the two fatty acids, at least one should be oleic acid.
- (c) The position of the oleic acid moiety is not critical.
- (d) Replacement of the oleic acid by stearic acid abolishes antigenicity.
- (e) The anomeric configuration has no influence on antigenicity: α and β -galactoside derivatives can be equally antigenic.

- (f) The galactose moiety may be linked to either O-1 or to O-3 of *sn*-glycerol to maintain antigenicity.

Based on the above observations, we prepared an 'unnatural' BBGL-2 analog, using the chemistry described in Section 2.1 having the structure 3-*O*-(β -D-galactopyranosyl)-1,2-di-*O*-oleoyl-DL-glycerol **46**, which is much easier and less expensive to synthesize than the native analogs, by starting from DL-glycerol. Figure 2. shows that the 'unnatural' congener is just as good an antigen as are the reactive native analogs, and is a candidate for a vaccine against LD.

There is some similarity between our results on antigenic specificity and those published recently by others demonstrating that iNKT cells react with BBGL-2 glycolipids in a FA structure-dependent manner, and that the difference in fatty acids influences the steric position of the sugar epitope.^{18,39} The highest response, assessed by the level of induced IL-2 was to compound **4** having Ole and Pal residues in the *sn*-1 and *sn*-2 positions, respectively, whereas the reverse arrangement produced less than half of that. In our present study, compound **4** also showed a high level of antigenicity. On the other hand, compound **1** produced only slightly more than baseline IL-2 levels, as shown by others.^{18,39} In a similar fashion, in our study compound **1** was not recognized by sera induced to *B. burgdorferi*, or to purified native BBGL-2.

3. Conclusions

We have described synthetic schemes to homogeneous BBGL-2 congeners that make up a distinctive group of major glycolipids on the surface of *B. burgdorferi*, the causative organism of Lyme disease, comprising α -galactopyranosyl-diacyl glycerols. We have presented evidence that the diastereomeric purity of these glycolipids can best be ascertained by examination of the carbonyl carbon resonances in their ¹³C NMR spectra, whereas the ¹H NMR spectra are less informative. We found that the antigenicity of the synthetic BBGL-2 glycolipids depends on the unsaturation of the lipid components, but is independent of the anomeric configuration of the *sn*-3 linked hydrophilic galactose head-group. The presence of two FAs in the glycerol part is necessary, of which at least one has to be an oleic acid residue for antigenicity, a property that is lost completely upon saturation.

4. Experimental

4.1. General methods

All chemicals were of commercial grade and used without purification. Solvents for chromatography were distilled prior to use. Anhydrous solvents were obtained from Aldrich. Column chromatography was performed on Silica Gel 60 (0.040–0.063 mm) and thin layer chromatography was performed on glass-supported silica gel layers obtained from Analtech (Uniplate) or on HPTLC plates from Merck. Visualization was carried out by inspection under UV light (254 nm), by iodine adsorption, and by charring using a solution of ammonium cerium(IV) sulfate and ammonium molybdate in sulfuric acid. 2-*O*-Oleoyl-1-*O*-palmitoyl-*sn*-glycerol (lot #160-181DG-22) was obtained from Avanti Polar Lipids Inc., Alabaster, AL. API-ES mass spectra were recorded on an Agilent Technologies LC MDS CL spectrometer.

NMR spectra were recorded at 300 K, using a Bruker DRX-500 spectrometer equipped with a 5 mm, HCN cryoprobe. Solutions containing ~20 mg of glycolipid in CDCl₃ unless indicated otherwise (0.5 mL) were used, except for compounds **1**, **5**, **9**, and **13** which were not soluble to this extent, and for which a saturated solution was employed. Tetramethylsilane was used as an internal chemical shift reference at $\delta = 0$, for ¹H and ¹³C

NMR spectra. The NMR data were acquired and processed by means of the Bruker Topspin program, version 1.3.

One-dimensional (1D) ^1H NMR spectra were acquired in 32,768 data points, zero-filled to 65,536 points, with use of a spectral width of 4.01 kHz, a 30° pulse (2.67 μs), and a recycle time of 6 s. For integration, the free induction decay (FID) was subjected to Gaussian multiplication, using a line-broadening of -0.5 Hz and a truncation fraction of 0.3. However, coupling constants were measured by using a line-broadening of -1.0 to -2.0 Hz. 1D ^{13}C NMR spectra were acquired in 65,536 data points, zero-filled to 131,072 points, by using a 25.15 kHz spectral width, a 45° pulse (7.7 μs), and a recycle time of 2 s. The free induction decays were subjected either to exponential multiplication using a line-broadening of 1.0 Hz, or to Gaussian multiplication with a line-broadening of -0.1 to -0.5 Hz.

2D COSY NMR spectra were acquired by using a 30° ^1H pulse, and 2048×512 point datasets, zero-filled to 2048×2048 points. The data were processed by using an unshifted sine-bell squared window in both dimensions, a magnitude calculation in F_1 , and symmetrization to remove artifacts. 2D TOCSY spectra were collected in $16,384 \times 512$ point datasets, zero-filled to $32,768 \times 2048$ points. Resolution enhancement by Gaussian multiplication was used in F_2 with a line-broadening of -1.5 Hz, and in F_1 , a sine-bell squared window, shifted by $\pi/2$ rad. The spectra were analyzed by means of extraction of 1D slices in the F_2 dimension. 2D HSQC spectra were recorded in 2048×512 point datasets, zero-filled to 8192×4096 points. A sine-bell squared window shifted by $\pi/4$ rad was used in both dimensions. 2D HMBC was conducted with an evolution delay of 83 ms, corresponding to optimization for $^{2,3}J_{\text{CH}}$ 6 Hz, together with 2048×512 or 2048 point datasets, zero-filled to 4096×4096 points. A magnitude calculation was used in F_1 , with sine-bell squared windows shifted by $\pi/2$ rad in both dimensions. ^1H -coupled 2D HSQC spectra were acquired in 8192×512 point datasets, zero-filled to 8192×2048 points, with a sine-bell squared window in both dimensions, shifted by $\pi/2$ rad. All 2D pulse sequences were field-gradient selected. ^1H and ^{13}C spectral widths were 3.005 and 25.15 kHz, respectively. The phase-sensitive, echo-anti-echo protocol was used for 2D TOCSY, HSQC, and ^1H -coupled HSQC.

The synthetic glycolipids were dissolved in $\text{CHCl}_3/\text{MeOH}$ 80:20 (v/v) at 1 mg/ml concentration. Then, 3 μL of the solution was directly pipetted onto stripes of a PVDF membrane (Millipore, Bedford, MA). After 10 min, the membranes were blocked with 5% BSA in PBS for 1 h at rt and washed three times with PBS. Next, the membranes were incubated for 2 h at rt with the following sera: (1) induced to heat killed *B. burgdorferi* cells (strain B31, ATCC 35210), diluted 1:100 in blocking buffer; (2) induced to purified BBGL-2, diluted 1:50 in blocking buffer; (3) serum from human patients diagnosed with Lyme disease, diluted 1:50 in blocking buffer; (4) control mouse and human sera. Mouse sera were prepared as described in Refs. 11 and 40. After washing as above, the membranes were incubated with goat anti-mouse or anti-human IgG conjugated to alkaline phosphatase (KPL, Gaithersburg, MD) for 1 h at rt, then washed and developed with BCIP/NBT phosphate substrate (KPL, Gaithersburg, MD).

4.2. 3-O- α -D-Galactopyranosyl-1,2-di-O-palmitoyl-*sn*-glycerol¹⁸ (1)

TLC: CH_2Cl_2 -MeOH (95:5). For the ^1H and ^{13}C NMR spectra see Tables 4–6. API-ES-MS: m/z calcd for $[\text{C}_{41}\text{H}_{78}\text{O}_{10}]\text{NH}_4^+$: 748.6. Found 748.4.

4.3. 3-O- α -D-Galactopyranosyl-1,2-di-O-oleoyl-*sn*-glycerol (2)

TLC: EtOAc-hexanes (1:1, 4:1). For the ^1H and ^{13}C NMR spectra see Tables 4–6. API-ES-MS: m/z calcd for $[\text{C}_{45}\text{H}_{82}\text{O}_{10}]\text{NH}_4^+$: 800.6. Found 800.6.

4.4. 3-O- α -D-Galactopyranosyl-2-O-oleoyl-1-O-palmitoyl-*sn*-glycerol¹⁷⁻²⁰ (3)

TLC: EtOAc–MeOH (99:1, 95:5), CH₂Cl₂–MeOH (95:5). For the ¹H and ¹³C NMR spectra see Tables 4–6. API-ES-MS: *m/z* calcd for [C₄₃H₈₀O₁₀]⁺NH₄: 774.6. Found 774.4.

4.5. 3-O- α -D-Galactopyranosyl-1-O-oleoyl-2-O-palmitoyl-*sn*-glycerol¹⁸⁻²⁰ (4)

TLC: EtOAc–MeOH (99:1, 95:5), CH₂Cl₂–MeOH (95:5). For the ¹H and ¹³C NMR spectra see Tables 4–6. API-ES-MS: *m/z* calcd for [C₄₃H₈₀O₁₀]⁺NH₄: 774.6. Found 774.6.

4.6. 1-O- α -D-Galactopyranosyl 2,3-di-O-palmitoyl-*sn*-glycerol (5)

TLC: CH₂Cl₂–MeOH (95:5); ¹H NMR (CDCl₃): δ 5.24 (m, 1H), 4.97 (d, 1H, *J* = 3.7 Hz), 4.29 (m, 1H, *J* = 4.1 Hz, *J* = 11.9 Hz), 4.20 (dd, 1H, *J* = 6.0 Hz, *J* = 11.9 Hz), 4.09 (br d, 1H, *J* = 2.1 Hz), 3.94 (dd, 1H, *J* = 6.3 Hz, *J* = 11.6 Hz), 3.87–3.81 (m, 4H), 3.74 (dd, 1H, *J* = 3.1 Hz, *J* = 9.8 Hz), 3.63 (dd, 1H, *J* = 4.8 Hz, *J* = 11.1 Hz); ¹³C NMR (CDCl₃): δ 173.55, 173.35, 99.29, 70.89, 70.28, 70.11, 69.92, 69.36, 66.89, 63.14, 62.17, 34.31, 34.10, 31.94, 29.71, 29.69, 29.67, 29.66, 29.51, 29.37, 29.30, 29.15, 29.12, 24.89, 22.70, 14.13. API-ES-MS: *m/z* calcd for [C₄₁H₇₈O₁₀]⁺NH₄: 748.6. Found 748.4.

4.7. 1-O- α -D-Galactopyranosyl-2,3-di-O-oleoyl-*sn*-glycerol (6)

TLC: EtOAc–hexanes (1:1, 4:1); ¹H NMR (CDCl₃): δ 5.39–5.29 (m, 4H), 5.25 (m, 1H), 4.97 (d, 1H, *J* = 3.6 Hz), 4.30 (dd, 1H, *J* = 3.9 Hz, *J* = 11.9 Hz), 4.19 (dd, 1H, *J* = 6.2 Hz, *J* = 11.9 Hz), 4.09 (d, 1H, *J* = 3.2 Hz), 3.93 (dd, 1H, *J* = 5.0 Hz, *J* = 11.6 Hz), 3.88–3.78 (m, 4H), 3.75 (dd, 1H, *J* = 3.3 Hz, *J* = 10.3 Hz), 3.62 (dd, 1H, *J* = 5.0 Hz, *J* = 11.1 Hz), 2.35–2.27 (m, 4H), 2.04–1.95 (m, 8H), 1.66–1.54 (m, 4H), 1.37–1.19 (m, 40H), 0.91–0.82 (m, 6H); ¹³C NMR (CDCl₃): δ 173.60, 173.34, 129.98, 129.64, 129.62, 99.37, 70.58, 70.18, 70.09, 69.95, 69.09, 66.63, 62.45, 62.41, 34.24, 34.04, 31.87, 29.73, 29.70, 29.70, 29.49, 29.28, 29.19, 29.13, 29.09, 29.06, 27.19, 27.16, 24.89, 24.82, 22.65, 14.08. API-ES-MS: *m/z* calcd for [C₄₅H₈₂O₁₀]⁺NH₄: 800.6. Found 800.4.

4.8. 1-O- α -D-Galactopyranosyl-2-O-oleoyl-3-O-palmitoyl-*sn*-glycerol (7)

TLC: EtOAc–MeOH (99:1, 95:5), CH₂Cl₂–MeOH (95:5); ¹H NMR (CDCl₃): δ 5.38–5.31 (m, 2H), 5.24 (m, 1H), 4.30 (dd, 1H, *J* = 4.9 Hz, *J* = 11.8 Hz), 4.19 (dd, 1H, *J* = 6.2 Hz, *J* = 11.8 Hz), 4.09 (d, 1H, *J* = 2.8 Hz), 3.93 (dd, 1H, *J* = 5.3 Hz, *J* = 11.8 Hz), 3.86–3.79 (m, 4H), 3.75 (dd, 1H, *J* = 2.8 Hz, *J* = 9.6 Hz), 3.62 (dd, 1H, *J* = 4.9 Hz, *J* = 10.9 Hz), 2.33 (t, 1H, *J* = 6.9 Hz), 2.31 (t, 2H, *J* = 7.4 Hz), 2.05–1.99 (m, 4H), 2.05–1.99 (m, 2H), 1.65–1.56 (m, 4H), 1.38–1.19 (m), 0.88 (t, 6H, *J* = 7.0 Hz); ¹³C NMR (CDCl₃): δ 173.69, 173.38, 130.02, 129.66, 99.39, 70.84, 70.28, 70.13, 69.94, 69.34, 66.83, 63.00, 62.23, 34.28, 34.10, 31.93, 31.91, 29.76, 29.71, 29.68, 29.67, 29.65, 29.54, 29.51, 29.37, 29.33, 29.32, 29.30, 29.21, 29.15, 29.09, 27.19, 27.18, 24.92, 24.88, 22.70, 22.69, 14.13. APIES-MS: *m/z* calcd for [C₄₃H₈₀O₁₀]⁺NH₄: 774.6. Found 774.6.

4.9. 1-O- α -D-Galactopyranosyl-3-O-oleoyl-2-O-palmitoyl-*sn*-glycerol (8)

TLC: EtOAc–MeOH (99:1, 95:5), CH₂Cl₂–MeOH (95:5); ¹H NMR (CDCl₃–MeOH-*d*₄): δ 5.30–5.23 (m, 2H), 5.15 (m, 1H), 4.85 (d, 1H, *J* = 3.4 Hz), 4.27 (dd, 1H, *J* = 3.7 Hz, *J* = 12.0 Hz), 4.11 (dd, 1H, *J* = 6.5 Hz, *J* = 12.0 Hz), 4.01 (d, 1H, *J* = 3.3 Hz), 3.48–3.65 (m, 5H), 3.54 (dd, 1H, *J* = 5.4 Hz, *J* = 10.8 Hz), 2.25 and 2.23 (2t, 2 \times 2H, *J* = 7.2 Hz for each), 1.99–1.91 (m, 4H), 1.60–1.48 (m, 4H), 1.31–1.14 (m), 0.86–0.77 (m, 6H); ¹³C NMR (CDCl₃–MeOH-*d*₄): δ 173.68, 173.35, 129.96, 129.6, 99.37, 70.51, 70.19, 69.96, 68.99, 66.56, 62.54, 62.16, 34.24, 34.07, 31.89, 31.86, 29.72, 29.71, 29.68, 29.63, 29.51, 29.49, 29.42, 29.33, 29.29, 29.28, 29.21, 29.14, 29.06, 28.18, 27.15, 24.88, 24.84, 22.65, 22.64, 14.07. APIES-MS: *m/z* calcd for [C₄₃H₈₀O₁₀]⁺NH₄: 774.6. Found 774.6.

4.10. 3-O- α -D-Galactopyranosyl-1,2-di-O-stearoyl-*sn*-glycerol¹⁷ (9)

TLC: EtOAc–MeOH (99:1, 95:5), CH₂Cl₂–MeOH (95:5); ¹H NMR (CDCl₃–MeOH-*d*₄): δ 5.26 (m, 1H), 4.88 (d, 1H, *J* = 3.7 Hz), 4.39 (dd, 1H, *J* = 3.6 Hz, *J* = 12.0 Hz), 4.16 (dd, 1H, *J* = 6.4 Hz, *J* = 12.0 Hz), 3.99 (d, 1H, *J* = 3.3 Hz), 3.85–3.75 (m), 3.73 (dd, 1H, *J* = 3.3 Hz, *J* = 10.0 Hz), 3.63 (dd, 1H, *J* = 5.9 Hz, *J* = 10.9 Hz), 2.53 (m, 1H), 2.45 (m, 1H), 2.35–2.30 (m, 4H), 1.65–1.56 (m, 4H), 1.36–1.19 (m, 56H), 0.88 (t, 6H, *J* ~6.8 Hz); ¹³C NMR (CDCl₃–MeOH-*d*₄): δ 173.82, 173.48, 99.34, 70.27, 70.03, 69.85, 69.62, 68.82, 66.15, 62.34, 61.71, 34.06, 33.89, 31.70–28.89, 25.62, 24.67, 24.65, 22.45, 13.79. API-ES-MS: *m/z* calcd for [C₄₅H₈₆O₁₀]NH₄⁺: 804.6. Found 804.7.

4.11. 3-O- α -D-Galactopyranosyl-2-O-palmitoyl-1-O-stearoyl-*sn*-glycerol (10)

TLC: EtOAc–MeOH (99:1, 95:5), CH₂Cl₂–MeOH (95:5); ¹H NMR (CDCl₃–MeOH-*d*₄): δ 5.28 (m, 1H), 4.87 (d, 1H, *J* = 3.6 Hz), 4.44 (dd, 1H, *J* = 3.2 Hz, *J* = 12.0 Hz), 4.19 (dd, 1H, *J* = 6.4 Hz, *J* = 12.0 Hz), 3.95 (d, 1H, *J* = 3.2 Hz), 3.86 (dd, 1H, *J* = 5.5 Hz, *J* = 10.8 Hz), 3.83–3.71 (m, 6H), 3.65 (dd, 1H, *J* = 5.7 Hz, *J* = 10.8 Hz), 2.35 and 2.34 (2t, 2 \times 2H, *J* ~7.1 Hz), 1.68–1.57 (m, 4H), 1.40–1.20 (52H), 0.89 (t, 6H, *J* ~7 Hz); ¹³C NMR (CDCl₃–MeOH-*d*₄): δ 173.54, 173.19, 99.03, 70.37, 69.64, 69.58, 69.19, 68.36, 65.53, 62.09, 61.06, 33.64, 33.47, 31.32, 29.07–28.49, 24.30, 22.03, 13.16. API-ES-MS: *m/z* calcd for [C₄₃H₈₂O₁₀]NH₄⁺: 776.6. Found 776.6.

4.12. 3-O- α -D-Galactopyranosyl-1-O-palmitoyl-*sn*-glycerol (11)

TLC: EtOAc–MeOH (95:5), CH₂Cl₂–MeOH (95:5, 9:1); ¹H NMR (CDCl₃–MeOH-*d*₄): δ 4.88 (d, 1H, *J* = 3.6 Hz), 4.17 (dd, 1H, *J* = 4.6 Hz, *J* = 11.3 Hz), 4.10 (dd, 1H, *J* = 5.9 Hz, *J* = 11.3 Hz), 4.06 (m, 1H), 3.95 (d, 1H, *J* = 3.1 Hz), 3.85–3.72 (m, 6H), 3.42 (dd, 1H, *J* = 7.6 Hz, *J* = 10.3 Hz), 2.36 (t, 2H, *J* = 7.5 Hz), 1.63 (m, 2H), 1.36–1.24 (m, 24H), 0.89 (t, 3H, *J* ~6.9 Hz); ¹³C NMR (CDCl₃–MeOH-*d*₄): δ 173.86, 99.01, 70.26, 69.61, 69.21, 68.97, 68.50, 67.85, 64.44, 61.07, 33.38, 31.27, 29.01, 28.98, 28.95, 28.82, 28.69, 28.63, 28.48, 24.22, 21.99, 13.11. API-ES-MS: *m/z* calcd for [C₂₅H₄₈O₉]NH₄⁺: 510.4. Found 510.5.

4.13. 3-O- α -D-Galactopyranosyl-1-O-oleoyl-*sn*-glycerol (12)

TLC: EtOAc–MeOH (95:5), CH₂Cl₂–MeOH (95:5, 9:1); ¹H NMR (CDCl₃–MeOH-*d*₄): δ 5.38–5.30 (m, 2H), 4.89 (d, 1H, *J* = 3.7 Hz), 4.17 (dd, 1H, *J* = 4.1 Hz, *J* = 11.3 Hz), 4.10 (dd, 1H, *J* = 5.8 Hz, *J* = 11.3 Hz), 4.05 (m, 1H), 3.95 (d, 1H, *J* = 3.4 Hz), 3.89–3.69 (m, 6H), 3.42 (dd, 1H, *J* = 7.6 Hz, *J* = 10.4 Hz), 2.37 (t, 2H, *J* = 7.4 Hz), 2.08–1.98 (m, 4H), 1.69–1.58 (m, 2H), 1.40–1.20 (m, 20H), 0.89 (t, 3H, *J* ~6.8 Hz); ¹³C NMR (CDCl₃–MeOH-*d*₄): δ 173.89, 129.36, 129.12, 99.07, 70.26, 69.67, 69.25, 69.10, 68.57, 67.93, 64.50, 61.13, 33.45, 31.32, 29.15, 29.11, 28.91, 28.72, 28.69, 28.61, 28.53, 28.51, 26.56, 26.54, 24.27, 22.06, 13.23. API-ES-MS: *m/z* calcd for [C₂₇H₄₈O₉]NH₄⁺: 536.4. Found 536.5.

4.14. 3-O- β -D-Galactopyranosyl-1,2-di-O-palmitoyl-*sn*-glycerol (13)

TLC: EtOAc–MeOH (95:5), CH₂Cl₂–MeOH (95:5, 9:1); ¹H NMR (CDCl₃–MeOH-*d*₄): δ 5.28 (m, 1H), 4.43 (dd, 1H, *J* = 3.0 Hz, *J* = 12.0 Hz), 4.25 (d, 1H, *J* = 7.5 Hz), 4.24 (dd, 1H, *J* = 6.8 Hz, *J* = 12.0 Hz), 3.99 (dd, 1H, *J* = 5.4 Hz, *J* = 12.0 Hz), 3.88 (dd, 1H, *J* = 1.1 Hz, *J* = 3.3 Hz), 3.81 (dd, 1H, *J* = 6.6, *J* = 11.5 Hz), 3.75 (dd, 1H, *J* = 5.4 Hz, *J* = 11.5 Hz), 3.74 (dd, 1H, 5.9 Hz, *J* = 11.0 Hz), 3.54 (dd, 1H, *J* = 7.5 Hz, *J* = 9.7 Hz), 3.89 (dd, 1H, *J* = 3.4 Hz, *J* = 9.7 Hz), 2.35–2.31 (m, 4H), 1.61 (m, 4H), 1.36–1.22 (m, 48H), 0.89 (t, 6H, *J* ~7.1 Hz); ¹³C NMR (CDCl₃–MeOH-*d*₄): 173.44, 173.11, 103.39, 74.60, 72.82, 70.47, 69.79, 68.19, 67.00, 62.17, 60.57, 33.50, 33.36, 31.21, 28.96–28.37, 24.19, 21.92, 12.98. API-ES-MS: *m/z* calcd for [C₄₁H₇₈O₁₀]NH₄⁺: 748.6. Found 748.7.

4.15. 3-O- β -D-Galactopyranosyl-1,2-di-O-oleoyl-*sn*-glycerol (14)

TLC: EtOAc–MeOH (95:5), CH₂Cl₂–MeOH (95:5, 9:1); ¹H NMR (CDCl₃–MeOH-*d*₄): δ 5.37–5.31 (m, 4H), 5.29 (m, 1H), 4.39 (dd, 1H, *J* = 3.1 Hz, *J* = 12.2 Hz), 4.27 (d, 1H, *J* = 7.6 Hz), 4.21 (dd, 1H, *J* = 6.6 Hz, *J* = 12.1 Hz), 4.02 (d, 1H, *J* = 2.6 Hz), 3.92 (dd, 1H, *J* = 5.4 Hz, *J* = 11.0 Hz), 3.84 (dd, 1H, *J* = 4.9 Hz, *J* = 12.0 Hz), 3.74–3.65 (m, 3H), 3.59 (dd, 1H, *J* = 2.9 Hz, *J* = 9.6 Hz), 3.54 (t, 1H, *J* ~4.6 Hz), 2.32 and 2.32 (2t, 2 \times 2H, *J* ~7.4 Hz), 2.08–1.94 (m, 8H), 1.65–1.54 (4H), 1.37–1.19 (m, 40H), 0.88 (t, 6H, *J* ~6.8 Hz); ¹³C NMR (CDCl₃–MeOH-*d*₄): δ 173.88, 173.47, 129.97, 129.64, 104.00, 74.52, 73.37, 71.11, 70.12, 68.88, 68.04, 62.87, 61.52, 34.24, 34.11, 31.88, 27.78–29.08, 27.20, 27.18, 24.86, 24.84, 22.65, 14.09. API-ES-MS *m/z* calcd for [C₄₅H₈₂O₁₀]NH₄⁺: 800.6. Found 800.4.

4.16. Phenyl 3,4,6-tri-O-benzoyl-2-O-benzyl-1-thio- β -D-galactopyranoside (17)

To a solution of compound **16** (32 g, 88.4 mmol) in a mixture of anhydrous CH₂Cl₂ (300 mL) and C₅H₅N (160 mL) containing a catalytic amount of DMAP was added benzoyl chloride (50 mL, 60.5 g, 430 mmol), dropwise at room temperature under stirring. After 24 h, the solution was treated with MeOH (70 mL). The solution was concentrated under reduced pressure. The residue was equilibrated between CHCl₃ and water. The organic layer was extracted with 4 N hydrochloric acid followed by water and saturated aq NaHCO₃. Column chromatographic purification of the product using a hexanes–EtOAc gradient 10:1 \rightarrow 4:1 afforded **17** (54.0 g, 91%) as a solid: ¹H NMR (CDCl₃): δ 8.03–7.13 (m, 25H), 5.92 (ddd, 1H, *J*_{3,4} = 3.5 Hz, *J*_{4,5} = 0.9 Hz, H-4), 5.50 (dd, *J*_{2,3} = 9.5 Hz, *J*_{3,4} = 3.5 Hz, H-3), 4.89 (d, 1H, *J*_{1,2} = 9.6 Hz, H-1), 4.84 and 4.61 [2d, 2 \times 1H *J* = 10.8 Hz each, CH₂ (Bn)], 4.62 (dd, 1H, *J*_{5,6} = 7.1 Hz, *J*_{6,6'} = 11.5 Hz, H-6), 4.41 (dd, 1H, *J*_{5,6'} = 5.7 Hz, *J*_{6,6'} = 11.5 Hz), 4.27 (ddd, 1H, *J*_{4,5} = 0.9 Hz, *J*_{5,6} = 7.1 Hz, *J*_{5,6'} = 5.7 Hz, H-5), 4.01 (t, 1H, *J*_{1,2} = *J* = 9.5 Hz); ¹³C NMR (CDCl₃): δ 166.01, 165.35, 137.36, 133.50, 133.22, 133.18, 132.81, 132.71, 129.92, 129.79, 129.64, 129.46, 129.29, 129.21, 128.99, 128.55, 128.41, 128.29, 128.27, 128.10, 127.93, 127.83, 87.51, 75.48, 75.22, 74.80, 74.77, 68.79, 62.58, 62.52. API-ES-MS: *m/z* calcd for [C₄₀H₃₄O₈S]NH₄⁺: 692.2. Found 692.2.

4.17. 3,4,6-Tri-O-benzoyl-2-O-benzyl- α -D-galactopyranosyl chloride (18)

To a solution of compound **17** (25.0 g, 41.6 mol) in anhydrous CH₂Cl₂ (250 mL) was added a saturated solution of chlorine in CCl₄ at 0 °C. After 30 min TLC (hexanes–EtOAc, 6:1) showed complete conversion to a slightly faster-moving compound. The reaction mixture was treated with 1-hexene (excess) followed by concentration. Column chromatographic purification of the residue (hexanes–EtOAc, 5:1) afforded amorphous **18** (15.4 g, 69%): ¹H NMR (CDCl₃, δ): 8.04–7.19 (m, 20H), 6.29 (d, 1H, *J*_{1,2} = 3.8 Hz, H-1), 5.99 (dd, 1H, *J*_{3,4} = 3.4 Hz, *J*_{4,5} = 1.2 Hz, H-4), 5.81 (dd, 1H, *J*_{2,3} = 10.3 Hz, *J*_{3,4} = 3.4 Hz, H-3), 4.84 (m, 1H, H-5), 4.70 and 4.64 [2d, 2 \times 1H *J* = 12.0 Hz each, CH₂ (Bn)], 4.56 (dd, 1H, *J*_{5,6} = 6.8 Hz, *J*_{6,6'} = 11.5 Hz, H-6), 4.34 (dd, 1H, *J*_{5,6} = 6.0 Hz, *J*_{6,6'} = 11.5 Hz, H-6'), 4.32 (dd, 1H, *J*_{1,2} = 3.8 Hz, *J*_{2,3} = 10.3 Hz, H-2); ¹³C NMR (CDCl₃): δ 165.90, 165.26, 165.21, 136.88, 133.55, 133.24, 133.18, 129.79, 129.76, 129.66, 129.33, 129.28, 129.08, 128.58, 128.51, 128.40, 128.28, 128.17, 128.04, 93.01, 73.09, 72.81, 69.89, 69.71, 68.59, 61.84.

4.18. (3,4,6-Tri-O-benzoyl-2-O-benzyl- α -D-galactopyranosyl)-1,2-O-isopropylidene-*sn*-glycerol (20) and (3,4,6-tri-O-benzoyl-2-O-benzyl- β -D-galactopyranosyl)-1,2-O-isopropylidene-*sn*-glycerol (21)

To a stirred mixture of chloride **18** (15.2 g, 25.3 mmol), 2,4,6-tri-*tert*-butylpyrimidine⁴¹ (22.5 g), crushed 4 Å molecular sieves (15 g), 1,2-O-isopropylidene-*sn*-glycerol **19** (4.5 mL, 4.8 g, 36.5 mmol), and anhydrous CH₂Cl₂ (125 mL) was added AgOTf (15 g, 58.2 mmol) at –40 °C. After 15 min TLC (3:1 hexanes–EtOAc) indicated the disappearance of **18** and the formation of two closely migrating products. To the reaction mixture were added Bu₄NBr

(11.5 g) and saturated aqueous NaHCO₃ and the resulting mixture was filtered through a layer of Celite. The organic layer was concentrated and the residue was chromatographed on silica gel using a 20:1→2:1 hexanes-EtOAc gradient to afford **20** (10.8 g, 61%) as an amorphous substance: ¹H NMR (CDCl₃, δ): 8.01–7.25 (m, 20H), 5.94 (dd, 1H, *J*_{3,4} = 3.0 Hz, *J*_{4,5} = 1.2 Hz, H-4), 5.76 (dd, 1H, *J*_{2,3} = 10.5 Hz, *J*_{3,4} = 3.0 Hz, H-3), 5.14 (d, 1H, *J*_{1,2} = 3.6 Hz, H-1), 4.71 and 4.63 [2d, 2 × 1H *J* = 12.3 Hz each, CH₂ (Bn)], 4.58 (m, 1H, H-5), 4.53 (dd, 1H, *J*_{7,5} Hz, *J* = 11.1), 4.36 (m, 1H, H-2), 4.32 (dd, 1H, *J* = 5.2 Hz, *J* = 11.0 Hz), 4.16 (dd, 1H, *J* = 3.5 Hz, *J* = 10.5 Hz, H-2), 4.09 (dd, *J* = 6.5 Hz, 8.4 Hz), 3.78 (dd, *J* = 6.5 Hz, *J* = 8.4 Hz), 3.74 (dd, 1H, *J* = 5.5 Hz, *J* = 10.5 Hz), 3.64 (dd, 1H, *J* = 5.5 Hz, *J* = 10.5 Hz), 1.42 and 1.37 (2s, 2 × 3H, 2CH₃); ¹³C NMR (CDCl₃): δ 166.02, 165.47, 165.46, 137.78, 133.40, 133.18, 133.02, 129.83, 129.72, 129.66, 129.53, 129.41, 128.54, 128.41, 128.25, 127.96, 127.93, 109.54, 97.85, 74.55, 73.44, 72.87, 70.05, 69.51, 69.16, 67.10, 66.85, 62.72, 26.81, 25.50. API-ES-MS: *m/z* calcd for [C₄₀H₄₀O₁₁]⁺NH₄: 714.3. Found 714.2.

Also isolated was (3,4,6-tri-*O*-benzoyl-2-*O*-benzyl-β-*D*-galactopyranosyl)-1,2-*O*-isopropylidene-*sn*-glycerol **21** (3.35 g, 19%): ¹H NMR (CDCl₃): δ 8.03–7.09 (m, 20H), 5.88 (br d, 1H, *J*_{3,4} = 3.5 Hz, H-4), 5.44 (dd, 1H, *J*_{2,3} = 10.1 Hz, *J*_{3,4} = 3.5 Hz, H-3), 4.87 and 4.69 [2d, 2 × 1H *J* = 11.8 Hz each, CH₂ (Bn)], 4.71 (d, [2d, 2 × 1H *J* = 12.3 Hz each, CH₂ (Bn)], 4.71 (d, 1H, *J*_{1,2} = 7.7 Hz, H-1), 4.39 (m, 1H, H-2'), 4.36 (dd, 1H, *J* = 6.5 Hz, *J* = 11.3 Hz), 4.10 (dd, 1H, *J* = 6.5 Hz, *J* = 8.3 Hz), 4.08 (dd, 1H, *J* = 5.0 Hz, *J* = 10.4 Hz), 3.94 (dd, 1H, *J* = 7.8 Hz, *J* = 10.4 Hz), 3.90 (dd, 1H, *J* = 6.0 Hz, *J* = 8.2 Hz), 3.72 (dd, 1H, *J* = 6.0 Hz, *J* = 10.4 Hz), 1.45 and 1.37 (2s, 2 × 3H, 2CH₃); ¹³C NMR (CDCl₃): δ 165.89, 165.41, 165.33, 137.65, 133.32, 133.13, 133.00, 129.80, 129.62, 129.60, 129.33, 129.23, 129.18, 128.41, 128.34, 128.11, 128.09, 127.97, 127.53, 109.43, 104.18, 76.08, 74.56, 74.24, 72.62, 70.93, 70.60, 68.39, 66.56, 61.97, 26.71, 25.21. API-ES-MS: *m/z* calcd for [C₄₀H₄₀O₁₁]⁺NH₄: 714.3. Found 714.3.

4.19. 3-*O*-(2-*O*-Benzyl-α-*D*-galactopyranosyl)-1,2-*O*-isopropylidene-*sn*-glycerol (**22**)

To a stirred solution of tribenzoate **20** (8.0 g, 11.4 mmol), anhydrous CH₂Cl₂ (50 mL) and MeOH (50 mL) was added a 25% solution of NaOMe in MeOH (10 mL) at rt. After 12 h, the solution was carefully treated with Dowex 50WX8 to approx. pH 5 as seen by indicator paper. The resin was removed by filtration and the filtrate was treated with an ethereal solution of CH₂N₂ until the yellow color persisted. Purification of the residue by silica gel chromatography using EtOAc as the eluant afforded solid **22** (5.0 g, 91%): ¹H NMR (CDCl₃): δ 7.38–7.29 (m, 5H), 4.95 (d, 1H, *J* = 3.5 Hz, H-1), 4.66 [s, 2H, CH₂ (Bn)] 4.32 (m, 1H), 4.08 (dd, 1H, *J* = 1.1 Hz, *J* = 3.4 Hz, H-4), 4.05 (dd, 1H, *J* = 6.5 Hz, *J* = 8.3 Hz), 3.99 (dd, 1H, *J* = 3.4 Hz, *J* = 9.8 Hz), 3.91 (dd, 1H, *J* = 5.2 Hz, *J* = 11.2 Hz), 3.85 (dt, 1H, *J* = 1.2 Hz, *J* = 4.3 Hz, *J* = 5.4 Hz), 3.81 (dd, 1H, *J* = 4.2 Hz, *J* = 11.2 Hz), 3.76 (dd, 1H, *J* = 3.5 Hz, *J* = 9.7 Hz), 3.73 (dd, 1H, *J* = 6.2 Hz, *J* = 8.2 Hz), 3.65 (dd, 1H, *J* = 4.9 Hz, *J* = 10.7 Hz), 3.46 (dd, 1H, *J* = 6.2 Hz, *J* = 10.7 Hz), 1.43 and 1.36 (2s, 2 × 3H, 2CH₃); ¹³C NMR (CDCl₃): δ 137.88, 128.63, 128.20, 128.15, 109.63, 97.11, 76.48, 74.63, 72.67, 70.29, 69.32, 68.98, 68.75, 66.47, 63.02, 26.78, 25.39. API-ES-MS: *m/z* calcd for [C₁₉H₂₈O₈]⁺NH₄: 402.2. Found 402.2.

4.20. 3-*O*-(α-*D*-Galactopyranosyl)-1,2-*O*-isopropylidene-*sn*-glycerol (**23**)

A mixture of compound **22** (480 mg), 2,4,6-tri-*tert*-butylpyrimidine (500 mg), 10% palladium-on-charcoal (0.5 g), EtOAc (10 mL) was stirred under hydrogen at 200 psi. After 10 min, the mixture was filtered and the solids were washed with MeOH. Concentration of the combined solutions under reduced pressure afforded a semisolid which was purified by silica gel column chromatography using 9:1 CH₂Cl₂–MeOH as the eluant to afford amorphous **23** (320 mg, 87%): ¹H NMR (CDCl₃): δ 4.91 (d, 1H, *J* = 3.5 Hz), 4.37 (m, 1H,

H-2'), 4.11 (dd, 1H, $J = 6.8$ Hz, $J = 8.5$ Hz), 3.97 (br d, 1H, $J = 3.2$ Hz), 3.85–3.73 (m, 7H), 3.50 (dd, 1H, $J = 6.8$ Hz, $J = 10.5$ Hz), 1.46 and 1.39 (2s, 2×3 H, 2CH_3); ^{13}C NMR (CDCl_3): δ 109.45, 98.89, 74.39, 70.25, 69.92, 69.39, 68.78, 68.68, 65.85, 61.37, 26.07, 24.72. API-ES-MS: m/z calcd for $[\text{C}_{12}\text{H}_{22}\text{O}_8]\text{NH}_4^+$: 312.2. Found 312.2.

4.21. 1,2-O-Isopropylidene-3-O-(2,3,4,6-tetra-O-levulinoyl- α -D-galactopyranosyl)-sn-glycerol (24)

To a stirred solution of tetraol **23** (320 mg, 1.09 mmol) in EtOAc (10 mL) were added sequentially levulinic acid (760 mg, 6.55 mmol), *N,N*-dicyclohexylcarbodiimide (1.6 g, 7.7 mmol) and a catalytic amount of 4-dimethylaminopyridine at rt. After 24 h, the solids were removed by filtration and washed with EtOAc). The solutions were combined and concentrated under reduced pressure. The residue was purified by silica gel column chromatography using 1:1→1:4 hexanes–EtOAc as the eluant to afford **24** (702 mg, 94%) as a syrup: ^1H NMR (CDCl_3): δ 5.43 (dd, 1H, $J = 1.2$ Hz, $J = 3.7$ Hz), 5.34 (dd, 1H, $J = 3.7$ Hz, $J = 10.6$ Hz), 5.13 (dd, 1H, $J = 3.7$ Hz, $J = 9.5$ Hz), 5.12 (br s, 1H), 4.30 (dd, 1H, $J = 5.8$ Hz, $J = 11.6$ Hz), 4.26 (t, $J = 6.6$ Hz), 4.11 (br s, 1H), 4.04 (br s, 1H), 4.08 (dd, 1H, $J = 6.4$ Hz, $J = 8.3$ Hz), 3.76 (dd, 1H, $J = 6.4$ Hz, $J = 8.3$ Hz), 3.72 (dd, 1H, $J = 5.2$ Hz, $J = 10.5$ Hz), 3.56 (dd, 1H, $J = 5.2$ Hz, $J = 10.5$ Hz), 2.82–2.41 (m, 16H), 2.19, 2.177, 2.172, 2.171 (4s, each 3H); ^{13}C NMR (CDCl_3): δ 206.51, 206.36, 206.31, 206.06, 172.18, 172.08, 171.98, 171.84, 109.52, 96.63, 74.38, 69.15, 68.20, 67.97, 67.52, 66.49, 66.45, 61.84, 37.80, 37.78, 37.75, 37.70, 29.79, 29.78, 29.76, 27.92, 27.77, 27.75, 26.71, 25.47. API-ES-MS: m/z calcd for $[\text{C}_{32}\text{H}_{46}\text{O}_{16}]\text{NH}_4^+$: 704.3. Found 704.2.

4.22. 3-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-sn-glycerol (25)

A stirred solution of **24** (1.25 g, 1.8 mmol) in a mixture of MeOH (5 mL) and AcOH (5 mL) was heated at reflux for 5 min when TLC (EtOAc) indicated disappearance of **24**. The solution was concentrated and the residue purified by column chromatography using a 1:1→1:0 EtOAc–hexanes gradient as the eluant to afford **25** (1.0 g, 83%) as a syrup: ^1H NMR (CDCl_3): δ 5.42 (dd, 1H, $J = 1.3$ Hz, $J = 3.8$ Hz), 5.37 (dd, 1H, $J = 3.8$ Hz, $J = 10.6$ Hz), 5.14 (d, 1H, $J = 3.8$ Hz), 5.09 (dd, 1H, $J = 3.8$ Hz, $J = 10.6$ Hz), 4.26 (br t, 1H, $J = 6.7$ Hz), 4.15–4.07 (m, 2H), 3.92 (m, 1H), 3.81 (dd, 1H, $J = 3.8$ Hz, $J = 10.7$ Hz), 3.72 (dd, 1H, $J = 4.3$ Hz, $J = 11.2$ Hz), 3.62 (dd, 1H, $J = 5.6$ Hz, $J = 11.2$ Hz), 3.53 (dd, 1H, $J = 6.5$ Hz, $J = 10.6$ Hz), 2.84–2.41 (m, 16H), 2.20 (s, 3H), 2.19 (s, 6H), 2.17 (s, 3H); ^{13}C NMR (CDCl_3): δ 207.77, 207.20, 206.62, 206.19, 172.18, 171.88, 171.79, 171.73, 96.40, 70.50, 69.93, 68.22, 68.10, 67.30, 66.37, 63.30, 61.92, 37.63, 37.58, 37.50, 29.65, 29.63, 29.60, 29.58, 29.50, 27.94, 27.76, 27.62, 27.59, 27.43. API-ES-MS: m/z calcd for $[\text{C}_{29}\text{H}_{42}\text{O}_{16}]\text{NH}_4^+$: 664.3. Found 664.2.

4.23. 1-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-sn-glycerol (27)

^1H NMR (CDCl_3): δ 5.42 (d, 1H, $J = 3.5$ Hz), 5.35 (dd, 1H, $J = 3.5$ Hz, $J = 10.8$ Hz), 5.17 (d, 1H, $J = 3.6$ Hz), 5.08 (dd, 1H, $J = 3.6$ Hz, $J = 10.8$ Hz), 4.24 (m, 1H), 4.16 (dd, 1H, $J = 5.2$ Hz, $J = 11.3$ Hz), 4.08 (dd, 1H, $J = 7.6$ Hz, $J = 11.3$ Hz), 3.88 (m, 1H), 3.79 (dd, 1H, $J = 5.7$ Hz, $J = 10.4$ Hz), 3.72 (d, 1H, $J = 4.5$ Hz), 3.62 (dd, 1H, $J = 5.1$ Hz, $J = 10.4$ Hz), 2.83–2.41 (m, 16H), 2.19 (s, 6H), 2.18 and 2.17 (2s, 3H each); ^{13}C NMR (CDCl_3): δ 207.75, 207.29, 206.46, 206.03, 172.31, 171.98, 171.93, 171.77, 96.19, 70.15, 69.33, 68.37, 68.28, 67.40, 66.65, 63.33, 62.31, 37.80, 37.76, 37.74, 37.66, 29.86, 29.78, 29.76, 27.87, 27.74, 27.70. API-ES-MS: m/z calcd for $[\text{C}_{29}\text{H}_{42}\text{O}_{16}]\text{NH}_4^+$: 664.3. Found 664.2.

4.24. 3-O-(2,3,4,6-Tetra-O-acetyl- β -D-galactopyranosyl)-1,2-O-isopropylidene-*sn*-glycerol (29)

To a stirred solution of imidate³⁵ **28** (4.0 g, 8 mmol) and alcohol **19** (2.1 g, 16 mmol) in anhydrous CH₂Cl₂ was added at 0 °C TMSOTf (20 μ L). After 1 h the solution was treated with a saturated aqueous solution of NaHCO₃. The usual processing followed by column chromatographic purification afforded **29** as a syrup (3.0 g, 80%): ¹H NMR (CDCl₃): δ 5.38 (dd, 1H, *J* = 1.1 Hz, *J* = 3.5 Hz), 5.21 (dd, 1H, *J* = 7.9 Hz, *J* = 10.4 Hz), 5.01 (dd, 1H, *J* = 3.5 Hz, *J* = 10.4 Hz), 4.58 (d, 1H, *J* = 7.9 Hz), 4.26 (m, 1H), 4.18 (dd, 1H, *J* = 6.7 Hz, *J* = 11.2 Hz), 4.13 (dd, 1H, *J* = 6.7 Hz, *J* = 11.2 Hz), 4.02 (dd, 1H, *J* = 6.5 Hz, *J* = 8.3 Hz), 3.92 (dt, 1H, *J* = 1.1 Hz, *J* = 6.8 Hz), 3.89 (dd, 1H, *J* = 4.4 Hz, *J* = 10.8 Hz), 3.81 (dd, 1H, *J* = 5.9 Hz, *J* = 8.3 Hz), 3.63 (dd, 1H, *J* = 6.0 Hz, *J* = 10.7 Hz), 2.15, 2.07, 2.05, 1.99, 1.42, 1.35 (6s, 6 \times 3H); ¹³C NMR (CDCl₃): δ 170.32, 170.17, 170.08, 169.35, 109.26, 101.35, 74.19, 70.82, 70.66, 69.03, 68.66, 66.96, 66.16, 61.21, 26.53, 25.09, 20.68, 20.60, 20.59, 20.51. API-ES-MS: *m/z* calcd for [C₂₀H₃₀O₁₂]NH₄⁺: 480.2. Found 480.2.

4.25. 1,2-O-Isopropylidene-3-O-(2,3,4,6-tetra-O-levulinoyl- β -D-galactopyranosyl)-*sn*-glycerol (30)

To a solution of compound **29** (3.5 g, 7.6 mmol) in anhydrous MeOH was added NaOMe in MeOH (excess) until the pH of the solution reached 12 as seen with an indicator paper. After 24 h, the solution was carefully treated with Dowex 50WX8 (H⁺) ion exchange resin until the pH reached approx. 7, then the solution was concentrated and the residue dried under vacuum to obtain a semisolid. A solution of this material in EtOAc (50 mL) was treated under stirring with levulinic acid (7.1 g, 61 mmol) followed by *N,N*-dicyclohexylcarbodiimide (12.5 g, 61 mmol). After 3 h, MeOH (5 mL) was added and stirring was continued for an additional 2 h. The solids were filtered and the filtrate concentrated followed by column chromatographic purification using a hexane–EtOAc 10:1→2:1 gradient to yield **30** (4.8 g, 93%) as a syrup: ¹H NMR (CDCl₃): δ 5.36 (d, 1H, *J* = 3.6 Hz), 5.17 (dd, 1H, *J* = 8.0 Hz, *J* = 10.4 Hz), 5.02 (dd, 1H, *J* = 3.6 Hz, *J* = 10.4 Hz), 4.57 (d, 1H, *J* = 8.2 Hz), 4.27 (m, 1H), 4.20 (dd, 1H, *J* = 6.7 Hz, *J* = 11.4 Hz), 4.11 (dd, 1H, *J* = 6.7 Hz, *J* = 11.6 Hz), 4.04 (dd, 1H, *J* = 6.7 Hz, *J* = 8.2 Hz), 3.91–3.87 (m, 2H), 3.83 (dd, 1H, *J* = 5.9 Hz, *J* = 8.2 Hz), 3.63 (dd, 1H, *J* = 6.3 Hz, *J* = 10.8 Hz), 2.83–2.42 (m, 16H), 2.19, 2.18, 2.17, 2.16, 1.41, 1.34 (6s, 6 \times 3H); ¹³C NMR (CDCl₃): δ 207.74, 206.82, 206.64, 205.30, 172.16, 171.89, 171.88, 171.41, 109.29, 101.26, 74.24, 70.82, 70.75, 69.39, 68.85, 67.23, 66.39, 61.38, 37.85, 37.78, 37.76, 37.68, 29.76, 29.75, 29.67, 27.79, 27.76, 26.67, 25.19. API-ES-MS: *m/z* calcd for [C₃₂H₄₆O₁₆]NH₄⁺: 704.3. Found 704.3.

4.26. 3-O-(2,3,4,6-Tetra-O-levulinoyl- β -D-galactopyranosyl)-*sn*-glycerol (31)

Compound **30** was treated with acetic acid as described for **24** to afford **31** as a syrup: ¹H NMR (CDCl₃): δ 5.36 (d, 1H, *J* = 3.7 Hz), 5.17 (dd, 1H, *J* = 7.9 Hz, *J* = 10.5 Hz), 5.04 (dd, 1H, *J* = 3.4 Hz, *J* = 10.5 Hz), 4.53 (d, 1H, *J* = 8.1 Hz), 4.21 (dd, 1H, *J* = 7.2 Hz, *J* = 11.4 Hz), 4.11 (dd, 1H, *J* = 6.0 Hz, *J* = 11.4 Hz), 3.94–3.90 (m, 2H), 3.86 (m, 1H), 3.77 (dd, 1H, *J* = 3.6 Hz, *J* = 10.3 Hz), 3.68 (dd, 1H, *J* = 4.3 Hz, *J* = 11.5 Hz), 3.65 (dd, 1H, *J* = 5.1 Hz, *J* = 11.5 Hz), 2.91–2.41 (m, 16H), 2.20, 2.19, 2.18, 2.16 (4s, 4 \times 3H); ¹³C NMR (CDCl₃): δ 207.71, 206.76, 206.56, 206.09, 172.18, 171.87, 171.72, 171.67, 101.49, 72.04, 70.83, 70.54, 70.34, 68.92, 67.24, 63.24, 61.54, 37.76, 37.74, 37.69, 37.57, 29.76, 29.73, 29.69, 29.61, 27.71, 27.68, 27.66. API-ES-MS: *m/z* calcd for [C₂₉H₄₂O₁₆]NH₄⁺: 664.3. Found 664.5.

4.27. General procedure for the preparation of singly Gro-substituted intermediates featuring tetra-levulinoylated Gal moieties

To a solution of the diol 3-*O*-(2,3,4,6-tetra-*O*-levulinoyl- α -D-galactopyranosyl)-*sn*-glycerol **25** (129 mg, 0.2 mmol) in anhydrous CH₂Cl₂ (3 mL) was added fatty acid (0.2 mmol) followed by DCC (60 mg, 0.29 mmol) and a catalytic amount of DMAP at room temperature. After 16 h MeOH (0.5 mL) was added followed by stirring for 1 h. TLC (CH₂Cl₂-MeOH 100:4) indicated that most of the starting diol had disappeared and two closely migrating mono-substituted products had formed. Some less polar material was also formed that was the doubly-substituted product by mass spectrometry. The mixture was concentrated, then CH₂Cl₂ was added, followed by removal of the solids by filtration. The residue was applied to a silica gel column made in CH₂Cl₂ which was then eluted with CH₂Cl₂-MeOH 100:1 to afford the mono-substituted product in 65–75% yield, free of the slower-migrating isomer.

4.28. General procedure for the preparation of fully substituted intermediates having two different fatty acyl groups in the Gro moiety

To a stirred solution of the singly Gro-substituted intermediate in anhydrous CH₂Cl₂ was added fatty acid (1.3 equiv) followed by DCC (1.5 equiv) and a catalytic amount of DMAP. After 3 h, the solution was treated with MeOH (excess). After 1 h, the mixture was concentrated. To the residue was added CH₂Cl₂ followed by removal of the solids by filtration. Column chromatographic purification of the residue with a CH₂Cl₂-MeOH mixture of appropriate polarity afforded a near quantitative yield of the doubly Gro-substituted product.

4.29. General procedure for the preparation of fully protected intermediates having two identical fatty acyl groups in the Gro moiety

To a stirred solution of 3-*O*-(2,3,4,6-tetra-*O*-levulinoyl- α -D-galactopyranosyl)-*sn*-glycerol (**25**) in anhydrous CH₂Cl₂ was added fatty acid (3 equiv) followed by DCC (6 equiv) and a catalytic amount of DMAP. After 3 h, the solution was treated with MeOH (excess). After 1 h, the mixture was concentrated and processed as described in Section 4.28.

4.30. General procedure for the removal of the levulinoyl groups

The levulinoylated material is dissolved in a 1 M solution of hydrazine hydrate in pyridine-AcOH (3:2) at rt. After 3 h, the solution is treated with 2,4-pentanedione (excess). The solution is concentrated under reduced pressure and the residue is purified by silica gel column chromatography. Depending on the acyl groups attached to the Gro moiety, the product may precipitate as a solid material during work-up.

4.31. 3-*O*-(2,3,4,6-Tetra-*O*-levulinoyl- α -D-galactopyranosyl)-1-*O*-palmitoyl-*sn*-glycerol (**32**)

¹H NMR (CDCl₃): δ 5.42 (dd, 1H, $J = 1.4$ Hz, $J = 3.5$ Hz), 5.36 (dd, 1H, $J = 3.5$ Hz, $J = 10.7$ Hz), 5.16 (d, 1H, $J = 3.7$ Hz), 5.09 (dd, 1H, $J = 3.7$ Hz, $J = 10.7$ Hz), 4.25 (ddd, 1H), 4.18 (dd, 1H, $J = 4.6$ Hz, $J = 11.3$ Hz), 4.13 (dd, 1H, $J = 6.1$ Hz, $J = 11.2$ Hz), 4.09 (dd, 1H, $J = 7.0$ Hz, $J = 11.2$ Hz), 3.81 (dd, 1H, $J = 3.4$ Hz, $J = 10.9$ Hz), 3.57 (d, 1H, $J = 4.4$ Hz), 3.51 (dd, 1H, $J = 7.0$ Hz, $J = 10.9$ Hz), 2.84–2.41 (m, 16H), 2.34 (dd, 1H, $J = 7.3$ Hz, $J = 7.8$ Hz), 2.189, 2.187, 2.181, 2.17 (4s, 4 \times 3H), 1.66–1.58 (m, 2H), 1.35–1.20 (m, 24H), 0.91–0.84 (m, 3H); ¹³C NMR (CDCl₃): δ 207.35, 206.63, 206.40, 205.97, 173.71, 172.11, 171.86, 171.72, 171.67, 96.76, 70.18, 68.68, 68.15, 68.07, 67.27, 66.51, 64.91, 61.89, 37.66, 37.62, 37.59, 37.55, 33.99, 31.79, 29.66, 29.63, 29.62, 29.56, 29.53, 29.49, 29.43, 29.36, 29.23,

29.17, 29.04, 27.84, 27.66, 27.64, 27.60, 24.78, 22.56, 14.00. API-ES-MS: m/z calcd for $[\text{C}_{45}\text{H}_{72}\text{O}_{17}]\text{NH}_4^+$: 902.5. Found 902.4.

4.32. 3-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-1-O-oleoyl-*sn*-glycerol (33)

^1H NMR (CDCl_3): δ 5.42 (d, 1H, $J = 3.4$ Hz), 5.36 (dd, 1H, $J = 3.6$ Hz, $J = 10.4$ Hz), 5.36–5.32 (m, 2H), 5.16 (d, 1H, $J = 3.7$ Hz), 5.09 (dd, 1H, $J = 3.7$ Hz, $J = 10.7$ Hz), 4.24 (ddd, 1H), 4.18 (dd, 1H, $J = 4.4$ Hz, $J = 11.3$ Hz), 4.13 (dd, 1H, $J = 5.8$ Hz, $J = 10.7$ Hz), 4.09 (dd, 1H, $J = 7.2$ Hz, $J = 11.3$ Hz), 3.81 (dd, 1H, $J = 3.4$ Hz, $J = 10.7$ Hz), 3.51 (dd, 1H, $J = 6.8$ Hz, $J = 10.7$ Hz), 2.84–2.41 (m, 16H), 2.34 (t, 2H, $J = 7.6$ Hz), 2.188 (s, 6H), 2.181 (s, 3H), 2.17 (s, 3H), 2.04–1.98 (m, 2H), 1.65–1.59 (m, 2H), 1.36–1.24 (m), 0.89–0.86 (m, 3H); ^{13}C NMR (CDCl_3): δ 206.50, 206.42, 206.22, 205.98, 173.78, 172.20, 171.95, 171.80, 171.76, 129.96, 129.72, 96.86, 70.31, 68.82, 68.25, 68.16, 67.33, 66.60, 65.03, 61.99, 53.42, 37.76, 37.72, 37.69, 37.66, 34.08, 31.88, 29.77, 29.76, 29.74, 29.72, 29.69, 29.49, 29.29, 29.18, 29.10, 27.94, 27.74, 27.72, 27.69, 27.19, 27.16, 24.86, 22.66, 14.09. API-ES-MS: m/z calcd for $[\text{C}_{47}\text{H}_{74}\text{O}_{17}]\text{NH}_4^+$: 928.5. Found 928.4.

4.33. 3-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-1-O-stearoyl-*sn*-glycerol (34)

^1H NMR (CDCl_3): δ 5.42 (d, 1H, $J = 3.6$ Hz), 5.36 (dd, 1H, $J = 3.6$ Hz, $J = 11.0$ Hz), 5.16 (d, 1H, $J = 3.7$ Hz), 5.09 (dd, 1H, $J = 3.7$ Hz, $J = 11.0$ Hz), 4.25 (t, 1H, $J = 6.6$ Hz), 4.19–4.07 (m, 5H), 3.81 (dd, 1H, $J = 3.1$ Hz, $J = 10.6$ Hz), 3.58 (d, 1H, $J = 3.9$ Hz), 3.51 (dd, 1H, $J = 6.7$ Hz, $J = 10.6$ Hz), 2.84–2.41 (m, 16H), 2.34 (t, 2H, $J \sim 7.6$ Hz), 2.19 (s, 6H), 2.18 and 2.17 (2s, 2 \times 3H), 1.62 (m, 2H), 1.38–1.18 (m, 28H), 0.88 (t, 3H, $J \sim 6.7$ Hz); ^{13}C NMR (CDCl_3): δ 207.31, 206.59, 206.37, 205.93, 173.67, 172.07, 171.83, 171.69, 171.63, 96.73, 70.16, 68.65, 68.13, 68.04, 67.25, 66.48, 64.88, 61.86, 37.63, 37.60, 37.57, 37.53, 33.97, 31.77, 29.69–29.02, 27.82, 27.63, 27.61, 27.57, 24.76, 22.54, 13.98. API-ES-MS: m/z calcd for $[\text{C}_{47}\text{H}_{76}\text{O}_{17}]\text{NH}_4^+$: 930.5. Found 930.4.

4.34. 1-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-3-O-palmitoyl-*sn*-glycerol (35)

^1H NMR (CDCl_3): δ 5.42 (dd, 1H, $J = 1.2$ Hz, $J = 3.4$ Hz), 5.34 (dd, 1H, $J = 3.4$ Hz, $J = 10.7$ Hz), 5.15 (d, 1H, $J = 3.8$ Hz), 5.10 (dd, 1H, $J = 3.8$ Hz, $J = 10.7$ Hz), 4.24 (t, 1H, $J \sim 6.5$ Hz), 4.18–4.14 (m, 2H), 4.08 (dd, 1H, $J = 7.5$ Hz, $J = 11.4$ Hz), 4.05–4.01 (m, 1H), 3.75 (dd, 1H, $J = 5.7$ Hz, $J = 11.0$ Hz), 3.60 (dd, 1H, $J = 4.4$ Hz, $J = 10.8$ Hz), 3.46 (d, 1H, $J = 5.6$ Hz), 2.83–2.41 (m, 16H), 2.34 (d, 2H, $J \sim 7.6$ Hz), 2.19 (s, 9H), 2.17 (s, 3H), 1.66–1.58 (m, 2H), 1.35–1.21 (m, 24H), 0.88 (t, 3H, $J \sim 7$ Hz); ^{13}C NMR (CDCl_3): δ 207.18, 206.98, 206.45, 206.02, 173.86, 172.23, 171.96, 171.94, 171.76, 96.39, 76.51, 69.53, 68.46, 6.31, 68.13, 67.39, 66.68, 65.12, 62.17, 37.76, 37.71, 37.69, 34.14, 31.92, 29.83, 29.75, 29.69, 29.66, 29.48, 29.35, 29.30, 29.18, 27.92, 27.75, 29.69, 24.91, 22.69, 14.12. API-ES-MS: m/z calcd for $[\text{C}_{45}\text{H}_{72}\text{O}_{17}]\text{NH}_4^+$: 902.5. Found 902.4.

4.35. 1-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-3-O-oleoyl-*sn*-glycerol (36)

^1H NMR (CDCl_3): δ 5.42 (d, 1H, $J = 3.3$ Hz), 5.37–5.30 (m, 3H), 5.14 (d, 1H, $J = 3.6$ Hz), 5.10 (dd, 1H, $J = 3.6$ Hz, $J = 10.6$ Hz), 3.24 (m, 1H), 4.20–4.11 (m, 3H), 4.07 (dd, 1H, $J = 7.4$ Hz, $J = 11.3$ Hz), 3.76 (dd, 1H, $J = 5.5$ Hz, $J = 10.8$ Hz), 2.83–2.41 (m, 16H), 2.32 (t, 2H, $J \sim 7.6$ Hz), 2.19 (s, 9H), 2.17 (s, 3H), 2.03–1.99 (m, 4H), 1.67–1.28 (m, 2H), 1.39–1.20 (m, 20H), 0.89–0.86 (m, 3H); ^{13}C NMR (CDCl_3): δ 207.13, 206.94, 206.42, 205.93, 173.74, 172.16, 171.90, 171.87, 171.69, 129.91, 129.67, 96.31, 69.44, 68.36, 68.22, 58.06, 67.32, 66.59, 65.05, 62.08, 37.69, 37.68, 37.64, 37.60, 34.04, 35.00, 31.81, 29.74, 29.68, 29.64, 29.43, 29.23, 29.18, 29.12, 29.06, 29.05, 27.85, 27.69, 27.67, 27.62, 27.13, 27.10, 24.82, 22.59, 14.04. API-ES-MS: m/z calcd for $[\text{C}_{47}\text{H}_{74}\text{O}_{17}]\text{NH}_4^+$: 928.5. Found 928.4.

4.36. 3-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-1,2-di-O-palmitoyl-*sn*-glycerol (37)

^1H NMR (CDCl_3): δ 5.43 (dd, 1H, $J = 1.2$ Hz, $J = 3.4$ Hz), 5.32 (dd, 1H, $J = 3.4$ Hz, $J = 10.8$ Hz), 5.22 (m, 1H), 5.12 (dd, 1H, $J = 3.7$ Hz, $J = 10.8$ Hz), 5.06 (d, 1H, $J = 3.7$ Hz), 4.37 (dd, 1H, 3.9 Hz, $J = 11.8$ Hz), 4.20 (ddd, 1H), 4.18–4.06 (m, 3H), 3.83 (dd, 1H, $J = 4.5$ Hz, $J = 11.2$ Hz), 3.65 (dd, 1H, $J = 5.0$ Hz, $J = 11.0$ Hz), 2.83–2.42 (m, 16H), 2.33 and 2.30 (2t, $2 \times 2\text{H}$, $J \sim 7.8$ Hz), 1.66–1.56 (m, 4H), 1.36–1.20 (m, 44H), 0.92–0.85 (m, 6H); ^{13}C NMR (CDCl_3): δ 206.40, 206.35, 206.30, 206.00, 173.25, 172.97, 172.09, 172.08, 171.90, 171.79, 96.58, 69.73, 67.99, 67.81, 67.44, 66.63, 66.47, 62.19, 61.65, 37.73, 37.68, 37.66, 34.20, 34.01, 31.88, 29.69, 29.66, 29.61, 29.49, 29.47, 29.31, 29.29, 29.27, 29.11, 29.10, 27.81, 27.74, 27.71, 27.65, 24.89, 24.84, 22.64, 14.07. API-ES-MS: m/z calcd for $[\text{C}_{61}\text{H}_{102}\text{O}_{18}]\text{NH}_4^+$: 1140.7. Found 1160.6.

4.37. 3-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-1,2-di-O-oleoyl-*sn*-glycerol (38)

^1H NMR (CDCl_3): δ 5.43 (d, 1H, $J = 3.7$ Hz), 5.38–5.30 (m 5H), 5.22 (m, 1H), 5.13 (dd, 1H, $J = 3.7$ Hz, $J = 10.8$ Hz), 5.06 (d, 1H, $J = 3.7$ Hz), 4.37 (dd, 1H, $J = 4.0$ Hz, $J = 11.8$ Hz), 4.22–4.06 (m, 4H), 3.83 (dd, 1H, $J = 4.5$ Hz, $J = 11.2$ Hz), 3.65 (dd, 1H, $J = 5.0$ Hz, $J = 11.2$ Hz), 2.83–2.42 (m, 16H), 2.33 and 2.30 (2t, $2 \times 3\text{H}$, $J \sim 7.7$ Hz), 2.18 (s, 3H), 2.177 (s, 3H), 2.170 (s, 6H), 2.05–1.97 (m, 8H), 1.66–1.56 (m, 4H), 1.38–1.20 (m, 42H), 0.88 (t, 6H, $J \sim 7.0$ Hz); ^{13}C NMR (CDCl_3): δ 206.35, 206.12, 205.90, 173.17, 172.89, 172.04, 172.03, 171.87, 171.75, 129.90, 129.63, 96.55, 69.70, 67.95, 67.77, 67.40, 66.58, 66.44, 62.15, 61.61, 37.68, 37.64, 37.62, 34.13, 33.94, 31.81, 29.67, 29.66, 29.65, 29.43, 29.22, 29.15, 29.13, 29.09, 29.06, 29.03, 29.02, 27.77, 27.70, 27.67, 27.62, 27.13, 27.11, 27.10, 24.82, 24.77, 22.59, 14.03. APIES-MS: m/z calcd for $[\text{C}_{65}\text{H}_{106}\text{O}_{18}]\text{NH}_4^+$: 1192.8. Found 1192.6.

4.38. 3-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-2-O-oleoyl-1-O-palmitoyl-*sn*-glycerol (39)

^1H NMR (CDCl_3): δ 5.43 (d, 1H, $J = 1.2$ Hz, $J = 3.5$ Hz), 5.38–5.30 (m, 3H), 5.22 (m, 1H), 5.12 (dd, 1H, $J = 3.7$ Hz, $J = 10.7$ Hz), 4.37 (dd, 1H, $J = 4.0$ Hz, 12.0 Hz), 4.22–4.06 (m, 4H), 3.83 (dd, 1H, $J = 4.6$ Hz, $J = 11.1$ Hz), 3.65 (4.9 Hz, $J = 11.3$ Hz), 2.82–2.42 (m, 16H), 2.33 and 2.30 (2t, $s \times 2\text{H}$, $J = 7.5$ Hz for each), 2.05–1.96 (m, 4H), 1.65–1.55 (m, 4H), 1.37–1.19 (m), 0.88 (t, 6H, $J = 6.7$ Hz); ^{13}C NMR (CDCl_3): δ 206.41, 206.18, 206.96, 173.25, 172.94, 172.09, 172.08, 171.91, 171.80, 129.94, 129.66, 96.58, 69.73, 67.99, 67.81, 67.44, 66.61, 66.47, 62.17, 61.65, 37.72, 37.67, 37.66, 34.17, 34.00, 31.87, 31.85, 29.71, 29.70, 29.69, 29.68, 29.65, 29.62, 29.60, 29.47, 29.46, 29.31, 29.26, 29.19, 29.12, 29.10, 29.06, 27.80, 27.73, 27.70, 27.65, 27.17, 27.15, 24.86, 24.83, 22.64, 22.63, 14.06. API-ES-MS: m/z calcd for $[\text{C}_{63}\text{H}_{104}\text{O}_{18}]\text{NH}_4^+$: 1166.8. Found 1166.6.

4.39. 3-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-1-O-oleoyl-2-O-palmitoyl-*sn*-glycerol (40)

^1H NMR (CDCl_3): δ 5.43 (d, 1H, $J = 3.7$ Hz), 5.36–5.29 (m, 3H), 5.22 (m, 1H), 5.12 (dd, 1H, $J = 3.9$ Hz, $J = 10.6$ Hz), 5.06 (d, 1H, $J = 3.5$ Hz), 4.36 (dd, 1H, $J = 4.1$ Hz, $J = 12.3$ Hz), 4.21–4.06 (m, 4H), 3.83 (dd, 1H, $J = 4.5$ Hz, $J = 11.3$ Hz), 3.65 (dd, 1H, $J = 5.0$ Hz, $J = 11.0$ Hz), 2.82–2.42 (m, 16H), 2.18 and 2.176 (2s, $2 \times 3\text{H}$), 2.172 (s, 3H), 2.33 and 2.30 (2t, $2 \times 2\text{H}$, $J = 7.5$ Hz), 2.01 (m, 4H), 1.62 (m, 4H), 1.37–1.21 (m), 0.88 (t, 6H, $J = 7.3$ Hz); ^{13}C NMR (CDCl_3): δ 206.45, 206.23, 206.22, 205.99, 173.29, 173.03, 172.15, 172.14, 171.97, 171.86, 130.01, 129.73, 96.64, 69.79, 68.05, 67.87, 67.50, 66.70, 66.53, 62.26, 61.71, 37.79, 36.74, 37.73, 34.25, 34.05, 31.93, 31.91, 31.79, 29.77, 29.75, 29.72, 29.71, 29.68, 29.55, 29.54, 29.46, 29.38, 29.35, 29.34, 29.33, 29.24, 29.17, 29.14, 27.86, 27.80, 27.76, 27.71, 27.23, 27.20, 24.95, 24.86, 22.70, 22.69, 14.13. API-ES-MS: m/z calcd for $[\text{C}_{63}\text{H}_{104}\text{O}_{18}]\text{NH}_4^+$: 1166.8. Found 1166.6.

4.40. 3-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-1,2-di-O-stearoyl-*sn*-glycerol (41)

$^1\text{H NMR}$ (CDCl_3): δ 5.43 (d, 1H, $J = 3.4$ Hz), 5.32 (dd, 1H, $J = 3.4$ Hz, $J = 10.7$ Hz), 5.22 (m, 1H), 5.12 (dd, 1H, $J = 3.6$ Hz, $J = 10.7$ Hz), 5.06 (d, 1H, $J = 3.6$ Hz), 4.37 (dd, 1H, $J = 3.9$ Hz, $J = 12.0$ Hz), 4.22–4.06 (m, 4H), 3.83 (d, 1H, $J = 4.6$ Hz, $J = 11.2$ Hz), 3.66 (dd, 1H, $J = 5.1$ Hz, $J = 11.2$ Hz), 2.86–2.41 (m, 16H), 2.33 and 2.30 (2t, $2 \times 2\text{H}$, $J \sim 7.5$ Hz), 2.18 and 2.17 (2s, $2 \times 3\text{H}$), 2.16 (s, 6H), 1.66–1.56 (m, 4H), 1.59–1.21 (m, 56H), 0.88 (t, 6H, $J \sim 7.0$ Hz); $^{13}\text{C NMR}$ (CDCl_3): δ 206.32, 206.10, 205.87, 173.18, 172.91, 172.03, 172.01, 171.85, 171.74, 96.52, 69.68, 67.94, 67.76, 67.39, 66.58, 66.42, 62.14, 61.60, 37.67, 37.62, 37.61, 34.14, 33.96, 31.82, 29.74–29.04, 27.76, 27.69, 27.66, 27.60, 24.84, 24.78, 22.59, 14.02. API-ES-MS: m/z calcd for $[\text{C}_{65}\text{H}_{110}\text{O}_{18}]\text{NH}_4^+$: 1196.8. Found 1197.0.

4.41. 1-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-2,3-di-O-palmitoyl-*sn*-glycerol (42)

$^1\text{H NMR}$ (CDCl_3): δ 5.42 (dd, 1H, $J = 1.3$ Hz, $J = 3.5$ Hz), 5.30 (dd, 1H, $J = 3.5$ Hz, $J = 10.7$ Hz), 5.20 (m, 1H), 5.12 (dd, 1H, $J = 3.7$ Hz, $J = 10.7$ Hz), 5.08 (d, 1H, $J = 3.7$ Hz), 4.32 (dd, 1H, $J = 3.7$ Hz, $J = 12.5$ Hz), 4.21–4.17 (m, 2H), 4.11 (s, 1H), 4.09 (m, 1H), 3.82 (dd, 1H, $J = 5.4$ Hz, $J = 11.1$ Hz), 3.63 (dd, 1H, $J = 5.4$ Hz, $J = 11.1$ Hz), 2.83–2.42 (m, 16H), 2.31 and 2.30 (2t, $2 \times 2\text{H}$, $J \sim 7.5$ Hz), 2.18, 2.176, 2.174, 2.171 (4s, $3 \times 3\text{H}$), 1.65–1.56 (m, 4H), 1.34–1.21 (m, 48H), 0.89–0.86 (m, 6H); $^{13}\text{C NMR}$ (CDCl_3): δ 206.37, 206.26, 206.20, 206.19, 173.21, 172.78, 172.07, 172.03, 171.85, 171.71, 96.61, 69.67, 67.97, 67.75, 67.31, 66.55, 66.28, 62.18, 61.63, 37.70, 37.68, 37.63, 37.61, 34.10, 33.97, 31.83, 29.66, 29.64, 29.61, 29.58, 29.57, 29.43, 29.42, 29.27, 29.23, 29.22, 29.06, 29.04, 27.77, 27.69, 27.66, 27.63, 24.81, 24.78, 22.60, 14.03. API-ES-MS: m/z calcd for $[\text{C}_{61}\text{H}_{102}\text{O}_{18}]\text{NH}_4^+$: 1140.7. Found 1140.6.

4.42. 1-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-2,3-di-O-oleoyl-*sn*-glycerol (43)

$^1\text{H NMR}$ (CDCl_3): δ 5.42 (dd, 1H, $J = 1.1$ Hz, $J = 3.4$ Hz), 5.37–5.31 (m, 4H), 5.30 (dd, 1H, $J = 3.4$ Hz, $J = 10.7$ Hz), 5.20 (m, 1H), 5.12 (dd, 1H, $J = 3.8$ Hz, $J = 10.7$ Hz), 5.08 (d, 1H, $J = 3.8$ Hz), 4.32 (dd, 1H, $J = 3.8$ Hz, $J = 12.0$ Hz), 4.1–4.17 (m, 2H), 4.09 (m, 1H), 3.82 (dd, 1H, $J = 5.3$ Hz, $J = 11.2$ Hz), 3.63 (dd, 1H, $J = 3.63$ Hz, $J = 11.2$ Hz), 2.82–2.42 (m, 16H), 2.31 and 2.30 (2t, $2 \times 2\text{H}$, $J \sim 7.5$ Hz), 2.183, 2.176, 2.173, 2.170 (4s, 3H each), 2.03–1.99 (m, 4H), 1.36–1.22 (m, 40H), 0.89–0.86 (m, 6H); $^{13}\text{C NMR}$ (CDCl_3): δ 206.48, 206.25, 206.20, 206.17, 173.21, 172.77, 172.09, 172.05, 171.87, 172.77, 172.09, 172.05, 171.88, 171.74, 129.92, 129.64, 96.63, 69.70, 67.99, 7.77, 67.33, 66.57, 66.30, 62.20, 61.64, 37.72, 37.70, 37.66, 37.63, 34.09, 33.96, 31.83, 29.69, 29.66, 29.45, 29.24, 29.15, 29.09, 29.07, 29.05, 29.02, 27.79, 27.70, 27.68, 27.65, 27.15, 27.12, 27.11, 24.80, 24.78, 22.60, 14.04. API-ES-MS: m/z calcd for $[\text{C}_{65}\text{H}_{106}\text{O}_{18}]\text{NH}_4^+$: 1192.8. Found 1192.6.

4.43. 1-O-(2,3,4,6-Tetra-O-levulinoyl- α -D-galactopyranosyl)-2-O-oleoyl-3-O-palmitoyl-*sn*-glycerol (44)

$^1\text{H NMR}$ (CDCl_3): δ 5.43 (d, 1H, $J = 3.4$ Hz), 5.37–5.27 (m, 3H), 5.20 (m, 1H), 5.12 (dd, 1H, $J = 3.7$ Hz, $J = 10.7$ Hz), 5.08 (d, 1H, $J = 3.7$ Hz), 4.32 (dd, 1H, $J = 3.7$ Hz, $J = 11.9$ Hz), 4.22–4.16 (m, 2H), 4.12–4.08 (m, 2H), 3.82 (dd, 1H, $J = 5.3$ Hz, $J = 11.1$ Hz), 3.63 (dd, 1H, $J = 5.3$ Hz, $J = 11.1$ Hz), 2.82–2.42 (m, 16H), 2.31 and 2.30 (2t, 2H, $J \sim 7.6$ Hz), 2.180 (s, 3H), 2.175 (s, 6H), 2.172 (s, 3H), 2.06–1.98 (m, 4H), 1.66–1.56 (m, 4H), 1.38–1.20 (m, 44H), 0.92–0.86 (m, 3H); $^{13}\text{C NMR}$ (CDCl_3): δ 206.44, 206.26, 206.18, 205.99, 173.33, 172.86, 172.18, 172.14, 171.96, 171.83, 130.01, 129.72, 96.73, 69.78, 68.07, 67.86, 67.41, 66.65, 66.38, 62.27, 61.73, 37.81, 37.79, 37.74, 37.72, 34.18, 34.06, 31.93, 31.91, 29.78, 29.76, 29.71, 29.69, 29.67, 29.54, 29.52, 29.37, 29.34, 29.33, 29.24, 29.18, 29.11, 27.87,

27.78, 27.76, 27.73, 27.24, 27.21, 24.89, 22.70, 14.13. API-ES-MS: m/z calcd for $[C_{63}H_{104}O_{18}]NH_4^+$: 1166.8. Found 1166.6.

4.44. 1-*O*-(2,3,4,6-Tetra-*O*-levulinoyl- α -*D*-galactopyranosyl)-3-*O*-oleoyl-2-*O*-palmitoyl-*sn*-glycerol (45)

1H NMR ($CDCl_3$): δ 5.42 (d, 1H, $J = 3.4$ Hz), 5.38–5.32 (m, 2H), 5.30 (dd, 1H, $J = 3.4$ Hz, $J = 10.6$ Hz), 5.20 (m, 1H), 5.12 (dd, 1H, $J = 3.6$ Hz, $J = 10.6$ Hz), 5.08 (dd, 1H, $J = 3.6$ Hz), 4.32 (dd, 1H, $J = 3.6$ Hz, $J = 11.8$ Hz), 4.22–4.17 (m, 2H), 4.12–4.08 (m, 2H), 3.82 (dd, 1H, $J = 5.3$ Hz, $J = 11.1$ Hz), 3.63 (dd, 1H, $J = 5.3$ Hz, $J = 11.1$ Hz), 2.83–2.42 (m, 16H), 2.307 and 2.305 (2t, $2 \times 2H$, $J \sim 7.4$ Hz each), 2.18 (s, 3H), 2.175 (s, 6H), 2.171 (s, 3H), 2.06–1.97 (m, 4H), 1.67–1.56 (m, 4H), 1.42–1.19 (m, 44H), 0.94–0.85 (m, 6H); ^{13}C NMR ($CDCl_3$): δ 206.44, 206.26, 206.18, 205.99, 173.29, 172.89, 172.18, 172.14, 171.96, 171.82, 130.01, 129.73, 96.72, 69.77, 68.07, 67.86, 67.41, 66.65, 66.39, 62.29, 61.73, 37.81, 37.79, 37.74, 37.72, 34.20, 34.05, 31.93, 31.91, 29.78, 29.75, 29.72, 29.70, 29.67, 29.54, 29.37, 29.33, 29.23, 29.16, 29.14, 27.87, 27.78, 27.76, 27.73, 27.23, 27.20, 24.91, 24.87, 22.70, 22.69, 14.13. API-ES-MS: m/z calcd for $[C_{63}H_{104}O_{18}]NH_4^+$: 1166.8. Found 1166.6.

4.45. 3-*O*- α -*D*-Galactopyranosyl)-1,2-di-*O*-oleoyl-*DL*-glycerol (46)

1H NMR ($CDCl_3$): δ 5.39–5.28 (m, 5H), 4.40 (dd, 1H, 3.6 Hz, $J = 12.5$ Hz), 4.28 (d, 1H, $J = 7.6$ Hz), 4.21 (dd, 1H, $J = 6.7$ Hz, $J = 12.0$ Hz), 4.14 (dd, 1H, $J = 6.0$ Hz, $J = 12.0$ Hz), 4.02–3.84 (m, 4H), 3.77 (dd, 1H, $J = 6.4$ Hz, $J = 11.5$ Hz), 3.75 (dd, 1H, $J = 6.4$ Hz, $J = 11.5$ Hz), 3.66 (t, 1H, $J = 8.4$ Hz), 3.61–3.57 (m, 1H), 3.55 (m, 2H), 2.35–2.29 (m, 4H), 2.06–1.96 (m, 8H), 1.67–1.54 (m, 6H), 1.38–1.19 (m, 38H), 0.88 (t, 6H, $J \sim 7.2$ Hz); ^{13}C NMR ($CDCl_3$): δ 173.76, 173.70, 173.52, 173.51, 130.06, 129.72, 104.06, 103.65, 74.53, 74.40, 73.46, 73.36, 71.76, 71.67, 70.19, 70.14, 69.58, 68.49, 68.26, 63.02, 62.96, 62.70, 62.44, 34.33, 34.31, 34.14, 34.13, 31.92, 29.79, 29.74, 29.55, 29.35, 29.34, 29.21, 29.15, 29.12, 29.08, 27.25, 27.20, 24.91, 24.86, 22.70, 14.13. API-ES-MS: m/z calcd for $[C_{45}H_{82}O_{10}]NH_4^+$: 800.6. Found 800.5.

5. Note added after the review process

An anonymous referee suggested the use of Lemieux's halideion catalysis protocol. Accordingly, tetra-*O*-benzyl-*D*-galactopyranosyl bromide, obtained in situ by the action of bromine on the corresponding ethylthio-galactoside, was treated with 1,2-*O*-isopropylidene-*sn*-glycerol (6 molar equiv) in the presence of Bu_4NBr and Hünig's base for 4 days at rt to afford the expected α -galactopyranosyl-glycerol derivative together with an inseparable minor product (<5%, NMR) in a combined yield of 78% yield (for two steps).

Acknowledgments

This work was supported by the intramural programs of the Eunice Kennedy Shriver National Institute of Child Health and Human Development, and the National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD.

Abbreviations

API-ES-MS	atmospheric pressure ionization electrospray mass spectrometry
BBGL	<i>Borrelia burgdorferi</i> glyco lipid
DCC	<i>N,N</i> -dicyclohexylcarbodiimide
DMAP	4-dimethylaminopyridine

FA	fatty acid
Gal	galactosyl
Gro	glycerol
IL	interleukin
iNKR	invariant natural killer T cell
LD	Lyme disease
Ole	oleoyl
Pal	palmitoyl
sn	stereochemical nomenclature
Ste	stearoyl
TLC	thin layer chromatography

References

- Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. *Science*. 1982; 216:1317–1319. [PubMed: 7043737]
- <http://www.cdc.gov>
- Bacon RM, Kugeler KJ, Mead PS. *MMWR*. 2008; 57:1–9. [PubMed: 18830214]
- Steere AC, Sikand VK, Meurice F, Parenti DL, Fikrig E, Schoen RT, Nowakowski J, Schmid CH, Laukamp S, Buscarino C, Krause DS. Lyme Dis Vaccine Study Grp. *N. Eng. J. Med.* 1998; 339:209–215.
- Gross DM, Forsthuber T, Tary-Lehmann M, Etling C, Ito K, Nagy ZA, Field JA, Steere AC, Huber BT. *Science*. 1998; 281:703–706. [PubMed: 9685265]
- Nigrovic LE, Thompson KM. *Epidemiol. Infect.* 2007; 135:1–8. [PubMed: 16893489]
- Ball R, Shadomy SV, Meyer A, Huber BT, Leffell MS, Zachary A, Belotto M, Hilton E, Bryant-Genevier M, Schriefer ME, Miller FW, Braun MM. *Arthritis Rheum.* 2009; 60:1179–1186. [PubMed: 19333928]
- Shen AK, Mead PS, Beard C. *Clin. Infect. Dis.* 2011; 52:S247–S252. [PubMed: 21217171]
- Plotkin SA. *Clin. Infect. Dis.* 2011; 52:S271–S275. [PubMed: 21217175]
- Takayama K, Rothenberg RJ, Barbour AG. *Infect. Immun.* 1987; 55:2311–2313. [PubMed: 3623705]
- Ben-Menachem G, Kubler-Kielb J, Coxon B, Yergey A, Schneerson R. *Proc. Natl. Acad. Sci. U.S.A.* 2003; 100:7913–7918. [PubMed: 12799465]
- Hossain H, Wellensiek HJ, Geyer R, Lochnit G. *Biochimie.* 2001; 83:683–692. [PubMed: 11522398]
- Schroder NWJ, Schombel U, Heine H, Gobel UB, Zähringer U, Schumann R. *Biol. Chem.* 2003; 278:33645–33653.
- Stubs G, Fingerle V, Wilske B, Gobel UB, Zähringer U, Schumann RR, Schroder NWJ. *J. Biol. Chem.* 2009; 284:13326–13334. [PubMed: 19307181]
- Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O, Ketchum KA, Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb JF, Fleischmann RD, Richardson D, Peterson J, Kerlavage AR, Quackenbush J, Salzberg S, Hanson M, van Vugt R, Palmer N, Adams MD, Gocayne J, Weidman J, Utterback T, Watthey L, McDonald L, Artiach P, Bowman C, Garland S, Fujii C, Cotton MD, Horst K, Roberts K, Hatch B, Smith HO, Venter JC. *Nature*. 1997; 390:580–586. [PubMed: 9403685]
- Jones KL, Seward RJ, Ben-Menachem G, Glickstein LJ, Costello CE, Steere AC. *Clin. Immunol.* 2009; 132:93–102. [PubMed: 19342303]

17. Wu D, Xing GW, Poles MA, Horowitz A, Kinjo Y, Sullivan B, Bodmer-Narkevitch V, Plettenburg O, Kronenberg M, Tsuji M, Ho DD, Wong CH. *Proc. Natl. Acad. Sci. U.S.A.* 2005; 102:1351–1356. [PubMed: 15665086]
18. Kinjo Y, Tupin E, Wu D, Fujio M, Garcia-Navarro R, Benhnia MREI, Zajonc DM, Ben-Menachem G, Ainge GD, Painter GF, Khurana A, Hoebe K, Behar SM, Beutler B, Wilson IA, Tsuji M, Sellati TJ, Wong CH, Kronenberg M. *Nat. Immunol.* 2006; 7:978–986. [PubMed: 16921381]
19. Du W, Kulkarni SS, Gervay-Hague J. *Chem. Commun.* 2007:2336–2338.
20. Schombs M, Park FE, Du WJ, Kulkarni SS, Gervay-Hague J. *J. Org. Chem.* 2010; 75:4891–4898. [PubMed: 20387787]
21. Xia J, Hui YZ. *Tetrahedron: Asymmetry.* 1997; 8:3019–3021.
22. Fraser BH, Perlmutter P, Wijesundera C. *J. Am. Oil Chem. Soc.* 2007; 84:11–21.
23. Andrews PC, Fraser BH, Junk PC, Massi M, Perlmutter P, Thienthong N, Wijesundera C. *Tetrahedron.* 2008; 64:9197–9202.
24. Fureby AM, Virto C, Adlercreutz P, Mattiasson B. *Biocatal. Biotransform.* 1996; 14:89–111.
25. Greimel P, Ito Y. *Tetrahedron Lett.* 2008; 49:3562–3566.
26. Martin SF, Josey JA, Wong YL, Dean DW. *J. Org. Chem.* 1994; 59:4805–4820.
27. Rosseto R, Bibak N, Hajdu J. *Org. Biomol. Chem.* 2006; 4:2358–2360. [PubMed: 16763679]
28. Pozsgay V, Kubler-Kielb J, Coxon B, Ekborg G. *Tetrahedron.* 2005; 61:10470–10481.
29. Pozsgay V, Kubler-Kielb J. *Carbohydr. Res.* 2007; 342:621–626. [PubMed: 17182019]
30. Khair N, Martin-Lomas M. *J. Org. Chem.* 1995; 60:7017–7021.
31. Osswald M, Lang U, Friedrich-Bochnitschek S, Pfrengle W, Kunz HZ. *Naturforsch., B.* 2003; 58:764–774.
32. van Boom JH, Burgers PMJ. *Tetrahedron Lett.* 1976:4875–4878.
33. Takatani M, Matsuo I, Ito Y. *Carbohydr. Res.* 2003; 338:1073–1081. [PubMed: 12706973]
34. Pozsgay V. *Tetrahedron: Asymmetry.* 2000; 11:151–172.
35. Eisele T, Windmüller R, Schmidt RR. *Carbohydr. Res.* 1998; 306:81–91. [PubMed: 9691441]
36. Cinco M, Banfi E, Balanzin D, Godeas C, Panfili E. *FEMS Microbiol. Immunol.* 1991; 3:33–38. [PubMed: 1711876]
37. Eiffert H, Lotter H, Jareckikhan K, Thomssen R. *Med. Microbiol. Immunol.* 1991; 180:229–237. [PubMed: 1722277]
38. Wheeler CM, Monco JCG, Benach JL, Golightly MG, Habicht GS, Steere AC. *J. Infect. Dis.* 1993; 167:665–674. [PubMed: 8440937]
39. Wang J, Li YL, Kinjo Y, Mac TT, Gibson D, Painter GF, Kronenberg M, Zajonc DM. *Proc. Natl. Acad. Sci. U.S.A.* 2010; 107:1535–1540. [PubMed: 20080535]
40. Chu C, Liu B, Watson D, Szu S, Bryla D, Shiloach J, Schneerson R, Robbins JB. *Infect. Immun.* 1991; 59:4450–4458. [PubMed: 1937803]
41. Crich D, Smith M, Yao QJ, Picione J. *Synthesis-Stuttgart.* 2001:323–326.





Figure 1. Immunoblotting of BBGL-2 derivatives 1–14 with mice sera induced against killed *B. burgdorferi* cells (A), against purified BBGL-II (B), and with LD patient serum (C). No reaction was observed with control serum (picture not shown).




3-*O*- α -D-Galp-1,2-di-*O*-Ole-*sn*-GRO (**2**)


1-*O*- α -D-Galp-2,3-di-*O*-Ole-*sn*-GRO (**6**)

3-*O*- β -D-Galp-1,2-di-*O*-Ole-DL-GRO (**46**)

Figure 2. Dot-blot of BBGL-2 derivatives **2**, **6**, and **46** with mice sera induced by native purified BBGL-2 injected in Freund's adjuvant.

**Scheme 1.**

Reagents and conditions: (a) 4.9 equiv BzCl, CH₂Cl₂, C₆H₅N, DMAP (cat), rt, 24 h, 91%; (b) Cl₂ in CCl₄ (excess), CH₂Cl₂, 0 °C, 30 min, hex-1-ene (excess), 69%; (c) 1.4 equiv 19, 2,4,6-tri-*tert*-butylpyrimidine (22.5 g, 184 mmol), 4 Å molecular sieves, AgOTf (15 g, 58.2 mmol), -40 °C, 15 min, 61%; (d) NaOMe (excess), CH₂Cl₂, MeOH, Dowex 50WX8, CH₂N₂, 91%; (e) H₂, Pd/C, 2,4,6-tri-*tert*-butylpyrimidine, EtOAc, 15 min, 87%; (f) levulinic acid (6 equiv), DCC (7 equiv), 4-dimethylaminopyridine (cat), EtOAc, 94%; (g) AcOH, MeOH, reflux, 5 min, 83%.

**Scheme 2.**

Reagents and conditions: (a) 2 equiv **19**, CH_2Cl_2 , TMSOTf (cat), 0°C , 1 h, 80%); (b) NaOMe (excess), MeOH, 24 h, Dowex 50WX8 (H+); (c) levulinic acid (8 equiv), DCC (8 equiv), EtOAc, 93%); (d) AcOH, MeOH, reflux, 5 min, 85%.




Chart 1.


Table 1

Synthetic BBGL-2 analogs

<i>3-O-α-D-Galp-1,2-di-O-Pal-sn-GRO</i>	1
<i>3-O-α-D-Galp-1,2-di-O-Ole-sn-GRO</i>	2
<i>3-O-α-D-Galp-2-O-Ole-1-O-Pal-sn-GRO</i>	3
<i>3-O-α-D-Galp-1-O-Ole-2-O-Pal-sn-GRO</i>	4
<i>1-O-α-D-Galp-2,3-di-O-Pal-sn-GRO</i>	5
<i>1-O-α-D-Galp-2,3-di-O-Ole-sn-GRO</i>	6
<i>1-O-α-D-Galp-2-O-Ole-3-O-Pal-sn-GRO</i>	7
<i>1-O-α-D-Galp-3-O-Ole-2-O-Pal-sn-GRO</i>	8
<i>3-O-α-D-Galp-1,2-di-O-Ste-sn-GRO</i>	9
<i>3-O-α-D-Galp-2-O-Pal-1-O-Ste-sn-GRO</i>	10
<i>3-O-α-D-Galp-1-O-Pal-sn-GRO</i>	11
<i>3-O-α-D-Galp-1-O-Ole-sn-GRO</i>	12
<i>3-O-β-D-Galp-1,2-di-O-Pal-sn-GRO</i>	13
<i>3-O-β-D-Galp-1,2-di-O-Ole-sn-GRO</i>	14

Table 2


Singly Gro-acylated protected BBGL-2 congeners



Compound	R ¹	R ²	R ³
32	H	OH	Pal
33	H	OH	Ole
34	H	OH	Ste
35	OH	H	Pal
36	OH	H	Ole

Table 3

Doubly Gro-acylated protected BBGL-2 congeners



Compound	R ¹	R ²	R ³
37	H	OPal	Pal
38	H	OOle	Ole
39	H	OOle	Pal
40	H	OPal	Ole
41	H	OSte	Ste
42	OPal	OH	Pal
43	OOle	OH	Ole
44	OOle	OH	Pal
45	OPal	OH	Ole

Table 4Selected ^1H NMR chemical shifts (ppm) of galactopyranosyl diglyceride analogs **1–4** in CDCl_3

	1	2	3	4
Gal H-1	4.954	4.925	4.940	4.937
H-2	3.829	3.868	3.850	3.856
H-3	3.758	3.788	3.774	3.776 ^a
H-4	4.098	4.087	4.095	4.093 ^a
H-5	3.824	3.798	3.808	3.806
H-6	3.962	3.833	3.914	3.905
H-6'	3.851	3.799	3.837	3.834
Gro H-1	4.379	4.3794	4.377	4.373
H-1'	4.117	4.148	4.132	4.135
H-2	5.254	5.250	5.252	5.254
H-3	3.856	3.812	3.834	5.254
H-3'	3.641	3.624	3.632	3.631
Ole H-2,2'	—	2.318, 2.309	2.321	2.313
H-8,8'	—	2.009, 2.008	2.010	2.009
H-9,10	—	5.342, 5.342	5.344	5.343
H-11,11'	—	2.009, 2.008	2.010	2.009
H-18	—	0.880, 0.880	0.880	0.880
Pal H-2,2'	2.324, 2.317	—	2.313	0.880
H-16	0.881, 0.881	—	0.880	0.880

^aOur assignments are at variance with those published in Ref. 19 for compound 4: H-3 of Gal was reported in the range of 4.03–4.12 ppm, while H-4 was in the 3.79–3.94 ppm range, that is, reversed from the data shown above.

Table 5Selected ^{13}C NMR chemical shifts (ppm) of galactopyranosyl diglyceride analogs **1–4** in CDCl_3

	1	2	3	4
Gal C-1	99.25	99.37	99.31	99.34
C-2	69.45	69.13	69.30	69.28
C-3	71.00	70.64	70.81	70.79
C-4	70.31	70.14	70.29	70.29
C-5	69.98	70.14	70.03	70.06
C-6	63.28	62.45	62.93	62.85
Gro C-1	61.95	62.42	62.17	62.24
C-2	69.87	69.88	69.87	69.87
C-3	66.88	66.46	66.67	66.65
Ole C-1	—	173.73, 173.35	173.29	173.63
C-2	—	34.28, 34.11	34.28	34.30
C-9	—	130.03	130.05	130.04
C-10	—	129.69	129.69	129.69
C-18	—	14.13, 14.13	14.13	14.13
Pal C-1	173.50, 173.25	—	173.65	173.34
C-2	34.29, 34.10	—	34.16	34.10
C-16	14.13, 14.13	—	14.13	14.13

Table 6

Selected two and three bond homonuclear H-H and one-bond heteronuclear C-H NMR coupling constants (Hz) of galactopyranosyl diglyceride analogs **1–4** in CDCl₃

	1	2	3	4
Gal $J_{1,2}$	3.8	3.7	3.8	3.8
$J_{2,3}$	9.7	10.3	9.8	9.9
$J_{3,4}$	3.3	3.2	3.2	3.3
$J_{4,5}$	0.9	<0.5	1.1	1.3
$J_{5,6}$	5.1	Nr ^a	5.0	5.0
$J_{5,6'}$	Nr	Nr	4.6	4.7
$J_{6,6'}$	11.6	nr	11.4	11.3
$J_{C-1,H-1}$	170.1	170.1	170.4	170.1
Gro $J_{1,1'}$	11.8	12.0	11.9	12.0
$J_{1,2}$	4.2	3.6	4.0	3.9
$J_{1',2}$	5.7	6.2	5.9	6.0
$J_{2,3}$	4.6	5.4	5.0	5.0
$J_{2,3'}$	6.3	5.8	6.1	6.0
$J_{3,3'}$	11.0	10.8	10.9	10.9

^aNr = not resolved.