

CORRIGENDUM

Mercedes Hernandez, Nieves Fernandez-Garcia, Pedro Diaz-Vivancos and Enrique Olmos

A different role for hydrogen peroxide and the antioxidative system under short and long salt stress in *Brassica oleracea* roots *J. Exp. Bot.* 2010 61: 521-535; doi:10.1093/jxb/erp321

The authors would like to apologize for the errors in figures 1, 4, and 7 in this paper. Control conditions are represented by filled circles and 80 mM of NaCl by inverted triangles. The correct legends are the following:

- Fig. 1. Time-course of hydrogen peroxide and MDA contents in Zone I (A, C) and Zone II (B, D) of *Brassica oleracea* roots grown under control conditions (filled circles), and with 40 mM of NaCl (open circles) and 80 mM of NaCl (inverted triangles). Values represent the means \pm SD of five different samples. Significant differences (P < 0.05) between days and treatments are indicated by different letters according to Tukey's test.
- Fig. 4. Time-course of SOD, APX, CAT, and POX enzyme activities in Zone I (A, C, E, G) and Zone II (B, D, F, H) of Brassica oleracea roots grown under control conditions (filled circles), and with 40 mM of NaCl (open circles) and 80 mM of NaCl (inverted triangles). Values represent the means \pm SD of five different samples. Significant differences (P < 0.05) between days and treatments are indicated by different letters according to Tukey's test.
- Fig. 7. Time-course of MDHAR, DHAR, and GR enzyme activities in Zone I (A, C, E) and Zone II (B, D, F) of *Brassica oleracea* roots grown under control conditions (filled circles), and with 40 mM of NaCl (open circles) and 80 mM of NaCl (inverted triangles). Values represent the means \pm SD of five different samples. Significant differences (P < 0.05) between days and treatments are indicated by different letters according to Tukey's test.