

Published in final edited form as:

*Alcohol*. 2011 November ; 45(7): 653–661. doi:10.1016/j.alcohol.2010.11.003.

## The cerebellar GABA<sub>A</sub>R $\alpha 6$ R100Q polymorphism alters ligand binding in outbred Sprague-Dawley rats in a similar manner as in selectively bred AT and ANT rats

Leena-Stiina Kontturi<sup>a</sup>, Asko J. Aalto<sup>a</sup>, Martin Wallner<sup>b</sup>, and Mikko Uusi-Oukari<sup>a,\*</sup>

<sup>a</sup> Department of Pharmacology, Drug Development and Therapeutics, University of Turku, FIN-20014 Turku, Finland <sup>b</sup> Department of Molecular and Medical Pharmacology, University of California, Los Angeles, California 90095, USA

### Abstract

The alcohol-tolerant AT and -non-tolerant ANT rat lines have been selectively bred for innate sensitivity to ethanol-induced motor impairment. The cerebellar GABA<sub>A</sub> receptor  $\alpha 6$  subunit alleles  $\alpha 6$ -100R and  $\alpha 6$ -100Q are segregated in the AT and ANT rats, respectively. This  $\alpha 6$  polymorphism might explain various differences in pharmacological properties and density of GABA<sub>A</sub> receptors between the rat lines. In the present study we have used non-selected outbred Sprague-Dawley rats homozygous for the  $\alpha 6$ -100RR (RR) and  $\alpha 6$ -100QQ (QQ) genotypes to show that these RR and QQ rats display similar differences between genotypes as AT and ANT rat lines. The genotypes differed in their affinity for [<sup>3</sup>H]Ro 15–513 and classical benzodiazepines to cerebellar “diazepam-insensitive” binding sites, in density of cerebellar [<sup>3</sup>H]muscimol binding and in the antagonizing effect of furosemide on GABA-induced inhibition of [<sup>3</sup>H]EBOB binding. The results suggest the involvement of  $\alpha 6$ -R100Q polymorphism in these line differences and in the differences previously found between AT and ANT rats. In addition, the  $\alpha 6$ -R100Q polymorphism induces striking differences in [<sup>3</sup>H]Ro 15-4513 binding kinetics to recombinant  $\alpha 6\beta 3\gamma 2s$  receptors as well as cerebellar diazepam-insensitive sites. Association of [<sup>3</sup>H]Ro 15-4513 binding was ~10 fold faster and dissociation was ~3–4 fold faster in diazepam-insensitive  $\alpha 6\beta \gamma 2$  receptors containing the  $\alpha 6$ -100Q allele, with a resulting change of ~2.5-fold in equilibrium dissociation constant ( $K_D$ ). The results indicate that in addition to the central role of the homologous  $\alpha 6$ -100R/Q ( $\alpha 1$ -101H) residue in benzodiazepine binding and efficacy, this critical benzodiazepine binding site residue has a major impact on benzodiazepine binding kinetics.

### Keywords

Cerebellar granule cell; Benzodiazepine; GABA<sub>A</sub> receptors; Ethanol sensitivity; Ro 15-4513; Selected rat lines

---

\*Corresponding author (Mikko Uusi-Oukari), Address: Itainen Pitkakatu 4, 20014 University of Turku, FINLAND; Tel: +358-2-3337607; fax: +358-2-3337216. mikko.uusi-oukari@utu.fi (M. Uusi-Oukari).

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Introduction

Selective breeding of rodent lines has been used as a tool to identify genes that mediate behavioral differences related to the selection criteria used. At the Biomedical Research Center, Alko Ltd, two rat lines, the AT and ANT rats, differing in their innate sensitivity to motor-impairing effects of ethanol have been produced (Eriksson and Rusi, 1981). The ethanol-sensitive ANT rats are also more sensitive than the ethanol-insensitive AT rats to GABA<sub>A</sub> receptor- (GABA<sub>A</sub>R) positive modulators lorazepam, diazepam and barbital (Hellevuo et al., 1989; Wong et al., 1996b) suggesting the involvement of GABA<sub>A</sub>ergic mechanisms mediating the sensitivity differences between the rat lines.

Various differences have been found between the AT and ANT lines in characteristics of cerebellar GABA<sub>A</sub>Rs. The density of high affinity GABA<sub>A</sub> agonist (<sup>3</sup>H]muscimol) binding sites is lower in ANT than in AT rats (Malminen and Korpi, 1988; Uusi-Oukari and Korpi, 1989). The affinity of “classical” benzodiazepine (BZ) agonists to  $\alpha 6$  subunit-containing, “diazepam-insensitive” (DZ-IS) binding sites in  $\alpha 6\beta 2$  receptors is about 100-fold higher in ANT than in AT rats (Uusi-Oukari and Korpi, 1990, 1991). This line difference is dependent on a single point mutation in the  $\alpha 6$  codon 100 (CGA→CAA) in ANT rats leading to an amino acid change from arginine (R) to glutamine (Q) (Korpi et al., 1993). The homologous position in  $\alpha 1,2,3,5$  receptors is a histidine residue which has been shown to be a critical residue for classical benzodiazepine binding at the  $\alpha\gamma 2$  subunit interface (Wieland et al., 1992). The  $\alpha 6$ -R100Q mutation increases the affinity of [<sup>3</sup>H]Ro 15-4513 to DZ-IS binding sites (Mäkelä et al., 1995; Uusi-Oukari and Korpi, 1990). The allosteric interaction between GABA and classical BZs in DZ-IS sites was found to be present only in ANT rats (Korpi et al., 1993; Uusi-Oukari and Korpi, 1992). Furthermore, furosemide, a GABA<sub>A</sub>R antagonist selective for  $\alpha 6$  subunit-containing receptors (Korpi et al., 1995) is less efficient in enhancing basal binding of the GABA<sub>A</sub>R-selective channel blocker [<sup>35</sup>S]TBPS and less efficient in antagonizing GABA-induced inhibition of [<sup>35</sup>S]TBPS binding in ANT rats than in AT rats (Mäkelä et al., 1996, 1999). The same  $\alpha 6$ -R100Q mutation was later found to be enriched in the Sardinian non-alcohol-preferring (sNP) rat line (Saba et al., 2001), and also in rats selected for alcohol (non)preference (Carr et al., 2003), and it was shown to be a rather frequent naturally occurring  $\alpha 6$  polymorphism in some Sprague-Dawley laboratory rat colonies (Hanchar et al., 2005).

The higher affinity and functional sensitivity of cerebellar  $\alpha 6\beta 2$  receptors in ANT than in AT rats to classic BZs suggest that the  $\alpha 6$ -100Q mutation determines the increased BZ-induced motor impairment in ANT rats (Korpi et al., 1993). However, the  $\alpha 6$ -R100Q mutation expressed in  $\alpha 6\beta 2/\gamma 2$  combination did not affect GABA or ethanol sensitivity (Hanchar et al., 2005; Korpi et al., 1993). In contrast, it was shown that  $\alpha 6$ -100Q mutation in the ethanol-sensitive receptor subtype  $\alpha 6\beta 3\delta$  (Wallner et al., 2003), while not affecting GABA sensitivity, dramatically increases ethanol sensitivity in the ethanol concentration range of 3–30 mM in recombinantly expressed receptors and in cerebellar granule cells in slices (Hanchar et al., 2005) suggesting the involvement of  $\alpha 6\beta 3\delta$  receptors in alcohol sensitivity differences between AT and ANT rats. The increased ethanol sensitivity seen with the R100Q mutation is consistent with earlier work using receptors reconstituted (by injection of cerebellar vesicles from  $\alpha 6$ -100RR and  $\alpha 6$ -100QQ rats) in oocytes (Sanna et al., 2003). Higher functional sensitivity to ethanol and BZs was also demonstrated in ANT when compared to AT cerebellar synaptoneurosomes (Schmid et al., 1999). However, using patch-clamp electrophysiological techniques in cerebellar slices of outbred Sprague-Dawley rats homozygous for the  $\alpha 6$ -100R and -100Q alleles, Botta et al., 2007 were not able to detect genotype dependent ethanol sensitivity differences. In addition, crossbreeding studies using AT and ANT animals did not support the conclusion that the  $\alpha 6$ R100Q polymorphism is important for differential ethanol sensitivity (Radcliffe et al., 2004). The controversy

surrounding the ethanol sensitivity of  $\alpha 4/6\beta\delta$ -GABA<sub>A</sub>Rs and whether this is a solution for the ethanol/GABA<sub>A</sub>R dilemma is discussed in detailed in a special issue of the Journal Alcohol, where the issue is introduced by an editorial by Lovinger and Homanics (2007).

In the present study we have compared cerebellar and hippocampal GABA<sub>A</sub>Rs of Sprague-Dawley rats homozygous for the  $\alpha 6$ -100R and -100Q alleles. We focused on receptor differences found between AT and ANT rats to reveal if these differences are related to  $\alpha 6$ -R100Q polymorphism. In addition, we characterized the effects of the polymorphism on cerebellar [<sup>3</sup>H]Ro 15-4513 binding kinetics to cerebellar DZ-IS binding sites and recombinant  $\alpha 6\beta 3\gamma 2$  receptors.

## Materials and methods

### Animals

Sprague-Dawley rats obtained from Charles River (Hollister, CA) were genotyped as described in Hanchar et al. (2005) to identify animals homozygous for the  $\alpha 6$ -100R and  $\alpha 6$ -100Q polymorphisms. A total of 14 RR and 17 QQ adult rats were used for the studies. The rats used in experiments of [<sup>3</sup>H]Ro 15-4513 binding kinetics were 6–12 month old males, while rats used in other experiments were 3–4 month old males. The animals were killed by decapitation, their brains were removed, and the cerebelli and hippocampi frozen on dry ice and stored at -70 °C. All procedures were in accordance with protocols approved by the University of California at Los Angeles (UCLA) Chancellor's Animal Research Committee and by the Institutional Animal Care and Use Committee of the University of Turku.

### Reagents

The radioligands [propyl-2,3-<sup>3</sup>H]EBOB (specific activity 48 Ci/mmol), [methylene-<sup>3</sup>H]muscimol (18 Ci/mmol), and [7,9-<sup>3</sup>H]Ro 15-4513 (28 Ci/mmol) were purchased from Perkin Elmer Life and Analytical Sciences (Boston, MA, USA). Flumazenil (Ro 15-1788) was a gift from F. Hoffmann-La Roche Ltd (Basel, Switzerland). Diazepam, GABA and picrotoxin were from Sigma Chemical Co. (St. Louis, MO, USA).

### Recombinant GABA<sub>A</sub> receptor expression in HEK 293 cells

Human embryonal kidney (HEK) 293 cells were transfected with rat cDNAs under the control of CMV promoter ( $\alpha 1$ ,  $\alpha 6$ -100R or  $\alpha 6$ -100Q: $\beta 3$ : $\gamma 2S$ , 1:1:2) as described in Meera et al. (1997) and the cells were harvested 48 h after transfection. The cells were washed with PBS, homogenized in PBS using an Ultra-Turrax and stored frozen at -70 °C. Before binding assays the suspensions were thawed, washed once with ice-cold assay buffer by resuspension and centrifugation, and homogenized in ice-cold assay buffer with an Ultra-Turrax.

### [<sup>3</sup>H]Ro 15-4513 binding assay

Cerebellar and hippocampal membranes were prepared and [<sup>3</sup>H]Ro 15-4513 binding assays performed essentially as described in Uusi-Oukari and Korpi (1990). Tris-HCl (50 mM, pH 7.4) containing 120 mM NaCl was used as the incubation buffer. Non-specific binding was determined in the presence of 10  $\mu$ M flumazenil. Triplicate samples were incubated in an ice-water bath in the dark with shaking for 1 h in a total volume of 300  $\mu$ l. Diazepam-insensitive (DZ-IS) [<sup>3</sup>H]Ro 15-4513 binding was determined in the presence of 100  $\mu$ M diazepam to differentiate between RR and QQ binding sites. Receptors containing the  $\alpha 6$ -100Q allele display 1  $\mu$ M diazepam affinity (Mäkelä et al., 1995; Uusi-Oukari and Korpi, 1992). Diazepam at 100  $\mu$ M displaces essentially all binding in QQ rats while hardly affecting  $\alpha 6$ -100R receptors (Mäkelä et al., 1995; Uusi-Oukari and Korpi, 1992). The

incubation was terminated by filtration of the samples with a Brandel Cell Harvester (model 48R, Gaithersburg, Maryland, USA) onto Whatman GF/B filters (Whatman International Ltd., Maidstone, UK). The samples were rinsed twice with 4–5 ml of ice-cold incubation buffer. Air-dried filters were immersed in 4 ml of Optiphase HiSafe 2 scintillation fluid (Wallac, Turku, Finland) and radioactivity determined in a Wallac model 1410 liquid scintillation counter (Wallac, Turku, Finland). Non-specific binding was subtracted from total binding and from binding in the presence of diazepam to get total specific binding and DZ-IS binding, respectively. DZ-IS binding was subtracted from total specific binding to get diazepam-sensitive (DZ-S) binding.

### Measurement of [<sup>3</sup>H]Ro 15-4513 binding kinetics

To measure association of [<sup>3</sup>H]Ro 15-4513 binding, cerebellar membranes of RR and QQ rats were incubated in incubation buffer with 5 nM [<sup>3</sup>H]Ro 15-4513 in the absence and presence of 1 μM diazepam (DZ-IS binding) or 10 μM flumazenil (non-specific binding) in conditions described above. Diazepam at 1 μM displaces essentially all DZ-S binding, but only 50% of DZ-IS binding in QQ membranes thus allowing measurement of DZ-IS binding selectively also in QQ cerebellum. The incubations were terminated at various time points. To measure dissociation of [<sup>3</sup>H]Ro 15-4513 binding triplicate samples of membranes were first preincubated in a total volume of 300 μl for 1 h with 5 nM [<sup>3</sup>H]Ro 15-4513 in the absence and presence of 1 μM diazepam or 10 μM flumazenil. The dissociation was started by adding 100 μl of 40 μM flumazenil to the incubation mixtures to reach a final 10 μM flumazenil concentration in all tubes. The tubes were mixed and incubations terminated at various time points as described above.

### [<sup>3</sup>H]Muscimol binding assay

Cerebellar membranes were prepared and saturation analysis of [<sup>3</sup>H]muscimol binding performed essentially as described in Uusi-Oukari and Korpi (1989). The radioligand was used at a concentration range of 0.5–20 nM. Non-specific binding was determined in the presence of 1 mM GABA. Triplicate reaction mixtures in 50 mM Tris-citrate buffer, pH 7.4, were incubated in ice-water bath in the dark with shaking for 30 min. The final incubation volume was 300 μl. The incubation was terminated as described for [<sup>3</sup>H]Ro 15-4513 binding.

### [<sup>3</sup>H]EBOB binding assay

Cerebellar membranes for binding of [<sup>3</sup>H]EBOB, a ligand binding to the same site as the more commonly used [<sup>35</sup>S]TBPS, were prepared and [<sup>3</sup>H]EBOB binding assay performed essentially as described in Uusi-Oukari and Maksay (2006). Triplicate samples of cerebellar membranes were incubated at room temperature with shaking for 2 h in 50 mM Tris-HCl, pH 7.4, containing 120 mM NaCl with 1 nM [<sup>3</sup>H]EBOB in a total volume of 400 μl in the absence and presence of 5 μM GABA with or without 300 μM furosemide. Non-specific binding was determined in the presence of 100 μM picrotoxin. The incubation was terminated as described for [<sup>3</sup>H]Ro 15-4513 binding.

### Protein measurement

In all ligand binding studies, protein concentrations of membranes were determined with the Bio-Rad Coomassie blue dye-based protein assay kit (Hercules, CA, USA) according to manufacturer's instructions.

### Data analysis

Saturation isotherms for estimation of  $K_D$  and  $B_{max}$  values, and association and dissociation curves for estimation of association and dissociation rate constants were analysed with Prism 5 software (Graph Pad, San Diego, CA, USA). Two-tailed unpaired *t*-test or one-way

ANOVA followed by Tukey's post hoc test was used to assess statistical significances of the differences between the groups. P-values of less than 0.05 were considered significant.

## Results

### [<sup>3</sup>H]Ro 15-4513 binding to brain membranes and to recombinant receptors

Cerebellar DZ-IS [<sup>3</sup>H]Ro 15-4513 binding was determined from all animals used in the study. The portion of 5 nM [<sup>3</sup>H]Ro 15-4513 binding in the presence of 100 μM diazepam was  $21.9 \pm 0.9\%$  and  $1.0 \pm 0.1\%$  of total binding in membranes of RR and QQ rats, respectively. The portion of DZ-IS binding was 20–25% in all RR samples, while QQ samples under these conditions displayed only small amounts of DZ-IS [<sup>3</sup>H]Ro 15-4513 binding. The values are consistent with previous studies on AT and ANT rat lines and confirm the GABA<sub>A</sub>R α6-100RR and α6-100QQ rat genotypes.

The association and dissociation rates of total, DZ-S and DZ-IS [<sup>3</sup>H]Ro 15-4513 binding were determined using cerebellar membranes from RR and QQ rats (Fig. 1 and 2, Table 1). Association and dissociation rates of total and DZ-S binding were very similar in RR and QQ membranes (Fig. 1, Table 1). In contrast, the  $K_{on}$  value of DZ-IS binding was 7.4-fold lower in RR than in QQ membranes indicating slower association of [<sup>3</sup>H]Ro 15-4513 to DZ-IS receptors in RR membranes. The same applies to dissociation of DZ-IS [<sup>3</sup>H]Ro 15-4513 binding. The  $K_{off}$  value was 2.9-fold lower in RR than in QQ membranes indicating slower [<sup>3</sup>H]Ro 15-4513 dissociation in RR membranes (Fig. 2, Table 1). The calculation of  $K_D$  values ( $K_{off}/K_{on}$ ) for [<sup>3</sup>H]Ro 15-4513 binding were 1.6 nM and 1.4 nM (DZ-S) and 2.3 nM and 0.9 nM (DZ-IS) for RR and QQ rats, respectively (Table 1).

Association rate of [<sup>3</sup>H]Ro 15-4513 binding to DZ-S α1β3γ2 recombinant receptors was comparable to that of [<sup>3</sup>H]Ro 15-4513 binding to DZ-S receptors in RR and QQ membranes (Table 1, Fig. 3A). Similarly, dissociation rate of the binding to α1β3γ2 receptors was indistinguishable from dissociation of DZ-S binding in cerebellar membranes (Table 1, Fig. 3B). Association and dissociation rates of [<sup>3</sup>H]Ro 15-4513 binding to α6-100R-β3γ2 recombinant receptors and DZ-IS binding to cerebellar membranes of RR rats were almost identical, while the association rate constant was lower in DZ-IS binding to QQ rat cerebellar membranes than to α6-100Q-β3γ2 recombinant receptors (Table 1, Fig. 3A, 3B). This difference between QQ recombinant receptors and membranes is likely due to 1 μM diazepam present with cerebellar membranes. The  $K_i$  of diazepam to DZ-IS sites in QQ animals is 1 μM (Uusi-Oukari and Korpi, 1992) indicating that diazepam competes with [<sup>3</sup>H]Ro 15-4513 in binding to these sites. The association rate constant of [<sup>3</sup>H]Ro 15-4513 binding was 11 fold lower in α6-100R-β3γ2 than in α6-100Q-β3γ2 receptors (Table 1, Fig. 3A) ( $p < 0.001$ , two-tailed unpaired *t*-test). The dissociation rate constant in α6-100Q-β3γ2 receptors was 4.6-fold larger than that in α6-100R-β3γ2 receptors (Table 1, Fig. 3B) ( $p < 0.05$ , two-tailed unpaired *t*-test). The calculated  $K_D$  values ( $K_{off}/K_{on}$ ) of [<sup>3</sup>H]Ro 15-4513 binding to recombinant receptors were 2.9 nM (α1β3γ2), 2.3 nM (α6-100R-β3γ2) and 1.0 nM (α6-100Q-β3γ2) (Table 1). The association and dissociation kinetics of [<sup>3</sup>H] Ro15-4513 binding were in a similar manner slower also in α6-100R-β2γ2 recombinant receptors as compared to α6-100Q-β2γ2 receptors (L.-S. Kontturi and M. Uusi-Oukari, unpublished results) indicating that the difference in binding kinetics is independent on β variant present in the receptor.

Since association of [<sup>3</sup>H]Ro 15-4513 binding to RR cerebellar membranes in the presence of diazepam (DZ-IS binding) and to α6-100R-β3γ2 recombinant receptors was very slow, we tested longer incubation periods. A 2 h incubation with 5 nM [<sup>3</sup>H] Ro15-4513 yielded 23% higher binding both to RR cerebellar membranes in the presence of diazepam and to α6-100R-β3γ2 recombinant receptors (Fig. 4).



Saturation analysis of [<sup>3</sup>H]Ro 15-4513 binding to hippocampal membranes revealed 23% higher B<sub>max</sub> value in RR than in QQ rats (Fig. 5). This difference, however, did not reach statistical significance (p=0.063).

### [<sup>3</sup>H]Muscimol binding to cerebellar membranes

The B<sub>max</sub> value of [<sup>3</sup>H]muscimol binding to cerebellar membranes was lower in QQ (1.46 ± 0.10 pmol/mg protein, mean ± SEM) than in RR (1.84 ± 0.14 pmol/mg protein) animals (p<0.05, two-tailed unpaired *t*-test; RR, n=7; QQ, n=8) (Fig. 6). The K<sub>D</sub> values of cerebellar [<sup>3</sup>H]muscimol binding did not differ between the rat lines (2.21 ± 0.13 and 2.52 ± 0.22 nM for RR and QQ rats, respectively).

### [<sup>3</sup>H]EBOB binding to cerebellar membranes

The GABA-antagonizing effect of furosemide on [<sup>3</sup>H]EBOB binding was studied in cerebellar membranes of RR and QQ rats. GABA (5 μM) inhibited basal binding of 1 nM [<sup>3</sup>H]EBOB by 65–70% (Fig. 7). The effect of GABA on the binding did not significantly differ between RR and QQ rats. However, furosemide (300 μM) significantly antagonized the effect of GABA on [<sup>3</sup>H]EBOB binding only in RR rats (p<0.05, one-way ANOVA followed by Tukey's post hoc test).

## Discussion

Using two groups of Sprague-Dawley rats homozygous for GABA<sub>A</sub>R alleles α6-100R and α6-100Q we found several differences between the groups in their cerebellar GABA<sub>A</sub>Rs previously found between the AT and ANT rat lines: the affinity of cerebellar DZ-IS [<sup>3</sup>H]Ro 15-4513 binding sites is higher to (1) [<sup>3</sup>H]Ro 15-4513 and to (2) classical BZs in QQ than in RR animals; (3) the number of cerebellar high-affinity [<sup>3</sup>H]muscimol binding sites is lower in QQ than in RR animals; and (4) the GABA-antagonizing effect of furosemide on GABA-induced inhibition of [<sup>3</sup>H]EBOB binding is lower in QQ than in RR animals. The affinity differences (1) and (2) have been previously published between alcohol-nonpreferring sNP rats homozygous for QQ and RR alleles (Sanna et al., 2003), between Sprague-Dawley rats with α6-100RR and α6-100QQ genotypes (Hanchar et al., 2005) and using α6-100R/Q-β2/3γ2 recombinant receptors (Hanchar et al., 2005; Korpi et al., 1993). The results clearly show that the differences (1, 2) are caused by the R to Q substitution in α6 residue 100.

As a surprise, this amino acid change affects association and dissociation kinetics of [<sup>3</sup>H]Ro 15-4513 binding. Residue 100 in α6 (and the homologous His residue in α1) reside in loop A of the BZ binding pocket (Tan et al., 2007). Association and dissociation rates of [<sup>3</sup>H]Ro 15-4513 binding appear to be very similar in α1-101H-β3γ2 and α6-100Q-β3γ2 receptors. It has been previously shown, that the α1-H101Q mutation in α1β2γ2 recombinant receptors increases [<sup>3</sup>H]Ro 15-4513 binding affinity (Davies et al., 1998). In fact, α1-H101 substitutions F, K, R, E, Q and C increase [<sup>3</sup>H]Ro 15-4513 affinity while drastically decreasing flunitrazepam affinity (Davies et al., 1998; Wieland et al., 1992). These substitutions also affect efficacy of BZs, including Ro 15-4513, in α1β2γ2 receptors indicating the central role of the residue in BZ binding and function (Dunn et al., 1999). We performed [<sup>3</sup>H]Ro 15-4513 association and dissociation measurements at 0 °C degrees. The association rates in vivo at 37 °C would be much faster and although obviously slower in α6-100R receptors, equilibrium would be reached within minutes. Our data suggest that in vitro binding studies using [<sup>3</sup>H]Ro 15-4513 as a ligand, 1 h incubation times at concentrations ≤ 5 nM at 0 °C have obviously led to an underestimation of the number of cerebellar [<sup>3</sup>H]Ro 15-4513 DZ-IS binding sites.

The other two AT/ANT line differences, (3) and (4), were present in RR/QQ rats, but the molecular basis of these differences is less clear. Cerebellar high affinity [<sup>3</sup>H]muscimol binding is mainly localized in cerebellar granule layer and associated with  $\alpha 6\beta \gamma 2$  and  $\alpha 6\beta \delta$  receptor subtypes (Jones et al., 1997; Korpi et al., 2002; Mihalek et al., 1999). The higher  $B_{\max}$  of high affinity [<sup>3</sup>H]muscimol binding therefore suggests a higher amount of  $\alpha 6\beta \gamma 2$  and/or  $\alpha 6\beta \delta$  receptors in AT/RR than in ANT/QQ cerebellar granule cells. The basal level  $\alpha 6$  mRNA expression has been shown to be higher in sNP rats with RR than with QQ genotype (Saba et al., 2005). This difference was suggested to be due to differences in  $\alpha 6$  promoter sequences between RR and QQ genotypes (Saba et al., 2005). However, no difference has been found between AT and ANT rats in the number of cerebellar [<sup>3</sup>H]Ro 15-4513 binding sites (Uusi-Oukari and Korpi, 1990) or in cerebral granule cell tonic GABAergic currents (Valenzuela et al., 2005), and only a tendency (~10%) to higher level of  $\alpha 6$  mRNA expression in AT than in ANT rats (Uusi-Oukari et al., 2000). However, the slow association of [<sup>3</sup>H]Ro 15-4513 to DZ-IS binding sites in RR rats suggests that the  $B_{\max}$  value of AT/RR rats might have been underestimated because the binding does not reach equilibrium within 1 h of incubation. Due to high BZ sensitivity it is not possible to determine  $B_{\max}$  of DZ-IS [<sup>3</sup>H]Ro 15-4513 binding in QQ animals. It might be also difficult to see a 30% difference corresponding to difference in [<sup>3</sup>H]muscimol binding in DZ-IS [<sup>3</sup>H]Ro 15-4513 binding by determining total [<sup>3</sup>H]Ro 15-4513 binding, because a 30% difference in 20–25% DZ-IS portion of total binding would make a 7% difference in  $B_{\max}$  of total binding. In addition to a difference in  $\alpha 6$  mRNA transcription there may be differences in the efficiency between  $\alpha 6$ -100R and  $\alpha 6$ -100Q proteins to combine with other subunits to form  $\alpha 6\beta \gamma 2$  and/or  $\alpha 6\beta \delta$  receptors.

The blunted furosemide action in antagonising GABA-induced inhibition of [<sup>3</sup>H]EBOB binding was also seen in QQ rats in the present study. The result suggests the involvement of residue 100Q on the mechanism(s) of blunted furosemide response in ANT/QQ rats. However, the inability to reproduce this line difference using  $\alpha 6$ -100Q- $\beta 3\gamma 2$  recombinant receptors (Mäkelä et al., 1996) suggests that in addition to  $\alpha 6$ -R100Q amino acid change there are other unidentified alteration(s) in ANT/QQ receptors necessary for the insensitivity to furosemide. There may also be GABA<sub>A</sub>R-associated component(s) present in cerebellar granule cells needed for the ANT/QQ furosemide insensitivity not present in HEK cells used for recombinant receptor expression. Granule cell-specific posttranslational modification is also possible. Anyway, the data shown here suggest that the furosemide-insensitive property seems to be linked to  $\alpha 6$ -100Q allele.

There was a tendency for lower [<sup>3</sup>H]Ro 15-4513 binding in hippocampal membranes of QQ rats as compared to RR rats. The result is in accordance with the hippocampal line difference between AT and ANT rats (Uusi-Oukari and Korpi, 1990). GABA<sub>A</sub>R subunits are clustered in genome as clusters of 3–4 subunits (Darlison et al., 2005). The  $\alpha 6$  gene is clustered with genes for  $\alpha 1$ ,  $\beta 2$ , and  $\gamma 2$  subunits, the latter three subunits forming the major GABA<sub>A</sub>R subtype  $\alpha 1\beta 2\gamma 2$  in the brain (Garrett et al., 1997; McKernan and Whiting, 1996). The  $\alpha 6$ -100Q genotype is strictly associated with various silent polymorphisms/single nucleotide polymorphisms (SNPs) in the other three GABA<sub>A</sub>R subunit genes of the cluster (Congeddu et al., 2003). Therefore, it remains possible that SNPs in  $\alpha 1$ ,  $\beta 2$  and  $\gamma 2$  subunit genes (that co-segregate with the  $\alpha 6$ -100Q genotype) might contribute to lower number of hippocampal [<sup>3</sup>H]Ro 15-4513 binding sites in ANT/QQ than in AT/RR rats.

While the results suggest that most of the differences in cerebellar GABA<sub>A</sub>Rs are dependent on  $\alpha 6$ -R100Q polymorphism, it is possible that other polymorphisms found in the genes of the  $\beta 2$ - $\alpha 6$ - $\alpha 1$ - $\gamma 2$  cluster and linked to  $\alpha 6$ -R100Q polymorphism might play a role in some of the differences between RR and QQ rats. The R100Q mutation is enriched in two alcohol-non-preferring rats lines (sNP and Alko Non-Alcohol, ANA) suggesting that in addition to

ethanol and BZ sensitivity to motor impairment the polymorphism and/or the associated SNPs of the cluster might be associated in alcohol preference and avoidance (Carr et al., 2003; Saba et al., 2001; Wong et al., 1996a). Genes in  $\beta 2$ - $\alpha 6$ - $\alpha 1$ - $\gamma 2$  cluster have also been associated to ethanol-related behaviors in humans. Several genetic association studies with human alcohol-dependent subjects suggest a role for genes in  $\beta 2$ - $\alpha 6$ - $\alpha 1$ - $\gamma 2$  cluster in the development of alcohol dependence (Chang et al., 2002; Loh et al., 1999, 2002; Radel et al., 2005; Sander et al., 1999).

## Acknowledgments

This study was supported by grants from the Finnish Society of Sciences and Letters (MU-O), Oskar Öflund Foundation (MU-O) and the Turku University Foundation (MU-O) and by NIH grant AA017891 (MW).


## References

- Botta P, Mameli M, Floyd KL, Radcliffe RA, Valenzuela CF. Ethanol sensitivity of GABAergic currents in cerebellar granule neurons is not increased by a single amino acid change (R100Q) in the  $\alpha 6$  GABA<sub>A</sub> receptor subunit. *J Pharmacol Exp Ther.* 2007; 323:684–691. [PubMed: 17704355]
- Carr LG, Spence JP, Eriksson CJ, Lumeng L, Li TK. AA and ANA rats exhibit the R100Q mutation in the GABA<sub>A</sub> receptor  $\alpha 6$  subunit. *Alcohol.* 2003; 31:93–97. [PubMed: 14615016]
- Chang YT, Sun HS, Fann CS, Chang CJ, Liao ZH, Huang JL, Loh EW, Yu WY, Cheng AT. Association of the  $\gamma$ -aminobutyric acid A receptor gene cluster with alcohol dependence in Taiwanese Han. *Mol Psychiatry.* 2002; 7:828–829. [PubMed: 12232773]
- Congeddu E, Saba L, Porcella A, Sanna A, Marchese G, Lobina C, Gessa GL, Pani L. Molecular characterization of new polymorphisms at the  $\beta 2$ ,  $\alpha 1$ ,  $\gamma 2$  GABA<sub>A</sub> receptor subunit genes associated to a rat nonpreferring ethanol phenotype. *Brain Res, Mol Brain Res.* 2003; 110:289–297. [PubMed: 12591165]
- Darlison MG, Pahal I, Thode C. Consequences of the evolution of the GABA<sub>A</sub> receptor gene family. *Cell Mol Neurobiol.* 2005; 25:607–624. [PubMed: 16075381]
- Davies M, Bateson AN, Dunn SM. Structural requirements for ligand interactions at the benzodiazepine recognition site of the GABA<sub>A</sub> receptor. *J Neurochem.* 1998; 70:2188–2194. [PubMed: 9572307]
- Dunn SM, Davies M, Muntoni AL, Lambert JJ. Mutagenesis of the rat  $\alpha 1$  subunit of the  $\gamma$ -aminobutyric acid<sub>A</sub> receptor reveals the importance of residue 101 in determining the allosteric effects of benzodiazepine site ligands. *Mol Pharmacol.* 1999; 56:768–774. [PubMed: 10496960]
- Eriksson, K.; Rusi, M. Finnish selection studies on alcohol-related behaviours: General outline. In: McClearn, GE.; Deitrich, RA.; Erwin, G., editors. *Development of Animal Models as Pharmacogenetic tools*, NIAAA Research Monograph No 6. U.S. Government Printing Office; Washington, DC: 1981. p. 87-117.
- Garrett KM, Haque D, Berry D, Niekrasz I, Gan J, Rotter A, Seale TW. The GABA<sub>A</sub> receptor  $\alpha 6$  subunit gene (*Gabra6*) is tightly linked to the  $\alpha 1$ - $\gamma 2$  subunit cluster on mouse chromosome 11. *Brain Res, Mol Brain Res.* 1987; 45:133–137. [PubMed: 9105679]
- Hanchar HJ, Dodson PD, Olsen RW, Otis TS, Wallner M. Alcohol-induced motor impairment caused by increased extrasynaptic GABA<sub>A</sub> receptor activity. *Nat Neurosci.* 2005; 8:339–345. [PubMed: 15696164]
- Hellevo K, Kiiianmaa K, Korpi ER. Effect of GABAergic drugs on motor impairment from ethanol, barbital and lorazepam in rat lines selected for differential sensitivity to ethanol. *Pharmacol Biochem Behav.* 1989; 34:399–404. [PubMed: 2516323]
- Jones A, Korpi ER, McKernan RM, Pelz R, Nusser Z, Mäkelä R, Mellor JR, Pollard S, Bahn S, Stephenson FA, et al. Ligand-gated ion channel subunit partnerships: GABA<sub>A</sub> receptor  $\alpha 6$  subunit gene inactivation inhibits delta subunit expression. *J Neurosci.* 1997; 17:1350–1362. [PubMed: 9006978]




- Korpi ER, Kleingoor C, Kettenmann H, Seeburg PH. Benzodiazepine-induced motor impairment linked to point mutation in cerebellar GABA<sub>A</sub> receptor. *Nature*. 1993; 361:356–359. [PubMed: 7678923]
- Korpi ER, Kuner T, Seeburg PH, Lüddens H. Selective antagonist for the cerebellar granule cell-specific  $\gamma$ -aminobutyric acid type A receptor. *Mol Pharmacol*. 1995; 47:283–289. [PubMed: 7870036]
- Korpi ER, Mihalek RM, Sinkkonen ST, Hauer B, Hevers W, Homanics GE, Sieghart W, Lüddens H. Altered receptor subtypes in the forebrain of GABA<sub>A</sub> receptor delta subunit-deficient mice: recruitment of  $\gamma$ 2 subunits. *Neuroscience*. 2002; 109:733–743. [PubMed: 11927155]
- Loh EW, Ball D. Role of the GABA<sub>A</sub>  $\beta$ 2, GABA<sub>A</sub>  $\alpha$ 6, GABA<sub>A</sub>  $\alpha$ 1 and GABA<sub>A</sub>  $\gamma$ 2 receptor subunit genes cluster in drug responses and the development of alcohol dependence. *Neurochem Int*. 2000; 37:413–423. [PubMed: 10871693]
- Loh EW, Smith I, Murray R, McLaughlin M, McNulty S, Ball D. Association between variants at the GABA<sub>A</sub>  $\beta$ 2, GABA<sub>A</sub>  $\alpha$ 6 and GABA<sub>A</sub>  $\gamma$ 2 gene cluster and alcohol dependence in a Scottish population. *Mol Psychiatry*. 1999; 4:539–544. [PubMed: 10578235]
- Lovinger DM, Homanics GE. Tonic for what ails us? High-affinity GABA<sub>A</sub> receptors and alcohol. *Alcohol*. 2007; 41:139–143. [PubMed: 17521844]
- Malminen O, Korpi ER. GABA/benzodiazepine receptor/chloride ionophore complex in brains of rat lines selectively bred for differences in ethanol-induced motor impairment. *Alcohol*. 1988; 5:239–249. [PubMed: 2458111]
- Mäkelä R, Lehtonen M, Wisden W, Lüddens H, Korpi ER. Blunted furosemide action on cerebellar GABA<sub>A</sub> receptors in ANT rats selectively bred for high alcohol sensitivity. *Neuropharmacology*. 1996; 35:1493–1502. [PubMed: 9014165]
- Mäkelä R, Uusi-Oukari M, Oja SS, Alho H, Anghelescu I, Klawe C, Lüddens H, Korpi ER. Furosemide action on cerebellar GABA<sub>A</sub> receptors in alcohol-sensitive ANT rats. *Alcohol*. 1999; 19:197–205. [PubMed: 10580508]
- Mäkelä R, Wong G, Lüddens H, Korpi ER. Phenotypic and genotypic analysis of rats with cerebellar GABA<sub>A</sub> receptors composed from mutant and wild-type  $\alpha$ 6 subunits. *J Neurochem*. 1995; 65:2401–2408. [PubMed: 7595533]
- McKernan RM, Whiting PJ. Which GABA<sub>A</sub>-receptor subtypes really occur in the brain? *Trends Neurosci*. 1996; 19:139–143. [PubMed: 8658597]
- Meera P, Wallner M, Song M, Toro L. Large conductance voltage- and calcium-dependent K<sup>+</sup> channel, a distinct member of voltage-dependent ion channels with seven N-terminal transmembrane segments (S0–S6), an extracellular N terminus, and an intracellular (S9–S10) C terminus. *Proc Natl Acad Sci USA*. 1997; 94:14066–14071. [PubMed: 9391153]
- Mihalek RM, Banerjee PK, Korpi ER, Quinlan JJ, Firestone LL, Mi ZP, Lagenaur C, Tretter V, Sieghart W, Anagnostaras SG, et al. Attenuated sensitivity to neuroactive steroids in  $\gamma$ -aminobutyrate type A receptor  $\delta$  subunit knockout mice. *Proc Natl Acad Sci USA*. 1999; 96:12905–12910. [PubMed: 10536021]
- Radel M, Vallejo RL, Iwata N, Aragon R, Long JC, Virkkunen M, Goldman D. Haplotype-based localization of an alcohol dependence gene to the 5q34  $\gamma$ -aminobutyric acid type A gene cluster. *Arch Gen Psychiatry*. 2005; 62:47–55. [PubMed: 15630072]
- Saba L, Porcella A, Congeddu E, Colombo G, Peis M, Pistis M, Gessa GL, Pani L. The R100Q mutation of the GABA<sub>A</sub>  $\alpha$ 6 receptor subunit may contribute to voluntary aversion to ethanol in the SNP rat line. *Brain Res, Mol Brain Res*. 2001; 87:263–270. [PubMed: 11245930]
- Saba L, Porcella A, Sanna A, Congeddu E, Marziliano N, Mongeau R, Grayson D, Pani L. Five mutations in the GABA<sub>A</sub>  $\alpha$ 6 gene 5' flanking region are associated with a reduced basal and ethanol-induced  $\alpha$ 6 upregulation in mutated Sardinian alcohol non-preferring rats. *Brain Res, Mol Brain Res*. 2005; 137:252–257. [PubMed: 15950783]
- Sander T, Ball D, Murray R, Patel J, Samochowiec J, Winterer G, Rommelspacher H, Schmidt LG, Loh EW. Association analysis of sequence variants of GABA<sub>A</sub>  $\alpha$ 6,  $\beta$ 2, and  $\gamma$ 2 gene cluster and alcohol dependence. *Alcohol Clin Exp Res*. 1999; 23:427–431. [PubMed: 10195814]


- Sanna A, Congeddu E, Porcella A, Saba L, Pistis M, Peis M, Marchese G, Ruiu S, Lobina C, Grayson DR, et al. Characterization of wild-type (R100R) and mutated (Q100Q) GABA<sub>A</sub>  $\alpha$ 6 subunit in Sardinian alcohol non-preferring rats (sNP). *Brain Res.* 2003; 967:98–105. [PubMed: 12650970]
- Schmid G, Bonanno G, Raiteri L, Sarviharju M, Korpi ER, Raiteri M. Enhanced benzodiazepine and ethanol actions on cerebellar GABA<sub>A</sub> receptors mediating glutamate release in an alcohol-sensitive rat line. *Neuropharmacology.* 1999; 38:1273–1279. [PubMed: 10471080]
- Tan KR, Baur R, Gonthier A, Goeldner M, Sigel E. Two neighboring residues of loop A of the  $\alpha$ 1 subunit point towards the benzodiazepine binding site of GABA<sub>A</sub> receptors. *FEBS Lett.* 2007; 581:4718–4722. [PubMed: 17854801]
- Uusi-Oukari M, Heikkilä J, Sinkkonen ST, Mäkelä R, Hauer B, Homanics GE, Sieghart W, Wisden W, Korpi ER. Long-range interactions in neuronal gene expression: evidence from gene targeting in the GABA<sub>A</sub> receptor  $\beta$ 2- $\alpha$ 6- $\alpha$ 1- $\gamma$ 2 subunit gene cluster. *Mol Cell Neurosci.* 2000; 16:34–41. [PubMed: 10882481]
- Uusi-Oukari M, Klein R, Mäkelä R, Lüddens H, Korpi ER. Quantification of GABA<sub>A</sub> receptor subunit mRNAs by non-radioisotopic competitive RT-PCR utilizing plate-based EIA methodology. *J Neurosci Methods.* 2000; 95:65–73. [PubMed: 10776816]
- Uusi-Oukari M, Korpi ER. Cerebellar GABA<sub>A</sub> receptor binding and function in vitro in two rat lines developed for high and low alcohol sensitivity. *Neurochem Res.* 1989; 14:733–739. [PubMed: 2554173]
- Uusi-Oukari M, Korpi ER. Diazepam sensitivity of the binding of an imidazobenzodiazepine, [<sup>3</sup>H]Ro 15-4513, in cerebellar membranes from two rat lines developed for high and low alcohol sensitivity. *J Neurochem.* 1990; 54:1980–1987. [PubMed: 2159982]
- Uusi-Oukari M, Korpi ER. Specific alterations in the cerebellar GABA<sub>A</sub> receptors of an alcohol-sensitive ANT rat line. *Alcohol Clin Exp Res.* 1991; 15:241–248. [PubMed: 1647706]
- Uusi-Oukari M, Korpi ER. Functional properties of GABA<sub>A</sub> receptors in two rat lines selected for high and low alcohol sensitivity. *Alcohol.* 1992; 9:261–269. [PubMed: 1318725]
- Uusi-Oukari M, Maksay G. Allosteric modulation of [<sup>3</sup>H]EBOB binding to GABA<sub>A</sub> receptors by diflunisal analogues. *Neurochem Int.* 2006; 49:676–682. [PubMed: 16884828]
- Valenzuela CF, Mameli M, Carta M. Letter to the Editor. *Alcohol, Clin Exp Res.* 2005; 29:1356–1357. [PubMed: 16046882]
- Wallner M, Hanchar HJ, Olsen RW. Ethanol enhances  $\alpha$ 4 $\beta$ 3 $\delta$  and  $\alpha$ 6 $\beta$ 3 $\delta$   $\gamma$ -aminobutyric acid type A receptors at low concentrations known to affect humans. *Proc Natl Acad Sci USA.* 2003; 100:15218–15223. [PubMed: 14625373]
- Wieland HA, Lüddens H, Seeburg PH. A single histidine in GABA<sub>A</sub> receptors is essential for benzodiazepine agonist binding. *J Biol Chem.* 1992; 267:1426–1429. [PubMed: 1346133]
- Wong G, Ovaska T, Korpi ER. Brain regional pharmacology of GABA<sub>A</sub> receptors in alcohol-preferring AA and alcohol-avoiding ANA rats. *Addict Biol.* 1996a; 1:263–272. [PubMed: 12893465]
- Wong G, Sarviharju M, Toropainen M, Matecka D, Korpi ER. Pharmacologic actions of subtype-selective and novel GABAergic ligands in rat lines with differential sensitivity to ethanol. *Pharmacol Biochem Behav.* 1996b; 53:723–730. [PubMed: 8866977]



**Fig. 1.** Association of total, DZ-S and DZ-IS [ $^3\text{H}$ ]Ro 15-4513 binding to cerebellar membranes from RR and QQ rats (mean  $\pm$  SEM of three independent experiments made in triplicate). DZ-IS binding was measured in the presence of 1  $\mu\text{M}$  diazepam.




**Fig. 2.** Dissociation of total, DZ-S and DZ-IS [<sup>3</sup>H]Ro 15-4513 binding from cerebellar membranes from RR and QQ rats (mean ± SEM of three independent experiments made in triplicate). DZ-IS binding was measured in the presence of 1 μM diazepam.




**Fig. 3.** A. Association kinetics of 5 nM [ $^3\text{H}$ ]Ro 15-4513 to  $\alpha 1\beta 3\gamma 2$ ,  $\alpha 6-100R-\beta 3\gamma 2$  and  $\alpha 6-100Q-\beta 3\gamma 2$  recombinant receptors expressed in HEK cells. 3B. Dissociation of [ $^3\text{H}$ ]Ro 15-4513 binding from  $\alpha 1\beta 3\gamma 2$ ,  $\alpha 6-100R-\beta 3\gamma 2$  and  $\alpha 6-100Q-\beta 3\gamma 2$  recombinant receptors expressed in HEK cells (mean  $\pm$  SEM, n=3 independent experiments made in triplicate).







**Fig. 4.** Association of [ $^3\text{H}$ ]Ro 15-4513 to  $\alpha 6-100R-\beta 3\gamma 2$  recombinant receptors and to cerebellar membranes from RR rats using longer incubation times (mean values from two independent experiments made in triplicate). DZ-IS binding was measured in the presence of 1  $\mu\text{M}$  diazepam.



**Fig. 5.** Saturation analysis of [<sup>3</sup>H]Ro 15-4513 binding to hippocampal membranes of RR and QQ rats (mean  $\pm$  SEM; RR, n= 4; QQ, n=5), Difference in  $B_{\max}$  between the rat lines:  $p=0.062$ , two-tailed unpaired  $t$ -test.



**Fig. 6.** Saturation analysis of [<sup>3</sup>H]muscimol binding to cerebellar membranes of RR and QQ rats (mean ± SEM; RR, n=7; QQ, n=8). \*p<0.05, significance of the difference in B<sub>max</sub> between the rat lines, two-tailed unpaired *t*-test.



**Fig. 7.** Antagonizing effect of furosemide on GABA-induced displacement of [<sup>3</sup>H]EBOB in cerebellar membranes from RR and QQ rats (mean  $\pm$  SEM; RR, n=11; QQ, n=13). \*\*p<0.01, significance of the difference between GABA-induced displacement and antagonism of this displacement in RR rats, one-way ANOVA followed by Tukey's post hoc test.

**Table 1**

Association and dissociation rate constants of [<sup>3</sup>H]Ro 15-4513 binding in cerebellar membranes of RR and QQ rats and in recombinant receptors expressed in HEK cells

	$K_{on}$ ( $M^{-1} \times min^{-1}$ )	$K_{off}$ ( $min^{-1}$ )	$K_D$ ( $K_{off}/K_{on}$ ) (nM)
Total binding			
RR	$4.6 \pm 0.4 \times 10^7$	$0.063 \pm 0.002$	$1.4 \pm 0.1$
QQ	$7.6 \pm 1.1 \times 10^7$	$0.080 \pm 0.006$	$1.1 \pm 0.3$
DZ-S binding			
RR	$6.0 \pm 0.2 \times 10^7$	$0.097 \pm 0.004$	$1.6 \pm 0.1$
QQ	$6.9 \pm 0.9 \times 10^7$	$0.090 \pm 0.008$	$1.4 \pm 0.3$
DZ-IS binding			
RR	$8.2 \pm 0.3 \times 10^6$	$0.019 \pm 0.001$	$2.3 \pm 0.1$
QQ	$6.1 \pm 0.8 \times 10^7^{##}$	$0.055 \pm 0.001^{###}$	$0.9 \pm 0.2^{###}$
Recombinant recept.			
$\alpha 1\beta 3\gamma 2$	$3.4 \pm 0.2 \times 10^7^{***}$	$0.098 \pm 0.010^{**}$	$2.9 \pm 0.3^*$
$\alpha 6-100R-\beta 3\gamma 2$	$8.0 \pm 0.8 \times 10^6^{***}$	$0.018 \pm 0.002$	$2.4 \pm 0.4^*$
$\alpha 6-100Q-\beta 3\gamma 2$	$8.7 \pm 0.2 \times 10^7^{***}$	$0.083 \pm 0.013^{**}$	$1.0 \pm 0.1$

DZ-S = diazepam-sensitive, DZ-IS = diazepam-insensitive,  $K_D$  = dissociation constant,  $K_{on}$  = association rate constant,  $K_{off}$  = dissociation rate constant. Total, DZ-S and DZ-IS binding to cerebellar membranes, mean  $\pm$  SEM values (n=3, measured in triplicate).

$###$  p<0.001,

$##$  p<0.01, significantly different from the corresponding RR value (unpaired *t*-test); Recombinant receptors, mean  $\pm$  SEM (n=3, measured in triplicate).

$***$  p<0.001, significantly different from two other groups;

$**$  p<0.01, significantly different from  $\alpha 6-100R$  value;

$*$  p<0.05, significantly different from  $\alpha 6-100Q$  value; one-way ANOVA followed by Tukey's post hoc test.