

Paranoid Schizophrenia is Characterized by Increased CB₁ Receptor Binding in the Dorsolateral Prefrontal Cortex

Victoria S Dalton^{1,2}, Leonora E Long^{2,3,4}, Cyndi Shannon Weickert^{2,3,5} and Katerina Zavitsanou^{*,1,2}

¹ANSTO Life Sciences, Australian Nuclear Science and Technology Organization, Sydney, New South Wales, Australia; ²Schizophrenia Research Institute, Sydney, New South Wales, Australia; ³Schizophrenia Research Laboratory, Neuroscience Research Australia, Sydney, New South Wales, Australia; ⁴School of Medical Sciences, University of New South Wales, Sydney, New South Wales, Australia; ⁵School of Psychiatry, University of New South Wales, Sydney, New South Wales, Australia

A number of studies suggest a dysregulation of the endogenous cannabinoid system in schizophrenia (SCZ). In the present study, we examined cannabinoid CB₁ receptor (CB₁R) binding and mRNA expression in the dorsolateral prefrontal cortex (DLPFC) (Brodmann's area 46) of SCZ patients and controls, post-mortem. Receptor density was investigated using autoradiography with the CB₁R ligand [³H] CP 55 940 and CB₁R mRNA expression was measured using quantitative RT-PCR in a cohort of 16 patients with paranoid SCZ, 21 patients with non-paranoid SCZ and 37 controls matched for age, post-mortem interval and pH. All cases were obtained from the University of Sydney Tissue Resource Centre. Results were analyzed using one-way analysis of variance (ANOVA) and *post hoc* Bonferroni tests and with analysis of covariance (ANCOVA) to control for demographic factors that would potentially influence CB₁R expression. There was a main effect of diagnosis on [³H] CP 55 940 binding quantified across all layers of the DLPFC ($F(2,71) = 3.740$, $p = 0.029$). *Post hoc* tests indicated that this main effect was due to patients with paranoid SCZ having 22% higher levels of CB₁R binding compared with the control group. When ANCOVA was employed, this effect was strengthened ($F(2,67) = 6.048$, $p = 0.004$) with paranoid SCZ patients differing significantly from the control ($p = 0.004$) and from the non-paranoid group ($p = 0.016$). In contrast, no significant differences were observed in mRNA expression between the different disease subtypes and the control group. Our findings confirm the existence of a CB₁R dysregulation in SCZ and underline the need for further investigation of the role of this receptor particularly in those diagnosed with paranoid SCZ.

Neuropsychopharmacology (2011) **36**, 1620–1630; doi:10.1038/npp.2011.43; published online 6 April 2011

Keywords: CB₁ receptor; paranoid schizophrenia; post-mortem; autoradiography; real time PCR

INTRODUCTION

Cannabis and cannabis-related drugs act principally through two seven-transmembrane-domain, G protein-coupled receptors termed the cannabinoid 1 (CB₁R) and cannabinoid 2 (CB₂R) receptors that are also activated by endogenous ligands termed endocannabinoids. The CB₁R is considered to mediate the majority of the psychoactive properties of cannabis (Ameri, 1999) and is expressed abundantly throughout the human brain (Glass *et al*, 1997). In the cortex, the CB₁R is expressed mainly on presynaptic terminals of GABAergic inhibitory interneurons (Eggen and Lewis, 2007).

Abnormalities in the CB₁R in cortical regions in schizophrenia (SCZ) have been reported in a number of studies. Dean *et al* (2001) found increased binding of the

CB₁R agonist [³H] CP 55 940, in the dorsolateral prefrontal cortex (DLPFC, Brodmann's area 9) in patients with SCZ that was independent of cannabis use before death and in caudate-putamen that appeared to be related to premorbid cannabis use in these patients. Urigüen *et al* (2009) reported that immunodensity of CB₁R in the frontal cortex was significantly decreased in antipsychotic-treated patients with SCZ but not in drug-free patients (Urigüen *et al*, 2009). Looking in other cortical regions, we have previously shown an increase in the binding of the selective cannabinoid antagonist [³H] SR141716A in the anterior cingulate cortex of patients with SCZ post-mortem (Zavitsanou *et al*, 2004), a finding that was confirmed in the posterior cingulate cortex (Newell *et al*, 2006). In contrast to the receptor binding studies, reduced cortical CB₁R mRNA and protein expression have been found in the post-mortem DLPFC (Brodmann's areas 9 and 46) in SCZ (Eggen *et al*, 2008; Eggen *et al*, 2010), whereas another study (Koethe *et al*, 2007) found no change in the density of CB₁R immunopositive cells in the anterior cingulate cortex in SCZ. Importantly, a recent imaging study using the novel positron emission tomography (PET) CB₁R tracer [¹¹C]

*Correspondence: Dr K Zavitsanou, ANSTO Life Sciences, Australian Nuclear Science and Technology Organisation, Building 1, Locked Bag 2001 Kirrawee DC, Sydney, NSW 2232, Australia, Tel: +61 29 717 3337, Fax: +61 29 717 9262, E-mail: kza@ansto.gov.au
Received 18 November 2010; revised 3 March 2011; accepted 3 March 2011

OMAR (JHU75528) showed an elevated mean binding of this tracer in patients with SCZ across all regions studied that reached significance in the pons (Wong *et al*, 2010a). Perhaps more importantly, the same study suggested that CB₁R binding in SCZ increases with severity of the positive symptoms and decreases with severity of negative symptoms (Wong *et al*, 2010a). Another imaging study using the selective high affinity PET radioligand [¹⁸F]MK-9470 showed a significant increase of CB₁R availability in the mesocorticolimbic circuitry, especially in the nucleus accumbens of both antipsychotic treated and untreated SCZ patients compared with controls (Ceccarini *et al*, 2010).

The discrepancies in the studies above may reflect methodological or regional differences, or may relate to other factors such as antipsychotic medication (eg, Urigüen *et al*, 2009) or cannabis consumption (Dean *et al*, 2001) or they may be related to cohort make up, as cohorts varied in SCZ subtype composition and the degree of symptoms varied among individuals in all of these studies. Recently, it has been suggested that different genetic and pathophysiological mechanisms may underlie different subtypes of SCZ (Chavarría-Siles *et al*, 2008). For example, genetic studies provide evidence that a variation of CNR1, the CB₁ receptor gene, confers risk for hebephrenic (disorganized) SCZ with no association to the more general phenotype of SCZ. The same authors suggested that inclusion of other subtypes may dilute the power to find association of SCZ with changes in CNR1. Similarly, the inclusion of different disease subtypes in the post-mortem studies described above may have an impact on the measures of CB₁R protein or mRNA levels and contributes to the discrepancies reported. Importantly, Giuffrida *et al* (2004) have shown that cerebrospinal fluid (CSF) levels of the endocannabinoid anandamide are profoundly and selectively elevated in paranoid-type SCZ, and that the levels of anandamide negatively correlate with the psychotic symptoms of the disease (Giuffrida *et al*, 2004).

In this study, we hypothesized that CB₁R density may vary according to the disease's subtype clusters. In the past decade, SCZ research on post-mortem human tissue has matured, largely due to the widespread availability of larger cohorts that allow for examination of specific disease subtypes. Thus, to understand further the changes in CB₁R in SCZ, we measured CB₁R binding and mRNA expression in the DLPFC (Brodmann's area 46) of a large cohort of control subjects and subjects with SCZ to determine (a) whether these changes were disease subtype specific with focus on the paranoid subtype and (b) whether changes in CB₁R binding were associated with changes in levels of mRNA expression. In our study close attention was paid to peri-mortem and demographic variables, which can impact studies of this kind (Mato and Pazos, 2004; Urigüen *et al*, 2009; Weickert *et al*, 2010).

MATERIALS AND METHODS

Human Post-Mortem Brain Samples

All research was approved and conducted under the guidelines of the Human Research Ethics Committees at the University of Wollongong (#HE99/222) and at the

University of New South Wales (#HREC07261). Tissue was provided by the New South Wales Tissue Resource Centre (University of New South Wales Human Research Ethic Committee #HREC07261). Characterization and tissue preparation for this Australian SCZ cohort has been described previously (Weickert *et al*, 2010). A post-mortem clinical diagnosis was determined for each case through careful examination of the donor's lifetime symptom profile by experienced clinicians. The Diagnostic Instrument for Brain Studies—Revised (DIBS) was then employed. DIBS is a semi-structured instrument specifically designed for post-mortem psychiatric assessment using medical records and informants that enables diagnosis at a sub-syndrome and symptom-based level (Sundqvist *et al*, 2008). The DIBS was applied to the clinical summary to generate a diagnosis of SCZ based on ICD-10, DSM-III-R, DSM-IV, Research Diagnostic Criteria, Schneider and Feighner criteria (Hill *et al*, 1996; Keks *et al*, 1997; Roberts *et al*, 1998). A diagnostic subtype was established as rated in individual items in the DIBS. For example according to the ICD-10, persons with paranoid SCZ must meet the general criteria for SCZ and must also experience prominent delusions or hallucinations. However, flattening of affect, catatonic symptoms, or incoherent speech must not dominate the clinical picture, although they may be present to a mild degree. Profiles were cross-matched with DSM-IV diagnostic criteria to determine the most appropriate subtype for the case. Normal controls had no history of significant psychological problems or psychological care, psychiatric admissions or drug detoxification, and no known history of psychiatric symptoms or substance abuse, as determined by both telephone screening and medical records, and no significant neuropathological changes upon examination of the brain (Weickert *et al*, 2010).

Tissue samples and sections were prepared from the large cohort of non-paranoid SCZ ($n=21$), paranoid SCZ ($n=16$), and control ($n=37$) cases matched for age, gender, pH, and post-mortem interval (Table 1; Weickert *et al*, 2010). The non-paranoid SCZ group included cases that met criteria for undifferentiated ($n=7$), residual ($n=2$), disorganized ($n=5$), and schizoaffective (bipolar and depressive subtype, $n=7$) type of SCZ (Table 1).

Tissue Dissection and Section Preparation

Tissue dissection has been described in detail previously (Weickert *et al*, 2010). Briefly, at autopsy, brain weight, and volume were determined (Harper *et al*, 1988). The fresh tissue was cut into ~1 cm coronal slices and various anatomical areas were dissected for separate freezing. For the DLPFC dissections, frozen tissue was dissected on a dry ice platform using a dental drill (Cat# UP500-UG33, Brasseler, USA). DLPFC tissue (average weight of tissue ~0.5 g gray matter tissue from the crown of the middle frontal gyrus) was obtained from the coronal slab corresponding to the middle one-third (rostral caudally) found anterior to the genu of the corpus callosum. The dissected tissue slices were immediately frozen and stored at -80°C ($\pm 5^{\circ}\text{C}$). Coronal tissue sections of the DLPFC (14 μm) were cut on a cryostat, thaw mounted onto microscope slides and stored at -80°C until use.

Table 1 Summary of Cohort Demographics

	n	pH	Age at death	PMI (h)	Freezer/ months	Brain weight (g)	RIN	Age of onset	Duration of illness in years	Lifetime chlorpromazine (mg)	[³ H]CP55940 binding (fmoles/mg TE)	Normalized CB ₁ mRNA expression
Control	37	6.7 ± 0.3	51.1 ± 14.6	248 ± 11.0	69.6 ± 42.7	1446.4 ± 127.1	7.3 ± 0.6				47.9 ± 12.6	4.6 ± 1.1
Paranoid schizophrenia	16	6.6 ± 0.3	51.6 ± 13.0	32.2 ± 16.0	83.6 ± 27.5	1364.4 ± 177.7	7.2 ± 0.6	23.6 ± 6.4	28.1 ± 12.8	7.7 × 10 ⁶ ± 7.8 × 10 ⁶	58.7 ± 13.4	4.9 ± 1.1
Non-paranoid schizophrenia	21	6.6 ± 0.3	51.1 ± 15.3	25.6 ± 11.4	77.1 ± 43.7	1417.1 ± 153.2	7.3 ± 0.6	23.8 ± 6.0	27.3 ± 14.9	8.1 × 10 ⁶ ± 8.2 × 10 ⁶	47.9 ± 16.2	4.1 ± 1.0
<i>Non-paranoid subtypes</i>												
Residual	2	6.1 ± 0.6	51.0 ± 0.0	16.5 ± 6.4	136.0 ± 4.2	1325.0 ± 233.3	6.6 ± 0.5	31.0 ± 5.7	20.0 ± 5.7	3.4 × 10 ⁶ ± 1.8 × 10 ⁶	29.6 ± 21.0	3.2 ± 0.2
Disorganized	5	6.7 ± 0.3	54.0 ± 13.1	25.7 ± 6.8	74.8 ± 42.8	1385.2 ± 184.2	7.2 ± 0.7	19.8 ± 1.8	34.2 ± 12.5	8.9 × 10 ⁶ ± 4.3 × 10 ⁶	47.6 ± 15.8	4.9 ± 1.4
Depressive	4	6.9 ± 0.1	52.3 ± 24.0	27.5 ± 15.8	46.8 ± 20.7	1490.0 ± 94.2	7.4 ± 0.5	24.5 ± 9.3	27.8 ± 21.3	4.4 × 10 ⁶ ± 5.5 × 10 ⁶	48.7 ± 7.3	4.4 ± 0.5
Bipolar	3	6.6 ± 0.3	50.7 ± 14.6	23.7 ± 5.9	71.7 ± 29.7	1396.7 ± 105.0	7.2 ± 0.3	29.3 ± 2.1	21.3 ± 12.5	1.1 × 10 ⁷ ± 1.0 × 10 ⁷	49.2 ± 10.3	3.3 ± 0.6
Undifferentiated	7	6.5 ± 0.2	48.6 ± 16.9	28.0 ± 14.9	82.0 ± 53.2	1433.0 ± 175.1	8.0 ± 0.5	21.9 ± 3.9	27.0 ± 16.4	9.9 × 10 ⁶ ± 1.2 × 10 ⁷	52.4 ± 21.0	3.9 ± 0.9

Abbreviations: h, hours; PMI, post-mortem interval; RIN, RNA integrity number; TE, tissue equivalent. Average values ± SD for continuous variables are shown.

In Vitro Autoradiography

All sections (three sections per case) were processed simultaneously to minimize experimental variance. On the day of the experiment, sections were pre-incubated for 30 min at room temperature in 50 mM Tris HCl (pH 7.4) buffer containing 5% bovine serum albumin (BSA). Two sections per case were then incubated for 2 h at room temperature in the same buffer with the addition of 10 nM [³H] CP 55 940 (specific activity 139.6 Ci/mmol, Perkin Elmer, USA). Non-specific binding was determined by incubating adjacent sections (one/case) in 10 nM [³H] CP 55 940 in the presence of 10 μM CP 55 940. After the incubation all sections were washed for 1 h at 4 °C in 50 mM Tris HCl (pH 7.4) containing 1% BSA followed by a second wash for 3 h and by third wash for 5 min in the same buffer. Sections were then dipped briefly in ice-cold distilled water and then air dried.

Dried sections were apposed to Kodak Biomax MR film, together with autoradiographic standards ([³H] microscales from Amersham), in X-ray film cassettes for 30 days.

Quantitative Analysis of Autoradiographic Images

Films were analyzed by using a computer-assisted image analysis system, Multi-Analyst, connected to a GS-690 Imaging Densitometer (Bio-Rad, USA). Two to three areas for quantification on each slide were previously defined by identifying the cyto-architectural characteristics of Brodmann's area 46 with neuronal nuclei (NeuN) immunostaining (Rajkowska and Goldman-Rakic, 1995; Yang *et al*, 2010). A rectangular box was drawn in each specified area from layers I-VI and density of receptor binding within these areas quantified. Quantification was performed blind to diagnosis by measuring the average optical density in three adjacent brain sections (two for the total binding and one for the non-specific binding). Non-specific binding (< 20–30% in the majority of cases) was subtracted from the total binding to determine the specific binding. Optical density measurements were then converted into fmoles [³H] CP 55 940 per mg tissue equivalent (fmoles/mg TE), according to the calibration curve obtained from the tritium standards.

Total RNA Isolation and RNA Quality Assessment

Total RNA was extracted from ~ 300 mg of frozen tissue per subject for qPCR analysis using Trizol (Invitrogen, Carlsbad, California) according to the manufacturer's instructions (Kozlovsky *et al*, 2004). The quality of extracted total RNA was determined using the Agilent Bioanalyzer 2100 (Agilent Technologies, Palo Alto, California). A volume of 100–200 ng RNA was applied to an RNA 6000 Nano LabChip, without heating before loading. The RNA Integrity Number (RIN) was used as an indicator of RNA quality, ranging from 1 (lowest quality) to 10 (highest quality). The cDNA was synthesized in three reactions of 3 μg of total RNA in a 26.25 μl reaction using the Superscript First-Strand Synthesis Kit (Invitrogen) according to the manufacturer's protocol.

Quantitative Real-Time PCR (qPCR)

CB₁R mRNA levels were measured using a pre-designed TaqMan Gene Expression Assay (Applied Biosystems) for CNR1 (Hs00275634_m1). Each 10 µl qPCR reaction contained FAM-labeled probe (250 nmol/l), primers (900 nmol/l), and 1.14 ng cDNA in 1x Taqman Universal Mastermix containing AmpliTaq Gold DNA polymerase, deoxynucleoside triphosphates, uracil-N-glycosylase, and passive reference. The PCR protocol used involved incubation at 50 °C for 2 min and 95 °C for 10 min, followed by 40 consecutive cycles of 95 °C for 15 s and 60 °C for 1 min. Serial dilutions of pooled cDNA (from all cases) were included on every qPCR plate and used by Sequence Detection Software (SDS; Applied Biosystems) to quantify sample expression by the relative standard curve method. Control wells containing no cDNA template displayed no amplification in any assay. Efficiencies of the qPCR reactions ranged from 77 to 100%, with r^2 values of between 0.95 and 1.00. All reactions were performed in triplicate. Expression levels were normalized to the geometric mean of four 'housekeeper' genes that did not change expression with diagnosis: ACTB (Hs99999903_m1), GAPDH (Hs99999905_m1), UBC (Hs00824723_m1), and TBP (Hs00427620_m1) (Weickert *et al*, 2010; Wong *et al*, 2010b). Population outliers were excluded if the normalized expression value was greater than two standard deviations from the group mean. As a result, RNA samples were unavailable for five members of the control cohort and 2 members of the SCZ cohort.

Statistical Analysis

Statistical analyses were conducted using SPSS software (version 14). The Kolmogorov–Smirnov test was employed to test for normal (Gaussian) distribution. Parametric tests were used in subsequent analysis as data were normally distributed. Mean values for binding and mRNA expression are reported \pm SD.

One-way analysis of variance (ANOVA) or Student's *t*-tests were used to compare the mean brain pH, age at death, PMI, freezer storage time, brain weight, RIN, mean age of illness onset, illness duration, and estimated lifetime exposure to antipsychotics (stated as chlorpromazine equivalent dose (mg)) between the diagnostic groups (Table 1).

For continuous, descriptive variables (brain pH, age at death, PMI, freezer storage time, brain weight, RIN, age of illness onset, illness duration, and estimated lifetime exposure to antipsychotics), we tested for significant Pearson correlations for the abnormal CB₁R binding and mRNA expression. Non-continuous descriptive variables such as gender (male/female), hemisphere (right/left), cause of death (suicide/other), presence of a cannabis use history (yes/no), agonal state (excellent: 1, good: 2, and poor: 3), daily alcohol intake (none: 0, low: 1, moderate: 2, high: 3, and unknown: 4), and tobacco smoking (unknown: 0, moderate: 1, and heavy: 2) were used as grouping variables with *t*-tests or one-way ANOVA to evaluate their effects on binding and mRNA expression.

The effects of the continuous and non-continuous variables were examined in all subjects (in both the control and SCZ group), in the control group alone, and the SCZ

group (not divided into paranoid and non-paranoid SCZ) alone. Effects of the continuous and non-continuous variables were then examined in the paranoid and non-paranoid SCZ groups to ensure that our measurements were not affected by a particular variable in each subgroup.

CB₁R binding and mRNA expression levels were compared between diagnostic groups (paranoid and non-paranoid SCZ and controls) using one-way ANOVA followed by *post hoc* Bonferroni tests to account for multiple comparisons. Separate analysis of covariance (ANCOVA) controlling for pH, age at death, freezer storage time, brain volume and RIN followed by Bonferroni *post hoc* tests were also calculated where appropriate.

Exploratory one-way ANOVAs were also performed to compare CB₁R binding and mRNA expression between the diagnostic SCZ subgroups (residual, disorganized, schizoaffective, undifferentiated, paranoid) and controls. Due to low subject numbers in some groups, LSD *post hoc* tests were used where appropriate to reduce the risk of type II error.

Results

The mean pH, age at death, PMI, freezer storage time, brain weight, and RIN did not differ between the groups studied (paranoid and non-paranoid SCZ, and controls; $0.008 \leq F \leq 2.153$, $df = 2$, $0.124 \leq p \leq 0.992$, Table 1). Also the age of onset of disease, the duration of illness, and lifetime exposure to antipsychotics (lifetime chlorpromazine) did not differ between paranoid and non-paranoid groups ($-0.167 \leq t \leq 0.167$, $df = 35$, $0.868 \leq p \leq 0.904$, Table 1) or between the six diagnostic subgroups ($F(5,31) = 1.848$, $p = 0.132$). In agreement with the literature, however, SCZ cases of the disorganized subtype had the earliest mean age of onset of disease (~ 20 years, Table 1).

Disease-Related Effects

One-way ANOVA revealed a statistically significant difference between the three diagnostic groups (paranoid SCZ, non-paranoid SCZ, and control) on [³H] CP 55 940 binding in the DLPFC ($F(2,71) = 3.740$, $p = 0.029$, Figures 1 and 2a). *Post hoc* analysis (Bonferroni) indicated that patients with paranoid SCZ had significantly (22%) higher levels of binding compared with controls (58.7 ± 13.4 vs 47.9 ± 12.6 fmoles/mg TE, $p = 0.036$, Figure 2a). Paranoid SCZ patients also had higher binding compared with the non-paranoid SCZ patients but this increase was not statistically significant (58.7 ± 13.4 vs 47.9 ± 16.2 fmoles/mg TE, $p = 0.068$). However, given that the sample size was smaller in the patient groups compared with the control group, we may have been underpowered to detect this increase. To have sufficient power (eg, 80%) to detect an increase in binding in paranoid patients compared with the non-paranoid patients, the groups would need to be increased from $n = 12$ – 17 to $n = 24$ per group.

Peri-mortem and demographic variables can impact CB₁R density and mRNA expression in post-mortem human studies (Mato and Pazos, 2004). Therefore although most of the correlations between the continuous variables and our data were between 0.3 and 0.5 (see below and also Table 2),




Figure 1 Typical autoradiographs showing [³H] CP55 940 binding in the dorsolateral prefrontal cortex in control, non-paranoid (NP SCZ), and paranoid (P SCZ) SCZ groups. Areas for quantification on each slide were previously defined by identifying the cyto-architectural characteristics of BA46 with NeuN immunostaining (Rajkowska and Goldman-Rakic, 1995; Yang *et al.*, 2010).




Figure 2 Cannabinoid CB₁ receptor density (a) and mRNA (b) expression in the dorsolateral prefrontal cortex in control, non-paranoid (NP SCZ), and paranoid (P SCZ) schizophrenia patients. (a) [³H] CP 55 940 binding density in fmoles/mg TE (tissue equivalent). (b) CB₁ mRNA expression normalized to the geometric mean of four housekeeping genes. #: A 22% increase in CB₁ receptor density was found in the P SCZ group that was statistically significant when compared with controls in a one-way analysis of variance ($F(2,71) = 3.740$, $p = 0.029$) and significantly different from controls ($p = 0.004$) following analysis of covariance (ANCOVA) ($F(2,67) = 6.048$, $p = 0.004$). *: The 22% increase in CB₁ receptor density in the P SCZ was statistically significant when compared with the NP SCZ group ($p = 0.016$) after ANCOVA ($F(2,67) = 6.048$, $p = 0.004$).

suggesting a moderate association (Cohen 1988), we proceeded with ANCOVA to account for continuous variables (pH, age at death, freezer storage time, and brain weight) that might impact on our data. Importantly, when ANCOVA was conducted, the significant difference between

the three diagnostic groups was retained ($F(2,67) = 6.048$, $p = 0.004$) with paranoid SCZ patients having significantly higher binding as compared with the control group ($p = 0.004$) and the non-paranoid SCZ group ($p = 0.016$) (Figure 2a).

In contrast, we failed to reveal a statistically significant difference between CB₁ mRNA expression in the three diagnostic groups by ANOVA ($F(2,63) = 2.183$, $p = 0.121$) (Figure 2b). This result remained unchanged after we controlled for continuous variables with ANCOVA ($F(2,57) = 2.239$, $p = 0.116$). A positive correlation was found, however, between CB₁R mRNA expression and binding ($r = 0.384$; $p = 0.001$).

We also carried out exploratory one-way ANOVAs treating each group (residual, disorganized, schizoaffective, undifferentiated, paranoid SCZ, and controls) as a single diagnosis to determine the strength of the complex variability on CB₁R binding and mRNA expression across different subtypes of SCZ. This analysis should be treated with caution, however, due to low subject numbers within some groups (Table 1). A significant effect of diagnosis on CB₁R binding was found ($F(5,68) = 2.392$, $p = 0.047$; Figure 3a). *Post hoc* tests indicated that this effect was due to patients with paranoid SCZ having higher CB₁R binding compared with controls ($p = 0.011$) and to the residual SCZ group ($p = 0.006$) and patients with residual SCZ having lower CB₁R binding compared with those with undifferentiated SCZ ($p = 0.042$; Table 1, Figure 3a). CB₁R mRNA did not differ between the diagnostic subgroups ($F(5,60) = 1.644$, $p = 0.162$; Table 1; Figure 3b).

Effects of Continuous and Non-Continuous Variables

Pearson's rank order correlations for continuous variables (brain pH, age at death, PMI, freezer storage time, brain weight, RIN, age of illness onset, illness duration, and estimated lifetime exposure to antipsychotics) are presented in Table 2. The *t*-tests and ANOVA for non-continuous variables (gender (male/female), hemisphere (right/left), cause of death (suicide/other), presence of a cannabis use history (yes/no), agonal state (excellent: 1, good: 2, and poor: 3), daily alcohol intake (none: 0, low: 1, moderate: 2, high: 3, and unknown: 4), and tobacco smoking (unknown: 0, moderate: 1, and heavy: 2)) were performed and are presented as Supplementary information (Figures S1 and S2).

Table 2 Pearson's Rank Order Correlations for Continuous Variables CB₁ Receptor Binding and mRNA Expression in the Dorsolateral Prefrontal Cortex of all Subjects, Controls only, SCZ Patients Alone and the Paranoid and Non-Paranoid SCZ Groups Alone

Variable	All subjects		Controls		SCZ		Paranoid SCZ		Non-paranoid SCZ	
	CB ₁ binding	mRNA	CB ₁ binding	mRNA	CB ₁ binding	mRNA	CB ₁ binding	mRNA	CB ₁ binding	mRNA
pH	r = 0.392 p = 0.001	r = 0.299 p = 0.015	r = 0.339 p = 0.040	r = 0.221 p = 0.209	r = 0.470 p = 0.003	r = 0.376 p = 0.034	r = 0.378 p = 0.149	r = 0.118 p = 0.689	r = 0.526 p = 0.014	r = 0.541 p = 0.021
Age at death	r = -0.358 p = 0.002	r = -0.356 p = 0.003	r = -0.358 p = 0.029	r = -0.401 p = 0.019	r = -0.374 p = 0.023	r = -0.303 p = 0.092	r = -0.464 p = 0.070	r = -0.246 p = 0.396	r = -0.373 p = 0.096	r = -0.337 p = 0.172
PMI (h)	r = 0.137 p = 0.244	r = 0.013 p = 0.915	r = -0.151 p = 0.372	r = -0.043 p = 0.809	r = 0.285 p = 0.087	r = 0.074 p = 0.685	r = 0.178 p = 0.509	r = 0.184 p = 0.528	r = 0.275 p = 0.228	r = 0.148 p = 0.556
Freezer storage time	r = 0.344 p = 0.003	r = -0.177 p = 0.154	r = 0.541 p = 0.001	r = 0.346 p = 0.045	r = 0.148 p = 0.382	r = 0.006 p = 0.976	r = 0.342 p = 0.295	r = 0.561 p = 0.037	r = 0.043 p = 0.852	r = -0.359 p = 0.144
Brain weight (g)	r = 0.291 p = 0.012	r = 0.108 p = 0.390	r = 0.463 p = 0.004	r = 0.161 p = 0.362	r = 0.246 p = 0.142	r = 0.047 p = 0.796	r = 0.291 p = 0.274	r = -0.192 p = 0.512	r = 0.356 p = 0.113	r = 0.446 p = 0.064
RIN		r = 0.285 p = 0.021		r = 0.254 p = 0.154		r = 0.314 p = 0.080		r = 0.379 p = 0.181		r = 0.361 p = 0.141
Age of disease onset					r = -0.137 p = 0.417	r = -0.325 p = 0.070	r = -0.108 p = 0.691	r = -0.246 p = 0.396	r = -0.305 p = 0.179	r = -0.450 p = 0.061
Duration of illness					r = -0.322 p = 0.052	r = -0.162 p = 0.375	r = -0.527 p = 0.036	r = -0.147 p = 0.616	r = -0.206 p = 0.255	r = -0.149 p = 0.555
Lifetime chlorpromazine					r = -0.084 p = 0.622	r = -0.042 p = 0.820	r = -0.210 p = 0.436	r = 0.177 p = 0.546	r = -0.002 p = 0.992	r = -0.154 p = 0.541

Abbreviation: SCZ, schizophrenia.
Significant correlations are shown in bold.

Within all subjects, [³H] CP 55 940 binding was correlated with pH ($r = 0.392$, $p = 0.001$), age at death ($r = -0.385$, $p = 0.002$), freezer storage time ($r = 0.344$, $p = 0.003$), and brain weight ($r = 0.291$, $p = 0.012$) (Table 2). However, there was a negative correlation of borderline significance between freezer time and age at death ($r = -0.208$, $p = 0.075$), which may explain the relationship between binding and freezer time. Binding levels did not correlate with PMI, or vary according to gender, hemisphere, agonal state, daily alcohol, and smoking (Table 2, Supplementary Figure S1). Significant correlations were observed between mRNA expression and pH ($r = 0.299$, $p = 0.015$), age at death ($r = -0.356$; $p = 0.003$), and RIN ($r = 0.285$; $p = 0.021$) (Table 2). CB₁ mRNA expression was unaffected by all other continuous (Table 2) and non-continuous variables (data not shown).

In the control group, there were significant positive correlations between CB₁ binding and pH ($r = 0.339$, $p = 0.040$), freezer storage time ($r = 0.541$, $p = 0.001$) and brain weight ($r = 0.463$, $p = 0.004$), and a negative correlation between binding and age at death ($r = -0.358$, $p = 0.029$) (Table 2). Binding was unaffected by all the other continuous and non-continuous variables (Table 2 and Supplementary Figure S1). A significant negative correlation was also found between mRNA expression and age at death ($r = -0.401$, $p = 0.019$) whereas the correlation between mRNA and freezer storage time was positive ($r = 0.346$, $p = 0.045$) (Table 2). The mRNA was not found to be affected by any of the non-continuous variables examined (data not shown).

In the SCZ group as a whole, binding was correlated with pH ($r = 0.470$, $p = 0.003$) and age at death ($r = -0.374$, $p = 0.023$), and correlations of borderline significance were found between binding and PMI ($r = 0.285$, $p = 0.087$) and duration of illness ($r = -0.322$, $p = 0.052$) (Table 2). CB₁ mRNA expression was positively correlated with pH ($r = 0.376$, $p = 0.034$) with a correlation of borderline significance between mRNA and age at death ($r = -0.303$, $p = 0.092$) (Table 2). CB₁ mRNA expression was unaffected by all other continuous variables (Table 2). The non-continuous variables considered were not shown to have an effect on CB₁ binding (Supplementary Figure S2) or mRNA expression (data not shown).

In the paranoid SCZ group alone, [³H] CP 55 940 binding was unaffected by all continuous and non-continuous variables except duration of illness ($r = -0.527$, $p = 0.036$) (Table 2, Supplementary Figure S2). A significant positive correlation was found between mRNA expression and freezer storage time ($r = 0.561$, $p = 0.037$) (Table 2). CB₁ mRNA expression was unaffected by all other continuous and non-continuous variables (Table 2, data not shown for non-continuous variables).

In the non-paranoid SCZ group alone, CB₁ receptor density was affected by pH ($r = 0.526$, $p = 0.014$) but was not affected by any other continuous and non-continuous variable (Table 2, Supplementary Figure S2). CB₁ mRNA expression was affected by pH ($r = 0.541$, $p = 0.021$) (Table 2) and was unaffected by all non-continuous variables (data not shown).




Figure 3 Cannabinoid CB₁ receptor density (a) and mRNA (b) expression in the DLPFC in diagnostic subtypes of schizophrenia and controls. (a) [³H] CP 55940 binding density in fmol/mg TE (tissue equivalent). In an exploratory one-way analysis of variance (ANOVA), where the bipolar and depressive subtypes were treated as one group (schizoaffective) and compared with controls along with the other four schizophrenia subtypes, a significant variation in CB₁R density was found ($F(5,68) = 2.392, p = 0.047$). * $p < 0.05$, $^{\$}p = 0.006$ in LSD *post hoc* tests. (b) CB₁ mRNA expression normalized to the geometric mean of four housekeeping genes. CB₁ mRNA expression did not differ between diagnostic subtypes ($F(5,60) = 1.644, p = 0.162$) when analyzed with one-way ANOVA.

DISCUSSION

Disease-Related Effects

Despite the widespread acknowledgement of the heterogeneity of SCZ, few studies have investigated whether indices of neuroreceptor densities/mRNA expression are associated with different clinical subtypes of the disorder. In the present study, we report a significant increase of 22% in cannabinoid CB₁R binding in the DLPFC in a subgroup of patients who suffered from paranoid SCZ compared with normal controls. The patients with paranoid SCZ also had elevated CB₁R binding compared with patients with non-paranoid SCZ.

Looking at a smaller cohort of 14 SCZ patients and matched controls, Dean *et al* (2001) found increased [³H]

CP 55940 binding in Brodmann's area 9 of the DLPFC. We have also previously shown an increase in binding sites for the selective antagonist [³H] SR141716A in the anterior cingulate cortex of 10 SCZ patients compared with their matched controls (Zavitsanou *et al*, 2004), a finding that was confirmed in the posterior cingulate cortex with [³H] CP 55940 (Newell *et al*, 2006). In contrast, Deng *et al* (2007) found no changes in both the binding of [³H] SR141716A or [³H] CP 55940 in the superior temporal gyrus in eight SCZ patients. Recently, the development of suitable radioligands that target the CB₁R *in vivo* in the living brain has allowed for the study of CB₁Rs in SCZ patients using PET. In agreement with the post-mortem studies, two recent PET studies (Wong *et al*, 2010a; Ceccarini *et al*, 2010) also reported elevated CB₁R binding in the pons and nucleus accumbens respectively in patients with SCZ.

The increase in CB₁R binding we observed in the present study was not accompanied by changes in CB₁R mRNA. Despite this lack of overall change in primary transcript levels of CB₁R mRNA, there was a positive correlation between mRNA and binding in SCZ. As this correlation was of a moderate effect, it is possible that the increase in CB₁R binding sites without a similar change in mRNA may arise from a change in post-translational processes such as a greater rate of translation per mRNA molecule or less receptor degradation/turnover. In agreement with our mRNA data and using the same methodology, Uriguen *et al* (2009) reported unchanged CB₁ mRNA expression in Brodmann's area 9 in SCZ. In contrast, Eggen *et al* (2008) reported decreases in both CB₁R immunoreactivity and mRNA expression in the DLPFC (Brodmann's area 9) and in CB₁R immunoreactivity in Brodmann's area 24 in the DLPFC in SCZ employing immunocytochemistry and *in situ* hybridization.

The discrepancy between ligand binding and immunocytochemistry approaches to CB₁R protein measurement needs to be understood and suggests that changes in the CB₁R system in SCZ are not simple or straightforward. Clearly, more studies will be needed to reach a consensus. [³H] CP 55940 has equal affinity for both CB₁R and CB₂R, however, binding of [³H] CP 55940 to CB₂R is unlikely to have influenced our results as this receptor is expressed at much lower levels in the mammalian brain than the CB₁R (Onaivi *et al*, 2006; Liu *et al*, 2009) and the cortex is among the regions that express the lowest levels of CB₂R in the human brain (Liu *et al*, 2009). Receptor binding studies are more likely to quantitatively reflect actual numbers of CB₁R binding sites in brain and the consistent effect of a broad range of ligands in both post-mortem and *in vivo* studies points to an increase in CB₁R in SCZ. In addition, CB₁R antibodies may not qualitatively or quantitatively stain receptors in all cell types or subcellular compartments (Eggen and Lewis, 2007) and may therefore fail to detect receptors that are detectable using radioligand binding.

In addition to methodological differences, the differences between the studies mentioned above may relate to differences in cohort make-up. Indeed, the wide range of SCZ subtypes that are included in post-mortem studies together with the lack of adequate numbers of cases in each subgroup may dilute the power to find neurochemical changes restricted to a specific diagnostic subtype. Indeed, we observed differences of only borderline significance

when binding in the entire SCZ cohort was compared with controls as a whole. Interestingly, in our cohort the average binding in the schizoaffective cases (depressive and bipolar subtypes pooled) was similar to the controls, whereas the disorganized and residual groups had lower binding compared with controls (Table 1). Furthermore, in an exploratory investigation, which should be treated with caution due to low subject numbers in some diagnostic subgroups, significant differences were found between binding in the residual group with that in the undifferentiated and paranoid groups. These results need to be confirmed in cohorts consisting of larger numbers. If confirmed, however, this evidence raises the possibility that CB₁Rs may show very specific adaptations in the cortical regions of different subtypes of SCZ. Overall, most studies support an increase in cortical CB₁R binding in patients with SCZ, and our study suggests that this increase may be especially evident in people suffering from paranoid SCZ.

Although in the present study the mean onset of the disease did not differ between the paranoid and non-paranoid group, several lines of evidence support that the paranoid subtype of SCZ is associated with later disease onset and better prognosis (McGlashan and Fenton, 1991; Zalewski *et al*, 1998). A connection between an abnormality in the endocannabinoid system and paranoid SCZ has been reported. In 1976, the paranoid psychosis associated with long term cannabis use was compared with the symptoms of paranoid SCZ in 25 psychiatric patients (Thacore and Shukla, 1976). Subjects with acute cannabis intoxication often display a SCZ-like syndrome with hallucinations, altered judgement, false beliefs, and cognitive impairment that are also features of paranoid SCZ (De Marchi *et al*, 2003; D'Souza *et al*, 2009). CSF levels of the endocannabinoid, anandamide, are profoundly and selectively elevated in paranoid-type SCZ as compared with dementia or affective disorder patients, negatively correlate with the psychotic symptoms of the disease and are normalized by treatment with typical but not atypical antipsychotic drugs (Giuffrida *et al*, 2004). Interestingly in two recent PET imaging studies, CB₁R binding (expressed as the distribution volume) in the frontal lobe and the middle and posterior cingulate cortex correlated with positive and inversely correlated with the negative symptoms (Wong *et al*, 2010a) whereas CB₁R uptake in the insula was positively associated with the positive PANSS subscale 'conceptual disorganization' and negatively correlated to psychomotor speed and attention in the amygdala, hippocampus, and putamen (Ceccarini *et al*, 2010). These results taken together with the present study suggest that it may be possible to relate CB₁R abnormalities to the severity of clinical symptoms in SCZ (Wong *et al*, 2010a) or to specific subtypes (present study) and also that subjects with SCZ with abnormal CB₁Rs may have an inherent tendency to a particular symptomatology (ie, paranoia).

As CB₁Rs in the human and monkey DLPFC are thought to be localized in inhibitory GABAergic interneurons of the CCK-expressing subtype (Glass *et al*, 1997; Eggan and Lewis, 2007), it is likely that the majority of CB₁R binding observed in the present study in the paranoid group comes from intrinsic sources. It is also possible that this increase reflects a compensatory response that, if assumed to result

in reduced inhibitory input from GABAergic interneurons (Bodor *et al*, 2005), could be related to tighter control of cognitive function in these patients and also to increased paranoid ideation. Post-mortem examination of other relevant brain regions such as the hippocampus and striatum, which contain high density of CB₁Rs would illuminate this issue.

Implications for Pharmacotherapy

In general, agents that bind to CB₁R and can act as antagonists display antipsychotic properties in animal models (Zuardi *et al*, 2006; Roser *et al*, 2010). The non-psychotomimetic cannabis constituent cannabidiol, which antagonizes the effects of THC (Pertwee 2008), as well as the synthetic CB₁R antagonist rimonabant (SR141716A) have been evaluated in humans as antipsychotics for the treatment of SCZ (Roser *et al*, 2010). Some conflicting results, however, have been obtained in these studies (Roser *et al*, 2010; Leweke *et al*, 2009). In a double-blind clinical trial involving 42 patients with paranoid SCZ, Leweke *et al* (2009) reported that cannabidiol possessed substantial antipsychotic properties with fewer side effects than amisulpride. Following cannabidiol administration, Zuardi *et al*, (1995) also reported an improvement in symptoms in one female with SCZ, however, no improvement was seen in three treatment-resistant males (Zuardi *et al*, 2006). Rimonabant improved psychiatric symptoms in some patients with SCZ (Kelly *et al*, 2011) but was shown to have no effect on psychopathology in comparison to placebo by Meltzer *et al* (2004) and was associated with a relapse to psychosis in another study (Ugur *et al*, 2008). The results of the current study may help to explain the conflicting findings on the effects of cannabinoid antagonists in humans; that is cannabinoid antagonists may be more effective for treating paranoid SCZ than other subtypes of the disease owing in part to the higher levels of CB₁ receptor we observed in this group.

Effects of Continuous and Non-Continuous Variables

Many demographic and peri-mortem factors influence CB₁R mRNA expression and/or binding in addition to any effects of SCZ (Mato and Pazos, 2004; Eggan *et al*, 2008; Ludányi *et al*, 2008; Urigüen *et al*, 2009). For example, CB₁R densities are influenced by aging, post-mortem delay and freezer storage time (our study and Mato and Pazos, 2004). We found negative correlations between CB₁R binding and mRNA with age suggesting that CB₁R density in the DLPFC decreases with age. Similar findings have been reported in the literature in the frontal cortex of normal individuals (Mato and Pazos, 2004). In agreement with Mato and Pazos (2004), we found no significant correlation for CB₁R with post-mortem delay but a positive correlation with freezer storage time that is in contrast to the findings of Mato and Pazos (2004) who reported that CB₁R density is reduced with freezer storage time. However, in our cohort, there was a negative correlation of borderline significance between freezer time and age that might explain the relationship between binding and freezer time. Brain weight was also found to positively correlate with CB₁R binding and whereas there is no convincing explanation in relation to

this finding, it is of interest that Harrison *et al* (2010) reported a positive correlation between human brain weight and the expression of two 'housekeeping genes' and suggested that brain weight should be added to the list of variables to be taken into account in post-mortem studies. Our study shows the retention of a statistically significant elevation in the [³H] CP 55940 binding in paranoid SCZ after all these variables were co-varied for. Along with careful matching of disease and control group, we used correlation and covariate analysis to identify and control for confounding variables, with stringent *post hoc* tests (Bonferroni tests) to account for multiple comparisons. Therefore, the significant difference associated with the diagnostic subtype that we observed here is robust.

Antipsychotic medication may also have had an effect on CB₁R binding and mRNA expression in our disease groups (Sundram *et al*, 2005; Cheng *et al*, 2008; Urigüen *et al*, 2009; Secher *et al*, 2010). Unfortunately, absence of adequate brain tissue from drug naïve persons with SCZ makes it difficult to overcome this problem. However, a number of points should be considered. First, we found no correlation between lifetime antipsychotic drug exposure and CB₁R density or mRNA expression. Second, the existing body of evidence suggests overall that most antipsychotic drugs do not bind the CB₁R *in vitro* (Theisen *et al*, 2007). In rats, antipsychotics do not change the CB₁R binding in the cortex and striatum (the regions with some of the highest density of CB₁R) (Sundram *et al*, 2005; Wiley *et al*, 2008) but may decrease the CB₁R in the brainstem and nucleus accumbens (Sundram *et al*, 2005; Weston-Green *et al*, 2008), or, in the case of risperidone, may increase CB₁R binding in the hypothalamus, hippocampus, and amygdala (Secher *et al*, 2010). Aripiprazole treatment is associated with upregulation in CB₁R mRNA in the rat frontal cortex (Cheng *et al*, 2008), however, Urigüen *et al* (2009) reported no effect of antipsychotic treatment on levels of CB₁R mRNA in human subjects with SCZ compared with controls. It should be noted that in the SCZ group as a whole, 31 of the 37 cases were treated mainly with typical antipsychotics and the remaining six cases received predominately atypical antipsychotic drugs. Within the paranoid group alone, 14 out of 16 cases were treated mainly with typical antipsychotics. Third, Ceccarini *et al* (2010) using the selective high affinity PET radioligand [¹⁸F] MK-9470 showed a significant increase of CB₁R availability in the mesocorticolimbic circuitry, especially in the nucleus accumbens of both antipsychotic treated and untreated (*n* = 5 drug naïve and *n* = 4 after drug washout) SCZ patients compared with controls, supporting the notion that CB₁Rs can be increased regardless of antipsychotic.

Finally, another potential influence on CB₁R binding and mRNA expression may be the effects of cannabis consumption in our SCZ group as chronic cannabinoid exposure has been shown to downregulate CB₁R binding in animal models (Dalton *et al*, 2009) and humans (Villares, 2007). In the current study, however, we observed no effect of cannabis exposure on CB₁R density or mRNA in the DLPFC, in agreement with other studies in the DLPFC (Dean *et al*, 2001; Eggen *et al*, 2010) and superior temporal gyrus (Deng *et al*, 2007).

CONCLUSIONS

Our finding of increased CB₁R binding in paranoid SCZ could reflect a greater involvement of the endocannabinoid system in the DLPFC in this subtype of patients with SCZ as may be suggested by their more marked positive thought disorders and delusional symptoms. An understanding between neurochemical deficits in the endocannabinoid system and SCZ subtypes may ultimately emerge from investigations that combine genetics with brain imaging approaches, biological assays, and neuropsychological techniques in specific SCZ subtypes. The present findings particularly if confirmed by further investigations, would suggest different levels of participation of elements of the endocannabinoid system in the different subtypes of SCZ, arguing for distinct neurochemical correlates of clinical subtypes and raising the possibility of instituting psychopharmacological treatment accordingly.

ACKNOWLEDGEMENTS

This work was supported by the Schizophrenia Research Institute, utilizing infrastructure funding from New South Wales Health, the Macquarie Group Foundation, Neuroscience Research Australia, and the University of New South Wales. Tissues were received from the New South Wales Tissue Resource Centre at the University of Sydney, which is supported by the National Health and Medical Research Council of Australia, Schizophrenia Research Institute, National Institute of Alcohol Abuse and Alcoholism (NIH (NIAAA) R24AA012725). We thank Nina Sundqvist of the New South Wales Tissue Resource Centre for her assistance.

DISCLOSURE

The authors declare no conflict of interest.

REFERENCES

- Ameri A (1999). The effects of cannabinoids on the brain. *Prog Neurobiol* 58: 315–348.
- Bodor AL, Katona I, Nyiri G, Mackie K, Ledent C, Hajos N *et al* (2005). Endocannabinoid signaling in rat somatosensory cortex: laminar differences and involvement of specific interneuron types. *J Neurosci* 25: 6845–6856.
- Ceccarini J, De Hert M, van Winkel R, Koethe D, Bormans G, Leweke M *et al* (2010). *In vivo* PET imaging of cerebral type 1 cannabinoid receptor availability in patients with schizophrenia. *Schizophrenia Res* 117: 170.
- Chavarria-Siles I, Contreras-Rojas J, Hare E, Walss-Bass C, Quezada P, Dassori A *et al* (2008). Cannabinoid receptor 1 gene (CNR1) and susceptibility to a quantitative phenotype for hebephrenic schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* 147: 279–284.
- Cheng MC, Liao DL, Hsiung CA, Chen CY, Liao YC, Chen CH (2008). Chronic treatment with aripiprazole induces differential gene expression in the rat frontal cortex. *Int J Neuropsychopharmacol* 11: 207–216.
- Cohen J (1988). *Statistical Power Analysis for the Behavioral Sciences*, 2nd edn. Lawrence Erlbaum Associates: Hillsdale, New Jersey.
- Dalton VS, Wang H, Zavitsanou K (2009). HU210-induced downregulation in cannabinoid CB1 receptor binding strongly

- correlates with body weight loss in the adult rat. *Neurochem Res* 34: 1343–1353.
- De Marchi N, De Petrocellis L, Orlando P, Daniele F, Fezza F, Di Marzo V (2003). Endocannabinoid signalling in the blood of patients with schizophrenia. *Lipids Health Dis* 2: 5.
- Dean B, Sundram S, Bradbury R, Scarr E, Copolov D (2001). Studies on [³H]CP-55940 binding in the human central nervous system: regional specific changes in density of cannabinoid-1 receptors associated with schizophrenia and cannabis use. *Neuroscience* 103: 9–15.
- Deng C, Han M, Huang X-F (2007). No changes in densities of cannabinoid receptors in the superior temporal gyrus in schizophrenia. *Neurosci Bull* 23: 341–347.
- D'Souza DC, Sewell RA, Ranganathan M (2009). Cannabis and psychosis/schizophrenia: human studies. *Eur Arch Psychiatry Clin Neurosci* 259: 413–431.
- Eggan SM, Hashimoto T, Lewis DA (2008). Reduced cortical cannabinoid 1 receptor messenger RNA and protein expression in schizophrenia. *Arch Gen Psychiatry* 65: 772–784.
- Eggan SM, Lewis DA (2007). Immunocytochemical distribution of the cannabinoid CB1 receptor in the primate neocortex: a regional and laminar analysis. *Cereb Cortex* 17: 175–191.
- Eggan SM, Stoyak SR, Verrico CD, Lewis DA (2010). Cannabinoid CB1 receptor immunoreactivity in the prefrontal cortex: Comparison of schizophrenia and major depressive disorder. *Neuropsychopharmacology* 35: 2060–2071.
- Giuffrida A, Leweke FM, Gerth CW, Schreiber D, Koethe D, Faulhaber J *et al* (2004). Cerebrospinal anandamide levels are elevated in acute schizophrenia and are inversely correlated with psychotic symptoms. *Neuropsychopharmacology* 29: 2108–2114.
- Glass M, Dragunow M, Faull RL (1997). Cannabinoid receptors in the human brain: a detailed anatomical and quantitative autoradiographic study in the fetal, neonatal and adult human brain. *Neuroscience* 77: 299–318.
- Harper CG, Kril JJ, Daly JM (1988). Brain shrinkage in alcoholics is not caused by changes in hydration: a pathological study. *J Neurol Neurosurg Psychiatry* 51: 124–127.
- Harrison PJ, Laatikainen LM, Tunbridge EM, Eastwood SL (2010). Human brain weight is correlated with expression of the 'housekeeping genes' beta-2-microglobulin (beta2M) and TATA-binding protein (TBP). *Neuropathol Appl Neurobiol* 36: 498–504.
- Hill C, Keks NA, Roberts S, Opeskin K, Dean B, MacKinnon A *et al* (1996). Problem of diagnosis in postmortem brain studies of schizophrenia. *Am J Psychiatry* 153: 533–537.
- Keks NA, Hill C, Roberts S, Dean B, Opeskin K, MacKinnon A *et al* (1997). Diagnostic instrument for brain studies (DIBS). *Schizophrenia Res* 24: 34.
- Kelly DL, Gorelick DA, Conley RR, Boggs DL, Linthicum J, Liu F *et al* (2011). Effects of the cannabinoid-1 receptor antagonist rimonabant on psychiatric symptoms in overweight people with schizophrenia: a randomized, double-blind, pilot study. *J Clin Psychopharmacol* 31: 86–91.
- Koethe D, Llenos IC, Dulay JR, Hoyer C, Torrey EF, Leweke FM *et al* (2007). Expression of CB1 cannabinoid receptor in the anterior cingulate cortex in schizophrenia, bipolar disorder, and major depression. *J Neural Transm* 114: 1055–1063.
- Kozlovsky N, Weickert CS, Tomaskovic-Crook E, Kleinman JE, Belmaker RH, Agam G (2004). Reduced GSK-3beta mRNA levels in postmortem dorsolateral prefrontal cortex of schizophrenic patients. *J Neural Transm* 111: 1583–1592.
- Leweke FM, Koethe D, Pahlisch F, Schreiber D, Gerth CW, Nolden BM *et al* (2009). Antipsychotic effects of cannabidiol. *Eur Psychiatry* 24(Suppl 1): 207.
- Liu Q-R, Pan C-H, Hishimoto A, Li C-Y, Xi Z-X, Llorente-Berzal A *et al* (2009). Species differences in cannabinoid receptor 2 (CNR2 gene): identification of novel human and rodent CB2 isoforms, differential tissue expression and regulation by cannabinoid receptor ligands. *Genes Brain Behav* 8: 519–530.
- Ludányi A, Eross L, Cziráj S, Vajda J, Halász P, Watanabe M *et al* (2008). Downregulation of the CB1 cannabinoid receptor and related molecular elements of the endocannabinoid system in epileptic human hippocampus. *J Neurosci* 28: 2976–2990.
- Mato S, Pazos A (2004). Influence of age, postmortem delay and freezing storage period on cannabinoid receptor density and functionality in human brain. *Neuropharmacology* 46: 716–726.
- McGlashan TH, Fenton WS (1991). Classical subtypes for schizophrenia: literature review for DSM-IV. *Schizophr Bull* 17: 609–632.
- Meltzer HY, Arvanitis L, Bauer D, Rein W (2004). Placebo-controlled evaluation of four novel compounds for the treatment of schizophrenia and schizoaffective disorder. *Am J Psychiatry* 161: 975–984.
- Newell KA, Deng C, Huang X-F (2006). Increased cannabinoid receptor density in the posterior cingulate cortex in schizophrenia. *Exp Brain Res* 172: 556–560.
- Pertwee RG (2008). The diverse CB₁ and CB₂ receptor pharmacology of three plant cannabinoids: (delta)9-tetrahydrocannabinol, cannabidiol and (delta)9-tetrahydrocannabivarin. *Br J Pharmacol* 153: 199–215.
- Onaivi ES, Ishiguro H, Gong J-P, Patel S, Perchuk A, Meozzi PA *et al* (2006). Discovery of the presence and functional expression of cannabinoid CB2 receptors in brain. *Ann N Y Acad Sci* 1074: 514–536.
- Rajkowska G, Goldman-Rakic PS (1995). Cytoarchitectonic definition of prefrontal areas in the normal human cortex: I. Remapping of areas 9 and 46 using quantitative criteria. *Cereb Cortex* 5: 307–322.
- Roberts S, Hill C, Dean B, Keks NA, Opeskin K, Copolov D (1998). Confirmation of the diagnosis of schizophrenia after death using DSM-IV: a Victorian experience. *Aust NZ J Psych* 32: 73–76.
- Roser P, Vollenweider FX, Kawohl W (2010). Potential antipsychotic properties of central cannabinoid (CB1) receptor antagonists. *World J Biol Psychiatry* 11(2 Part 2): 208–219.
- Secher A, Husum H, Holst B, Egerod KL, Møllerup E (2010). Risperidone treatment increases CB1 receptor binding in rat brain. *Neuroendocrinology* 91: 155–168.
- Sundram S, Copolov D, Dean B (2005). Clozapine decreases [³H] CP 55940 binding to the cannabinoid 1 receptor in the rat nucleus accumbens. *Naunyn Schmiedebergs Arch Pharmacol* 371: 428–433.
- Sundqvist N, Garrick T, Bishop I, Harper C (2008). Reliability of post-mortem psychiatric diagnosis for neuroscience research. *Aust N Z J Psychiatry* 42: 221–227.
- Thacore VR, Shukla SR (1976). Cannabis psychosis and paranoid schizophrenia. *Arch Gen Psychiatry* 33: 383–386.
- Theisen FM, Haberhausen M, Firnges MA, Gregory P, Reinders JH, Remschmidt H *et al* (2007). No evidence for binding of clozapine, olanzapine and/or haloperidol to selected receptors involved in body weight regulation. *Pharmacogenomics J* 7: 275–281.
- Ugur T, Bartels M, Kis B, Scherbaum N (2008). Psychosis following anti-obesity treatment with rimonabant. *Obes Facts* 1: 103–105.
- Urigüen L, García-Fuster MJ, Callado LF, Morentin B, La Harpe R, Casadó V *et al* (2009). Immunodensity and mRNA expression of A2A adenosine, D2 dopamine, and CB1 cannabinoid receptors in postmortem frontal cortex of subjects with schizophrenia: effect of antipsychotic treatment. *Psychopharmacology (Berl)* 206: 313–324.
- Villares J (2007). Chronic use of marijuana decreases cannabinoid receptor binding and mRNA expression in the human brain. *Neuroscience* 145: 323–334.
- Weickert CS, Sheedy D, Rothmond DA, Dedova I, Fung S, Garrick T *et al* (2010). Selection of reference gene expression in a schizophrenia brain cohort. *Aust N Z J Psychiatry* 44: 59–70.

- Weston-Green K, Huang XF, Han M, Deng C (2008). The effects of antipsychotics on the density of cannabinoid receptors in the dorsal vagal complex of rats: implications for olanzapine-induced weight gain. *Int J Neuropsychopharmacol* 11: 827–835.
- Wiley JL, Kendler SH, Burston JJ, Howard DR, Selley DE, Sim-Selley LJ (2008). Antipsychotic-induced alterations in CB₁ receptor-mediated G-protein signaling and *in vivo* pharmacology in rats. *Neuropharmacology* 55: 1183–1190.
- Wong DF, Kuwabara H, Horti AG, Raymond V, Brasic J, Guevara M *et al* (2010a). Quantification of cerebral cannabinoid receptors subtype 1 (CB₁) in healthy subjects and schizophrenia by the novel PET radioligand [¹¹C]OMAR. *Neuroimage* 52: 1505–1513.
- Wong J, Hyde TM, Cassano HL, Deep-Soboslay A, Kleinman JE, Weickert CS (2010b). Promoter specific alterations of brain-derived neurotrophic factor mRNA in schizophrenia. *Neuroscience* 169: 1071–1084.
- Yang Y, Fung SJ, Rothwell A, Tianmei S, Weickert CS (2010). Increased interstitial white matter neuron density in the dorsolateral prefrontal cortex of people with schizophrenia. *Biol Psychiatry* 69: 63–70.
- Zalewski C, Johnson-Selfridge MT, Ohriner S, Zarrella K, Seltzer JC (1998). A review of neuropsychological differences between paranoid and nonparanoid schizophrenia patients. *Schizophr Bull* 24: 127–145.
- Zavitsanou K, Garrick T, Huang X-F (2004). Selective antagonist [³H]SR141716A binding to cannabinoid CB₁ receptors is increased in the anterior cingulate cortex in schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 28: 355–360.
- Zuardi AW, Crippa JA, Hallak JE, Moreira FA, Guimaraes FS (2006). Cannabidiol, a Cannabis sativa constituent, as an antipsychotic drug. *Braz J Med Biol Res* 39: 421–429.
- Zuardi AW, Morais SL, Guimaraes FS, Mechoulam R (1995). Antipsychotic effect of cannabidiol. *J Clin Psychiatry* 56: 485–486.

Supplementary Information accompanies the paper on the Neuropsychopharmacology website (<http://www.nature.com/npp>)