



Published in final edited form as:

*J Clin Psychiatry*. 2010 February ; 71(2): 201–206. doi:10.4088/JCP.08m04663yel.

## Neurocognitive impairment and psychosis in bipolar disorder during early remission from an acute episode of mood disturbance

Boaz Levy, Ph.D. and Roger D. Weiss, MD.

McLean Hospital/Harvard Medical School, Belmont, MA USA

### Abstract

**Objective**—Recent studies have reported greater neurocognitive impairment in euthymic bipolar disorder patients with a history of psychosis relative to patients without such a history. To further explore the relation between psychosis and cognitive dysfunction in bipolar disorder, the current study examined the cognitive functioning of patients during early remission from a discrete episode of mood disturbance. The study aimed to determine whether the presence of psychosis during inpatient hospitalization was associated with greater cognitive impairment at the time of hospital discharge.

**Method**—Fifty-nine inpatients who met DSM-IV criteria for bipolar disorder (24 admitted with psychosis, 35 admitted without psychosis), ages 18–59, completed a neuropsychological battery and mood measures 24–48 hours before discharge. The cognitive battery included standardized tests of IQ, attention and working memory, visual memory, verbal memory and executive functioning.

**Results**—A multivariate analysis of variance detected group differences on measures of verbal memory ( $p < 0.001$ ) and executive functioning ( $p < 0.003$ ), using mood measures and previous number of psychiatric admissions as covariates. Post-hoc analysis of between-subjects effects revealed significantly poorer performance on the California Verbal Learning Test–Second Edition, Logical Memory test from Wechsler Memory Scale–Revised, Stroop–Word/Color Interference test and the Wisconsin Card Sorting Test for patients who were admitted to the hospital with psychosis. These results remained significant after matching the groups for past psychosis, with the exception of the Logical Memory test.

**Conclusions**—The results of this study indicate that patients with bipolar disorder who were admitted to the hospital due to psychosis exhibited significantly more severe cognitive impairment at the time of discharge than patients admitted for an acute mood disturbance without psychosis. These findings may be important for improving discharge planning and the development of more effective outpatient services.

### Keywords

bipolar disorder; cognitive impairment; psychosis; inpatients

## 1. Introduction

Patients with bipolar disorder often suffer from debilitating cognitive impairment that persists beyond the remission of mood symptoms (1). Studies show that cognitive deficits

are more likely to be present in patients who suffer from a more severe course of illness, as indicated by longer durations of mood disturbance, younger age of onset, and a higher number of inpatient admissions (2, 3). More recently, researchers have expanded this line of investigation to explore the impact of other clinical variables that are associated with poor illness outcome on the cognitive functioning of patients. In this respect, psychosis has come into focus, as it often constitutes an important clinical marker for a more severe course of illness in bipolar disorder (4, 5).

The few studies that have investigated the impact of psychosis on the cognitive functioning of patients with bipolar disorder have shown mixed results. Several authors found no differences between patients with and without a history of psychotic episodes (6, 7, 8, 9,10); in contrast, Albus et al. (11) reported an overall more compromised performance on cognitive measures in first-break patients who presented with psychotic symptoms. Similarly, Zubieta et al. (12) found that patients with a history of psychosis obtained lower scores on measures of attention, verbal memory and executive functioning than healthy controls. Consistent with these findings, Martinez-Aran et al. (2, 13, 14) observed more severe deficits on related measures in bipolar disorder patients who presented with previous psychotic episodes relative to patients without a history of psychosis. Specific to executive functioning, Bora et al. (15) concluded that deficits in cognitive flexibility may be the trait marker of psychotic features among patients with bipolar disorder. In the nonverbal domain of cognitive functioning, Glahn et al. (16) reported significantly weaker performance on measures of spatial working memory in patients with a history of psychosis compared with patients without such a history. Although the findings to date appear inconclusive, given the inconsistency across studies, the multiple, independent reports associating psychosis with a more severe course of illness and greater cognitive impairment warrant additional studies in this area.

Further investigation may benefit from expanding the direct implications of studies to patient care. Most of the studies to date have focused primarily on the long-term impact of previous psychotic episodes on the cognitive functioning of euthymic patients. However, in a clinical view, it may be equally important to explore the impact of a discrete psychotic episode on the cognitive functioning of patients during the early phases of recovery from acute disturbance. Understanding how an acute psychotic episode affects the cognitive functioning of patients during early remission may be critical for improving patient care following discharge from inpatient admission. The important clinical question here is whether patients with bipolar disorder who were admitted to the hospital due to psychosis suffer more debilitating cognitive impairment at the time of discharge than patients who required an inpatient admission for an acute mood disturbance without psychosis. The current study examined this question. We hypothesized that patients who were admitted to the hospital with psychosis would perform more poorly upon discharge on neuropsychological tests than would patients who did not experience psychotic symptoms over the course of hospitalization.

## 2. Method

### 2.1 Subjects

Fifty-nine inpatients at McLean Hospital who met DSM-IV diagnostic criteria for bipolar I disorder completed the neuropsychological battery. In this sample, 41 patients were admitted with manic symptoms, 8 were admitted for depression, and 10 patients were hospitalized with a mixed mood presentation. The sample consisted of 33 men and 26 women, of whom 10 identified an affiliation with an ethnic minority group. The age range was 18 to 59. Participants were classified into two groups based on the presence (n=24) or absence (n=35) of psychotic symptoms during the current admission. In the former group, 7 participants

experienced their first psychotic episode in the context of the current admission, and in the latter group 11 patients reported previous psychotic episodes. Consistent with previous studies (14), psychosis was defined here by the presence of either delusions or hallucinations. Patients did not present psychotic symptoms at the time of testing. The presence of psychosis was determined through cross-validating clinical information among three sources: diagnostic interviews, medical record notes, and direct communications with the treatment team. Agreement in reporting psychotic symptoms (delusions or hallucinations) prior to or during admission determined the presence of a psychotic condition. The absence of psychosis at the time of discharge was determined by failure to detect psychotic symptoms by the research clinician who interviewed the patients and the treatment team.

## 2.2 Inclusion/Exclusion Criteria

All of the participants in this study were adult (age  $\geq 18$ ) inpatients who met DSM-IV diagnostic criteria for bipolar I disorder. To reduce the impact of severe mood symptoms on participants' test performance, inclusion criteria also required a Beck Depression Inventory – Second Edition (BDI-II) score  $< 15$  (17), a Beck Hopelessness Scale (BHS) score  $< 10$  (18), and a Young Mania Rating Scale (YMRS) score  $< 15$  (19) at the time of testing. Exclusion criteria included electroconvulsive therapy during the 12 months prior to admission, a history of neurological illness or injury, and lifetime diagnoses of substance abuse or dependence. This information was obtained through structured diagnostic interviews and a review of medical records.

## 2.3 Diagnosis and Procedure

After the treatment team on the unit determined that a potential participant had reached sufficient stability for discharge, the patient was approached by a trained clinician to sign an informed consent form for the study. The clinician administered the Structured Clinical Interview for DSM-IV (20) – Part I to confirm a bipolar I diagnosis. The diagnostic procedure also integrated information from family members and outpatient treatment providers that was available in the medical record or through verbal communications with the treatment team. After the diagnostic procedure was completed, a trained examiner administered the neuropsychological battery and mood measures. This procedure occurred 24–48 hours prior to discharge.

## 2.4 The Neuropsychological Battery

In this study we employed a neuropsychological test battery with well-documented norms and satisfactory estimates of reliability and validity. The battery was designed to assess IQ, attention and working memory, visual memory, verbal memory and executive functioning. IQ estimates were obtained with the Wechsler Abbreviated Scale of Intelligence (WASI; 21). Measures of attention included the Digit Span subtest from the Wechsler Adult Intelligence Scale – Third Edition (22), Trails Making Test parts A and B (23), and the Letter and Symbol Cancellation Task (24). We assessed verbal memory with the Logical Memory subtest from Wechsler's Memory Scale-Revised (25), and the California Verbal Learning Test II – Short Form (26). To assess perceptual organization and visual memory, we administered the Rey Complex Figure test (27). The battery also consisted of several measures of executive functioning, including the Stroop Color-Word Interference Test (28), the Controlled Oral Word Association Test (COWAT) - FAS letters format, and Animal Naming Task (29) - and the Wisconsin Card Sorting Test – 64 Card Version (30). Administration of the battery required approximately 2 hours.

## 2.5 Statistical Analysis

Preliminary analyses of demographic and clinical data applied the Pearson's chi-square and t-tests. Because the neuropsychological test battery included multiple measures in several cognitive domains, group differences in test scores were determined by a multivariate analysis of variance. To control for residual mood symptoms and illness severity, analyses used the mood measures administered at the time of testing (i.e., BDI-II and YMRS scores) and previous number of psychiatric hospitalizations as covariates. The same procedure was repeated for each cognitive domain separately: attention and working memory, visual memory, verbal memory, executive functioning and IQ. ANOVA of post-hoc between-subjects effects in test performance on individual measures applied the same covariates. To control for group differences in past psychotic episodes, subsequent analysis repeated previously significant comparisons after excluding patients without a history of psychosis from the group that presented to the current admission without psychosis. This analysis compared patients who were admitted to the hospital with (n=24) and without (n=11) psychosis, where the latter group consisted only of patients who reported a history of psychosis. Analyses applied standardized scaled scores for the cognitive measures (mean = 50, SD = 10; California Verbal Learning Test II immediate and delayed recall and recognition scores had a mean of 0 and SD of 1), with lower values representing poorer performance.

## 3. Results

### 3.1 Clinical and Demographic Variables

Table 1 presents means, standard deviations, and test statistics of the clinical and demographic variables. As Table 1 indicates, no group differences emerged in clinical variables with respect to residual mood symptoms, previous number of psychiatric admissions, or age at first hospitalization. Further analysis also failed to detect differences in the number of medications patients were prescribed on the day of testing, or diagnostic subtype upon admission. Group comparisons of demographic variables also did not reveal significant differences in age, gender, years of formal education, marital status, or ethnic minority status.

### 3.2 Cognitive Measures

**Attention, Working Memory and IQ**—The Multivariate ANOVA procedure revealed no group differences on measures of attention and working memory (Wilk's Lambda,  $F_{(7,48)}=0.28$ ,  $p<0.95$ ). Consistent with this result, none of the ANCOVA tests for differences on the individual measures approached significance. The multivariate and ANCOVA procedures also did not detect significant group differences on the WASI IQ measures (Wilk's Lambda;  $F_{(7,48)}=0.71$ ,  $p<0.65$ ), which included Full Scale IQ, Verbal IQ, Performance IQ, Block Design, Matrix Reasoning, Vocabulary, and Similarities.

**Memory**—Analysis detected highly significant group differences in measures of verbal memory (Wilk's Lambda,  $F_{(7,48)}=9.11$ ,  $p<0.001$ ). As Table 2 reveals, the ANCOVA procedure indicates that the bipolar disorder group without psychosis performed significantly better than the group with psychosis on immediate and delayed recall of passages (Logical Memory) and on the acquisition, immediate and delayed recall of a word-list (CVLT-II). When patients without a history of psychosis were excluded from analysis, results remained significant only on immediate and delayed recall of the word-list (see Table 3). No group differences emerged in visual memory (Wilks' Lambda,  $F_{(7,48)}=0.31$ ,  $p<0.87$ ), as measured by the Rey Complex Figure test.

**Executive Functioning**—The multivariate procedure yielded highly significant group differences in measures of executive functioning (Wilk's Lambda,  $F_{(7,48)} = 3.67$ ,  $p < 0.003$ ). The ANCOVA procedure revealed poorer performance for the bipolar disorder group with psychosis on the Stroop and the Wisconsin Card Sorting tests (see Table 2). These results remained significant after the exclusion of patients without a history of psychosis (see Table 3).

#### 4. Discussion

The results of the current study indicated more severe cognitive deficits in patients with bipolar disorder who were admitted to the hospital with psychotic symptoms than in patients who were hospitalized for an acute mood disturbance without psychosis. Analyses revealed that the presence of psychosis on admission was associated with greater impairment in verbal memory at the time of discharge, as indicated by more compromised performance on a passage memory test and a list-learning task. Patients who presented to the hospital with psychosis also exhibited more severe impairment in executive functioning, especially on tasks that measure cognitive flexibility (i.e. ability to adjust a course of action in response to corrective feedback) and the ability to inhibit dominant or automatic reactions in favor of responses relevant to task demands. Group differences in executive functioning and verbal memory remained significant after matching for past psychosis. However, in the domain of verbal memory, this was the case only on measures of the list-learning task, which tends to be more sensitive to executive dysfunction.

The results of this study are consistent with several previous reports that have emphasized greater impairment in verbal memory and executive functioning among euthymic patients with a history of psychosis (2, 12, 13, 14, 15). In fact, it is possible that previous psychotic episodes contributed to the group differences in cognitive functioning observed in the current sample. At the same time, it is notable that the differences remained highly significant after controlling for certain variables that would be related to the overall severity of illness (i.e. previous number of inpatient admissions, and the presence of residual mood symptoms). This result may advance speculation about the degree to which the observed group differences relate more directly to the nature of the current episode. It is thus possible that early remission from psychosis may be accompanied by greater cognitive impairment relative to early remission from an acute mood disturbance without psychosis.

Understanding the potential impact of a discrete psychotic episode on cognitive functioning in the early stage of remission may be important for clinical care. In the hospital, the presence and effects of cognitive impairment may remain largely inconspicuous due to the minimal functional demands that this environment poses on the brain. Upon discharge, however, patients may face the challenge of functioning in an unsupervised environment at a time when their cognitive abilities are most compromised. Previous reports highlight the relationship between cognitive impairment and psychosocial disability in bipolar disorder (2). In particular, executive and consequent memory impairments can compromise patients' ability to negotiate the demands of everyday life in unprotected settings (31, 32, 33). After discharge, cognitively impaired patients may encounter major difficulties in keeping appointments, managing medication, sustaining proper nutritional and fluid intake, and maintaining hygiene. These difficulties may increase stress, trigger affective instability, and ultimately result in rapid decompensation and readmission. Longitudinal evidence suggests that about 60% of bipolar disorder patients experience poor post-hospital adjustment (6). This may be partly due to the untoward effects of cognitive impairment, exacerbated by a recent psychotic episode. The results of the current study warrant careful discharge planning with regard to the assessment of cognitive functioning for bipolar patients who were admitted for psychosis.

Several limitations of the current study deserve mention. The study reports a correlation between psychosis and cognitive functioning, which may limit conclusions about the degree to which acute psychosis contributes to poor cognitive outcome upon discharge. A follow-up assessment several months after discharge would allow for the examination of the relationship between psychosis, cognitive recovery, and functional outcome. In the context of the naturalistic environment of the hospital setting, the current study also did not control for the cognitive effects of medication, although the groups did not differ in terms of number of prescribed medications on the day of testing. The exclusion criterion of lifetime diagnoses of substance abuse or dependence was quite strict and limited the generalizability of findings. In terms of cognitive measurement, the current study may have failed to detect group differences in visual memory, as the Rey-Osterrieth is particularly sensitive to the adverse effects of executive dysfunction, which were present in both groups. Finally, the current sample did not allow for the analysis of ethnicity, which again may limit the generalization of results.

Despite these limitations, the current study indicates a potential link between the presence of psychosis and cognitive impairment during the early phase of remission from acute disturbance. These results may inform clinical care and may highlight the need to increase the accommodations and supports that bipolar disorder patients receive after discharge from the hospital.

## Acknowledgments

The authors are grateful for the technical assistance of the treatment teams on AB2 unit at McLean Hospital. In particular, we are thankful for the invaluable clinical support of Matt Bernstein, MD, Liz Liebson, MD, Katherine Healey, MSW, Amy Carlson, MSW, Amy Burch, MSW and Kevin Aubrey, MSW.

### Funding source

The study was supported by the Kaplen Award on Depression (granted by the Harvard Medical School, Department of Psychiatry), NARSAD Young Investigator Award, and grants R01 DA15968 and K24 DA022288 from the National Institute on Drug Abuse.

## References

1. Mur M, Portella MJ, Martinez-Aran A, Pifarre J, Vieta E. Persistent neuropsychological deficit in euthymic bipolar patients: Executive function as a core deficit. *J Clin Psychiatry*. 2007; 68:1078–1086. [PubMed: 17685745]
2. Martinez-Aran A, Vieta E, Colom F, Torrent C, et al. Cognitive impairment in euthymic bipolar patients: Implications for clinical and functional outcome. *Bipolar Disord*. 2004; 6:224–232. [PubMed: 15117401]
3. Thompson JM, Gallagher P, Hughes JH, Watson S, Gray JM, Ferrier IN, Young AH. Neurocognitive impairment in euthymic patients with bipolar affective disorder. *Br J Psychiatry*. 2005; 186:32–40. [PubMed: 15630121]
4. Milkowitz DJ. Longitudinal outcome and medication non-compliance among manic patients with and without mood-incongruent psychotic features. *J Nerv Ment Dis*. 1992; 180:703–711. [PubMed: 1359003]
5. Tohen M, Hennen J, Zarate CM, et al. Two-year syndromal and functional recovery in 219 cases of first-episode major affective disorder with psychotic features. *Am J Psychiatry*. 2000; 157:220–228. [PubMed: 10671390]
6. Goldberg JF, Harrow M, Grossman LS. Course and outcome in bipolar affective disorder: a longitudinal follow up study. *Am J Psychiatry*. 1995; 152:379–384. [PubMed: 7864263]
7. Selva G, Salazar J, Balanza-Martinez V, et al. Bipolar I patients with and without a history of psychotic symptoms: do they differ in their cognitive functioning? *J Psychiatr Res*. 2007; 41:265–272. [PubMed: 16762369]

8. Donaldson S, Goldstein LH, Landau S, et al. The Maudsley Bipolar Disorder Project: the effects of medication, family history, and duration of illness on IQ and memory in bipolar disorder. *J Clin Psychiatry*. 2003; 64:86–93. [PubMed: 12590629]
9. Frangou S, Donaldson S, Hadjulic M, et al. The Maudsley Bipolar Disorder Project: executive dysfunction in bipolar disorder I and its clinical correlates. *Bio Psychiatry*. 2005; 58:859–569. [PubMed: 16039620]
10. Altshuler LL, Ventura J, Van Gorp WG, Green MF, Theberge DC, Mintz J. Neurocognitive function in clinically stable men with bipolar I disorder or schizophrenia and normal control subjects. *Biol Psychiatry*. 2004; 56:560–569. [PubMed: 15476685]
11. Albus M, Hubbman W, Wahlheim C, et al. Contrasts in neuropsychological test profiles between patients with first first-episode schizophrenia and first episode affective disorder. *Acta Psychiatr Scand*. 1996; 94:87–93. [PubMed: 8883568]
12. Zubieta JK, Huguelet P, O'Neil RL, Giordani BJ. Cognitive function in euthymic bipolar I disorder. *Psychiatry Res*. 2001; 102:9–20. [PubMed: 11368835]
13. Martinez-Aran A, Vieta E, Reinares M, et al. Cognitive function across manic or hypomanic, depressed, and euthymic states in bipolar disorder. *Am J Psychiatry*. 2004; 16:262–270. [PubMed: 14754775]
14. Martinez-Aran A, Torrent C, Tabares-Seisdedos R, et al. Neurocognitive impairment in bipolar patients with and without a history of psychosis. *J clin Psychiatry*. 2008:233–239. [PubMed: 18232725]
15. Bora E, Vahip S, Akdeniz F. The effects of previous psychotic mood episodes on cognitive impairment in euthymic bipolar patients. *Bipolar Disord*. 2007; 9:468–477. [PubMed: 17680917]
16. Glahn DC, Bearden CE, Cakir S, et al. Differential working memory impairment in bipolar disorder and schizophrenia: effects of lifetime history of psychosis. *Bipolar Disord*. 2006; 8:117–123. [PubMed: 16542181]
17. Dozois DJ, Dobson KS, Ahnberg LJ. A psychometric evaluation of the Beck depression Inventory – II. *Psychol Assess*. 1998; 10:83–89.
18. Beck AT, Weissman A, Lester D, Trexler L. The measurement of pessimism: the hopelessness scale. *J Consult Clin Psychol*. 1974; 42:861–865. [PubMed: 4436473]
19. Young RC, Biggs JT, Ziegler VE, Meyer DA. A rating scale for mania: Reliability, validity and sensitivity. *Br J Psychiatry*. 1978; 133:429–435. [PubMed: 728692]
20. First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JBW. Biometric Research. New York: New York State Psychiatric Institute; 1994. Structured clinical interview for Axis I DSM-IV disorders-patients edition (SCID-I/P), version 2.0.
21. Wechsler, D. Wechsler Abbreviated Scale of Intelligence. San Antonio, TX: The Psychological Cooperation; 1999.
22. Wechsler, D. Wechsler Adult Intelligence Scale – Third Edition. San Antonio, TX: The Psychological Cooperation; 1997.
23. Spreen, O.; Strauss, E. A compendium of neuropsychological tests: Administration, norms, and commentary. 2nd. New York: Oxford University Press; 1998.
24. Lockwood KA, Marcotte AC, Stern C. Differentiation of attention deficit/hyperactivity disorder subtypes: application of neuropsychological model of attention. *J Clin Exp Neuropsychol*. 2001; 23:317–330. [PubMed: 11404810]
25. Wechsler, D. Wechsler Memory Scale – Revised. San Antonio, TX: The Psychological Cooperation; 1987.
26. Delis, D.; Kramer, J.; Kaplan, E.; Ober, B. The Psychological Cooperation. Australia: Harcourt Assessment, Inc; 1999. California Verbal Learning Test—Second Edition (CVLT-II) Manual.
27. Meyers, JE.; Meyers, KR. Rey Complex Figure Test and Recognition Trial. Florida: Psychological Assessment Resources, Inc; 1995.
28. Golden, C.; Freshwater, SM. Stroop Color Word Test. Illinois: Stoelting Co; 2002.
29. Tombaugh TH, Kozak J, Rees L. Normative data stratified by age and education for two measures of verbal fluency: FAS and animal naming. *Arch Clin Neuropsychol*. 1999; 14:167–177. [PubMed: 14590600]

30. Heaton, KH.; Chelune, GJ.; Talley, JL.; Kay, GG.; Curtis, G. Psychological Assessment resources. Texas: Harcourt Assessment, Inc; 2003. Wisconsin Card Sorting Test: Computer Version 4 (WCST: CV4™) Research Edition.
31. Martinez-Aran A, Penades R, Vieta E, et al. Executive function in patients with remitted bipolar disorder and schizophrenia and its relationship with functional outcome. *Psychother Psychosom.* 2002; 71:39–46. [PubMed: 11740167]
32. Goswami U, Sharma A, Khastigir U, Ferrier IN, Young AH, Gallagher P, Thompson JM, Moore PB. Neuropsychological dysfunction, soft neurological signs and social disability in euthymic patients with bipolar disorder. *Br J Psychiatry.* 2006; 188:366–373. [PubMed: 16582064]
33. Laes JR, Sponheim SR. Does cognition predict community function only in schizophrenia?: A study of schizophrenia patients, bipolar affective disorder patients, and community control subjects. *Schizophr Res.* 2006; 84:121–131. [PubMed: 16443348]



**Table 1**  
Means, Standard Deviations, and Group Comparisons of Demographic and Mood Measures

Measures	Bipolar Patients with Psychosis		Bipolar Patients without Psychosis		t	P Value
	Mean	SD	Mean	SD		
Demographic						
Education	14.2	1.78	14.3	2.44	-0.05	0.96
Age	37.2	12.5	37.5	13.4	-0.08	0.93
Onset	24.9	9.6	28.7	11.8	1.20	0.20
Admissions	5.50	5.31	5.82	5.92	-0.19	0.85
YMRS	7.25	4.60	6.63	3.60	0.57	0.57
BDI – II	8.33	3.29	7.91	3.58	0.45	0.65
BHS	3.88	4.06	4.50	2.99	0.67	0.50

Note: BDI=Beck Depression Inventory, BHS=Beck Hopelessness Scale, YMRS=Young Mania Rating Scale, Admissions=previous number of psychiatric admissions, Onset=age of first psychiatric admission for mood disturbance.

**Table 2**  
Group Comparison of Scaled Scores of Verbal Memory and Executive Functioning Measures

Measures	Test of Between-Subjects Effects				Means and Standard deviations of measures				
	MD	MS	df	F	Sig.	Mean	SD	Mean	SD
Verbal Memory									
LM-I	7.5	249.5	4.0	2.46*	0.05	40.5	10.1	48.0	10.3
LM-D	8.2	154.5	4.0	3.49*	0.05	38.9	6.80	47.1	11.3
CVLT-AQ	11.6	506.0	4.0	4.51**	0.003	34.1	12.5	45.7	8.70
CVLT-SD	1.77	11.4	4.0	13.1***	0.0001	-2.27	0.67	-0.5	1.05
CVLT-LD	1.15	4.97	4.0	4.48**	0.003	-2.12	0.85	-0.97	1.13
CVLT-REC	0.95	4.22	4.0	2.05	0.10	-1.77	1.42	-0.82	1.41
Executive Functioning									
FAS	5.05	195.1	4.0	1.34	0.26	50.8	15.3	55.8	9.11
Animals	3.80	61.4	4.0	0.73	0.57	41.8	8.93	45.6	8.93
Stroop-INT	9.90	396.6	4.0	5.12***	0.001	41.0	8.09	50.9	9.21
WCST-C	1.45	8.67	4.0	5.19***	0.001	0.95	0.85	2.40	1.51
WCST-N	9.20	352.3	4.0	3.12*	0.02	30.0	8.7	39.2	11.5
WCST-P	3.00	147.1	4.0	7.82	0.13	37.5	10.2	40.5	8.35

Note: MD = Mean Difference, MS = Mean Square, SD = Standard Deviation, FAS = Controlled Oral Association Test, LM = Logical Memory (I = Immediate Recall, D = Delayed Recall), Stroop (CW = Color-Word Condition, INT = Interference), WCST = Wisconsin Card Sorting Test (C = number of categories completed, N = non-perseverative errors, P = perseverative errors), Sig. = Significance level (P value),

\* P<0.05,

\*\* P<0.01=

\*\*\* P<0.001.

Group Comparison of Scaled Scores of Verbal Memory and Executive Functioning Measures, Excluding Patients without a History of Psychosis

**Table 3**

Measures	Test of Between-Subjects Effects				Means and Standard deviations of measures				
	MD	MS	df	F	Sig.	Mean	SD	Mean	SD
Verbal Memory									
LM-I	7.91	202.2	4.0	1.68	0.18	40.5	10.1	48.4	12.4
LM-D	8.10	161.3	4.0	1.73	0.16	38.9	6.80	47.0	13.6
CVLT-AQ	10.5	268.3	4.0	2.13	0.10	34.1	12.5	44.6	6.46
CVLT-SD	1.96	7.50	4.0	12.5***	0.0001	-2.27	0.67	-0.31	0.92
CVLT-LD	1.40	4.20	4.0	5.43**	0.002	-2.12	0.85	-0.72	0.95
CVLT-REC	0.87	3.46	4.0	1.68	0.17	-1.77	1.42	-0.90	1.51
Executive Functioning									
FAS	5.50	144.1	4.0	0.75	0.56	50.8	15.3	56.3	8.01
Animals	3.68	44.5	4.0	0.48	0.74	41.8	8.93	45.5	9.96
Stroop-INT	6.79	160.4	4.0	2.91*	0.03	41.0	8.09	47.8	6.66
WCST-C	0.95	3.65	4.0	3.20**	0.01	0.95	0.85	1.90	0.36
WCST-N	9.72	242.5	4.0	3.02*	0.03	30.0	8.7	39.7	9.54
WCST-P	2.72	134.7	4.0	1.61	0.13	37.5	10.2	39.9	7.43

Note: MD = Mean Difference, MS = Mean Square, SD = Standard Deviation, FAS = Controlled Oral Association Test, LM = Logical Memory (I = Immediate Recall, D = Delayed Recall), Stroop (CW = Color-Word Condition, INT = Interference), WCST = Wisconsin Card Sorting Test (C = number of categories completed, N = non-perseverative errors, P = perseverative errors), Sig. = Significance level (P value).

\* P<0.05,

\*\* P<0.01=

\*\*\* P<0.001.