

ORIGINAL ARTICLE

Metatranscriptomic analyses of chlorophototrophs of a hot-spring microbial mat

Zhenfeng Liu¹, Christian G Klatt², Jason M Wood², Douglas B Rusch³, Marcus Ludwig¹, Nicola Wittekindt¹, Lynn P Tomsho¹, Stephan C Schuster¹, David M Ward² and Donald A Bryant¹

¹Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University Park, PA, USA; ²Department of Land Resources and Environmental Sciences, Montana State University, Bozeman, MT, USA and ³The J Craig Venter Institute, 9704 Medical Center Drive, Rockville, MD, USA

The phototrophic microbial mat community of Mushroom Spring, an alkaline siliceous hot spring in Yellowstone National Park, was studied by metatranscriptomic methods. RNA was extracted from mat specimens collected at four timepoints during light-to-dark and dark-to-light transitions in one diel cycle, and these RNA samples were analyzed by both pyrosequencing and SOLiD technologies. Pyrosequencing was used to assess the community composition, which showed that ~84% of the rRNA was derived from members of four kingdoms *Cyanobacteria*, *Chloroflexi*, *Chlorobi* and *Acidobacteria*. Transcription of photosynthesis-related genes conclusively demonstrated the phototrophic nature of two newly discovered populations; these organisms, which were discovered through metagenomics, are currently uncultured and previously undescribed members of *Chloroflexi* and *Chlorobi*. Data sets produced by SOLiD sequencing of complementary DNA provided >100-fold greater sequence coverage. The much greater sequencing depth allowed transcripts to be detected from ~15 000 genes and could be used to demonstrate statistically significant differential transcription of thousands of genes. Temporal differences for *in situ* transcription patterns of photosynthesis-related genes suggested that the six types of chlorophototrophs in the mats may use different strategies for maximizing their solar-energy capture, usage and growth. On the basis of both temporal pattern and transcript abundance, intra-guild gene expression differences were also detected for two populations of the oxygenic photosynthesis guild. This study showed that, when community-relevant genomes and metagenomes are available, SOLiD sequencing technology can be used for metatranscriptomic analyses, and the results suggested that this method can potentially reveal new insights into the ecophysiology of this model microbial community.

The ISME Journal (2011) 5, 1279–1290; doi:10.1038/ismej.2011.37; published online 23 June 2011

Subject Category: integrated genomics and post-genomics approaches in microbial ecology

Keywords: metatranscriptome; photosynthesis; phototrophy; cyanobacteria; chlorophyll; reaction centers

Introduction

The chlorophototrophic microbial mats of alkaline siliceous hot springs in Yellowstone National Park have served for decades as model systems for understanding the composition, structure and function of microbial communities (Brock, 1978; Ward *et al.*, 1998, 2002, 2006, 2011; Ward and Castenholz, 2000; Klatt *et al.*, 2011). Members of the kingdoms *Cyanobacteria* (mostly *Synechococcus* spp.) and *Chloroflexi* (family *Chloroflexales*; filamentous anoxygenic phototrophs (FAPs), mostly *Roseiflexus*

and *Chloroflexus* spp.) (Pierson and Castenholz, 1974; van der Meer *et al.*, 2010; Ward *et al.*, 2011) are the dominant organisms in these mat communities (Table 1). Cultivation and DNA sequencing methods have produced complete or nearly complete consensus genome sequences for several chlorophototrophic organisms that occur in these mats, including *Synechococcus* spp. strains A and B', which are close relatives of indigenous populations that co-occur in the 60 °C mat (Ferris and Ward, 1997; Allewalt *et al.*, 2006; Bhaya *et al.*, 2007); *Roseiflexus* sp. strain RS-1 (van der Meer *et al.*, 2010; Bryant *et al.*, 2011); and *Chloroflexus aurantiacus* strains Y-400-fl and 396-1 (Madigan and Brock, 1977; Bryant *et al.*, 2011). The application of metagenomic methods, in combination with traditional microbial cultivation methods, recently led to the identification of a previously unknown chlorophototroph, Candidatus Chloracidobacterium

Correspondence: DA Bryant, Department of Biochemistry and Molecular Biology, The Pennsylvania State University, S-235 Frear Building, University Park, PA 16802, USA.
E-mail: dab14@psu.edu

Received 10 June 2010; revised 26 June 2010; accepted 11 February 2011; published online 23 June 2011

Table 1 Relationship between predominant phototrophic phylogenetic groups and functional guilds

Phylogenetic group	Chlorophylls	Carbon metabolism	Reaction center gene expression ^a	Chl gene expression ^a	Functional guild
<i>Synechococcus</i> spp. A-like	Chl <i>a</i>	Autotrophy	Day	Day	Oxygenic phototrophs (600–700 nm)
<i>Synechococcus</i> spp. B'-like	Chl <i>a</i>	Autotrophy	Day	Day	Oxygenic phototrophs (600–750 nm)
<i>Roseiflexus</i> spp.	BChl <i>a</i>	Mixotrophy	Morning low light	Evening	850–950 nm mixotrophs
<i>Anaerolineae</i> -like <i>Chloroflexi</i>	BChl <i>a</i> ^b	Unknown	Evening	Evening	BChl <i>a</i> /near-IR mixotrophs
<i>Chloroflexus</i> spp.	Major: BChl <i>c</i> , Minor: BChl <i>a</i>	Mixotrophy	Morning low light	Evening	700–750 nm mixotrophs
<i>C. thermophilum</i>	Major: BChl <i>c</i> , Minor: BChl <i>a</i> , Chl <i>a</i>	Heterotrophy	Evening	Day	700–750 nm heterotrophs
<i>Chlorobiales</i>	Major: BChl <i>c</i> , Minor: BChl <i>a</i> , Chl <i>a</i>	Mixotrophy	Evening	Day	700–750 nm mixotrophs

^aPeriod of maximal gene expression among times sampled to date.

^bInsufficient evidence currently exists to determine whether this organism can synthesize other chlorophylls.

thermophilum, from the kingdom *Acidobacteria* as an additional member of these mats (Bryant *et al.*, 2007). The genome of this organism has also recently been completed (Bryant *et al.*, 2011; AM Garcia Costas, Z Liu, LP Tomsho, SC Schuster, DM Ward, and DA Bryant, in preparation). As documented in the accompanying paper (Klatt *et al.*, 2011) and in previous studies (Bryant *et al.*, 2007), metagenomics analyses additionally suggested the existence of other chlorophototrophic members of this community, which were previously undescribed and are currently uncultivated organisms from the kingdoms *Chloroflexi* and *Chlorobi* (Table 1). Collectively, the availability of these genomes was instrumental in analyzing metagenomes from these mat communities (Klatt *et al.*, 2011).

Understanding transcription and its regulation in natural microbial communities has been an intriguing, although technically challenging goal, mainly because of the broad biodiversity of most such communities and the inherent difficulties in working with RNA. Microarrays (Parro *et al.*, 2007) and cloning of randomly amplified complementary DNAs (cDNAs) (Poretsky *et al.*, 2005) have been successfully applied to metatranscriptomic studies. Because of recent developments in sequencing technologies, metatranscriptomic analyses by intensive RNA sequencing have recently emerged as a much more powerful approach to the study of gene transcription in microbial communities. Moreover, there is no prerequisite to obtain genome sequences as required for microarray analyses, and such approaches can rapidly generate enormous amounts of information. Several pioneering studies using pyrosequencing technology on marine and soil microbial communities (Frias-Lopez *et al.*, 2008; Urich *et al.*, 2008; Gilbert *et al.*, 2008) have not only

demonstrated the feasibility of such an approach, but have also demonstrated its advantages in defining community structure (Urich *et al.*, 2008) and its power in identifying mRNAs of known or novel genes (Gilbert *et al.*, 2008). Several studies (Gilbert *et al.*, 2008; Hewson *et al.*, 2009; Poretsky *et al.*, 2009) compared metatranscriptomes obtained under different environmental conditions, and these studies were able to detect differential gene expression. However, compared with the number of genes occurring in a microbial community (at least tens of thousands, if not millions), the number of mRNA sequences generated in the above-mentioned pyrosequencing studies (from ~5000 to ~160 000) were insufficient to provide satisfactory coverage or to infer statistically meaningful transcription patterns under different conditions for most genes occurring in the members of a complex community. For this reason previous studies have mainly focused on the most dominant members of a given microbial community, and comparisons of transcriptional behaviors of different phylogenetic and physiological groups within the same community have not yet been described. Therefore, one of the main objectives of this study was to test the feasibility of using so-called Next-Generation (NextGen) sequencing methodologies in metatranscriptomic studies.

In a complementary study to that of Klatt *et al.* (2011), two different approaches to transcription profiling of the chlorophototrophic microbial mat community of Mushroom Spring were compared. In the first approach, pyrosequencing (Roche GS FLX, 454 Life Sciences, Branford, CT, USA) of cDNAs produced from RNA extracted from mat samples collected at 60 °C for four different time points in a diel cycle was carried out. Samples were collected

from light-to-dark and dark-to-light transition periods, because these have been shown to be important times for changes in mat physiology and metabolism, such as changes in anoxygenic photosynthesis (van der Meer *et al.*, 2005) and nitrogen fixation (Steunou *et al.*, 2006, 2008). The ~225-bp sequences obtained by this approach primarily provided information on the composition of the community based on stable (that is, rRNA) sequences. In a second approach, and as a pilot study to determine the feasibility of using NextGen sequencing to conduct a full hourly diel-cycle metatranscriptomic analysis, extensive SOLiD-3 (Applied Biosystems, Carlsbad, CA, USA) sequencing was carried out using cDNAs produced from the same four RNA samples used in the pyrosequencing analysis. After *in silico* filtering to remove stable rRNA sequences, this approach provided 50-bp sequences derived from expressed sequences (that is, mRNAs) that provided detailed information about the global patterns of transcription for predominant members of the mat community. The combination of these two approaches validated and refined conclusions about the mat composition derived from metagenomic analyses (Klatt *et al.*, 2011), and accurately reproduced previous observations concerning the expression of *Synechococcus* spp. genes for photosynthesis and nitrogen fixation (Steunou *et al.*, 2006, 2008). This study further showed that, if suitable reference genome sequences or extensive metagenomic data are available, NextGen sequencing can be used to obtain unprecedented insights into the *in situ* physiological properties of community members as inferred from transcription profiles.

Materials and methods

Sample collection

Samples of microbial mats were collected from Mushroom Spring in the Yellowstone National Park on 10–11 July 2008 at a temperature of 61 to 64 °C. All samples were collected from the same, localized region of the mat, but the ambient temperature was slightly lower at sunset (2100 hours) and slightly higher in the morning because of wind and flow conditions. Mat core samples (~0.5 cm² area, #4 cores) were collected at 2100 hours (at sunset), 0515 hours (just before sunrise), 0640 hours (indirect

sunlight on the mat) and 0840 hours (direct sunlight on the mat) at the same site. The light intensities at these four sampling times were ~15, 0.05, 46 and 960 μmol photons m⁻²s⁻¹, respectively. The top ~2-mm thick layer containing most mat chlorophototrophs (Bauld and Brock, 1973; Ruff-Roberts *et al.*, 1994) was removed with a razor blade, placed in a 2.0 ml screw-cap tube containing 0.5 g of sterile glass beads (~100–200 μm diameter), and immediately immersed in liquid nitrogen. Samples were kept frozen until used for RNA extraction.

Methods for the extraction of total RNA, synthesis of cDNA, preparation of cDNA libraries, construction of rRNA and gene, and protein databases, analytical methods for the pyrosequencing and SOLiD databases, mapping validation and other statistical analyses are described in detail in the Supplementary Material. The sequences collected in this study have been deposited under accession number SRA018112 in the NCBI Sequence Read Archive.

Results and discussion

Pyrosequencing and composition analysis of the microbial mat samples

Table 2 summarizes the data obtained by pyrosequencing of cDNAs derived from the phototrophic microbial mat samples collected at four different times in a single diel cycle from Mushroom Spring. The total number of sequences ranged from 59 283 for the sample collected at sunset (2100 hours) to 128 208 for the sample collected in the low-light morning period (0640 hours) before the mat was exposed to full sunlight. As expected, most sequences were derived from rRNAs, which accounted for 88.9–94.4% of the total sequences. The percentage of sequences assignable as mRNAs differed for the four samples. The percentage of mRNA sequences was lowest (2.4%) just before sunrise (0515 hours) and was highest (8.1%) when the mat was fully illuminated (0840 hours). The percentage of sequences that were not identified as rRNA or mRNA was essentially constant in the four samples at 2.4–3.0%.

Figure 1 summarizes the kingdom-level composition of the sequences that were identified as rRNAs (both large subunit and small subunit). These data show that ~84% of the rRNA sequences were

Table 2 Summary of cDNA sequences using pyrosequencing

	Sunset 2100 hours	Pre-dawn 0515 hours	Morning, low light 0640 hours	Morning, high light 0840 hours
Total cDNA sequences	59 283	108 948	128 208	113 128
Ribosomal RNAs	54 660 (92.2%)	102 829 (94.4%)	119 654 (93.3%)	100 566 (88.9%)
Messenger RNAs	3006 (5.1%)	3154 (2.4%)	5536 (4.3%)	9167 (8.1%)
Unidentified RNAs	1617 (2.7%)	2965 (2.7%)	3018 (2.4%)	3395 (3.0%)




Figure 1 Kingdom-level composition of the mat community based on rRNA sequences. Only RNA sequences generated by pyrosequencing were analyzed. Data shown here are the sum of the aligned sequences for the large subunit and small subunit rRNA sequences.

derived from members of only four kingdoms: *Chloroflexi* (~40%), *Cyanobacteria* (~27%), *Chlorobi* (~11%) and *Acidobacteria* (~6%). The remaining sequences (~16%) assigned as rRNAs were widely distributed across the domain Bacteria (Supplementary Figure S1). No sequences obtained from the upper mat layer could definitively be assigned to members of the domain Archaea. In agreement with Klatt *et al.* (2011), the next most abundant sequences were assigned to the kingdoms *Firmicutes* and *Bacteroidetes*, which represented less than 2% of the rRNA sequences. Overall, the best matches to these remaining sequences were distributed across hundreds of different bacterial sequence entries. As can be seen in Figure 1, only small differences in the percentage composition values were observed for the samples collected at four different times during the light-to-dark and dark-to-light transition periods. Although mat *Synechococcus* spp. exhibit phototaxis (Ramsing *et al.*, 1997) and members of *Chloroflexi* have exhibited the most dramatic evidence of vertical mobility in field studies (Boomer *et al.*, 2000), the movements of such organisms are relatively slow. It took 24 h of shading the mat to observe ‘puffs’ of filaments migrating to the surface in Octopus Spring (Boomer *et al.*, 2000). The small differences in sequence composition are unlikely to arise because of the movements of organisms up or down between the upper 2 mm and deeper mat layers (not sampled) during the diel cycle. However, it is possible that some of these differences reflect minor variations in the rRNA content of the cells as a function of the time of day. Assuming that the cells of all samples were lysed with equal efficiency in all samples, it is more likely that these differences merely reflect the slight compositional variation of the mat at the nearby but obviously slightly different sampling locations. Whatever the case, the composition of the rRNA sequences recovered from the mat at the kingdom level was essentially constant across the

diel cycle within the narrow temperature range (61–64 °C) and localized region of the mat that was sampled. These observations allow the samples to be compared in order to analyze changes in transcripts as a function of the diel cycle.

The conclusion that the upper photic layer of the mat is mostly comprised of organisms from four kingdoms agrees with results from metagenomic analyses (Klatt *et al.*, 2011). On the basis of recruitment of sequences to relevant reference genomes, metagenomic analyses of high- and low-temperature metagenomes of Mushroom Springs suggested that *Cyanobacteria*, *Chloroflexi*, *Acidobacteria* and *Chlorobi* accounted for about 41.7, 32.1, 7.6 and 3.4% of the sequences, respectively (Klatt *et al.*, 2011). Several factors might be responsible for the observed abundance differences inferred from the metagenomic versus the metatranscriptomic data. First, these methods are not quantitative and very little is known about the relative cellular contents of DNA, RNA and proteins of the organisms comprising the mat. Second, the samples were not collected from the same mat location, and moreover, the samples were collected at different times of the year (mid-summer versus late fall). These differences could have influenced the percentage of metabolically active cells, which would in turn affect the yields of DNA and rRNA. Finally, although the *Synechococcus* spp. cells are much larger in volume than those of the members of the *Chloroflexi* in the mat, the exact effects of these differences on DNA and ribosome content per cell are unknown. For all of these reasons, the precise composition of the mat cannot accurately be determined by either method.

Because of the diversity exhibited by the members of the four major kingdoms, especially *Chloroflexi* and *Acidobacteria*, which include phototrophs and non-phototrophs (Ward *et al.*, 2009; Bryant *et al.*, 2011), in-depth analyses of the 16S rRNA sequences of these kingdoms were necessary. These analyses were carried out using reference sequences with clearer taxonomic affiliations and with higher resolving power to discriminate among similar sequence alignments (see Supplementary Information for details). Figure 2 shows the results of one such analysis for the kingdom *Chloroflexi* for the sequences derived from the sample collected at sunset (2100 hours). The majority (83.5%) of the sequences in this analysis were assigned as *Roseiflexus* spp. and 6.3% of the sequences were assigned to *Chloroflexus* spp. *Roseiflexus* sp. RS-1 and *Chloroflexus* sp. 396–1 were the closest sequenced representatives from these two genera, respectively. Sequences derived from close relatives of the aerobic chemo-organotroph, *Thermomicrobium roseum*, which was reported to be present in similar mat systems in Yellowstone (Wu *et al.*, 2009; Klatt *et al.*, 2011), were also detected, although at a very low level (0.3%). As shown in Figure 2a, about 10% of the remaining sequences were assigned to three




Figure 2 In-depth composition analyses of *Chloroflexi* 16S rRNA sequences based on the high-resolution *Chloroflexi* 16S rRNA database (**a**) and after addition of partial 16S rRNA sequence of the putatively phototrophic, currently uncultivated *Chloroflexi* population/cluster/scaffolds from Mushroom Spring (**b**) in one sample (sunset). Only RNA sequences generated by pyrosequencing were analyzed. The sizes of the circles are proportional to the relative abundances of rRNA sequences assigned to a particular taxon. Figure generated by MEGAN (Huson *et al.*, 2007) (absolute score cutoff = 71, relative cutoff = 0% of best hit).

different families, mainly *Anaerolineaceae*. To probe the phylogenetic affiliation of these remaining sequences in greater detail, the 16S rRNA sequence of a novel, unclassified member of the *Chloroflexi* from the mat metagenome was used as additional reference. Figure 2b shows that almost all of the remaining sequences not assigned as *Roseiflexus* spp. or *Chloroflexus* spp. were assigned to this added sequence derived from the metagenome (Klatt *et al.*, 2011). Thus, this unclassified member of the *Chloroflexi* and its close relatives probably accounted for ~10% of *Chloroflexi* or ~4% of the total rRNA sequences. On the basis of analyses of the metagenome of Mushroom and Octopus Springs, Klatt *et al.* (2011) have found strong evidence that

this high-abundance, unclassified and uncultivated *Chloroflexi* population (most closely, but still distantly related to members of the class *Anaerolineae*) is highly likely to be a previously unrecognized chlorophototroph. This is consistent with the observation that all other high-abundance populations in this mat community are also chlorophototrophs.

As shown in Supplementary Figure S2 and as judged by the sequence similarity distribution of the 16S rRNA sequences derived from the sunset sample, members of the genus *Synechococcus* spp. accounted for 99.3% of the rRNA sequences assigned to the kingdom *Cyanobacteria* (Supplementary Figure S2A), and not surprisingly, the

previously characterized, mat-derived *Synechococcus* sp. strain A and *Synechococcus* sp. strain B' best represented those sequences (Bhaya *et al.*, 2007). Because no mat representative from the kingdom *Chlorobi* has yet been cultured or had its genome completely sequenced, the situation for this taxon was more complex. Among well-characterized *Chlorobiales* strains, the closest relative of the *Chlorobiales*-like mat population is *Chp. thalassium* (Gibson *et al.*, 1984). Approximately 45% of the rRNA sequences were assigned to this genus, and the remaining sequences assigned to the *Chlorobiales* were widely distributed across the other three genera included in the analysis (Supplementary Figure S2B). However, when the 16S rRNA sequences of the *Chlorobiales* population was recovered from the mat metagenome and used in a similar analysis, all of the *Chlorobiales*-like rRNA sequences were assigned to this single sequence type (Supplementary Figure S2C). *Candidatus* C. thermophilum accounted for 99.8% of the sequences assigned to the kingdom *Acidobacteria* (Supplementary Figure S2D). The same in-depth analysis was carried out on all samples, and the results for all four samples were similar (data not shown for the other three samples). Collectively, the rRNA sequence data suggested that the organisms belonging to the kingdom *Chloroflexi* are significantly more phylogenetically heterogeneous than those that are affiliated with the *Cyanobacteria*, *Chlorobi* and *Acidobacteria*. More importantly, the organisms affiliated with these four kingdoms, which collectively accounted for ~84% of the rRNA sequences detected from the mat community, are known to be, or are highly likely to be, chlorophototrophs.

Pyrosequencing and mRNA analysis of the microbial mat samples

Figure 3a shows a compositional analysis of the mRNA sequences derived from pyrosequencing. These data have been compared with the average percentage composition value for each major taxon based on rRNA, which were derived from the data in Figure 1. For each of the major kingdoms, the highest percentage of mRNA sequences assigned by comparison with the reference metagenome for the four samples was comparable to their average percent composition based on the rRNA sequences. The percentage of sequences identified as mRNAs but assigned to the 'Others' category was consistently higher than the average rRNA content. The main reason behind this observation was the incomplete identification of sequences of the seven major chlorophototrophic populations in the reference metagenome. Small scaffolds in the metagenome were assigned as 'Others' (see Supplementary Materials for more Supplementary Information) because it is very difficult to assign them to a phylogenetic group with high confidence. Many of these small scaffolds probably belong to the seven




Figure 3 Composition of mRNA sequences at the kingdom level (major organisms only) in both data sets, pyrosequencing (a) and SOLiD-3 (b) compared with respective average rRNA sequence percentages. Mean \pm s.d. is shown for rRNA sequence percentages for all four samples. mRNA sequences of seven predominant phototrophic phylogenetic groups and functional guilds are counted. Percentages were calculated based on total mRNA sequences aligned to reference metagenome.

major populations. For example, some of these scaffolds contain fragments of highly expressed genes, such as photosynthetic reaction centers and nitrogenase genes. Another observation was that the percentages of *Chloroflexi* mRNA sequences were higher than the average rRNA percentage for this kingdom, whereas the opposite was true for the *Cyanobacteria*. This is expected because the *Chloroflexi* populations have larger genomes (more than 4000 genes each) than the *Synechococcus* spp. (~2800 genes each). (Table 3) Although the relationship between mRNA sequence abundances and genome sizes is probably not linear, it is likely that cells with larger genomes will have more mRNA sequences than cells with smaller genomes when both are compared under similar metabolic activity levels.

After careful consideration of the factors described above, large changes in the mRNA percentages among different timepoints were nevertheless observed for two of the four kingdoms. The mRNAs assigned to *Cyanobacteria* were significantly less abundant than their average rRNA content at sunset and pre-dawn, but the mRNA levels for the

Table 3 Comparison of cDNA sequences assigned to major reference genomes using two sequencing methods

Phylogenetic groups ^a	Number of genes in reference metagenome	Number of genes in a comparable genome ^b	Number of mRNA sequences assigned ^c		Number of genes covered ^d		Number of genes significantly regulated ^e	
			454	SOLiD	454	SOLiD	454	SOLiD
Rosei	4733	4517	6446	784 685	1879	4575	28	2123
Chlor	2003	4044	210	36 549	126	1736	0	265
Anaer	2614	Not Available	778	77 958	358	2326	2	535
SyneA + SyneB'	5433	5622	2908	356 910	1204	5226	13	2695
Cab	2847	3285	1025	193 815	465	2776	5	1781
GSB	2775	Not Available	1560	274210	513	2553	7	1708
Sum of above	20 405	Not Available	12 927	1 724 127	4545	19 192	55	9107

^aRosei, *Roseiflexus* sp.; Chlor, *Chloroflexus* sp.; Anaer, novel *Anaerolineae*-like *Chloroflexi* population; SyneA, *Synechococcus* sp. strain A; SyneB', *Synechococcus* sp. strain B'; Cab, *Candidatus Chloracidobacterium thermophilum*; GSB, a *Chlorobiales* population.

^bGenomes of strains isolated in the same or similar microbial communities: *Roseiflexus* sp. RS-1 (complete); *Synechococcus* sp. strain A (complete); *Synechococcus* sp. strain B' (complete); *Candidatus Chloracidobacterium thermophilum* (complete); *Chloroflexus* sp. 396-1 (draft).

^cTotal of uniquely assigned cDNA sequences in the four samples.

^dGenes with at least one assigned cDNA sequence in the four samples.

^eGenes, expression levels of which are statistically different ($P < 0.001$) between at least one pair of samples.

Cyanobacteria increased dramatically when the mat was fully illuminated in the morning. The sequences identified as mRNAs from the *Acidobacteria* were highest at sunset and in high-intensity light in the morning but were significantly lower in the other two samples. The percentages of mRNA sequences were often dictated by expression patterns for the most highly expressed genes. The gene with the most mRNA sequences usually accounted for 5–10% of the mRNA sequences assigned to a particular taxon, and sometimes accounted for up to 25% of the mRNA sequences in a single sample. For all of the chlorophototrophs in the community, genes related to photosynthesis, especially genes encoding reaction center polypeptides, were consistently at or near the top of the lists of genes with most mRNA sequences (Supplementary Table 1). Thus, the change in the percentages of mRNA sequences of *Cyanobacteria* and *Acidobacteria* reflected how their photosynthesis-related genes are regulated and expressed (see further discussion below).

The total numbers of sequences assigned to specific coding sequences (that is, as derived from mRNAs) in the pyrosequencing approach were rather low (~3000–9000; Table 2), especially in relationship to the number of genes collectively encoded in the reference genomes of the seven major taxa (> 20 000; Table 3). Although the coverage and resolution power of the data obtained through this approach were obviously limited, it was still possible to identify statistically meaningful changes in gene expression across the four time points and to draw meaningful conclusions from a few highly expressed genes. For example, it was previously shown that the *nifH* and *nifD* genes, encoding nitrogenase in *Synechococcus* spp. populations in this mat system, exhibited maximal transcription levels at sunset and a secondary rise in transcription levels in the low-light early morning period (Steunou et al., 2008). This was clearly evident in the pattern




Figure 4 Relative transcript abundances of two *Synechococcus* spp. nitrogenase genes in two different data sets. Number of mRNA sequences assigned to *Synechococcus* sp. strain A-like and *Synechococcus* sp. strain B'-like *nifH* and *nifD* genes or partial genes were summed, normalized and compared.

of expression of the *Synechococcus* spp. *nifH* and *nifD* genes detected in this analysis (Figure 4). The mRNA abundances for the *nif* genes was highest in the sunset sample, were equivalently low in the pre-dawn and morning high-light samples, and showed a smaller but nevertheless significant increase in the morning low-light sample. Although these data helped to validate the cDNA sequencing approach to transcription profiling, the low number of total mRNA sequences led us to explore NextGen sequencing of cDNAs by using SOLiD-3 methodology to determine whether this method could be used to assess global transcription patterns in the members of the mat community.

SOLiD-3 sequencing and transcription of photosynthesis genes

Table 4 summarizes the data obtained from the sequencing of cDNA samples prepared from the

Table 4 Summary of cDNA sequences using SOLiD technology

	Sunset 2100 hours	Pre-dawn 0515 hours	Morning, low light 0640 hours	Morning, high light 0840 hours
Total mapped sequences	12 520 192	10 840 071	9 974 349	10 574 300
Sequences mapped to CDS	743 902 (5.9%)	318 205 (2.9%)	477 362 (4.8%)	1 057 433 (10.0%)
Sequences uniquely mapped to CDS	623 239 (5.0%)	261 838 (2.4%)	393 655 (3.9%)	883 799 (8.4%)

same four RNA samples described above, but by using the SOLiD-3 methodology. Overall, about 10–12 million sequences were obtained for each of the four cDNA samples. Obviously, because of the large amount of sequence information, the BLAST-based approach used for the data generated from pyrosequencing became computationally untenable, and instead the BWA rapid alignment tool was used (Li and Durbin, 2009). The 50-bp sequences mapped to rRNA databases were undoubtedly much less informative phylogenetically than their ~225-bp counterparts generated by pyrosequencing. Thus, the rRNA sequences obtained from the SOLiD-3 data were not analyzed further and will not be discussed further here. Simulations performed by the method of Richter *et al.* (2008) showed that the upper limit for unique mapping errors, when allowing three to five mismatches for 50-bp sequences, was ~0.007 and 3.12% for non-uniquely mapped sequences (Supplementary Material and Supplementary Table S3). These are estimated upper limits because the simulated data includes rRNA sequences and other repeat sequences (for example, insertion sequences, transposases and other repeated sequences) that were actually removed (or were not counted) by the transcript mapping procedure used in this study.

Although the sequences in SOLiD data sets were different in length from the ones in pyrosequencing data sets and were aligned using different softwares, the results were generally very similar in multiple ways. In all, 2.9–10.0% of the cDNA sequences (that is, 318 205 to 1 057 433 total cDNA sequences) could be mapped to coding sequences in the reference metagenomes. For a few sequences, it was not possible to map these sequences uniquely to a single gene (for example, for some transcripts encoding members of the *psbA* multigene family found in *Synechococcus* spp.). Relatively few instances of this type were encountered, and in general these few cases did not interfere with the global interpretation of the results. Similar to the results obtained by pyrosequencing (Table 2), the sequences that were assigned as mRNAs were lowest in abundance in the pre-dawn sample and were highest in abundance in the morning sample obtained after the mat received full sunlight. Figure 3b shows the compositions of mRNA sequences among four kingdoms. When compared with the pyrosequencing results in Figure 3a, it was very obvious that the pattern of changes of mRNA sequences of each kingdom across the four time-

points was almost identical, although some differences in the actual percentages were observed. For example, the percentage of *Chlorobi* and *Acidobacteria* mRNA sequences were consistently higher in SOLiD data sets than that in pyrosequencing data sets. These differences might have arisen from different biases in sequence amplification procedures, because the libraries were prepared using different kits as prescribed by the manufacturers of the two platforms. However, these biases would not affect gene expression patterns among four time-points inferred from those two data sets because the biases should be consistently present within each platform among the four different timepoint samples. The expression patterns of genes derived from the two data sets were indeed very similar. As shown in Figure 4, the SOLiD-3 and pyrosequencing methods produced essentially identical expression patterns for *nifH* and *nifD*, and as mentioned above, this pattern also matches that obtained by reverse transcription-PCR (Steunou *et al.*, 2008). For 55 genes that were detected as significantly regulated genes in the pyrosequencing data sets (Table 3), the normalized expression patterns for the two data sets were compared to determine how well the two data sets were correlated. The correlation coefficients for 49 of the 55 genes were 0.8 or greater for the two expression profiles. These observations established the consistency and the credibility of both data sets and validated the analytical approaches used.

It was also obvious that the SOLiD-3 data sets were superior in providing insights into gene regulation. As a result of the much greater sequencing depth at a similar cost (>100-fold greater sequencing depth), the SOLiD-3 data allowed transcripts to be detected for more than 90% of the >20 000 genes in the reference data set. Furthermore, the resolution power resulting from the large numbers of mRNA sequences for each gene allowed statistically meaningful comparisons to be made for a much larger number of genes. The SOLiD-3 sequencing approach detected transcripts for 19 192 genes, of which nearly half (9107, 47%) were differentially expressed in a statistically meaningful manner (Table 3).

Because the mat community members showing highest abundances were chlorophototrophs, photosynthesis genes were analyzed first. Photosynthetic reaction center genes, which have tentatively been assigned by phylogenetic analyses to the unclassified and uncultured *Anaerolineae*-like members of




Figure 5 Relative transcript abundances of photosynthetic reaction center genes of six different phototrophs. The relative transcript abundances of the indicated photosynthetic reaction center genes with most assigned mRNA sequences (SOLiD data set only) are plotted for six different phototrophs. Syne, *Synechococcus* sp. strain A and B'; Cab, *Candidatus Chloracidobacterium thermophilum*; GSB, a *Chlorobiales* population; Rosei, *Roseiflexus* sp.; Chlor, *Chloroflexus* sp.; Anaer, novel *Anaerolineae*-like *Chloroflexi* population.

the *Chloroflexi* (Supplementary Information), were among the top genes with most abundant mRNA sequences (Supplementary Table S1). This suggests that these genes have an important role in the biological processes of this population, and thus provides further very strong support for the hypothesis that this population represents a previously unidentified and uncultured chlorophototroph. This is also the case for the novel *Chlorobiales* spp. Although it was previously expected to be a chlorophototroph because all other known *Chlorobiales* spp. are photolithoautotrophs (Bryant *et al.*, 2011), the recent discovery of a non-phototrophic member of the *Chlorobi*, *Ignavibacterium album* (Iino *et al.*, 2009), made this assumption questionable. These findings about its photosynthesis genes being highly transcribed strongly suggest that *Chlorobiales* spp. in the mat are chlorophototrophs.

Figure 5 shows the expression patterns for the genes encoding the subunits of the type-1 or type-2 photochemical reaction centers of six major chlorophototrophs (Table 1) in the mat community (for this analysis we did not differentiate here between A-like and B'-like *Synechococcus* spp. sequences; Table 1). Consistent with the chlorophototrophic lifestyles of the organisms belonging to these four kingdoms, in each case the genes encoding reaction centers were either the most highly expressed, or among the group of most highly expressed, genes (Supplementary Table S1). For the *Synechococcus* spp., the mRNA levels for *psaA*, which encodes one of the two large core subunits of the photosystem I reaction center (Grotjohann and Fromme, 2005), were lowest in the pre-dawn sample but rose very steeply as the light intensity increased in the morning. Overall, the transcript abundance for *psaA* increased more than 50-fold in the morning, and




Figure 6 Relative transcript abundances for bacteriochlorophyll synthesis genes of six different phototrophs. Mean \pm s.e.m. are shown here for all bacteriochlorophyll synthesis genes of six different phototrophs, expression levels of which are statistically different ($P < 0.001$) between at least a pair of samples. Syne, *Synechococcus* sp. strain A and B'; Cab, *Candidatus Chloracidobacterium thermophilum*; GSB, a *Chlorobiales* population; Rosei, *Roseiflexus* sp.; Chlor, *Chloroflexus* sp.; Anaer, novel *Anaerolineae*-like *Chloroflexi* population.

similar increases were observed for other *psa* genes, as well as the *psb* genes encoding subunits of photosystem II (data not shown; Steunou *et al.*, 2008). In contrast, the transcript abundance for photosynthetic reaction center genes of all anoxygenic phototrophs decreased when the mat was fully illuminated. The transcription patterns for the *pscA* genes, which encode subunits of type-1 reaction centers of *Candidatus C. thermophilum* and the *Chlorobiales*-like populations in the mat (Bryant *et al.*, 2007), were very similar. For these two organisms, the *pscA* transcripts were most abundant at sunset and were least abundant when the mat was fully illuminated in the morning. Transcription patterns for the *pufM*, *pufL* and *pufC* genes, associated with the type-2 reaction centers of three different *Chloroflexi* populations, were inconsistent, but transcript levels for these genes seemed to decrease as the light intensity increased in the morning. Because the changes in the transcript levels for these genes in the *Chloroflexi* were small (\sim two-fold changes) for the time points sampled, we are currently unsure whether these differences reflect real changes in transcript levels or merely statistical variation (Figure 5).

With the obvious caveat that patterns of transcription obviously do not necessarily correlate with protein levels and enzyme activity patterns, these differences in transcription patterns for reaction center proteins nevertheless strongly imply that the organisms belonging to the major taxa in the mats likely use different strategies for light capture and usage (Table 1). Because it is unlikely that the specific absorption properties of individual cells in the mat change much during the diel cycle, the observed differences in transcription of the various chlorophototrophs suggest that it is the removal of

inhibitory factors (for example, oxygen) or the availability of required limiting nutrients for photoheterotrophs (for example, organic acids or amino acids) that ultimately controls when transcription of a given population will be maximal. Thus, for each type of organism the specific light-harvesting strategy must be to supply an adequate energy supply when the light intensity is not necessarily maximal. This could explain why chlorosomes, with their very large numbers of chromophores and low energy costs for production (Bryant and Frigaard, 2006), are the preferred antenna complexes for most of the anoxygenic phototrophs in the mats.

The biogenesis of the photosynthetic apparatus requires cells to synthesize chlorophylls and/or bacteriochlorophylls in addition to the proteins that bind these prosthetic groups. Figure 6 summarizes the sequence results for bacteriochlorophyll biosynthesis genes, expression patterns of which were statistically meaningful in the four major taxa. The pattern of expression of the chlorophyll biosynthesis genes for *Synechococcus* spp. was very similar to the pattern of expression for *psaA* (Figure 5) and other *psa* and *psb* genes. Transcript levels increased very sharply as the light intensity increased in the morning. However, for *Roseiflexus* spp. the expression of genes for bacteriochlorophyll *a* biosynthesis was highest at sunset and lowest in the morning when the mat received full sunlight. This pattern was consistent with transcripts involved in bacteriochlorophyll biosynthesis from *Chloroflexus* spp. and the novel *Anaerolineae*-like population as well, and was generally similar to that for expression of the reaction center gene *pufM* except that *pufM* transcription was lower at sunset (compare Figures 5 and 6). This pattern is consistent with the known physiology of characterized *Chloroflexi* strains, which are facultatively phototrophic. The biosynthesis of bacteriochlorophyll and components of the photosynthetic apparatus is strongly suppressed by oxygen in *Chloroflexus aurantiacus* (Bryant *et al.*, 2011). The transcription patterns for the bacteriochlorophyll biosynthesis genes for the populations of *Candidatus C. thermophilum* and the mat *Chlorobiales* spp. were essentially identical (Figure 6). Transcripts were high at sunset, declined overnight and rose sharply again in the morning and were at the highest levels when the mat received full sunlight (Figure 6). For the organisms belonging to these two populations, transcript levels for genes encoding enzymes of bacteriochlorophyll biosynthesis increased when the mat was fully illuminated, whereas transcript levels for reaction centers decreased. Interestingly, both of these types of organisms produce chlorosomes as light-harvesting antenna structures (Bryant and Frigaard, 2006; Bryant *et al.*, 2007), and it is possible that the synthesis of reaction centers (dependent on bacteriochlorophyll *a* and chlorophyll *a*, and maximal at sunset) and chlorosomes, which are dependent on bacteriochlorophyll *c* biosynthesis (Bryant and

Frigaard, 2006; Gomez Maqueo Chew and Bryant, 2007), may occur at different times of the day.

To study intra-guild gene expression for A-like and B'-like *Synechococcus* spp. transcripts, we first identified pairs of orthologous genes (blastp, reciprocal best hit), for which transcripts were both detectable and clearly assignable to one or the other genome type. The data were then examined to identify examples in that transcript levels for the two populations (i) differed in abundance in a statistically significant manner (*t*-test, $P < 0.05$); or (ii) had different expression patterns (correlation coefficient < 0). Examples of both types can be seen in Supplementary Table S2. For example, transcripts for genes encoding several chaperones were much lower in abundance in A-like than in B'-like populations. Moreover, transcripts of several genes for quinone biosynthesis and those for a glycolate oxidase subunit and glycerate kinase were more abundant in B'-like than in A-like populations. The temporal pattern of transcription of several of these genes also differed (Supplementary Table S2, for example, sensor histidine kinase and the Fe/S subunit of glycolate oxidase). Differences in both the abundance and temporal expression pattern were also observed for sensory kinases and response regulators, which suggested that entire regulons might be differentially expressed within members of these two populations. These examples illustrate that substantial transcriptional differences, which presumably reflect metabolic and physiological differences, occur with the A-like and B'-like *Synechococcus* populations within the guild of oxygenic phototrophs.

Conclusions

The data shown here showed that pyrosequencing and SOLiD-3 sequencing of cDNA samples provided complementary information concerning the composition and physiology/metabolism of the phototrophic microbial mats of alkaline siliceous hot springs. The longer sequences derived from pyrosequencing (~225 bp) provided detailed information concerning the composition of the microbial mat community. These data showed clearly that the phototrophic mat community is principally comprised of chlorophototrophic organisms derived from four kingdoms: *Chloroflexi*, *Cyanobacteria*, *Chlorobi* and *Acidobacteria*. The analysis of rRNA sequences provided more evidence for the presence of a previously unknown, unclassified and uncultured, and deeply branching chlorophototrophic member of the *Chloroflexi* in the mat. Within-guild differences in expression patterns were observed for the oxygenic photosynthesis guild, suggesting that with the current approach, the ability to distinguish the origin of transcripts is at a higher level of phylogenetic resolution than cluster analysis of metagenomic assemblies (Klatt *et al.*, 2011). The

analysis of mRNA sequences further demonstrated the feasibility and accuracy of metatranscriptomics by SOLiD NextGen sequencing for the first time. Although only four timepoints were analyzed here, this study suggested that the different phototrophic populations in the mat exhibit temporal differences in their transcriptional patterns (Table 1). Because of the ability to achieve dramatically greater sequencing depth at a reasonable cost, SOLiD-3 sequencing can provide unprecedented details concerning the global, diel patterns of transcription that occur for the organisms of this microbial mat community. Samples have been collected and sequencing has been completed for a complete, hour-by-hour analysis of the diel transcription pattern for this chlorophototrophic mat community. The description of these global diel patterns of transcription, which will provide novel insights into the *in situ* physiology of this community, will be presented elsewhere.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

This research was supported by the National Science Foundation Frontiers in Integrative Biology Research Program (EF-0328698), the National Aeronautics and Space Administration Exobiology Program (NAG5-8807), and the US Department of Energy (DOE), Office of Biological and Environmental Research (BER) as part of BER's Genomic Science Program (GSP). This contribution originates from the GSP Foundational Scientific Focus Area (FSFA) at the Pacific Northwest National Laboratory (PNNL). DAB additionally and gratefully acknowledges support from the National Science Foundation (MCB-0523100), Department of Energy (DE-FG02-94ER20137), and the Joint Genome Institute. The work conducted by the US Department of Energy Joint Genome Institute is supported by the Office of Science of the US Department of Energy under Contract No. DE-AC02-05CH11231. Finally, We gratefully acknowledge the support and assistance of National Park Service personnel at Yellowstone National Park.

References

Allewalt JP, Bateson MM, Revsbech NP, Slack K, Ward DM. (2006). Effect of temperature and light on growth of and photosynthesis by *Synechococcus* isolates typical of those predominating in the octopus spring microbial mat community of Yellowstone National Park. *Appl Environ Microbiol* **72**: 544–550.

Bauld J, Brock TD. (1973). Ecological studies of *Chloroflexis*, a gliding photosynthetic bacterium. *Arch Microbiol* **92**: 267–284.

Bhaya D, Grossman AR, Steunou A, Khuri N, Cohan FM, Hamamura N *et al.* (2007). Population level functional

diversity in a microbial community revealed by comparative genomic and metagenomic analyses. *ISME J* **1**: 703–713.

Boomer SM, Pierson BK, Austinhirst R, Castenholz RW. (2000). Characterization of novel bacteriochlorophyll-*a*-containing red filaments from alkaline hot springs in Yellowstone National Park. *Arch Microbiol* **174**: 152–161.

Brock TD. (1978). *Thermophilic Microorganisms and Life at High Temperatures*. Springer Verlag: New York.

Bryant DA, Costas AM, Maresca JA, Chew AG, Klatt CG, Bateson MM *et al.* (2007). *Candidatus Chloracidobacterium thermophilum*: an aerobic phototrophic Acidobacterium. *Science* **317**: 523–526.

Bryant DA, Frigaard N-U. (2006). Prokaryotic photosynthesis and phototrophy illuminated. *Trends Microbiol* **14**: 488–496.

Bryant DA, Klatt CG, Frigaard N-U, Liu Z, Li T, Zhao F *et al.* (2011). Comparative and functional genomics of anoxygenic green bacteria from the taxa *Chlorobi*, *Chloroflexi*, and *Acidobacteria*. In: Burnap RL, Vermaas W (eds), *Advances in Photosynthesis and Respiration*, Vol. 33 Functional Genomics and Evolution of Photosynthetic Systems, Springer: Dordrecht, The Netherlands, in press

Ferris MJ, Ward DM. (1997). Seasonal distributions of dominant 16S rRNA-defined populations in a hot spring microbial mat examined by denaturing gradient gel electrophoresis. *Appl Environ Microbiol* **63**: 1375–1381.

Frias-Lopez J, Shi Y, Tyson GW, Coleman ML, Schuster SC, Chisholm SW *et al.* (2008). Microbial community gene expression in ocean surface waters. *Proc Natl Acad Sci USA* **105**: 3805–3810.

Gibson J, Pfennig N, Waterbury JB. (1984). *Chloroherpeton thalassium* gen. nov. et spec. nov., a non-filamentous, flexing and gliding green sulfur bacterium. *Arch Microbiol* **138**: 96–101.

Gilbert JA, Field D, Huang Y, Edwards R, Li W, Gilna P *et al.* (2008). Detection of large numbers of novel sequences in the metatranscriptomes of complex marine microbial communities. *PLoS One* **3**: e3042.

Gomez Maqueo Chew A, Bryant DA. (2007). Chlorophyll biosynthesis in bacteria: the origins of structural and functional diversity. *Annu Rev Microbiol* **61**: 113–129.

Grotjohann I, Fromme P. (2005). Structure of cyanobacterial Photosystem I. *Photosynth Res* **85**: 51–72.

Hewson I, Poretsky RS, Dyhrman ST, Zielinski B, White AE, Tripp HJ *et al.* (2009). Microbial community gene expression within colonies of the diazotroph, *Trichodesmium*, from the Southwest Pacific Ocean. *ISME J* **3**: 1286–1300.

Huson DH, Auch AF, Qi J, Schuster SC. (2007). MEGAN analysis of metagenomic data. *Genome Res* **17**: 377–386.

Iino T, Mori K, Uchino Y, Nakagawa T, Harayama S, Suzuki KI. (2009). *Ignavibacterium album* gen. nov., sp. nov., a moderately thermophilic anaerobic bacterium isolated from microbial mats at a terrestrial hot spring, and proposal of *Ignavibacteria* classis nov. for a novel lineage at the periphery of the green sulfur bacteria. *Int J Syst Evol Microbiol* **60**(part 6): 1376–1382. PMID 19671715.

Klatt CG, Wood JM, Rusch DB, Bateson MM, Hamamura N, Heidelberg JF *et al.* (2011). Community ecology of hot

- spring cyanobacterial mats: predominant populations and their functional potential. *ISME J* **5**: 1262–1278.
- Li H, Durbin R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**: 1754–1760.
- Madigan MT, Brock TD. (1977). 'Chlorobium-type' vesicles of photosynthetically-grown *Chloroflexus aurantiacus* observed using negative staining techniques. *J Gen Microbiol* **102**: 279–285.
- van der Meer MTJ, Klatt CG, Wood J, Bryant DA, Bateson MM, Lammerts L *et al.* (2010). Cultivation and genomic, nutritional and lipid biomarker characterization of *Roseiflexus* sp. strains closely related to predominant *in situ* populations inhabiting Yellowstone hot spring microbial mats. *J Bacteriol* **192**: 3033–3042.
- van der Meer MTJ, Schouten S, Bateson MM, Nübel U, Wieland A, Kühl M *et al.* (2005). Diel variations in carbon metabolism by green nonsulfur-like bacteria in alkaline silicious hot spring microbial mats from Yellowstone National Park, USA. *Appl Environ Microbiol* **71**: 3978–3986.
- Parro V, Moreno-Paz M, González-Toril E. (2007). Analysis of environmental transcriptomes by DNA microarrays. *Environ Microbiol* **9**: 453–464.
- Pierson BK, Castenholz RW. (1974). A phototrophic gliding filamentous bacterium of hot springs, *Chloroflexus aurantiacus*, gen and sp nov. *Arch Microbiol* **100**: 5–24.
- Poretsky RS, Bano N, Buchan A, LeClerc G, Kleikemper J, Pickering M *et al.* (2005). Analysis of microbial gene transcripts in environmental samples. *Appl Environ Microbiol* **71**: 4121–4126.
- Poretsky RS, Hewson I, Sun S, Allen AE, Zehr JP, Moran MA. (2009). Comparative day/night metatranscriptomic analysis of microbial communities in the North Pacific subtropical gyre. *Environ Microbiol* **11**: 1358–1375.
- Ramsing NB, Ferris MJ, Ward DM. (1997). Light-induced motility of thermophilic *Synechococcus* isolates from Octopus Spring, Yellowstone National Park. *Appl Environ Microbiol* **63**: 2347–2354.
- Richter DC, Ott F, Auch AF, Schmid R, Huson DH. (2008). MetaSim—A sequencing simulator for genomics and metagenomics. *PLoS One* **3**: e3373.
- Ruff-Roberts AL, Kuenen JG, Ward DM. (1994). Distribution of cultivated and uncultivated cyanobacteria and *Chloroflexus*-like bacteria in hot spring microbial mats. *Appl Environ Microbiol* **60**: 697–704.
- Steunou AS, Bhaya D, Bateson MM, Melendrez MC, Ward DM, Brecht E *et al.* (2006). *In situ* analysis of nitrogen fixation and metabolic switching in unicellular thermophilic cyanobacteria inhabiting hot spring microbial mats. *Proc Natl Acad Sci USA* **103**: 2398–2403.
- Steunou AS, Jensen SI, Brecht E, Becraft ED, Bateson MM, Kilian O *et al.* (2008). Regulation of *nif* gene expression and the energetics of N₂ fixation over the diel cycle in a hot spring microbial mat. *ISME J* **2**: 364–378.
- Urich T, Lanzén A, Qi J, Huson DH, Schleper C, Schuster SC. (2008). Simultaneous assessment of soil microbial community structure and function through analysis of the meta-transcriptome. *PLoS One* **3**: e2527.
- Ward DM, Bateson MM, Ferris MJ, Kühl M, Wieland A, Koeppl A *et al.* (2006). Cyanobacterial ecotypes in the microbial mat community of Mushroom Spring (Yellowstone National Park, Wyoming) as species-like units linking microbial community composition, structure and function. *Phil Trans R Soc Lond B Biol Sci* **361**: 1997–2008.
- Ward DM, Castenholz RW. (2000). Cyanobacteria in geothermal habitats. In: Whitton BA, Potts M (eds), *Ecology of Cyanobacteria*. Kluwer Academic Publishers: Dordrecht, The Netherlands, pp 37–59.
- Ward DM, Ferris MJ, Nold SC, Bateson MM. (1998). A natural view of microbial biodiversity within hot spring cyanobacterial mat communities. *Microbiol Mol Biol Rev* **62**: 1353–1370.
- Ward DM, Klatt CG, Wood J, Cohan FM, Bryant DA. (2011). Functional genomics in an ecological and evolutionary context: maximizing the value of genomes in systems biology. In: Burnap RL, Vermaas W (eds), *Advances in Photosynthesis and Respiration*, Vol. 33 Functional Genomics and Evolution of Photosynthetic Systems, Springer: Dordrecht, The Netherlands, in press.
- Ward DM, Papke RT, Nübel U, McKittrick MC. (2002). Natural history of microorganisms inhabiting hot spring microbial mat communities: clues to the origin of microbial diversity and implications for microbiology and macrobiology. In: JT Staley and A.-L. Reysenbach (eds), *Biodiversity of Microbial Life: Foundations of Earth's Biosphere*. John Wiley and Sons: New York, pp 27–48.
- Ward NL, Challacombe JF, Janssen PH, Henrissat B, Coutinho PM, Wu M *et al.* (2009). Three genomes from the phylum *Acidobacteria* provide insight into the lifestyles of these microorganisms in soils. *Appl Environ Microbiol* **75**: 2046–2056.
- Wu D, Raymond J, Wu M, Chatterji S, Ren Q, Graham JE *et al.* (2009). Complete genome sequence of the aerobic CO-oxidizing thermophile *Thermomicrobium roseum*. *PLoS ONE* **4**: e4207.
- Yamada M, Zhang H, Hanada S, Nagashima KV, Shimada K, Matsuura K. (2005). Structural and spectroscopic properties of a reaction center complex from the chlorosome-lacking filamentous anoxygenic phototrophic bacterium *Roseiflexus castenholzii*. *J Bacteriol* **187**: 1702–1709.

Supplementary Information accompanies the paper on The ISME Journal website (<http://www.nature.com/ismej>)