

Published in final edited form as:

J Am Chem Soc. 2011 August 3; 133(30): 11482–11484. doi:10.1021/ja2048072.

Deoxyfluorination of Phenols

Pingping Tang, Weike Wang, and Tobias Ritter


Department of Chemistry and Chemical Biology, Harvard University, 12 Oxford Street, Cambridge, MA 02138

Tobias Ritter: ritter@chemistry.harvard.edu

Abstract

An operationally simple *ipso*-fluorination of phenols with a new deoxyfluorination reagent is presented.

While several arene fluorination reactions have been reported within the past five years,^{1,2} efficient fluorination of fine and bulk chemicals is still an unsolved problem.³ Simple aryl fluorides can be accessed by conventional fluorination reactions such as the Balz-Schiemann reaction,⁴ but currently, only two fluorination reactions are available to make functionalized fluoroarenes with a large variety of substitution patterns: Buchwald's palladium-catalyzed nucleophilic fluorination of aryl triflates,^{1j} and a silver-catalyzed electrophilic fluorination of aryl stannanes reported by our group.¹¹ Both reactions currently lack practicality: The palladium-catalyzed reaction requires anhydrous conditions and can give mixtures of constitutional isomers, and the silver-catalyzed reaction requires the stepwise synthesis of toxic aryl stannanes. Our goal was to develop a reliable and operationally simple fluorination reaction that provides more straightforward access to a wide variety of aryl fluorides. Here we present a fluorinating reagent that delivers aryl fluorides from phenols by a one-step *ipso* substitution.




(1)

The new deoxyfluorination reagent **1** (eq. 1) was prepared by treatment of *N,N*-diaryl-2-chloroimidazolium chloride,⁵ readily available in >100 g quantities (see Supporting Information), with CsF. Reagent **1** can be handled in air as a solid, but hydrolyzes upon prolonged storage in a wet atmosphere. It can be stored in a dry toluene solution for at least two months without detectable decomposition. Reaction of 4-methoxyphenol, **1**, and CsF in toluene or dioxane afforded 4-fluoroanisole in 82% and 88% yield, respectively (Table 1). The commercially available deoxyfluorination reagents⁶ shown in Table 1 did not afford any detectable fluoroanisole under identical reaction conditions. Reactions performed in polar (MeCN) or protic solvents did not afford any detectable aryl fluoride with any of the evaluated commercial fluorinating reagents, and less than 10% yield with fluorinating reagent **1**.

Correspondence to: Tobias Ritter, ritter@chemistry.harvard.edu.

Supporting Information Available: Experimental procedures and spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

A variety of phenols was evaluated for fluorination (Table 2). Phenols with electron withdrawing groups reacted faster than phenols substituted with electron-releasing functional groups, and typically afforded >90% yield of fluorinated arene after heating at 80 °C for 3 hours. In this study we focused on electron rich arenes because electron poor arenes can typically be prepared by nucleophilic aromatic substitution.⁷ Electron-rich phenols can be fluorinated by **1** when heated at 110 °C for 20 hours. It is noteworthy that even 4-fluoroaniline (**15**) can be prepared by the presented method. Strong hydrogen bond donors such as alcohols are not tolerated, unless the hydrogen atom of the alcohol OH group is engaged in intramolecular hydrogen bonding, such as in **28**. Amides derived from ammonia and primary amines, which are potential hydrogen bond donors, are challenging substrates for the fluorination reaction. For example, 4-fluorobenzamide was isolated in only 20% yield. Phenols featuring amides derived from secondary amines such as **17** can be fluorinated, as can 2- and 4- pyridones, (eq 2, see also structures **23** and **25** in Table 1). Both compound classes afford products that lack N–H hydrogen bond donors. Importantly, deoxyprotonated products and constitutional isomers were not observed throughout our study; all substitutions are *ipso*-substitutions with fluoride. The palladium-catalyzed fluorination of aryl triflates^{1j} can give constitutional isomers; for example, fluorination of the triflate derived from 4-methoxyphenol afforded 3-fluoroanisole as the major product (58%), and only 25% of **12**.



(2)

We propose that the mechanism for fluorination proceeds via a 2-phenoxy-imidazolium bifluoride salt such as **30**, formed by condensation of a phenol with **1** (Scheme 1). Salt **30** was formed upon mixing 4-methoxyphenol and **1**, and could be isolated in 91% yield. A hydrogen bond between one hydrogen atom of the imidazolium heterocycle and the bifluoride counteranion was identified in the crystal structure of **30** (C–F distance of C–H···F–H–F: 3.0 Å). We postulate that a similar hydrogen bond is also present in solution in apolar solvents: when the hexafluorophosphate salt **33** was treated with bifluoride, a shift of the ¹H NMR resonance of the hydrogen atoms of the imidazolium heterocycle was observed from 7.56 ppm to 7.98 ppm (see Supporting Information). Bifluoride is a strong hydrogen bond acceptor.⁹ Hydrogen bonding could facilitate fluorination because it renders the uronium a better nucleofuge or because the bifluoride salts such as **30** exist as soluble tight ion pairs in apolar solvents, with close proximity of fluoride to the ipso-carbon. In agreement with both hypotheses, imidazolium salts **31** and **32**, which lack the opportunity for similar hydrogen bonding, did not afford detectable aryl fluoride when heated. Likewise, exchange of the bifluoride for the less hydrogen bond-accepting counteranion PF₆ lowered the reaction yield to 2%. Moreover, the commercially available deoxyfluorinating reagents shown in Table 1 lack the ability to form intermediates suitable for C(sp²)···F–H–F hydrogen bonding as found in **30**, which may explain the difference in deoxyfluorinating reactivity between the known reagents and **1**. Quantitative rate measurements, for example for a Hammett analysis with different substituents in the *para* position of the phenol, were not obtained due to the fact that the reaction mixture is heterogeneous. The heterogeneous mixture may also explain the difference in yield of fluorination when fluorination is performed with **30** isolated (67%, reaction incomplete), rather than formed in situ in the presence of excess CsF (82%).

In conclusion we have developed a practical deoxyfluorination of phenols that can be used conveniently from milligram to multigram scale. Whereas in conventional nucleophilic fluorination chemistry hydrogen bonding is undesirable because it reduces the nucleophilicity of fluoride, hydrogen bonding appears to be crucial for the fluorination

reaction presented here. A disadvantage of the transformation is the generation of a stoichiometric amount of waste, the urea byproduct. However, the deoxyfluorination proceeds in one step from phenols. Ultimately, a fluorination reaction that could combine the operational simplicity of the transformation presented here, the catalysis approach of the Buchwald fluorination, and the ability to fluorinate complex molecules late stage as accomplished by the silver-catalyzed fluorination would significantly advance the field of fluorination.

Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Dr. Shao-Liang Zheng for his help with X-ray data collection and structure determination. This work was supported by NSF (CHE-0952753) and NIH-NIGMS (GM088237).

REFERENCES

1. For transition-metal-mediated C–F bond formation, see: (a) Hull KL, Anani WQ, Sanford MS. *J. Am. Chem. Soc.* 2006; 128:7134. [PubMed: 16734446] (b) Akana JA, Bhattacharyya KX, Müller P, Sadighi JP. *J. Am. Chem. Soc.* 2007; 129:7736. [PubMed: 17547409] (c) Kaspi AW, Yahav-Levi A, Goldberg I, Vigalok A. *Inorg. Chem.* 2008; 47:5. [PubMed: 18052157] (d) Furuya T, Kaiser HM, Ritter T. *Angew. Chem. Int. Ed.* 2008; 47:5993. (e) Furuya T, Ritter T. *J. Am. Chem. Soc.* 2008; 130:10060. [PubMed: 18616246] (f) Furuya T, Strom AE, Ritter T. *J. Am. Chem. Soc.* 2009; 131:1662. [PubMed: 19191693] (g) Ball ND, Sanford MS. *J. Am. Chem. Soc.* 2009; 131:3796. [PubMed: 19249867] (h) Wang X, Mei TS, Yu JQ. *J. Am. Chem. Soc.* 2009; 131:7520. [PubMed: 19435367] (i) Furuya T, Ritter T. *Org. Lett.* 2009; 11:2860. [PubMed: 19507870] (j) Watson DA, Su M, Teverovskiy G, Zhang Y, García-Fortanet J, Kinzel T, Buchwald SL. *Science.* 2009; 325:1661. [PubMed: 19679769] (k) Furuya T, Benitez D, Tkatchouk E, Strom AE, Tang P, Goddard WA III, Ritter T. *J. Am. Chem. Soc.* 2010; 132:3793. [PubMed: 20196595] (l) Tang P, Furuya T, Ritter T. *J. Am. Chem. Soc.* 2010; 132:12150. [PubMed: 20695434] (m) Tang P, Ritter T. *Tetrahedron.* 2011; 67:4449. [PubMed: 21691436]
2. For other recent arene fluorination reactions, see: (a) Sun H, DiMugno SG. *Angew. Chem. Int. Ed.* 2006; 45:2720. (b) Yamada S, Gavryushin A, Knochel P. *Angew. Chem. Int. Ed.* 2010; 49:2215. (c) Anbarasan P, Neumann H, Beller M. *Angew. Chem. Int. Ed.* 2010; 49:2219.
3. Furuya T, Kamlet AS, Ritter T. *Nature.* 2011; 473:470. [PubMed: 21614074]
4. (a) O'Hagan D. *Chem. Soc. Rev.* 2008; 37:308. [PubMed: 18197347] (b) Furuya T, Klein JEMN, Ritter T. *Synthesis.* 2010; 11:1804. [PubMed: 20953341]
5. Mendoza-Espinosa D, Donnadiu B, Bertrand G. *J. Am. Chem. Soc.* 2010; 132:7264. [PubMed: 20443633]
6. (a) Middleton WJ. *J. Org. Chem.* 1975; 40:574. (b) Lal GS, Pez GP, Pesaresi RJ, Prozonc FM. *Chem. Commun.* 1999; 2:215. (c) Hayashi H, Sonoda H, Fukumura K, Nagata T. *Chem. Commun.* 2002:1618. (d) Beaulieu F, Beauregard LP, Courchesne G, Couturier M, LaFlamme F, L'Heureux A. *Org. Lett.* 2009; 11:5050. [PubMed: 19799406]
7. For selected nucleophilic aromatic fluorination reactions, see: (a) Bunnett JF, Zahler RE. *Chem. Review.* 1951; 49:273. (b) Adams DJ, Clark JH. *Chem. Soc. Rev.* 1999; 28:225. (c) Kuduk SD, DiPardo RM, Bock MG. *Org. Lett.* 2005; 7:577. [PubMed: 15704898] (d) Sun H, DiMugno SG. *J. Am. Chem. Soc.* 2005; 127:2050. [PubMed: 15713075]
8. The role of Ph₂Zn has not yet been established.
9. Kollman PA, Allen LC. *J. Am. Chem. Soc.* 1970; 92:6101.



Scheme 1.
Studies of Hydrogen bonding
Ar = 2,6-diisopropylphenyl, Ar' = 4-methoxyphenyl

Table 1


Evaluation of different fluorination reagents^a.





























Fluorinating reagent:	MeCN	dioxane	toluene
 1	<1%	88%	82%
	<1% ^b		
 Xtalfluor-E [⊕]	<1% ^b		
 Xtalfluor-M [⊕]	<1% ^b		
 DAST [⊕] DEOXYFLUOR	<1% ^b		<1% ^b

^a) Yields were determined by ¹⁹F NMR with internal standard.

^b) The commercially available deoxyfluorination reagents afforded no detectable 4-fluoroanisole in MeCN, 1,4-dioxane, or toluene.

Table 2

Deoxyfluorination of phenols with reagent **1**.^a


 2, 93% (80 °C, 3 h)	 3, 98% ^b (80 °C, 3 h)	 4, 95% (80 °C, 3 h)	 5, 95% ^b (80 °C, 3 h)	 6, 92% ^b (80 °C, 3 h)	 7, 82% ^b (80 °C, 18 h)	 8, 90% (80 °C, 18 h)
 9, 91% (80 °C, 18 h)	 10, 88% ^b (80 °C, 18 h)	 11, 88% ^b (80 °C, 18 h)	 12, 82% ^b (110 °C, 20 h)	 13, 81% ^b (110 °C, 20 h)	 14, 87% ^{b,c,d} (110 °C, 20 h)	 15, 75% ^{b,c} (110 °C, 20 h)
 16, 74% ^b (110 °C, 20 h)	 17, 91% (110 °C, 20 h)	 18, 86% (110 °C, 20 h)	 19, 88% ^c (110 °C, 20 h)	 20, 90% (110 °C, 20 h)	 21, 92% (110 °C, 20 h)	 22, 95% (110 °C, 20 h)
 23, 90% ^b (80 °C, 18 h)	 24, 84% ^b (110 °C, 20 h)	 25, 50% ^{b,c,e} (110 °C, 20 h)	 26, 88% (110 °C, 20 h)	 27, 90% (110 °C, 20 h)	 28, 75% (110 °C, 20 h)	 29, 95% (110 °C, 20 h)

^a) Reaction conditions: 1.2 equiv of **1**, concentration = 0.1 M. Toluene was used as standard solvent due to its higher boiling point (111 °C) compared to 1,4-dioxane (101 °C).

^b) Yields were determined by ¹⁹F NMR with 1-fluoro-3-nitrobenzene as a standard.

^c) 1,4-dioxane was used as solvent.

^d) 20 mol% ZnPh₂ was added;⁷ 55% yield in the absence of ZnPh₂.

^e) 20 mol% ZnPh₂ was added; 25% yield in the absence of ZnPh₂.