

Published in final edited form as:

Nature. ; 476(7361): 446–449. doi:10.1038/nature10305.

A chronological framework for the British Quaternary based on *Bithynia opercula*

Kirsty E.H. Penkman¹, Richard C. Preece², David R. Bridgland³, David H. Keen^{4,†}, Tom Meijer⁵, Simon A. Parfitt^{6,7}, Tom S. White², and Matthew J. Collins¹

¹BioArCh, Departments of Archaeology & Chemistry, University of York, York YO10 5DD, UK

²Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ UK

³Department of Geography, University of Durham, South Road, Durham DH1 3LE ⁴Institute of Archaeology and Antiquity, University of Birmingham, Birmingham B15 2TT, UK ⁵Cainozoic Mollusca, Netherlands Centre for Biodiversity, Naturalis, P.O. Box 9517, 2300 RA Leiden, The Netherlands ⁶Institute of Archaeology, University College London, 31-34 Gordon Square, London WC1H 0PY, UK ⁷Department of Palaeontology, The Natural History Museum, Cromwell Road, London SW7 5BD, UK

Abstract

Marine and ice-core records show that the Earth has experienced a succession of glacial and interglacials during the Quaternary (last ~2.6 million years), although it is often difficult to correlate fragmentary terrestrial records with specific cycles. Aminostratigraphy is a method potentially able to link terrestrial sequences to the marine isotope stages (MIS) of the deep-sea record^{1,2}. We have used new methods of extraction and analysis of amino acids, preserved within the calcitic opercula of the freshwater gastropod *Bithynia*, to provide the most comprehensive dataset for the British Pleistocene based on a single dating technique. A total of 470 opercula from 74 sites spanning the entire Quaternary are ranked in order of relative age based on the extent of protein degradation, using aspartic acid (Asx), glutamic acid (Glx), serine (Ser), alanine (Ala) and valine (Val). This new aminostratigraphy is consistent with the stratigraphical relations of stratotypes, sites with independent geochronology, biostratigraphy and terrace stratigraphy³⁻⁶. The method corroborates the existence of four interglacial stages between the Anglian (MIS 12) and the Holocene in the terrestrial succession. It establishes human occupation of Britain in most interglacial stages after MIS 15, but supports the notion of human absence during the Last Interglacial (MIS 5e)⁷. Suspicions that the treeless 'optimum of the Upton Warren interstadial' at Isleworth pre-dates MIS 3 are confirmed. This new aminostratigraphy provides a robust framework against which climatic, biostratigraphical and archaeological models can be tested.

Despite the importance of the terrestrial record for climate models, the difficulties of assigning specific sedimentary sequences to individual climate cycles restricts the use of these data in climate modelling. The British Quaternary is exceptional for the number of

Correspondence and requests for materials should be addressed to K.E.H.P. (kirsty.penkman@york.ac.uk)..

[†]Deceased 16 April 2006

Supplementary Information Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Author contributions K.E.H.P., M.J.C., R.C.P. and D.H.K. designed the study. R.C.P., D.H.K., T.M., D.R.B., S.A.P., T.S.W. and K.E.H.P. supplied samples. K.E.H.P. undertook the analyses and processed the data. K.E.H.P., M.J.C., T.S.W. and R.C.P. wrote the paper. All authors discussed the results and commented on the manuscript.

Author information Reprints and permissions information is available at www.nature.com/reprints.

The authors declare no competing financial interests.

recorded sites and their biodiversity, which has fluctuated markedly due to the mid-latitude situation of this ephemeral island. A consensus has emerged from attempted differentiation between interglacials in Britain using river terrace stratigraphy^{6,8} and biostratigraphy³⁻⁵ (Table 1).

This study revisits research pioneered by Miller et al. (predominantly on bivalves)¹ and Bowen et al. (predominantly on gastropod shells)² who used the extent of racemization in the amino acid L-isoleucine (to its diastereomer D-alloisoleucine, yielding an A/I value) in non-marine mollusc shells to build an aminostratigraphy of terrestrial sequences that could be linked to the marine oxygen isotope stratigraphy. Following debate concerning certain correlations, we developed a revised method of extraction and analysis⁹. Shells of freshwater gastropods (*Bithynia* and *Valvata*) from many of the original sites¹⁰ have been re-analysed, confirming much of the A/I stratigraphy. However, it emerged that within-site and within-stage variability increases in shells from older sites. This variability probably results from diagenetic alteration of the biomineral carbonate from aragonite to the more thermodynamically stable calcite^{10,11}.

Our new method has five significant revisions, three of which reduced within-site variability. First, inter-species variation was minimised by analysing only a single genus of freshwater gastropod (*Bithynia*). Second, variability in amino acid concentration and D/L values was significantly lowered when samples were crushed to 500 µm and exposed to prolonged wet chemical oxidation (48 hours, 12% wt./vol. NaOCl, room temperature), destroying any contamination and leaving a functionally closed-system protein fraction defined as 'intra-crystalline'^{9,12}. Third, the calcitic operculum, which in life closes the aperture of the shell, was analysed instead of the aragonitic shell. Opercula display less within-site variation and greater stability than shells^{10,11}, and show subtle but minimal intraspecific differences in racemization (Supplementary Data 1). Further modifications included the analysis of a range of different amino acids¹³ (rather than only a single measure of racemization), which allows an estimate of 'Intra-crystalline Protein Decomposition' (IcPD). This integrates data from amino acids with differing rates of racemization (Asx >> Ala > Val ~ Glx) with the extent of dehydration of serine ([Ser]/[Ala]) to estimate age (Fig. 1). Finally, comparisons were made between free amino acids (FAA), liberated by diagenesis, and the total extent of racemization (THAA), in order to test closed-system conditions¹⁴.

The intra-crystalline fraction maintains constant chemical conditions so that the extent of protein degradation can be attributed to the thermal history of the sample. Within the study area differences in thermal history have been minor during the past century and chiefly related to burial depth and the thermal diffusivity of the overburden, mediated by the presence/absence of vegetation and/or snow cover (Supplementary Figure 1 and Discussion). Opercula with similar levels of protein degradation are therefore thought to be of equivalent age, assuming that similarly small thermal gradients existed during past interglacials, when most of the racemization would have occurred.

The consistency of our method has been tested by measuring opercula from British interglacial stratotypes and/or sites with independent geochronology (Table 1; Fig. 1, Supplementary Data 1). All stratotypes yielding *Bithynia* have been analysed, but stratotypes have not been formally defined for all stages¹⁵. Sites with independent geochronology can be used to calibrate IcPD results but, as only 7 of our pre-Holocene sites have associated dates, we have not done this here. Nevertheless, an age-dependent increase in the level of IcPD is seen from the Holocene to the Early Pleistocene, using a combination of fast (e.g. Asx) and slow (e.g. Val) racemizing amino acids to span this range.

Increasing protein decomposition within opercula is also consistent with increasing river terrace elevation (and therefore age) in Quaternary fluvial archives (Table 1 and Supplementary Data 1 and 2, Figure 2). The formation of river terraces is attributed to climatic forcing and uplift, linking aggradation and incision phases with climatically induced changes in sediment and water supply^{6,8,16}. Assemblages of amino acid values from the Thames sequence, the most complete British fluvial archive¹⁷⁻¹⁹, clearly correspond with discrete terrace aggradations (Fig. 2 and Supplementary Data 2). Four interglacials after the Anglian (MIS 12) stage are represented in this system, assigned to MIS 11, 9, 7 and 5e on the basis that each aggradation formed during a complete glacial/interglacial cycle^{6,16}. Although this relationship might not hold for all rivers²⁰, a similar pattern exists between Ala D/L values and terraces in other systems, such as the Severn/Avon²¹, Trent/Witham²² and Nene/Welland²³. Support for these interpretations comes from integrating several lines of evidence (e.g. biostratigraphy⁴⁻⁵ and some of the original A/I data¹⁶), not all of which are wholly independent. However, the ability of the method to differentiate between terrace aggradations (Supplementary Data 2) is not reliant on other data and sites where the age attribution is based fundamentally on amino acid data are not assigned ‘consensus MIS ages’ in Table 1.

Bithynia is generally rare in cold-stage contexts, although it occurred commonly in the ‘Upton Warren interstadial’ deposit at Isleworth. This was originally thought to fall within the ‘Middle Devensian’ (MIS 3) on the basis of a radiocarbon date of ~43 ka BP²⁴. The Isleworth opercula IcPD is consistently higher than from Cassington, a site tentatively correlated with MIS 5a²⁵, but lower than Last Interglacial opercula. Radiocarbon therefore provides only a minimum age for the Isleworth deposits. Our IcPD data indicate an earlier age and suggest that this new method can potentially be used to distinguish marine isotope sub-stages beyond the limits of radiocarbon dating.

Aminostratigraphic data also provide independent support for biostratigraphic age models developed for the Middle Pleistocene^{5,26,27}. In the early Middle Pleistocene, the water vole *Arvicola* is thought to have replaced its ancestor *Mimomys savini* within a relatively short time over large regions of Europe^{3,27}. This hypothesis gains support from our new data, which show that opercula from sites yielding *Arvicola* show less protein degradation than those containing *M. savini* (Table 1).

The occurrence of *Corbicula* (a bivalve) and *Hippopotamus* in the British Pleistocene is mutually exclusive^{5,26}. At British sites securely attributable to the Last Interglacial (MIS 5e) *Corbicula* is absent; conversely, after MIS 12, *Hippopotamus* is only known from the Last Interglacial, and is therefore widely regarded as an ‘indicator species’ for MIS 5e⁴. Our data support these conclusions, as post-Anglian sites with *Hippopotamus* show less protein breakdown than sites yielding *Corbicula*, with levels of protein degradation consistent with attribution to MIS 5e.

The comparisons above demonstrate the remarkable consistency of our new method with independent lines of evidence. This comprehensive dating framework enables us to explore the British archaeological record. Our data show that human occupation occurred within at least two distinct pre-Anglian stages, the older (Pakefield) associated with *Mimomys* and the younger (Waverley Wood) with *Arvicola*. The conclusion that Waverley Wood is younger than the Cromerian stratotype at West Runton supports the biostratigraphic age model²⁷ and contradicts a conclusion reached in an earlier aminostratigraphical study². Our data can provide age constraints for other archaeological assemblages, enabling attribution to specific marine isotope stages in younger deposits. The development of Levallois technology, characterized by the removal of flakes from specifically prepared cores, is unknown in Britain before MIS 9²⁸. Archaeological evidence from MIS 9 is sparse, but the far better

record from sites attributed to MIS 7 shows that Levallois industries had become dominant in southern England; our data support this view (Table 1; Fig. 3).

In recent years it has become clear that humans were absent from Britain during the Last Interglacial; earlier claims to the contrary have been shown to be based on misinterpretation of archaeological sites previously thought to be of Last Interglacial age (such as Aveley, Crayford, Grays, Purfleet and Stutton), invariably now assigned to earlier stages²⁹ (Supplementary Data 1). Our results confirm that no British archaeological site can be attributed to the Last Interglacial (Fig. 3), a conclusion consistent with human absence during this stage^{7,29}.

This stratigraphical framework provides a secure basis for relating the British terrestrial British sequence to global Quaternary climate records. This is fundamental to geological and archaeological research but, as importantly, it enables the rich British record to be used to test the ability of climate models to simulate pre-late Quaternary palaeoclimates. This dating technique offers a means of correlating terrestrial with marine and ice-core records, thereby increasing the confidence of model predictions³⁰. Moreover, the calcitic opercula of bithyniid (or similar) gastropods occur commonly in many Quaternary sequences, offering potential for development of regional aminostratigraphies around the world.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank P. Allen, N. Ashton, D. Bain, M. Bates, S. Boreham, D. Bowen, R. Briant, C. Buckingham, J. Clayden, G. R. Coope, A. Cruickshanks, P. Dark, B. Demarchi, M. Greenwood, T. van Kolfschoten, H. Langford, S. Lewis, D. Maddy, R. Markham, D. Mayhew, H. Roe, J. Rose, D. Schreve, R. Scott, K. Scott, C. Stringer, R. Waghorne, M. Warren and F. Wenban-Smith for providing some of the material analysed and general discussion. C. Helmsley and J. Todd released material from the Natural History Museum, London, for destructive analysis. R. Allen provided technical support. G. Peeters gave permission to reproduce images of *Bithynia*. The analyses were funded by English Heritage, NERC and the Wellcome Trust (grant GR076905MA). This is a contribution to the Ancient Human Occupation of Britain (AHOB) project funded by the Leverhulme Trust.

References

1. Miller GH, Hollin JT, Andrews JT. Aminostratigraphy of UK Pleistocene deposits. *Nature*. 1979; 281:539–543.
2. Bowen DQ, Hughes S, Sykes GA, Miller GH. Land-sea correlations in the Pleistocene based on isoleucine epimerization in non-marine molluscs. *Nature*. 1989; 340:49–51.
3. von Koenigswald, W.; van Kolfschoten, T. The early Middle Pleistocene in Europe. Turner, C., editor. Balkema; 1996. p. 211–226.
4. Schreve DC. Differentiation of the British late Middle Pleistocene interglacials: the evidence from mammalian biostratigraphy. *Quaternary Science Reviews*. 2001; 20:1693–1705.
5. Keen DH. Towards a late Middle Pleistocene non-marine molluscan biostratigraphy for the British Isles. *Quaternary Science Reviews*. 2001; 20:1657–1665.
6. Bridgland DR. The record from British Quaternary river systems within the context of global fluvial archives. *Journal of Quaternary Science*. 2010; 25:433–446.
7. Ashton N, Lewis SG. Deserted Britain: declining populations in the British late Middle Pleistocene. *Antiquity*. 2002; 76:388–396.
8. Bridgland DR, Maddy D, Bates M. River terrace sequences: templates for Quaternary geochronology and marine-terrestrial correlation. *Journal of Quaternary Science*. 2004; 19:203–218.

9. Penkman KEH, Kaufman DS, Maddy D, Collins MJ. Closed-system behaviour of the intracrystalline fraction of amino acids in mollusc shells. *Quaternary Geochronology*. 2008; 3:2–25. [PubMed: 19684879]
10. Penkman KEH, et al. Testing the aminostratigraphy of fluvial archives: the evidence from intracrystalline proteins within freshwater shells. *Quaternary Science Reviews*. 2007; 26:2958–2969. [PubMed: 19684880]
11. Penkman KEH, Preece RC, Keen DH, Collins MJ. Amino acid geochronology of the type Cromerian of West Runton, Norfolk, UK. *Quaternary International*. 2010; 228:25–37. [PubMed: 21217810]
12. Sykes GA, Collins MJ, Walton DI. The significance of a geochemically isolated intracrystalline fraction within biominerals. *Organic Geochemistry*. 1995; 23:1059–1065.
13. Kaufman DS, Manley WF. A new procedure for determining DL amino acid ratios in fossils using reverse phase liquid chromatography. *Quaternary Science Reviews*. 1998; 17:987–1000.
14. Preece RC, Penkman KEH. New faunal analyses and amino acid dating of the Lower Palaeolithic site at East Farm, Barnham, Suffolk. *Proceedings of the Geologists' Association*. 2005; 116:363–377.
15. Bowen DQ. A revised correlation of Quaternary deposits in the British Isles. *Geological Society of London, Special Report No. 23*. 1999:174.
16. Bridgland DR. River terrace systems in north-west Europe: an archive of environmental change, uplift and early human occupation. *Quaternary Science Reviews*. 2000; 19:1293–1303.
17. Gibbard, P. *The Pleistocene History of the Lower Thames Valley*. Cambridge University Press; 1994.
18. Bridgland, DR. *Quaternary of the Thames*. Chapman and Hall; 1994.
19. Bridgland DR. The Middle and Upper Pleistocene sequence in the Lower Thames: a record of Milankovitch climatic fluctuation and early human occupation of southern Britain. *Proceedings of the Geologists' Association*. 2006; 117:281–305.
20. Bridgland DR, et al. The Palaeolithic occupation of Europe as revealed by evidence from the rivers: data from IGCP 449. *Journal of Quaternary Science*. 2006; 21:437–455.
21. Maddy D, Keen DH, Bridgland DR, Green CP. A revised model for the Pleistocene development of the River Avon, Warwickshire. *Journal of the Geological Society of London*. 1991; 148:473–484.
22. White, TS.; Bridgland, DR.; Howard, AJ. *The Quaternary of the Trent Valley and Adjoining Regions: Field Guide*. White, TS.; Bridgland, DR.; Howard, AJ.; White, MJ., editors. Quaternary Research Association; 2007. p. 10-23.
23. Boreham S, White TS, Bridgland DR, Howard AJ, White MJ. The Quaternary history of the Wash fluvial network. *Proceedings of the Geologists' Association*. 2010; 121:393–409.
24. Coope GR, Angus RB. An ecological study of a temperate interlude in the middle of the last glaciation, based on fossil Coleoptera from Isleworth, Middlesex. *Journal of Animal Ecology*. 1975; 44:365–391.
25. Maddy D, et al. The Upper Pleistocene deposits at Cassington, near Oxford, England. *Journal of Quaternary Science*. 1998; 13:205–231.
26. Meijer T, Preece RC. A review of the occurrence of *Corbicula* in the Pleistocene of North-west Europe. *Netherlands Journal of Geosciences – Geologie en Mijnbouw*. 2000; 79:241–255.
27. Preece RC, et al. Biostratigraphic and aminostratigraphic constraints on the age of the Middle Pleistocene glacial succession in North Norfolk, UK. *Journal of Quaternary Science*. 2009; 24:557–580.
28. White MJ, Ashton N. Lower Palaeolithic core technology and the origins of the Levallois method in North-Western Europe. *Current Anthropology*. 2003; 44:598–609.
29. Lewis, SG.; Ashton, N.; Jacobi, R. *The Ancient Human Occupation of Britain*. Ashton, N.; Lewis, SG.; Stringer, C., editors. Elsevier; 2010. p. 125-164.
30. Jansen, E., et al. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Solomon, S., et al., editors. Cambridge University Press; 2007.

Site number (SI Fig. 1)	Site	Bifurca species	Number analysed	D/L		Independent Dendrochronology	Terrace stratigraphy Hypogonitimus / Coriulica Arvicola / Mimomys	Mesolithic Upper Palaeolithic Lepidolite Acheulian Clactonian Undiagnostic	Consensus MIS
				Ala	Val				
				mean standard deviation	mean standard deviation				
1	Acle (modern)	tr	3	0.19	0.00	0.10	0.00		
1	Acle (modern)	tr	2	0.04	0.00	0.02	0.00		
2	Enfield Lock	tr	4	0.05	0.01	0.03	0.01		1
3	Quadenham Mere, 250-260cm	tr	4	0.05	0.01	0.03	0.00		1
3	Quadenham Mere, 640-650cm	tr	3	0.05	0.01	0.03	0.00		1
4	Newby Wiske	tr	3	0.08	0.01	0.04	0.00		1
5	Aston-upon-Trent	tr	4	0.06	0.00	0.04	0.00		1
6	Star Carr, 245-250cm	tr	2	0.07	0.01	0.04	0.00		1
7	Sproughton	tr	4	0.09	0.00	0.05	0.00		2
6	Star Carr, 524-528cm	tr	3	0.10	0.02	0.04	0.00		2
8	Cassington	tr	4	0.12	0.01	0.07	0.00		5c-4
9	Ialeworth	tr	4	0.14	0.01	0.07	0.01		
10	East Mersea (R. site)	tr	2	0.16	0.02	0.08	0.01		5c
11	Coston	tr	3	0.17	0.00	0.08	0.00		5c
12	Masey	tr	4	0.17	0.03	0.08	0.02		
13	Woolpack Farm	tr	1	0.17	0.00	0.09	0.00		
14	Jaywick Sands	tr	4	0.17	0.01	0.09	0.01		
15	Saham Toney, 52	tr	4	0.17	0.01	0.10	0.01		
16	Itteringham	tr	1	0.18	0.00	0.09	0.01		
17	Tattershall Castle	tr	4	0.18	0.01	0.09	0.01		5c
18	Bobbitshole	tr	3	0.18	0.01	0.10	0.00		5c
19	Shrotham	tr	4	0.18	0.01	0.09	0.01		5c
20	Trafalgar Square	tr	6	0.19	0.01	0.10	0.01		5c
15	Saham Toney, 94	tr	2	0.19	0.00	0.11	0.01		
21	Bardon Quarry, Area 3	tr	3	0.19	0.02	0.09	0.01		
15	Saham Toney, 93	tr	4	0.19	0.01	0.10	0.01		
16	Itteringham, Bed d	tr	4	0.19	0.01	0.10	0.00		
22	Eckington	tr	4	0.20	0.01	0.10	0.01		5c
23	Crophorne New Inn	tr	4	0.20	0.01	0.09	0.00		5c
21	Bardon Quarry, Area 2	tr	3	0.21	0.03	0.11	0.02		
24	Furtham's Lane East	tr	25	0.22	0.01	0.12	0.01		
25	Stanton Harcourt	tr	4	0.23	0.01	0.11	0.00		7
26	Crayford	tr	4	0.24	0.00	0.12	0.01		7
27	Barnwell	tr	2	0.24	0.01	0.12	0.01		
28	Alibone-on-Stour	tr	4	0.24	0.01	0.12	0.01		
29	Histon Road	tr	2	0.24	0.01	0.12	0.01		
30	Strensham	tr	4	0.25	0.02	0.12	0.02		
31	Stutton	tr	4	0.25	0.01	0.14	0.01		
32	Cotonation Farm	tr	1	0.25	0.00	0.13	0.00		7
33	Somersham	tr	4	0.25	0.01	0.12	0.00		
34	Norton Bottoms	tr	15	0.25	0.01	0.13	0.01		7
35	Block Fen	tr	4	0.25	0.02	0.13	0.01		
33	Somersham	tr	4	0.25	0.01	0.13	0.01		
36	Ebbesfleet, U. loam, U. part	tr	1	0.26	0.01	0.14	0.00		
36	Ebbesfleet, U. loam, L. part	tr	2	0.26	0.01	0.14	0.01		
37	Brough	tr	6	0.27	0.01	0.16	0.01		7
38	Ilford (Uphall Pit)	tr	2	0.27	0.01	0.14	0.02		7
39	Mastenhall	tr	2	0.28	0.02	0.15	0.00		
40	West Thurrock (Lion Pit), 4	tr	25	0.29	0.01	0.15	0.01		7
40	West Thurrock (Lion Pit), 2000	tr	3	0.29	0.01	0.16	0.00		7
36	Ebbesfleet, Lmst Brickearth	tr	3	0.30	0.01	0.15	0.01		
41	Stoke Tunnel	tr	2	0.30	0.01	0.15	0.01		
40	West Thurrock (Lion Pit), 3	tr	25	0.30	0.01	0.16	0.02		7
40	West Thurrock (Lion Pit), 6	tr	5	0.31	0.01	0.16	0.01		7
42	Setley	tr	3	0.31	0.01	0.16	0.00		
43	Aweley	tr	4	0.32	0.00	0.16	0.00		7
44	West Wittering	tr	3	0.32	0.00	0.16	0.00		
45	Bushley Green	tr	1	0.34	0.00	0.16	0.00		
46	Cudmore Grove	tr	4	0.34	0.02	0.19	0.01		9
47	Hackney Downs	tr	2	0.34	0.01	0.17	0.01		9
46	Cudmore Grove, sample PE	tr	2	0.36	0.00	0.20	0.00		9
48	Bethus Park	tr	1	0.36	0.02	0.20	0.02		9
49	Purfleet, 1	tr	23	0.38	0.01	0.18	0.01		9
50	Bairing	tr	4	0.38	0.01	0.20	0.01		9
49	Purfleet, 6	tr	24	0.38	0.01	0.21	0.01		9
51	Shoeburyness	tr	5	0.39	0.01	0.20	0.01		
52	Grays	tr	2	0.40	0.01	0.20	0.00		9
53	Marks Tey	tr	2	0.40	0.01	0.22	0.01		11
48	Bethus Park, BP18	tr	4	0.40	0.01	0.21	0.01		9
54	Trimingham	tr	13	0.40	0.01	0.22	0.01		
55	Hoxne, Stratum B2, 64	tr	4	0.40	0.02	0.22	0.01		11
56	Barnham, BEF92	tr	3	0.40	0.02	0.21	0.01		11
57	Elveden, ELV 96	tr	2	0.41	0.01	0.20	0.01		
57	Elveden, ELV 95	tr	5	0.41	0.01	0.21	0.01		
58	Swanscombe (Barnfield Pit)	tr	6	0.42	0.01	0.24	0.03		11
56	Barnham, BEF93	tr	6	0.42	0.02	0.20	0.02		11
59	Woodston	tr	4	0.42	0.01	0.23	0.01		11
55	Hoxne, Stratum E	tr	4	0.42	0.01	0.23	0.01		11
60	West Stow (Beeches Pit)	tr	4	0.42	0.01	0.25	0.01		11
61	Dierden's Pit (Ingress Vale)	tr	2	0.43	0.01	0.22	0.01		11
55	Hoxne, Stratum B2, 50	tr	4	0.43	0.01	0.24	0.01		11
62	Southfleet Road	tr	7	0.43	0.01	0.22	0.01		11
61	Dierden's Pit (Ingress Vale)	tr	4	0.45	0.01	0.21	0.01		11
63	Clacton-on-Sea	tr	4	0.46	0.01	0.25	0.02		11
64	Wilverley Wood	tr	6	0.48	0.03	0.27	0.03		137
65	Sidestrand, Upper Unio-Bed	tr	7	0.54	0.02	0.29	0.01		137
65	Sidestrand, Lower Unio-Bed	tr	5	0.55	0.01	0.29	0.01		137
66	Sugworth	tr	6	0.56	0.01	0.28	0.01		157
67	Little Oakley	tr	2	0.59	0.00	0.28	0.00		157
68	Pakefield, PaCl	tr	4	0.57	0.03	0.30	0.01		17/157
68	Pakefield, PaCl	tr	3	0.58	0.01	0.35	0.02		17/157
69	West Runton	tr	11	0.61	0.02	0.33	0.01		17/157
70	Bavel	tr	4	0.61	0.02	0.34	0.04		317
71	Weybourne (Weybourne Crag)	tr	2	0.89	0.01	0.69	0.03		
72	Tegelen	tr	4	0.90	0.01	0.75	0.03		
71	Weybourne (Weybourne Crag)	tr	1	0.92	0.00	0.79	0.01		
73	Thorpe Aldringham (N. Crag)	sp	1	0.94	0.00	0.84	0.04		
74	Frechen (Pliocene)	tr	4	0.93	0.01	0.91	0.02		




Figure 1. Amino acid decomposition at sites of known age

Comparison of racemization in *Bithynia* opercula for FAA aspartic acid (Asx), THAA alanine (Ala) and valine (Val) for all sites with independent geochronology. Y-axis error bars indicate one standard deviation about the mean for the site. X-axis error bars are not shown for the Early Pleistocene and Pliocene samples, as these estimates of age are not based on numerical methods. Note the rapid racemization in Asx at relatively young sites and the plateau beyond ~0.5 Ma; Asx is therefore most valuable for separating sites younger than MIS 9. In contrast, Val racemizes more slowly and provides poorer resolution for younger sites, but is able to differentiate between sites back to the Pliocene. Utilising several amino acids with different rates of degradation therefore enables greater age resolution. The increase in racemization is not linear with time, but slows during cold stages. Inset: shell (a) and operculum (b) of *Bithynia tentaculata*.




Figure 2. THAA vs FAA D/L Ala for individual samples in relation to terrace stratigraphy (a) Idealised transverse sections through the Lower Thames terrace sequence¹⁹ and (b) THAA vs FAA D/L Ala. Data from Thames sites (coloured) are superimposed on the full dataset. Note the concordance of terrace aggradations with the extent of protein degradation. This finding is consistent in all four river systems tested (Table 1, Supplementary Table 2). The relation of FAA:THAA is a useful control on the reliability of the closed-system. For example, opercula distorted following deposition showed levels of racemization of THAA far lower than predicted from FAA¹⁴, suggesting that during demineralization and distortion smaller, more mobile, amino acids were lost whereas larger peptides were retained during re-mineralization. Only ~1% of samples analysed here were compromised, such as the outlier from Sugworth (circled).




Figure 3. THAA vs FAA D/L Ala in relation to the occurrence of archaeology

Data from archaeological sites (coloured) are superimposed on the full dataset. Error bars represent two standard deviations about the mean for each site. Symbols as in Table 1. The earliest Palaeolithic archaeology (Mode 1, i.e. flake tools made with hard-hammers) comes from pre-Anglian contexts yielding *Mimomys savini*. Sites with unequivocal Clactonian (core-and-flake) archaeology all fall within MIS 11. Sites yielding Levallois artefacts form a temporally discrete cluster after MIS 9 but before MIS 5e. Archaeology is unknown at British sites attributed to the Last Interglacial (MIS 5e). Opercula have also been analysed from one Upper Palaeolithic and one Mesolithic site.

Table 1
Intra-crystalline amino acid data from the opercula of *Bithynia* from sites in southern Britain

Sites were chosen because they: (i) are stratotypes (in bold) of various interglacial stages; (ii) have independent geochronology; (iii) occur within a fluvial terrace sequence; or (iv) have relative ages based on biostratigraphy (Supplementary Data 1). Some additional sites, including three key continental Plio-Pleistocene localities, are listed to show how they fit into this general framework. Sites are listed in rank order based on the THAA Ala D/L value, the most useful single measurement covering the timescales under discussion but full interpretation requires consideration of the overall IcPD (Supplementary Data 1). Our data are shown alongside terrace stratigraphy (NW: Nene/Welland; SA: Severn/Avon; Th: Thames; TW: Trent/Witham), occurrences of important biostratigraphic indicator species (? = indirect association), *in situ* archaeology and existing consensus views on correlation with the MIS record. Age attributions reliant on amino acid dating alone are excluded. The occurrence of the water vole *Arvicola* is only shown for pre-Anglian sites (i.e. pre-MIS 12). ▲ archaeology found in overlying sediments; ▼ archaeology found in underlying sediments; × archaeology from the same horizon as the opercula analysed; ◆ indirect association (i.e. archaeology recorded from the site but not this profile). *Bithynia tentaculata* (L.) does not occur

throughout the British Pleistocene, so where necessary we have used other species of *Bithynia* (*b/t* = *B. bavelensis/tentaculata*; *le* = *B. leachii*; *te* = *B. tentaculata*; *tr* = *B. troscheli*).

Site number (SI Fig. 1)	Site	<i>Bithynia</i> species	Number analysed	Ala D/L		Val D/L		Independent geochronology	Terrace stratigraphy	Biostrat.		Archaeology					Consensus MIS	
				mean	standard deviation	mean	standard deviation			<i>Hippopotamys/Corbicula</i>	<i>Arvicola</i> (pre-Anglian) / <i>Mimomys</i>	Mesolithic	Upper Palaeolithic	Levallois	Acheulian	Clactonian		Undiagnostic
1	Acle (modern)	<i>le</i>	3	0.04	0.00	0.02	0.00	•										
1	Acle (modern)	<i>te</i>	2	0.04	0.00	0.02	0.00											
2	Enfield Lock	<i>te</i>	4	0.05	0.01	0.03	0.01											1
3	Quidenham Mere, 250-260cm	<i>te</i>	4	0.05	0.01	0.03	0.00											1
3	Quidenham Mere, 640-650cm	<i>te</i>	3	0.05	0.01	0.03	0.00											1
4	Newby Wiske	<i>te</i>	3	0.06	0.01	0.04	0.00	•										1
5	Aston-upon-Trent	<i>te</i>	4	0.06	0.00	0.04	0.00	•										1
6	Star Carr, 245-250cm	<i>te</i>	2	0.07	0.01	0.04	0.00	•			◇							1
7	Sproughton	<i>te</i>	4	0.09	0.00	0.05	0.00	•				◇						2
6	Star Carr, 524-528cm	<i>te</i>	3	0.10	0.02	0.04	0.00	•										2
8	Cassington	<i>te</i>	4	0.12	0.01	0.07	0.00	•	Th									5c-4
9	Isleworth	<i>te</i>	4	0.14	0.01	0.07	0.01		Th									
10	East Mersea (R. site)	<i>te</i>	2	0.16	0.02	0.08	0.01			H								5e
11	Coston	<i>te</i>	3	0.17	0.00	0.08	0.00			H								5e
12	Maxey	<i>te</i>	4	0.17	0.03	0.08	0.02		NW									
13	Woolpack Farm	<i>te</i>	1	0.17	0.00	0.09	0.00											
14	Jaywick Sands	<i>te</i>	4	0.17	0.01	0.09	0.01											
15	Saham Toney, 92	<i>te</i>	4	0.17	0.00	0.10	0.01											
16	Itteringham	<i>te</i>	1	0.18	0.00	0.09	0.01											
17	Tattershall Castle	<i>te</i>	4	0.18	0.01	0.09	0.01	•	TW									5e
18	Bobbitshole	<i>te</i>	3	0.18	0.01	0.10	0.00											5e
19	Shropham	<i>te</i>	4	0.18	0.01	0.09	0.01			(H)								5e
20	Trafalgar Square	<i>te</i>	6	0.19	0.01	0.10	0.01		Th	H								5e
15	Saham Toney, 94	<i>te</i>	2	0.19	0.00	0.11	0.01											
21	Bardon Quarry, Area 3	<i>te</i>	3	0.19	0.02	0.09	0.01		TW									
15	Saham Toney, 93	<i>te</i>	4	0.19	0.01	0.10	0.01											
16	Itteringham, Bed d	<i>te</i>	4	0.19	0.01	0.10	0.00											
22	Eckington	<i>te</i>	4	0.20	0.01	0.10	0.01		SA	H								5e
23	Crothorne New Inn	<i>te</i>	4	0.20	0.01	0.09	0.00		SA	H								5e
21	Bardon Quarry, Area 2	<i>te</i>	4	0.21	0.03	0.11	0.02											
24	Funtham's Lane East	<i>te</i>	25	0.22	0.01	0.12	0.01		NW	C								
25	Stanton Harcourt	<i>te</i>	4	0.23	0.01	0.11	0.00		Th	(C)					◇			7
26	Crayford	<i>te</i>	4	0.24	0.00	0.12	0.01		Th	C			◇					7
27	Barnwell	<i>te</i>	2	0.24	0.01	0.12	0.01			C								
28	Ailstone-on-Stour	<i>te</i>	4	0.24	0.01	0.12	0.01		SA	C								
29	Histon Road	<i>te</i>	2	0.24	0.01	0.12	0.01			(C)								
30	Strensham	<i>te</i>	4	0.25	0.02	0.12	0.02		SA									
31	Stutton	<i>te</i>	4	0.25	0.01	0.14	0.01			C			◇					
32	Coronation Farm	<i>te</i>	1	0.25	0.00	0.13	0.00		TW									7
33	Somersham	<i>te</i>	4	0.25	0.01	0.12	0.00			C			◇					
34	Norton Bottoms	<i>te</i>	15	0.25	0.01	0.13	0.01		TW	C								7

35	Block Fen	<i>te</i>	4	0.25	0.02	0.13	0.01			(C)								
33	Somersham	<i>tr</i>	4	0.26	0.01	0.13	0.01			C			◇					
36	Ebbsfleet, U. loam, U. part	<i>te</i>	1	0.26	0.01	0.14	0.00						↓					
36	Ebbsfleet, U. loam, L. part	<i>te</i>	2	0.26	0.01	0.14	0.01						↓					
37	Brough	<i>te</i>	6	0.27	0.01	0.15	0.01		TW									7
38	Ilford (Uphall Pit)	<i>te</i>	2	0.27	0.01	0.14	0.02		Th	C								7
39	Maidenhall	<i>te</i>	2	0.28	0.02	0.15	0.00						◇					
40	West Thurrock (Lion Pit), 4	<i>te</i>	25	0.29	0.01	0.15	0.01		Th	C			↓					7
40	West Thurrock (Lion Pit), 2000	<i>te</i>	3	0.29	0.01	0.16	0.00		Th	C			↓					7