

## ERRATA

# Mechanisms of Antibiotic Resistance Determined by Resistance-Transfer Factors

JOEL UNOWSKY AND MARTIN RACHMELER

*Department of Microbiology, Northwestern University Medical School, Chicago, Illinois*

Volume 92, no. 2, pages 361 and 362: Change the legends to Fig. 2, 3, 4, and 5 to read as follows.

FIG. 2. Uptake of CM by CM-resistant cells, CSH-2-F<sup>-</sup>-fi<sup>+</sup> (○), and sensitive cells, CSH-2-F<sup>-</sup> (●).

FIG. 3. Uptake of TC by TC-resistant strains, CSH-2-F<sup>-</sup>-fi<sup>-</sup> (▲), CSH-2-F<sup>-</sup>-fi<sup>+</sup> (■), *S. flexneri* 2b-fi<sup>+</sup> (○), and a TC-sensitive strain, CSH-2-F<sup>-</sup> (●). TC-resistant strain CSH-2-F<sup>-</sup>-fi<sup>-</sup> was preincubated in TC and then tested for TC uptake (□).

FIG. 4. Uptake of SM by *str-s* strains, CSH-2-F<sup>-</sup> (●) and 15·51 (■), and *str-r* strains, CSH-2-F<sup>-</sup>-fi<sup>+</sup> (○), CSH-2-F<sup>-</sup>-fi<sup>-</sup> (△), W-677 (▲), and 15·49 (□).

FIG. 5. Effect of CM on SM uptake by an SM- and

CM-sensitive strain, CSH-2-F<sup>-</sup>, and an SM- and CM-resistant strain, CSH-2-F<sup>-</sup>-fi<sup>+</sup>. Log-phase cultures were divided in half; to one portion was added 5 μg/ml of tritiated SM; to the other, 5 μg of tritiated SM plus 25 μg/ml of CM. The cultures were grown in nutrient broth on a shaker at 37 C; 1-ml samples were withdrawn, filtered, and washed and their radioactivity was counted. (○) SM uptake by CSH-2-F<sup>-</sup>; (●) SM uptake by CSH-2-F<sup>-</sup> in presence of CM; (△) SM uptake by CSH-2-F<sup>-</sup>-fi<sup>+</sup>; (▲) SM uptake by CSH-2-F<sup>-</sup>-fi<sup>+</sup> in the presence of CM.

## Effect of Bromouracil-containing Deoxyribonucleic Acid on *Bacillus subtilis*

DIXIE M. GIMLIN, SUE D. HARDMAN, BETTY N. KELLEY, GRACE C. BUTLER,  
AND FRANKLIN R. LEACH

*Department of Biochemistry, Oklahoma State University, Stillwater, Oklahoma*

Volume 92, no. 2, page 369, column 2: In the second line of the footnote to Table 3, change "75 μg per ml of DNA" to "2.5 μg per ml of DNA."

## Relation of Polysaccharide Content to Some Biological Properties of Endotoxins from Mutants of *Salmonella typhimurium*

R. W. I. KESSEL, HENRY H. FREEDMAN, AND WERNER BRAUN

*Institute of Microbiology, Rutgers, The State University, New Brunswick, New Jersey*

Volume 92, no. 3, page 594: Add to legend of Fig. 2, "Triangles, white circles, and black circles indicate data from separate experiments. The abscissa indicates endotoxin dose in μg/mouse."