

Published in final edited form as:


Org Lett. 2010 January 15; 12(2): 336–339. doi:10.1021/ol902681t.

Regioselective Intramolecular Dipolar Cycloaddition of Azides and Unsymmetrical Alkynes

Ryan A. Brawn, Morgan Welzel, Jason T. Lowe, and James S. Panek*

Department of Chemistry and Center for Chemical Methodology and Library Development, Metcalf Center for Science and Engineering, 590 Commonwealth Avenue, Boston University, Boston, Massachusetts 02215

Abstract



Enantioenriched allenylsilanes are used in three component propargylation reactions with aldehydes and silyl ethers to form *syn*-homopropargylic ethers that contain an imbedded azide. These materials then undergo thermally induced intramolecular 1,3-dipolar cycloaddition reactions, resulting in unique fused ring systems containing 1,2,3-triazoles. The ability to modify all three components of the reaction allows for expedient access to compounds containing significant structural and stereochemical variation.

Recent advancements in the Huisgen 1,3-dipolar cycloaddition reaction have led to a renewed interest in its application in the synthesis of 1,2,3-triazoles.¹ Small molecules containing a triazole functionality have been shown to exhibit a range of biological function, including antitumor, antibacterial, antiparasitic, and antiviral activity (Figure 1).²

Historically terminal and symmetrically substituted alkynes have been used predominantly in these reactions, since unsymmetrical alkynes often result in diminished reaction rates and poor regioselectivity.³ One solution to this problem is a tandem dipolar cycloaddition/cross coupling reaction, which can access fully substituted triazoles.⁴ Another approach is an intramolecular cycloaddition, where the regioselectivity is determined by the initial positions of the azide and alkyne in the starting material.⁵ While both of these strategies have proven to be useful, methods for the formation of fully functionalized triazoles remain underdeveloped.

We have recently reported a three component reaction of enantioenriched allenylsilanes with aldehydes and silyl ethers, resulting in highly functionalized *syn*-homopropargylic ethers.⁶ In our continued interest in developing these allenes as chiral carbon nucleophiles,⁷ we sought to further expand the use of the resulting homopropargylic ethers by incorporating a pendant azide moiety which will further react to provide heterocycles containing fused 1,2,3-triazoles. In this paper we describe the development of a three component propargylation/cycloaddition reaction to give access to densely functionalized fused triazole ring systems.

panek@bu.edu.

 Supporting Information Available Experimental data and selected spectral data for all new compounds. This material is available free of charge via the internet at <http://pubs.acs.org>.

The propargylation of enantioenriched allenylsilane (*R_a*)-**4a** with 2-azidobenzaldehyde⁸ and methoxytrimethylsilane (TMSOMe), promoted by TMSOTf, resulted in formation of alkyne **5** in 71% yield as a single observed diastereomer (Scheme 1). Heating alkyne **5** in toluene at 110 °C promoted the desired dipolar cyclization reaction, resulting in the formation of triazole **6a** in 90% yield. As anticipated, a single regioisomer of the desired triazole was observed.

A similar experiment was conducted with 2-azidoacetaldehyde dimethylacetal (Scheme 1). Due to concerns with volatility and stability, the preformed acetal was used instead of the aldehyde. The propargylation reaction provided homopropargylic ether **7**, and heating this product in toluene gave desired triazole **8** in 90% isolated yield. This reaction sequence demonstrates that the methodology can be expanded to aliphatic systems using acetals with imbedded azides.

When this strategy was applied to more substituted azidobenzaldehydes we observed that some of the triazole product was formed in the crude propargylation reaction mixture.⁸ Attempts to purify these propargylation products proved difficult, as the triazole product would be observed in post-chromatographic NMR spectra. This led to the development of a two step sequence where the propargylation reaction was quenched, and the crude product was heated to 70 °C in toluene, to directly produce the desired triazole product.

This modified procedure gave the desired tricyclic system with an imbedded triazole **6** in good yield and high diastereoselectivity (Scheme 2). The reactions were tolerant to a range of functionality on the aromatic ring, including electron donating and withdrawing groups. When achiral allenylsilane **4b** was subjected to the same reaction conditions, products **6f** and **6g** were formed in moderate yield.

Further structural variation was achieved by using silyl ether reaction partners containing an azide functionality (Table 1). The silyl protected 2-azidoethanol could be prepared from 2-bromoethanol in two steps using a known procedure.⁹ The TBS ether was explored after the TMS version proved to be difficult to work with due to its inherent volatility.

The propargylation reaction of allenylsilane (*R_a*)-**4** with aldehydes and silyl protected 2-azidoethanol resulted in the formation of *syn*-homopropargylic ethers **9a-i**. High yields and selectivities were obtained for a variety of aromatic aldehydes (entries 1–6). The use of aliphatic aldehydes in this reaction sequence resulted in lower yields (entries 7–9), and lower selectivity for the unbranched system (entry 8).

For all of the examples studied, heating the reaction products to 130 °C in toluene gave the desired triazoles **10a-i** in good yield.¹⁰ None of the triazole products were observed after the propargylation reaction, as the alkyne products were stable at room temperature, and as expected, the dipolar cyclizations proceeded with complete regioselectivity.

To expand the scope of this methodology, we synthesized four additional azidosilyl ethers (Scheme 3) which would allow access to fused triazole systems with greater complexity.¹¹ Silyl ether **11a** was prepared in four steps from (*S*)-ethyl lactate, and contains a chiral center on the carbon with the silyl ether. Silyl ether **11b**, derived from the epoxide opening of (*R*)-styrene oxide, has a chiral center on the carbon containing the azide. TBS protected 3-azidopropanol **11c** was prepared from 3-bromopropanol using the same procedure as TBS protected 2-azidoethanol. Finally, the 3-carbon silyl ether containing a chiral center at C2 (**11d**) was accessed using commercially available (*S*)-methyl-3-hydroxy-2-methylpropanoate in four steps. These procedures are all general and relatively straightforward, allowing for the rapid incorporation of stereochemical variation into the synthesis of the desired fused triazole systems.

Propargylation reactions with **11a** proceeded in good yield, but with a significant loss of diastereoselectivity when the (*R_a*) enantiomer of allenylsilane **4a** was used (Scheme 4, **12a+b**). When the (*S_a*) enantiomer of the allenylsilane was used the diastereoselectivity was substantially higher (Scheme 4, **12c+d**), exhibiting a “matched pair” of reaction partners in a double stereodifferentiating reaction. The cycloadditions proceeded with moderate to high yield, although the products of matched cases (**13c+d**) showed higher yields than the mismatched cases (**13a+b**).¹²

When the silyl ether's chiral center was located on the carbon bearing the azide the diastereoselectivity of the reaction products was not effected by switching the enantiomer of the allene. Propargylations with silyl ether **11b** gave high yields and diastereoselectivities with both enantiomers (Scheme 4, **12e-h**). The cycloadditions also proceeded in similar yields for the different diastereomers (Scheme 4, **13e-h**), with moderate yields obtained in all cases.

Propargylations with the three-carbon silyl ether **11c** proceeded with good yield and excellent diastereoselectivity (Scheme 4, **12i+j**). While the reactions were sluggish in toluene, switching the solvent to chlorobenzene at 150 °C resulted in significantly higher yields for **13i+j**. Use of silyl ether **11d**, which contained a stereocenter at C2, had no effect on the selectivity, as the products were formed in good yields as a single observed diastereomer regardless of which enantiomer of allenylsilane **4a** was used (Scheme 4, **12k+l**).

In conclusion we have reported a tandem propargylation/dipolar cycloaddition sequence to rapidly construct structurally and stereochemically diverse fused ring systems containing 1,2,3-triazoles. Continued exploration of these ring systems, along with preliminary biological evaluation, is currently in progress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support was obtained from the NIH CA 53604 and through an AstraZeneca graduate fellowship to RAB.

References

- (a) Huisgen, R. 1,3-Dipolar Cycloaddition Chemistry. Padwa, A., editor. Wiley; New York: 1984. (b) Kolb HC, Finn MG, Sharpless KB. *Angew Chem Int Ed.* 2001; 40:2004–2021. (c) Rostovtsev VV, Green LG, Fokin VV, Sharpless KB. *Angew Chem Int Ed.* 2002; 41:2596–2599. (d) Kelly AR, Wei J, Kesavan S, Marié JC, Windmon N, Young DW, Marcaurelle LA. *Org Lett.* 2009; 11:2257–2260. [PubMed: 19473044] (e) Spiteric, C.; Moses, JE. *Angew Chem Int Ed.* 2009. p. 48
- (a) Lewis WG, Green LG, Grynszpan F, Radić Z, Carlier PR, Taylor P, Finn MG, Sharpless KB. *Angew Chem Int Ed.* 2002; 41:1053–1057. (b) Tron GC, Piralí T, Billington RA, Canonico PL, Sorba G, Genazzani AA. *Med Res Rev.* 2008; 28:278–308. [PubMed: 17763363] (c) Lee T, Cho M, Ko SY, Youn HJ, Baek DJ, Cho WJ, Kang CY, Kim S. *J Med Chem.* 2007; 50:585–589. [PubMed: 17266209] (d) Maurya SK, Gollapalli DR, Kirubakaran S, Zhang M, Johnson CR, Benjamin NN, Hedstrom L, Cuny GD. *J Med Chem.* 2009; 52:4623–4630. [PubMed: 19624136] (e) Kolb HC, Sharpless KB. *Drug Discovery Today.* 2003; 8:1128–1137. [PubMed: 14678739] (f) Puig-Basagoiti F, Qing M, Dong H, Zhang B, Zou G, Yuan Z, Shi PY. *Antiviral Research.* 2009; 83:71–79. [PubMed: 19501258]
- (a) Zhang L, Chen X, Xue P, Sun HHY, Williams ID, Sharpless KB, Fokin VV, Jia G. *J Am Chem Soc.* 2005; 127:15998–15999. [PubMed: 16287266] (b) Majireck MM, Weinreb SM. *J Org Chem.* 2006; 71:8680–8683. [PubMed: 17064059]

4. (a) Ackermann L, Potukuchi HK, Landsberg D, Vicente R. *Org Lett.* 2008; 10:3081–3084. [PubMed: 18549230] (b) Chowdhury C, Mukherjee S, Das B, Achari B. *J Org Chem.* 2009; 75:3612–3615. [PubMed: 19344130]
5. (a) Pearson WH, Bergmeier SC, Chytra JA. *Synthesis.* 1990:156–159. (b) Oliva AI, Christmann U, Font D, Cuevas F, Ballester P, Buschmann H, Torrens A, Yenes S, Pericás MA. *Org Lett.* 2008; 10:1617–1619. [PubMed: 18348569] (c) Balducci E, Bellucci L, Petricci E, Taddei M, Tafi A. *J Org Chem.* 2008; 74:1314–1321. [PubMed: 19138079]
6. Brawn RA, Panek JS. *Org Lett.* 2007; 9:2689–2692. [PubMed: 17559219]
7. For a recent review on chiral organosilanes in diversity oriented synthesis see: Shaw J. *Nat Prod Rep.* 2009; 26:11–26. [PubMed: 19374120]
8. For the synthesis of functionalized 2-azidobenzaldehydes see: (a) Pelkey ET, Gribble GW. *Tetrahedron Lett.* 1997; 38:5603–5606. (b) Main CA, Petersson HA, Rahman SS, Hartley RC. *Tetrahedron.* 2008; 64:901–914. (c) Cuevas JC, De Mendoza J, Prados P. *J Org Chem.* 1988; 53:2055–2066.
9. Smith RH, Mehl AF, Shantz DL, Chmurny GN, Michejda CJ. *J Org Chem.* 1988; 53:1467–1471.
10. All of the dipolar cycloaddition reactions were run in a sealed tube. No reactivity was observed using the copper (I) catalyst systems which have been previously reported.
11. For the synthesis of these silyl ethers: see Supporting Information.
12. Masamune S, Choy W, Petersen JS, Sita LR. *Angew Chem, Int Ed.* 1985; 24:1.






Figure 1. Representative examples of biologically active molecules containing 1,2,3-triazoles.

**Scheme 1.**


Two step triazole formation

^a Isolated yields after purification over silica gel. ^b Diastereomeric ratios determined by ¹H NMR analysis on crude material.


**Scheme 2.**

One-pot propargylation/cycloaddition

^a Isolated yields after purification over silica gel. ^b Diastereomeric ratios determined by ¹H NMR analysis on crude material. ^c Reaction run using achiral allenylsilane **4b**.



Scheme 3.
Silyl ether synthesis

**Scheme 4.**


Propargylations and dipolar cycloadditions with substituted silyl ethers

^a All yields refer to isolated products after purification over silica gel. ^b All diastereomeric ratios were determined by ¹H NMR analysis on crude material.


^c The (*S_a*) enantiomer of the allenylsilane **1** was used for the propargylation reaction. ^dCycloaddition reaction run in chlorobenzene at 150 °C.

Table 1

Propargylations with TBS 2-azidoethanol



entry	aldehyde (R)	product 9 (yield ^a ; dr ^b)	product 10 (yield ^a)
1		9a (72%; 18:1)	10a (71%)
2		9b (89%; >20:1)	10b (75%)
3		9c (76%; 15:1)	10c (76%)
4		9d (87%; >20:1)	10d (74%)
5		9e (74%; >20:1)	10e (66%)
6		9f (76%; 15:1)	10f (74%)



entry	aldehyde (R)	product 9 (yield ^a ; dr ^b)	product 10 (yield ^a)
7		9g (60%; >20:1)	10g (75%)
8		9h (37%; 5:1)	10h (57%)
9		9i (16%, >20:1)	10i (80%)

^a Isolated yields after purification over silica gel.

^b Diastereomeric ratios determined by ¹H NMR analysis on crude material.