

Isoleucine 309 acts as a C₄ catalytic switch that increases ribulose-1,5-bisphosphate carboxylase/oxygenase (rubisco) carboxylation rate in *Flaveria*

Spencer M. Whitney^{a,1}, Robert E. Sharwood^{a,2}, Douglas Orr^a, Sarah J. White^a, Hernan Alonso^a, and Jeroni Galmés^b

^aResearch School of Biology, Australian National University, Canberra ACT 2601, Australia; and ^bGrup de Recerca en Biologia de les Plantes en Condicions Mediterrànies Universitat de les Illes Balears, 07122 Mallorca, Spain

Edited by George H. Lorimer, University of Maryland, College Park, MD, and approved July 27, 2011 (received for review June 13, 2011)

Improving global yields of important agricultural crops is a complex challenge. Enhancing yield and resource use by engineering improvements to photosynthetic carbon assimilation is one potential solution. During the last 40 million years C₄ photosynthesis has evolved multiple times, enabling plants to evade the catalytic inadequacies of the CO₂-fixing enzyme, ribulose-1,5-bisphosphate carboxylase/oxygenase (rubisco). Compared with their C₃ ancestors, C₄ plants combine a faster rubisco with a biochemical CO₂-concentrating mechanism, enabling more efficient use of water and nitrogen and enhanced yield. Here we show the versatility of plastome manipulation in tobacco for identifying sequences in C₄-rubisco that can be transplanted into C₃-rubisco to improve carboxylation rate (V_C). Using transplastomic tobacco lines expressing native and mutated rubisco large subunits (L-subunits) from *Flaveria pringlei* (C₃), *Flaveria floridana* (C₃-C₄), and *Flaveria bidentis* (C₄), we reveal that Met-309-Ile substitutions in the L-subunit act as a catalytic switch between C₄ (³⁰⁹Ile; faster V_C , lower CO₂ affinity) and C₃ (³⁰⁹Met; slower V_C , higher CO₂ affinity) catalysis. Application of this transplastomic system permits further identification of other structural solutions selected by nature that can increase rubisco V_C in C₃ crops. Coengineering a catalytically faster C₃ rubisco and a CO₂-concentrating mechanism within C₃ crop species could enhance their efficiency in resource use and yield.

CO₂ assimilation | *rbcL* mutagenesis | gas exchange | chloroplast transformation

The future uncertainties of global climate change and estimates of unsustainable population growth have increased the urgency of improving crop yields (1). One possible solution is to “supercharge” photosynthesis by improving the C₃ cycle (2, 3). Although a simple idea, this is a complex challenge that involves several possible alternatives. Many of these alternatives focus on enhancing the performance of the CO₂-fixing enzyme ribulose-1,5-bisphosphate (RuBP) carboxylase/oxygenase (rubisco), which catalyzes the first step in the synthesis of carbohydrates. Despite its pivotal role, rubisco is a slow catalyst, completing only one to four carboxylation reactions per catalytic site per second in plants (4, 5). Moreover CO₂ not only is fixed through a complex catalytic process but also must compete with O₂. The oxygenation of RuBP produces 2-phosphoglycolate, whose recycling by photorespiration requires energy and results in the futile loss of fixed carbon [~30% of fixed CO₂ in many C₃ plants (6)].

To compensate for rubisco’s catalytic limitations, plants invest as much as 25% of their leaf nitrogen in rubisco (7). This value is much lower in C₄ plants, where a biochemical CO₂-concentrating mechanism (CCM) elevates CO₂ around rubisco. This optimized microenvironment allows rubisco to operate close to its maximal activity, reducing O₂ competition. This CCM has enabled C₄ plants to evolve rubiscos with substantially improved carboxylation rates (V_C) relative to their C₃ ancestors, albeit at the expense of reducing CO₂ affinity [i.e., a higher apparent K_m for CO₂ (K_C)] (8, 9). As a consequence, C₄ plants require less rubisco, thereby enhancing nitrogen use with improvements in

V_C correlating with improved efficiency in nitrogen use (10). In addition, the high concentration of CO₂ around rubisco allows C₄ plants to operate at lower CO₂ pressures within their leaf air spaces, thereby reducing their stomatal conductance requirements and the associated H₂O loss by transpiration. Indicative of these growth advantages, C₄ photosynthesis has evolved independently several times from multiple C₃ lineages during the last 20–40 million years (11, 12).

Following nature’s example, a number of CO₂ transgenic approaches have been designed to emulate CCM strategies in C₃ plastids and improve rubisco performance (2). These approaches include elevating the CO₂ concentration within chloroplasts using recombinant CO₂/HCO₃[−] transporters from cyanobacteria or engineering alternative pathways to bypass photorespiration and release CO₂ within the stroma (13, 14). Although each strategy faces continuous challenges in its fine tuning and integration into crops, further improvements in yield and in the efficiency of water and nitrogen use are likely by concurrently “speeding up” rubisco (9).

Identifying the natural changes that result in the faster C₄ rubiscos is far from simple, given the complex structure and biogenesis pathway of the hexadecameric rubisco (L₈S₈) in vascular plants, whose assembly requirements cannot be met by conventional bacterial or in vitro expression systems (15). The catalytic core of L₈S₈ rubisco comprises four 52-kDa large (L)-subunit pairs which are capped by two sets of 15-kDa small (S)-subunit tetramers that provide structural stability and influence catalysis (16, 17). Although supplementing rice rubisco with S-subunits from the C₄ plant sorghum was found to improve V_C of the heterologous L₈S₈ enzyme (16), crosses between C₃ and C₄ *Flaveria* and *Atriplex* species showed C₄ catalysis to be maternally inherited (18, 19) and hence defined by the chloroplast-encoded L-subunit gene (*rbcL*). Therefore, changes in both Rubisco L- and S-subunits can influence catalysis.

Although phylogenetic studies have identified potential L-subunit residues involved in the transition from C₃-like to C₄-like rubisco, it is uncertain which residues are catalytically determinant (11, 20). Here we undertake a transplastomic approach to identify such residues in vivo. By manipulating the *rbcL* gene in tobacco to produce hybrid L₈^FS₈^T rubiscos (containing variant *Flaveria* L- and tobacco S-subunits), we demonstrate that Met-309-Ile substitutions in the L-subunit act as

Author contributions: S.M.W. designed research; S.M.W., R.S., D.O., S.J.W., H.A., and J.G. performed research; S.M.W. and J.G. analyzed data; and S.M.W., R.S., H.A., and J.G. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: Spencer.Whitney@anu.edu.au.

²Present address: Hawkesbury Institute for the Environment, University of Western Sydney, Richmond NSW 2753, Australia.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1109503108/-DCSupplemental.

a catalytic switch between C₄ (³⁰⁹Ile) and C₃ (³⁰⁹Met) catalysis in *Flaveria rubisco*.

Results

Flaveria Rubisco L-Subunit Expression in Tobacco Chloroplasts. The *rbcL* genes from *Flaveria pringlei* (C₃), *Flaveria floridana* (a C₃-C₄ intermediate), and *Flaveria bidentis* (C₄) were chosen for transformation into tobacco plastids because of their diverse catalytic properties (ref. 21 and below) despite their high sequence similarity (Fig. 1A). As in all *Flaveria rbcL* genes, nonsilent nucleotide changes occur only at residues 149, 265, and 309 (20). The *Flaveria* L-subunits show >95% identity with the tobacco L-subunit, with 22–24 amino acid differences in addition to a highly charged TDKDKDKKR extension at the C terminus (Fig S1). The *Flaveria rbcL* genes were cloned into the plastome-transforming plasmid pLEV4, where the expression of the transgene is regulated by the native tobacco *rbcL* gene regulatory sequences [i.e., its promoter, 5', and 3'-untranslated sequences (22)]. The transforming plasmids, including the control pLEV4, were introduced biolistically into ^{cm}trL, a tailor-made tobacco master line for integrating *rbcL* transgenes into the tobacco chloroplast by homologous recombination (Fig. 1B) (23). Two independent transplastomic lines for each *rbcL* transgene were grown to maturity in soil in air supplemented with 0.5% (vol/vol) CO₂. The transformed tobacco lines that incorporated the *F. floridana*, *F. bidentis*, and *F. pringlei rbcL* genes were called “*tob^{flo}*,” “*tob^{bid}*,” and “*tob^{pring}*,” respectively.

Nondenaturing PAGE (ndPAGE) analysis of the soluble leaf protein was used to confirm the production of the hybrid L₈^FS₈^t rubisco (comprising *Flaveria* L-subunits and tobacco S-subunits) and to assess the homoplasmy of the *tob^{pring}*, *tob^{flo}*, and *tob^{bid}* lines (Fig. 1C). Each of the transformed lines examined was deemed to be homoplasmic, because no L₂ *Rhodospirillum rubrum* rubisco from the parental ^{cm}trL line was detected; homoplasmy was further confirmed by DNA blot analysis (Fig S2).

Differential Expression of the Hybrid L₈^FS₈^t Rubiscos. Differences in the intensities of the L₈^FS₈^t bands in ndPAGE indicated that the amounts of rubisco produced in the T₀ lines varied (Fig. 1C). This variation was confirmed by quantitative [¹⁴C]2-carboxyarabinitol-1,5-bisphosphate (CABP) binding. The amount of L₈^FS₈^t produced in the *tob^{flo}*, *tob^{bid}*, and *tob^{pring}* leaves was reduced by approximately 50%, 65%, and 75% relative to wild-type, respectively (Fig. 1C). In contrast, rubisco content in the *tob^{LEV4}* control transformants matched that in wild-type, indicating that the additional genome changes around *rbcL* were not the cause of the reduced L₈^FS₈^t expression. SDS PAGE-immunoblot analysis showed no unassembled *Flaveria* L-subunits accumulated as insoluble aggregates.

Contrary to the reduced L₈^FS₈^t content in the T₀ leaves, there was little or no difference in the *Flaveria rbcL* mRNA levels in the same leaves relative to wild type (Fig. 1D), indicating that L₈^FS₈^t synthesis probably is perturbed posttranscriptionally. As shown previously (23–25), a less abundant *rbcL-aadA* dicistronic mRNA (~10% that of the *rbcL* mRNA) was produced in all transformants as a result of inefficient transcription termination by the tobacco *rbcL* 3'UTR. The stages that hinder L₈^FS₈^t expression during rubisco biogenesis or degradation remain to be identified fully.

Plant Growth and Leaf Photosynthesis. The disparity in L₈^FS₈^t levels in leaves of the *tob^{flo}*, *tob^{bid}*, and *tob^{pring}* lines persisted to the T₂ progeny and correlated with their relative differences in photosynthesis and growth rates. For the *tob^{flo}* and *tob^{bid}* plants, the higher L₈^FS₈^t levels produced were sufficient for them to survive through to maturity in air (without CO₂ enrichment), although they grew more slowly than wild-type plants (Fig. 2A). In contrast, the *tob^{pring}* transformants grew poorly in air. As seen




Fig. 1. *Flaveria* rubisco L-subunit sequence and expression in tobacco chloroplasts. (A) Comparison of *Flaveria rbcL* sequences that differ only in L-subunit substitutions at amino acids 149, 265, and 309 (20). (B) The transforming plasmid pLEV4 contains a homologous plastome flanking sequence (indicated by dashed lines; numbering indicates region of sequence integration relative to plastome sequence; GenBank ID Z00444) that directed integration of the *rbcL* transgenes and a promoterless *aadA*-selectable marker gene into the plastome of the tobacco master line, ^{cm}trL (23). The L-subunit amino acid differences at residues 149 (black triangle), 265 (black triangle), and 309 (gray triangle) in the *tob^{LEV4}* (tobacco *rbcL* control), *tob^{pring}* (*F. pringlei rbcL*), *tob^{flo}* (*F. floridana rbcL*), *tob^{bid}* (*F. bidentis rbcL*), *tob^{flo-bid}* (chimeric *F. floridana*-*F. bidentis rbcL*), *tob^{bid-flo}* (chimeric *F. bidentis*-*F. floridana rbcL*), and *tob^{149A,265I}* (mutated tobacco *rbcL*) transplastomic tobacco (*tob*) lines are shown. Annealing locations of primers LSH, LSE, and the 5' $\Delta rbcL$ probe (24) are shown. N, NheI; S, Sall cloning sites. (C) Nondenaturing PAGE analysis of soluble protein from comparable leaves of independent T₀ transplastomic lines, ^{cm}trL, and wild-type tobacco (protein from 1.5 mm² of leaf was loaded per lane). Homoplasmic transformants produce only L₈S₈ rubisco (~520 kDa) and not the ~100-kDa *R. rubrum* L₂ rubisco produced in ^{cm}trL (23). (D) Detection of *rbcL* and *rbcL-aadA* mRNA in total RNA from 3 mm² (for wild type) or 6 mm² (other samples) of the leaves sampled in C.

previously in tobacco ^{Rst} lines producing hybrid L₈^FS₈^t rubisco (comprising sunflower L- and tobacco S-subunits) (24), the juvenile *tob^{pring}* plants displayed a pale green leaf phenotype with marginal curling and dimpling (Fig. 2A Right). This phenotype is




Fig. 2. Measurements of growth and leaf gas exchange in the transformants producing the variant L₈^FS₈^T rubiscos. (A) (Left) Comparatively slower growth in air of the tob^{Flaveria} transformants as a function of plant height relative to wild-type. (Right) The tob^{pring} lines grew extremely poorly in air. pce, post-coty ledon emergence. (B) Phenotype of the plants from A at age pce as shown. (Scale bars: 4 cm.) Air supplemented with 0.5% (vol/vol) CO₂ was used to grow tob^{pring} plants to maturity. (C) Comparative differences in gas-exchange measurements of CO₂ assimilation rates at 25 °C under varying chloroplast CO₂ pressures (C_c) at growth illumination (400 μmol quanta m⁻²·s⁻¹). Measurements were made on young mature leaves located at similar canopy positions (fifth leaf from the apical meristem) of physiologically comparable mature plants analogous to those shown in B. Leaf rubisco contents were 25.0 and 30.5; 12.4 and 11.9; 10.1 and 10.8; 4.1 and 4.3 μmol rubisco sites m⁻² in the independent wild-type (circles), tob^{flor} (squares; line 1 white, line 2 black), tob^{bid} (triangles; line 1 white, line 2 black), and tob^{pring} (diamonds; line 1 white, line 2 black) plants analyzed, respectively. The lines show the rubisco limited CO₂ assimilation rates for wild-type (---), tob^{flor} (—), tob^{bid} (···), and tob^{pring} (— · —) modeled according to ref. 26 using the catalytic parameters for the respective hybrid L₈^FS₈^T rubiscos in Fig. 3 and assuming rubisco was fully activated and a value of 0.3 mol m⁻²·s⁻¹·bar⁻¹ for mesophyll conductance.

likely a consequence of the very low L₈^FS₈^T content during early vegetative growth (<3 μmol sites m⁻²·s⁻¹) and in the young mature leaves (<6 μmol sites m⁻²·s⁻¹; Fig. 2C) of the tob^{pring} plants. Consistent with their different growth rates and varied L₈^FS₈^T contents, the leaf photosynthetic CO₂ assimilation rates at varying CO₂ partial pressures (pCO₂) were slowest for tob^{pring} [but still slightly higher than in tobacco^{Rst} (24)] and were successively better for the tob^{bid} and tob^{flor} lines, albeit still slower

than in wild-type tobacco (Fig. 2C). Measurements of the ratio of variable fluorescence to maximal fluorescence (F_v/F_m) in wild-type leaves (0.82 ± 0.01) were identical to those in the three tob^{Flaveria} genotypes (0.83 ± 0.01), indicating no difference in photochemical efficiency under the growth conditions.

Catalysis by Each L₈^FS₈^T Rubisco Matches the Source *Flaveria* Enzyme.

The catalytic properties of the recombinant L₈^FS₈^T were compared with the source L₈^FS₈^F enzymes from the corresponding *Flaveria* species (Fig. 3). As seen previously for the L₈^SS₈^T rubisco produced in tobacco^{Rst} (24), the L₈^FS₈^T and equivalent *Flaveria* L₈^FS₈^F enzymes were catalytically comparable with respect to their carboxylation (V_C) and oxygenation (V_O) rates, their apparent Michaelis constants (K_m) for CO₂ (K_C) and O₂ (K_O), and CO₂/O₂ specificities (S_{C/O}). Of particular interest was the faster carboxylation rate (~35% higher V_C relative to the C₃ rubiscos) and lower CO₂ affinity (~50% higher K_C) of the hybrid L₈^FS₈^T rubisco from tob^{bid} that matched the *F. bidentis* L₈^FS₈^F enzyme (Fig. 3). As expected, when the catalytic properties and content of each L₈^FS₈^T hybrid enzyme were used to model CO₂ assimilation rates according to Farquhar et al (26), the final values closely matched those measured by whole-leaf gas exchange (Fig. 2C). These results provide confidence both in the accuracy of the measured catalytic properties of the L₈^FS₈^T hybrid enzymes and the rubisco-limited CO₂-assimilation model.

Interchanging C₃-C₄ Catalysis via ³⁰⁹Met-³⁰⁹Ile Substitutions in *Flaveria* Rubisco.

The catalytic similarity between native *F. bidentis* rubisco and the hybrid tob^{bid} L₈^FS₈^T enzyme indicated that the introduced L-subunit determined the catalytic phenotype. Because the *F. bidentis* and *F. floridana* L-subunits differ only at residues 149 and 309 but show significant differences in V_C and K_C (Fig. 3), domain swapping of their *rbcL* was used to identify which residue(s) imparted the C₄ catalysis of *F. bidentis* rubisco. The chimeric *rbcL* gene in the transforming plasmid pLEV^{flor-bid} introduced a Met-309-Ile substitution into the *F. floridana* *rbcL* gene, whereas in pLEV^{bid-flor} the chimeric *rbcL* gene coded an Ile-309-Met substitution in the *F. bidentis* *rbcL* gene (Fig. 1B). Both plasmids were transformed into ^{cm}trL, and

Plant line	Rubisco structure	L-subunit residue			K _C (μM)						V _C (s ⁻¹)						K _O (μM)		S _{C/O} (mol·mol ⁻¹)		V _O (s ⁻¹)	
		149	265	309	0	10	20	0	1	2	3	4	5	0	1	0	1					
<i>N. tabacum</i>	L ₈ ^S S ₈	Q	V	V	12.6 ± 0.2	12.6 ± 0.2	12.6 ± 0.2	3.2 ± 0.2	3.2 ± 0.2	3.2 ± 0.2	3.2 ± 0.2	3.2 ± 0.2	3.2 ± 0.2	3.2 ± 0.2	274 ± 18	81 ± 1	0.8	0.8				
tob ^{149A,265I}	L ₈ ^S S ₈	A	I	I	12.2 ± 0.5	12.2 ± 0.5	12.2 ± 0.5	3.4 ± 0.2	3.4 ± 0.2	3.4 ± 0.2	3.4 ± 0.2	3.4 ± 0.2	3.4 ± 0.2	3.4 ± 0.2	306 ± 33	82 ± 1	1.1	1.1				
<i>F. pringlei</i> (C ₃)	L ₈ ^F S ₈ ^F	E	V	M	13.7 ± 0.5	13.7 ± 0.5	13.7 ± 0.5	3.5 ± 0.3	3.5 ± 0.3	3.5 ± 0.3	3.5 ± 0.3	3.5 ± 0.3	3.5 ± 0.3	376 ± 35	81 ± 1	1.0	1.0					
tob ^{pring}	L ₈ ^F S ₈				13.0 ± 0.4	13.0 ± 0.4	13.0 ± 0.4	3.5 ± 0.3	3.5 ± 0.3	3.5 ± 0.3	3.5 ± 0.3	3.5 ± 0.3	3.5 ± 0.3	317 ± 30	80 ± 2	1.1	1.1					
<i>F. floridana</i> (C ₃ -C ₄)	L ₈ ^F S ₈ ^F	D	I	M	14.4 ± 0.5	14.4 ± 0.5	14.4 ± 0.5	3.6 ± 0.1	3.6 ± 0.1	3.6 ± 0.1	3.6 ± 0.1	3.6 ± 0.1	3.6 ± 0.1	374 ± 33	82 ± 2	1.1	1.1					
tob ^{flor}	L ₈ ^F S ₈				14.5 ± 0.3	14.5 ± 0.3	14.5 ± 0.3	3.7 ± 0.2	3.7 ± 0.2	3.7 ± 0.2	3.7 ± 0.2	3.7 ± 0.2	3.7 ± 0.2	359 ± 22	81 ± 2	1.2	1.2					
<i>F. bidentis</i> (C ₄)	L ₈ ^F S ₈ ^F	A	I	I	20.4 ± 0.5	20.4 ± 0.5	20.4 ± 0.5	4.8 ± 0.3	4.8 ± 0.3	4.8 ± 0.3	4.8 ± 0.3	4.8 ± 0.3	4.8 ± 0.3	420 ± 37	81 ± 1	1.2	1.2					
tob ^{bid}	L ₈ ^F S ₈				19.9 ± 0.6	19.9 ± 0.6	19.9 ± 0.6	4.7 ± 0.2	4.7 ± 0.2	4.7 ± 0.2	4.7 ± 0.2	4.7 ± 0.2	4.7 ± 0.2	408 ± 28	79 ± 2	1.2	1.2					
tob ^{flor-bid}	L ₈ ^F S ₈ ^T	D	I	I	19.3 ± 0.3	19.3 ± 0.3	19.3 ± 0.3	4.6 ± 0.2	4.6 ± 0.2	4.6 ± 0.2	4.6 ± 0.2	4.6 ± 0.2	4.6 ± 0.2	393 ± 20	80 ± 1	1.2	1.2					
tob ^{bid-flor}	L ₈ ^F S ₈ ^T	A	I	M	13.8 ± 0.3	13.8 ± 0.3	13.8 ± 0.3	3.7 ± 0.2	3.7 ± 0.2	3.7 ± 0.2	3.7 ± 0.2	3.7 ± 0.2	3.7 ± 0.2	372 ± 24	81 ± 1	1.2	1.2					

Fig. 3. Comparative catalysis at 25 °C of the wild-type and Q¹⁴⁹A-V²⁶⁵I mutated tobacco (L₈^SS₈) rubisco, the source *Flaveria* (L₈^FS₈^F) rubiscos, the hybrid (L₈^FS₈^T, comprising *Flaveria* L- and tobacco S-subunits), and chimeric (L₈^FS₈^T, comprising chimeric *Flaveria* L- and tobacco S-subunits) rubisco variants produced in the transplastomic tobacco plants. The L-subunit amino acid residues at codons 149, 265, and 309 in each rubisco type are shown. Values shown are the average ± SD of independent assays (n = 4–24; see Materials and Methods for details). The maximal oxygenation rate (V_O) was calculated using the equation S_{C/O} = (V_C/K_C)/(V_O/K_O).

sertion of a more hydrophobic ³⁰⁹Ile this far from the active site might influence V_C (20).

Despite the improvements in V_C imparted by ³⁰⁹Ile in hybrid $L_8^F S_8^T$ rubisco, the accompanying reductions in CO_2 affinity (i.e., increased K_C) precluded gains in carboxylation efficiency. At 25 °C under ambient oxygen levels, the carboxylation efficiency (i.e., $V_C/K_C^{21\%O_2}$) of the C_4 -like ³⁰⁹Ile-containing rubiscos in $to b^{bid}$ and $to b^{flo-bid}$ ($145 \text{ mM}^{-1} \cdot \text{s}^{-1}$) were poorer than the C_3 -like ³⁰⁹Met- $L_8^F S_8^T$ enzymes in $to b^{pring}$ and $to b^{flo}$ (150 and $161 \text{ mM}^{-1} \cdot \text{s}^{-1}$, respectively). Thus, because of the low CO_2 levels within (unstressed) C_3 chloroplasts ($<10 \mu\text{M}$), the faster C_4 -like enzymes probably provide no advantage to plant growth within a C_3 plant (at least at 25 °C), as shown recently in rice (16). As modeled recently, optimal CO_2 concentrations required for C_4 rubisco are substantially higher ($\sim 80 \mu\text{M}$) (8), indicating that taking full advantage of a faster rubisco in a C_3 plant will require the combinatorial effect of a suitable CCM, for which a number of strategies are being pursued (2). Some of these approaches already have demonstrated that elevating CO_2 pressures within C_3 plastids can improve the capacity for CO_2 assimilation by reducing the energy costs of photorespiration (13, 14).

Our results suggest that the carboxylation rate of rubisco in a C_3 plant might be increased either by direct replacement with L-subunits sourced from C_4 plants (as in the $to b^{bid}$ plants; Fig. 3) or by tailoring appropriate sequence mutations into related C_3 rubisco L-subunits (as in the $to b^{flo-bid}$ plants; Fig. 3). Although the first approach suffers from our inability to predict a priori the assembly properties of foreign rubiscos within the chloroplast of the recipient transplastomic line, the second approach would require knowing the catalytic/structural effect of every possible mutation within the context of a particular rubisco enzyme, an understanding that we are still far from achieving. Indeed, the finding that tobacco rubisco encodes ³⁰⁹Ile but shows C_3 catalysis highlights the complex natural variation in the sequence–structure–function relationships among plant L-subunits. Even the sequence diversity at position 309 among C_4 rubiscos (Fig. 4A) indicates that this residue is not the only one that can impart C_4 catalysis. This result is consistent with the polyphyletic evolution of C_4 photosynthesis (12) and with predictions that at least eight L-subunit residues (including residue 309 but not residue 149) have been selected for positively by C_4 catalysis (11). Experimentally testing these predictions, identifying other catalytically determinant L-subunit residues, and exploring which particular rubiscos are affected by these changes have been hampered by the preferential location of *rbcl* in the plastome (27) and the small range of species whose plastomes can be transformed stably (28). However, as shown in this and previous studies (24, 29, 30), these experimental limitations may be circumvented by expressing hybrid rubiscos in tobacco plastids. The generality of this system for examining sequence–performance relationships within otherwise inaccessible, catalytically diverse foreign L-subunits remains to be explored fully.

Although this study demonstrates the pervasive role of the L-subunit in shaping catalysis in plant rubisco, the important role of the S-subunits on catalysis cannot be overlooked. The apparent catalytic neutrality of the tobacco S-subunit when assembled with heterologous L-subunits (Fig. 3) (24, 29, 30) contrasts with the recent success in shaping rice rubisco toward C_4 -like catalysis using heterologous S-subunits from C_4 sorghum (16). Similarly, structural changes to the S-subunit have improved *Chlamydomonas* rubisco catalysis (17). As highlighted recently (20), differences in rubisco S-subunit sequence also may account for the catalytic deviation of *Flaveria palmeri* rubisco, whose L-subunit sequence matches that of $to b^{flo-bid}$ (coding ³⁰⁹Ile) but shows C_3 catalysis.

The similar $S_{C/O}$ values determined for rubisco from *F. bidentis*, *F. floridana*, and *F. pringlei* (Fig. 3) in this study contrast with the slightly varying values determined previously ($S_{C/O} =$

76 ± 1 , 84 ± 1 , and 81 ± 1 respectively) (21). The reason for this variation is unknown but may lie in alterations in the catalytic competence as a result of different purification processes (ion exchange chromatography versus ammonium sulfate fractionation), the final enzyme purity, and the length of ultra-cold storage (24). By using fresh rubisco rapidly purified to $>95\%$ homogeneity by ion exchange chromatography, our measured $S_{C/O}$ values were highly reproducible between preparations from independent biological replicates.

Here we present an *rbcl* engineering approach involving hybrid rubisco production in tobacco plastids to unravel the sequence and catalytic diversity of related C_3 and C_4 rubiscos from *Flaveria*. Future applications of this experimental system are focused on identifying sequence changes that account for the natural diversity of rubisco performance and testing the feasibility of transplanting these catalytic improvements into the rubisco L-subunits of agriculturally relevant crops. In particular, when coengineered with biotechnological strategies to elevate CO_2 around rubisco in C_3 plants, a faster rubisco may translate into improved efficiency in water and nitrogen use and the enhanced yields currently enjoyed by C_4 plants.

Materials and Methods

RuBP and [¹⁴C]-CABP were synthesized as described (31, 32). Protein content was measured using a dye-binding assay (Pierce) and BSA as a protein-standard.

Tobacco Plastome Transformation and Growth. The transforming plasmid pLEV4 directs the insertion of an *rbcl* gene and a promoterless *aadA* gene (coding spectinomycin resistance) into the tobacco plastome in place of the L_2 *Rhodospirillum rubrum* rubisco-coding ^{cm}*rbclM* gene in the plastome of the tobacco master line ^{cm}trL (Fig. 1B) (23). The *rbcl* gene from *F. bidentis*, *F. pringlei*, and *F. floridana* was PCR amplified from leaf genomic DNA [isolated using the DNeasy Plant Mini Kit (Qiagen)] with the primers 5' Nhelrbcl (5'-AGCTAGCGTTGGATTCAAAGCTGGTGT-3' [the NheI site that spans the *rbcl* codons 9 (Ala) and 10 (Ser) is shown in italics] and 3' Sallrbcl (5'-TGTCGACTGTTTTATCTCTCTTATCTTATCT-3' [the reverse complement of the *rbcl* stop codon is shown in bold, and the Sall site is shown in italics]). The 1,439-bp NheI–Sall *rbcl* products were cloned into pLEV4 to give the transforming plasmids pLEV^{pring}, pLEV^{flo}, and pLEV^{bid}. The plasmids pLEV^{flo-bid} and pLEV^{bid-flo} were made by interchanging the 569-bp SphI–Sall fragments of the *F. bidentis* and *F. floridana* *rbcl* genes (Fig. 1B). Mutations coding substitutions Gln-149-Ala and Val-265-Ile in the tobacco *rbcl* gene in pLEV4 were introduced using the QuikChange Site-Directed Mutagenesis Kit (Stratagene) to produce plasmid pLEV^{tob}^{149A,265I}. All plasmids were sequenced using BigDye terminator sequencing at the Biomolecular Resource Facility, Australian National University (Canberra, Australia).

Each of the pLEV-derived plasmids was introduced biolistically into three leaves of ^{cm}trL1 as described (23), and three to seven independent spectinomycin-resistant plants were obtained for each mutant. Two independent plastome-transformed lines for each introduced *rbcl* gene were grown to maturity in soil in a growth atmosphere supplemented with 0.5% (vol/vol) CO_2 as described (24). At each generation the plants were fertilized artificially with wild-type pollen.

PAGE and Nucleotide Blot Analyses. The preparation and analysis of soluble leaf protein by SDS/PAGE, nondenaturing PAGE, and immunoblot analysis was performed as described (33). Total leaf genomic DNA was isolated using the DNeasy Plant Mini Kit and used to PCR amplify and sequence the transformed plastome region using primers 5'-CTATGGAATTCGAACCT-GAACTT-3' (LSH) and 5'-GAGGTGTGATACTGGCTTGATTC-3' (LSE) (Fig. 1B) (24). DNA blot analysis of the genomic DNA was used to confirm homoplasmy (Fig S1). Total RNA was extracted from 0.5 cm² of leaf in 0.8 mL TRIzol (Invitrogen). Six per cent or 12% of the RNA was separated on denaturing formaldehyde gels (34). The RNA was blotted onto Hybond-N nitrocellulose membrane (GE Healthcare) and probed with a ³²P-labeled 5' Δ *rbcl* probe (Fig. 1B) as described (24).

Rubisco Content and Catalytic Assessments. Rates of rubisco ¹⁴ CO_2 fixation using soluble leaf protein extract were measured in 7-mL septum-capped scintillation vials in reaction buffer [50 mM Hepes-NaOH (pH 7.8), 10 mM $MgCl_2$, 0.5 mM RuBP] containing varying concentrations of $NaH^{14}CO_3$ (0–67 μ M)

and O₂ (0–25%) (vol/vol), accurately mixed with nitrogen using Wostoff gas-mixing pumps as described (24, 33). Assays (0.5 mL total volume) were started by the addition of activated leaf protein, and the Michaelis constants (K_m) for CO₂ (K_C) and O₂ (K_O) were determined from the fitted data. Replicate measurements ($n = 4–8$) were made using protein preparations from two to four different leaves of independently transformed lines. For each sample the maximal rate of carboxylation (V_C) was extrapolated from the Michaelis–Menten fit and then normalized by dividing the rate by the number of rubisco-active sites quantified by [¹⁴C]2-CABP binding (35, 36). Rubisco CO₂/O₂ specificity (S_{CO}) was measured as described (37), using freshly extracted rubisco, quickly purified by ion exchange chromatography (24), from at least two separate plants for each independently transformed line.

Growth and Photosynthesis Analysis. Wild-type (*Nicotiana tabacum* L. Petit Havana) and transplastomic tobacco lines were grown in growth chambers at 25 °C and $400 \pm 50 \mu\text{mol photons m}^{-2}\text{s}^{-1}$ as described (24) in air or 0.5% (vol/vol) CO₂-enriched air. Plant height from the soil surface to the apical meristem was measured until the first floral apertures emerged. Leaf photosynthesis and dark respiration rates in plants of comparable physiological development (45–50 cm in height; 14 or 15 leaves) were made in the growth chamber using an LI-6400 gas-exchange system (LI-COR) (24). The maximum quantum efficiency of PSII in dark-adapted leaves [variable fluorescence (F_v)/maximum fluorescence (F_m)] was measured in the same leaves using an LI-6400–40 Leaf Chamber Fluorometer.

ACKNOWLEDGMENTS. This research was supported by Australian Research Council Discovery Grant FT0991407 (to S.M.W.) and by project AGL2009-07999 (Plan Nacional, Spain) (to J.G.).

- Edgerton MD (2009) Increasing crop productivity to meet global needs for feed, food, and fuel. *Plant Physiol* 149:7–13.
- Raines CA (2011) Increasing photosynthetic carbon assimilation in C₃ plants to improve crop yield: Current and future strategies. *Plant Physiol* 155:36–42.
- von Caemmerer S, Evans JR (2010) Enhancing C₃ photosynthesis. *Plant Physiol* 154:589–592.
- Andersson I, Backlund A (2008) Structure and function of Rubisco. *Plant Physiol Biochem* 46:275–291.
- Parry M, Madgwick P, Carvalho J, Andralojc P (2007) Prospects for increasing photosynthesis by overcoming the limitations of Rubisco. *J Agric Sci* 145:31–43.
- Zhu X-G, Portis AR, Long SP (2004) Would transformation of C₃ crop plants with foreign Rubisco increase productivity? A computational analysis extrapolating from kinetic properties to canopy photosynthesis. *Plant Cell Environ* 27:155–165.
- Evans JR (1989) Photosynthesis and nitrogen relationships in leaves of C₃ plants. *Oecologia* 78:9–19.
- Savir Y, Noor E, Milo R, Tlustý T (2010) Cross-species analysis traces adaptation of Rubisco toward optimality in a low-dimensional landscape. *Proc Natl Acad Sci USA* 107:3475–3480.
- Whitney SM, Houtz RL, Alonso H (2011) Advancing our understanding and capacity to engineer nature's CO₂-sequestering enzyme, Rubisco. *Plant Physiol* 155:27–35.
- Ghannoum O, et al. (2005) Faster Rubisco is the key to superior nitrogen-use efficiency in NADP-malic enzyme relative to NAD-malic enzyme C₄ grasses. *Plant Physiol* 137:638–650.
- Christin P-A, et al. (2008) Evolutionary switch and genetic convergence on *rbcl* following the evolution of C₄ photosynthesis. *Mol Biol Evol* 25:2361–2368.
- Sage RF, Christin PA, Edwards EJ (2011) The C₄ plant lineages of planet Earth. *J Exp Bot* 62:3155–3169.
- Kebeish R, et al. (2007) Chloroplastic photorespiratory bypass increases photosynthesis and biomass production in *Arabidopsis thaliana*. *Nat Biotechnol* 25:593–599.
- Liemman-Hurwitz J, Rachmilevitch S, Mittler R, Marcus Y, Kaplan A (2003) Enhanced photosynthesis and growth of transgenic plants that express *ictB*, a gene involved in HCO₃⁻ accumulation in cyanobacteria. *Plant Biotechnol J* 1:43–50.
- Mueller-Cajar O, Whitney SM (2008) Directing the evolution of Rubisco and Rubisco activase: First impressions of a new tool for photosynthesis research. *Photosynth Res* 98:667–675.
- Ishikawa C, Hatanaka T, Misoo S, Miyake C, Fukayama H (2011) Functional incorporation of sorghum small subunit increases the catalytic turnover rate of rubisco in transgenic rice. *Plant Physiol* 156:1603–1611.
- Spreitzer RJ, Peddi SR, Satagopan S (2005) Phylogenetic engineering at an interface between large and small subunits imparts land-plant kinetic properties to algal Rubisco. *Proc Natl Acad Sci USA* 102:17225–17230.
- Berry JA (1983) Genetic control of the kinetic parameters of RuP2 carboxylase: Studies of a C₃ and a C₄ species and their F hybrid. *Carnegie Institution of Washington Annual Report* 82:96–99.
- Hudson GS, et al. (1990) Comparisons of *rbcl* genes for the large subunit of ribulose-bisphosphate carboxylase from closely related C₃ and C₄ plant species. *J Biol Chem* 265:808–814.
- Kapralov MV, Kubien DS, Andersson I, Filatov DA (2011) Changes in Rubisco kinetics during the evolution of C₄ photosynthesis in *Flaveria* (Asteraceae) are associated with positive selection on genes encoding the enzyme. *Mol Biol Evol* 28:1491–1503.
- Kubien DS, Whitney SM, Moore PV, Jesson LK (2008) The biochemistry of Rubisco in *Flaveria*. *J Exp Bot* 59:1767–1777.
- Alonso H, Blayney MJ, Beck JL, Whitney SM (2009) Substrate-induced assembly of *Methanococcoides burtonii* D-ribulose-1,5-bisphosphate carboxylase/oxygenase dimers into decamers. *J Biol Chem* 284:33876–33882.
- Whitney SM, Sharwood RE (2008) Construction of a tobacco master line to improve Rubisco engineering in chloroplasts. *J Exp Bot* 59:1909–1921.
- Sharwood RE, von Caemmerer S, Maliga P, Whitney SM (2008) The catalytic properties of hybrid Rubisco comprising tobacco small and sunflower large subunits mirror the kinetically equivalent source Rubiscos and can support tobacco growth. *Plant Physiol* 146:83–96.
- Whitney SM, Andrews TJ (2003) Photosynthesis and growth of tobacco with a substituted bacterial Rubisco mirror the properties of the introduced enzyme. *Plant Physiol* 133:287–294.
- Farquhar G, vonCaemmerer S, Berry J (1980) A biochemical-model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* 149:78–90.
- Kanevski I, Maliga P (1994) Relocation of the plastid *rbcl* gene to the nucleus yields functional ribulose-1,5-bisphosphate carboxylase in tobacco chloroplasts. *Proc Natl Acad Sci USA* 91:1969–1973.
- Maliga P, Bock R (2011) Plastid biotechnology: Food, fuel, and medicine for the 21st century. *Plant Physiol* 155:1501–1510.
- Sharwood R, Whitney S (2010) *The Chloroplast: Basics and Applications*, ed Ca R (Springer Science, Dordrecht, The Netherlands), pp 285–306.
- Zhang XH, et al. (2011) Hybrid Rubisco of tomato large subunits and tobacco small subunits is functional in tobacco plants. *Plant Sci* 180:480–488.
- Kane HJ, Wilkin JM, Portis AR, Andrews TJ (1998) Potent inhibition of ribulose-bisphosphate carboxylase by an oxidized impurity in ribulose-1,5-bisphosphate. *Plant Physiol* 117:1059–1069.
- Pierce J, Tolbert NE, Barker R (1980) Interaction of ribulosebisphosphate carboxylase/oxygenase with transition-state analogues. *Biochemistry* 19:934–942.
- Whitney SM, Sharwood RE (2007) Linked Rubisco subunits can assemble into functional oligomers without impeding catalytic performance. *J Biol Chem* 282:3809–3818.
- Sambrook J, Fritsch EF, Maniatis T (2000) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).
- Ruuska S, et al. (1998) The interplay between limiting processes in C₃ photosynthesis studied by rapid-response gas exchange using transgenic tobacco impaired in photosynthesis. *Aust J Plant Physiol* 25:859–870.
- Whitney SM, Andrews TJ (2001) Plastome-encoded bacterial ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO) supports photosynthesis and growth in tobacco. *Proc Natl Acad Sci USA* 98:14738–14743.
- Kane HJ, et al. (1994) An improved method for measuring the CO₂/O₂ specificity of ribulosebisphosphate carboxylase-oxygenase. *Aust J Plant Physiol* 21:449–461.
- Christin P-A, et al. (2011) Complex evolutionary transitions and the significance of C₃-C₄ intermediate forms of photosynthesis in *Molluginaceae*. *Evolution* 65:643–660.