

# Cell Polarity in *Saccharomyces cerevisiae* Depends on Proper Localization of the Bud9 Landmark Protein by the EKC/KEOPS Complex

Yu Kato,<sup>1</sup> Hiroshi Kawasaki,<sup>2</sup> Yoshifumi Ohyama, Takashi Morishita,<sup>3</sup> Hiroshi Iwasaki,<sup>4</sup> Tetsuro Kokubo, and Hisashi Hirano

Department of Supramolecular Biology, Graduate School of Nanobioscience, Yokohama City University, Tsurumi-ku, Yokohama 230-0045, Japan

**ABSTRACT** In diploid *Saccharomyces cerevisiae* cells, bud-site selection is determined by two cortical landmarks, Bud8p and Bud9p, at the distal and proximal poles, respectively. Their localizations depend on the multigenerational proteins Rax1p/Rax2p. Many genes involved in bud-site selection were identified previously by genome-wide screening of deletion mutants, which identified *BUD32* that causes a random budding in diploid cells. Bud32p is an atypical kinase involved in a signaling cascade of Sch9p kinase, the yeast homolog of Akt/PKB, and a component of the EKC/KEOPS (endopeptidase-like, kinase, chromatin-associated/kinase, putative endopeptidase, and other proteins of small size) complex that functions in telomere maintenance and transcriptional regulation. However, its role in bipolar budding has remained unclear. In this report, we show that the Sch9p kinase cascade does not affect bipolar budding but that the EKC/KEOPS complex regulates the localization of Bud9p. The kinase activity of Bud32p, which is essential for the functions of the EKC/KEOPS complex but is not necessary for the Sch9p signaling cascade, is required for bipolar bud-site selection. *BUD9* is necessary for random budding in each deletion mutant of EKC/KEOPS components, and *RAX2* is genetically upstream of EKC/KEOPS genes for the regulation of bipolar budding. The asymmetric localization of Bud9p was dependent on the complex, but Bud8p and Rax2p were not. We concluded that the EKC/KEOPS complex is specifically involved in the regulation of Bud9p localization downstream of Rax1p/Rax2p.

**T**HE generation of cell polarity is important for the function of many cell types and underlies various processes such as cell division, differentiation, cell migration, cell–cell signaling, and fertilization (Freifelder 1960; Mooseker 1985; Hyman and White 1987; Bedinger *et al.* 1994; Chant and Pringle 1995; Kraut *et al.* 1996). In the budding yeast *Saccharomyces cerevisiae*, such polarization induces asym-

metric growth to form a bud that becomes the daughter cell. Yeast cells polarize and divide by budding in two patterns: the axial pattern of haploid  $a/\alpha$ -cells and the bipolar pattern of diploid  $a/\alpha$ -cells. Axial budding in the haploid  $a/\alpha$ -cells forms chain-like buds adjacent to the immediately preceding bud site, while bipolar budding in diploid  $a/\alpha$ -cells forms cluster-like buds at either the proximal pole or its opposite distal pole. Alternate bud-site selections between proximal and distal poles usually occur in bipolar budding.

The establishment of yeast cell polarity undergoes three basic steps (Drubin and Nelson 1996). First, site selection occurs on the cell surface, a spatial landmark where cells will polarize. The bud sites during axial budding are marked with the axial landmarks Axl1p, Axl2p, Bud3p, and Bud4p (Fujita *et al.* 1994; Chant *et al.* 1995; Halme *et al.* 1996; Sanders and Herskowitz 1996). The bipolar landmarks Bud8p and Bud9p of the distal and proximal poles, respectively, determine the polarization axis of budding in diploid cells (Harkins *et al.* 2001). These cortical landmarks provide recognition sites for the Rsr1p/Bud2p/Bud5p GTPase

Copyright © 2011 by the Genetics Society of America  
doi: 10.1534/genetics.111.128231

Manuscript received March 1, 2011; accepted for publication May 17, 2011

Supporting information is available online at <http://www.genetics.org/content/suppl/2011/05/30/genetics.111.128231.DC1>.

<sup>1</sup>Present address: Laboratory for Immunochaperons, Research Center for Allergy and Immunology (RCAI), RIKEN Yokohama Institute, Suehiro-cho 1-7-22, Tsurumi-ku, Yokohama 230-0045, Japan

<sup>2</sup>Corresponding author: Department of Supramolecular Biology, Graduate School of Nanobioscience, Yokohama City University, Suehiro-cho 1-7-29, Tsurumi-ku, Yokohama 230-0045, Japan. E-mail: [kawasaki@yokohama-cu.ac.jp](mailto:kawasaki@yokohama-cu.ac.jp)

<sup>3</sup>Present address: Genome Damage and Stability Centre, University of Sussex, Falmer, Brighton, East Sussex BN1 9RQ, United Kingdom.

<sup>4</sup>Present address: Department of Life Science, Graduate School of Bioscience and Biotechnology, Tokyo Institute of Technology, Nagatsuda-cho 4259 B-8, Midori-ku, Yokohama 256-8501, Japan.

signaling module (Kang *et al.* 2001; Park and Bi 2007). Next, the GTPase module transmits positional information from the axial and bipolar cortical markers to the protein Cdc42p GTPase and its guanine nucleotide-exchange factor Cdc24p for polarity establishment. Finally, Cdc42p recruits the machinery that organizes and polymerizes actin, actin-associated proteins, and septins to the selected site of growth.

Numerous proteins have been found to be involved in bud-site selection (Park and Bi 2007). A genome-wide screening of homozygous deletion mutants identified 127 mutants representing three different bud-site phenotypes: unipolar, axial-like, and random (Ni and Snyder 2001). Among them, 112 mutants displayed strong or weak random budding phenotypes. The homozygous deletion mutant of the gene *BUD32* displayed a random budding pattern (Ni and Snyder 2001). *BUD32* was originally found as a gene encoding an atypical protein kinase that belongs to the piD261 family of atypical Ser/Thr protein kinases found in virtually all eukaryotic and archaeal organisms (Stocchetto *et al.* 1997). Recently, two independent studies of haploid cells have shown that Bud32p is a component of the KEOPS (kinase, putative endopeptidase, and other proteins of small size) or the EKC (endopeptidase-like, kinase, chromatin-associated) complexes with functions of telomere maintenance and transcriptional regulation, respectively (Downey *et al.* 2006; Kisseleva-Romanova *et al.* 2006). Cgi121p, a component of the KEOPS complex, was identified by a genome-wide screen as a suppressor of *cdc13-1*, an allele of the gene encoding the telomere-capping protein Cdc13p (Downey *et al.* 2006). The components of the KEOPS complex containing Cgi121p are the protein kinase Bud32p, the putative peptidase Kae1p, and the uncharacterized protein Gon7p. Deletion of *BUD32* or genes encoding other KEOPS components produced phenotypes with short telomeres and failed to synthesize *de novo* telomeres to DNA double-stranded breaks (Downey *et al.* 2006). The EKC complex contains a homolog of cancer-testis antigens (Pcc1p) and four additional proteins found in the KEOPS complex. Pcc1p is a transcription factor, and its deletion affects the expression of several genes regulated by  $\alpha$ -factor and galactose (Kisseleva-Romanova *et al.* 2006). Bud32p is phosphorylated by Sch9p kinase, a yeast member of the Akt/PKB subfamily. The phosphorylation of Bud32p positively regulates its ability to interact with Grx4p and to phosphorylate it (Peggon *et al.* 2008). The signaling pathway from Akt/PKB (homolog of Sch9p) to PRPK (homolog of Bud32p) is evolutionarily conserved. Sch9p kinase regulates the nutrient signaling pathway in yeast (Roosen *et al.* 2005). The function of Bud32p in this pathway is independent of the role in the EKC/KEOPS complex (Peggon *et al.* 2008).

Genes that are specifically linked to the random budding of diploid cells encode factors important for the localization of bipolar cortical landmarks (Casamayor and Snyder 2002). Diploid cells select two poles for bipolar budding at the poles proximal or distal to the birth scar, which are deter-

mined by two cortical landmarks, Bud8p and Bud9p, respectively. *BUD8* and *BUD9* were originally identified as specific genes essential for normal bipolar budding. The *bud8 $\Delta$*  and *bud9 $\Delta$*  mutants bud at the proximal and distal poles, respectively (Harkins *et al.* 2001). In addition, Bud8p and Bud9p are localized at the distal and proximal poles, respectively (Harkins *et al.* 2001). Bud9p is also localized at the distal pole, whose localization may inhibit the function of Bud8p through an interaction between the two (Taheri *et al.* 2000). Previous study suggests that many of the genes that specifically cause random budding, when deleted, are associated with the localization of Bud8p (Ni and Snyder 2001). Rax1p and Rax2p are interdependently related in the maintenance of bipolar budding (Kang *et al.* 2004). The genes encoding these proteins were identified by screening genes that restored bipolar budding patterns in the haploid *axl1 $\Delta$*  mutant (Fujita *et al.* 1994). Bud8p and Bud9p interact with and depend on Rax1p and Rax2p, respectively, for their localization (Kang *et al.* 2004). The delivery of Bud8p and Bud9p to the presumptive bud site and mother-bud neck, respectively, is dependent on actin. The delivery of Bud9p is also dependent on septin (Schenkman *et al.* 2002).

In this article, we report that the Sch9p kinase cascade does not associate with bipolar budding, but that the EKC/KEOPS complex is involved in bipolar budding. The deletion mutants of each EKC/KEOPS component displayed a random budding pattern, and the phenotype was suppressed by the deletion of *BUD9*, suggesting that Bud9p is a cause of random budding in these mutants. Consistent with phenotypic analysis, the asymmetric localization of GFP-Bud9p was dependent on the EKC/KEOPS complex, but GFP-Bud8p was not. The localization of Bud9p is also regulated interdependently by Rax1p/Rax2p. The phenotype of double mutants of each EKC/KEOPS gene and *RAX2* was changed to one similar to the *rax2 $\Delta$*  mutant, which showed a random budding pattern specifically in diploid cells. The localization of Rax2p was normal even in *bud32 $\Delta$*  and other mutant cells with deleted EKC/KEOPS components. These results suggest that the EKC/KEOPS complex is required for the localization of Bud9p and acts as a downstream factor of Rax2p.

## Materials and Methods

### Yeast strains, culture medium, plasmids, and recombinant DNA methods

Homozygous deletion mutants were generated by mating *MATa* cells with *MAT $\alpha$*  cells or by transformation of the plasmid expressing the HO (Homothallic switching) endonuclease in haploid cells, which is expected to cause mating-type switching of *MATa* cells to *MAT $\alpha$*  or *MAT $\alpha$*  cells to *MATa* and hence subsequent mating between *MATa* and *MAT $\alpha$*  cells to form a homozygous diploid. In the supporting information, Table S1 summarizes the yeast strains used in this study. Yeast cells were grown in yeast extract peptone dextrose [YPD: 2% (w/v) bacto-peptone, 1% (w/v) yeast extract, and 2% (w/v) glucose] or in synthetic defined

(SD) medium [0.2% (w/v) yeast nitrogen base without amino acids and ammonium sulfate, 0.5% (w/v) ammonium sulfate, and 2% (w/v) glucose] or in synthetic complete (SC) medium [0.2% (w/v) yeast nitrogen base without amino acids and ammonium sulfate, 0.5% (w/v) ammonium sulfate, 2% (w/v) glucose, and complete amino acids] lacking appropriate amino acids to maintain various plasmids. Plasmid preparation was conducted with *Escherichia coli* XL10 GOLD (Stratagene, La Jolla, CA) grown in Luria-Bertani medium [0.5% (w/v) NaCl, 1% (w/v) yeast extract, and 1% (w/v) tryptone] containing 50  $\mu\text{g}\cdot\text{ml}^{-1}$  ampicillin. Solid media were made with 2% agar. Table S2 summarizes the plasmids used in this study. Standard methods of yeast genetics and DNA manipulation (Gietz *et al.* 1992; Ausubel *et al.* 1995; Longtine *et al.* 1998) were used, except where noted. Polymerase chain reaction (PCR) was performed using KOD FX DNA polymerase (Nacalai Tesque, Kyoto, Japan). Oligonucleotide primers shown in Table S3 were synthesized and purchased from Invitrogen (Tokyo).

### Plasmids

The pYC01 was created with pRS425 (Christianson *et al.* 1992) by inserting a fragment containing an upstream sequence of 808 bp, an open reading frame of *BUD8*, and a downstream sequence of 517 bp. PCR was performed to amplify the *BUD8* region with the primer set BUD8a/BUD8b and the genomic DNA of the BY4743 yeast strain as a template. The resulting *XhoI/SpeI* fragment was cloned into *XhoI/SpeI*-digested pRS425. Detailed procedures for the preparation of pYC03 to insert epitope tags using a *MluI* cleavable site (5'-ACGCGT-3') just downstream of the start codon of *BUD8* have been reported (Kato *et al.* 2009). For green fluorescent protein (GFP) tagging, PCR was performed using the primer set GFPa/GFPb and pFA6a-GFP<sup>S65T</sup>-kanMX6 (Longtine *et al.* 1998) as a template. The resulting *MluI* fragment was incubated with T4 polynucleotide kinase (T4 PNK) (Takara Shuzo, Kyoto, Japan) and then ligated with *EcoRV*-digested pBluescript (SK-) to create pYC05. *MluI*-digested pYC05 was separated using 1.5% agarose gel electrophoresis and gel-extracted. The resulting insert was ligated with *MluI*-digested pYC03 to generate pYC14. The GFP-Bud9p fragment was cloned into pRS426 (Christianson *et al.* 1992) using the same procedure to generate the pYC06.

The pYC09 was constructed to delete *GON7*. PCR was first performed to amplify the *GON7* region with the primer set GON7a/GON7b, with genomic DNA of the BY4743 yeast strain as the template. The resulting fragment containing an upstream 570 bp, an open reading frame of *GON7*, and a downstream 519 bp was phosphorylated with T4 PNK and then cloned in *EcoRV*-digested pBluescript (SK-) to create pYC07. Deletion of the open reading frame of *GON7* and creation of a *MluI* restriction site to generate pYC08 were performed by PCR using the primer set GON7c/GON7d and pYC07 as a template. For cloning the *URA3* region as a selective marker, PCR was next performed using the primer set

URA3a/URA3b and genomic DNA of the B8032 strain as the template. The resulting *MluI* fragment containing an upstream 871 bp, an open reading frame of *URA3*, and a downstream 1000 bp was digested with *MluI* and then cloned into *MluI*-digested pYC08 to create pYC09.

For FLAG tagging, PCR was performed using a primer set, FLAGa/FLAGb, and pTM55 as the template. The resulting fragment was digested with *MluI* and then ligated with *MluI*-digested pYC04 to create pYC10.

The pYC11 was created with pBluescript (SK-) by inserting a fragment containing an upstream sequence of 520 bp, an open reading frame of *BUD32*, and a downstream sequence of 789 bp. PCR was performed to amplify the *BUD32* region with the primer set BUD32a/BUD32b and the genomic DNA of the BY4743 yeast strain as a template. The resulting *SpeI/XhoI* fragment was cloned into *SpeI/XhoI*-digested pBluescript (SK-) to generate pYC11. Site-directed mutagenesis was performed to insert a *MluI* cleavable site (5'-ACGCGT-3') just upstream of the stop codon of *BUD32* with primer set BUD32c/BUD32d and pYC11 as a template to generate pYC12.

Plasmids pYC13 and pYC15 were created with pBluescript (SK-) by replacing base pairs of *BUD32* to mutate to alanine at Lys52 and Ser258 of *Bud32p*, respectively. PCR was performed to replace base pairs with the primer sets BUD32e/BUD32f, BUD32g/BUD32h, and pYC12 as a template to generate pYC13 and pYC15, respectively. The pYC13 was then digested with *SpeI/XhoI*, and the insert of pYC13 was transferred to pRS306 to generate pYC16.

*MluI*-digested pYC05 was separated using 1.5% agarose gel electrophoresis and gel-extracted. The resulting insert was ligated with *MluI*-digested pYC12, pYC13, and pYC15 to generate pYC17, pYC18, and pYC19, respectively. The resulting plasmids were then digested with *SpeI/XhoI*, and the inserts of these plasmids were transferred to pRS306 to generate pYC20, pYC21 and pYC22, respectively. All sequences were verified using an ABI 3010 sequencer (Applied Biosystems, Foster City, CA).

### Strains

Strains expressing *Rax2p*-GFP, CCY032, and CCY033 were created by a one-step tagging method using PCR product (Longtine *et al.* 1998). A cassette of GFP<sup>S65T</sup>-kanMX6 just upstream of the stop codon of *RAX2* using the primer set *RAX2*-GFPa/*RAX2*-GFPb and pFA6a-GFP<sup>S65T</sup>-kanMX6 as a template was amplified by PCR. The BY4741 and BY4742 strains were transformed with the resulting fragments to create strains CCY032 and CCY033, respectively. The transformants were incubated on YPD plates at 30° for 1 day and then selected to YPD plates with G418 (100  $\mu\text{g}\cdot\text{ml}^{-1}$ ). The correct insertion of GFP<sup>S65T</sup>-kanMX6 was verified by PCR using the genomic DNA of candidates as the template. The haploid strains CCY032 and CCY033 were mated on YPD plates for 1 day at 30° and then selected on SC plates not containing Lys and Met for 2 days at 30° to obtain the diploid strain expressing *Rax2p*-GFP (CCY034).

The *GON7* deletion strains (CCY015, CCY016, CCY 022, and CCY025) were created by homologous recombination with PCR products from pYC09. PCR was performed using the primer set GON7a/GON7b and pYC09 as the template. The resulting fragment was transformed in *bud32Δ* (CCY001), *cgi121Δ* (CCY004), *bud8Δ* (CCY010), and *bud9Δ* (CCY012) mutants, respectively, and then the transformants were grown on SC-Ura for 3 days at 25°. Alternatively, the one-step disruption method using PCR products was used to delete *GON7* in *RAX2-GFP* (CCY034) or *rax2Δ* (CCY044) strains. PCR was performed using the primer set GON7e/GON7f and pFA6a-His3MX6 as the template. The resulting fragment was transformed in the diploid strains CCY034 and CCY044, and then the transformants were grown in SC-His at 25° for 3 days. The resulting heterozygous strains CCY039 and CCY049 were then confirmed by PCR using these genomic DNAs as templates to verify disruption of *GON7*. After dissection and selection of spores from these strains, PCR was used to confirm the deletion of *GON7* and *RAX2* using genomic DNA of these segregates as the template to obtain the haploid strains CCY040 and CCY050, respectively. *MATa* or *MATα* was identified by mating with reference strains on SD plates for 3 days at 25°.

The strain expressing *Bud32p-GFP* (CCY052) was created by homologous recombination with a linearized fragment of pYC20, respectively. The pYC20 fragment was digested with *SphI* and purified by a gel extraction. The resulting linearized fragments were transformed in BY4741 strain. Transformants were incubated on YPD plate at 25° for 3 days and then transferred to an SC plate with 5-FOA (1 mg·ml<sup>-1</sup>) at 25° for 3 days. After the single isolation of colonies on YPD plate, the resulting strain (CCY052) was confirmed by PCR using these genomic DNAs as templates to verify the correct insertion of *BUD32-GFP*. The diploid *BUD32-GFP* strain (CCY053) was generated by transformation of the plasmid expressing the *HO* endonuclease in the CCY052 strain. The growth rate of the resulting strain (CCY053) was confirmed to be identical to that of the BY4743 strain by a spotting assay.

Strains expressing *Bud32p* (K52A) (CCY064), *Bud32p* (K52A)-GFP (CCY054 and CCY055), and *Bud32p* (S258A)-GFP (CCY067) were created by homologous recombination with a linearized fragment of pYC16, pYC21, and pYC22, respectively, using the same procedure to generate CCY052. The diploid *bud32* (K52A)-GFP strain (CCY056) was generated by mating CCY054 with CCY055, and the diploid *bud32* (S258A)-GFP strain (CCY068) was generated by transformation of the plasmid expressing the *HO* endonuclease in CCY067.

Strains expressing *Bud32p-13myc* (CCY062) and *Bud32p* (K52A)-13myc (CCY065) were created by the one-step tagging method using PCR product (Longtine *et al.* 1998). A cassette of 13Myc-kanMX6 just upstream of the stop codon of *BUD32* using the primer set BUD32i/BUD32j and pFA6a-13Myc-kanMX6 as a template was amplified by PCR. The BY4741 and CCY064 strains were transformed with the resulting fragment to generate strains CCY062 and


CCY065, respectively. The transformants were incubated on YPD plates at 30° for 1 day and then selected to YPD plates with G418 (100 μg·ml<sup>-1</sup>). The correct insertion of 13Myc-kanMX6 was verified by PCR using genomic DNA of candidates as the template. Diploid strains (CCY063 and CCY066) were generated by transformation of the plasmid expressing the *HO* endonuclease in CCY062 and CCY065, respectively. The growth rate of resulting the strain (CCY063) was confirmed to be identical to that of the BY4743 strain by the spotting assay.

Strains expressing *Cdc11p-GFP* (CCY071 and CCY072) were created by the one-step tagging method using PCR product (Longtine *et al.* 1998). A cassette of GFP<sup>S65T</sup>-kanMX6 just upstream of the stop codon of *CDC11* using the primer set CDC11a/CDC11b and pFA6a-GFP<sup>S65T</sup>-kanMX6 as a template was amplified by PCR. The BY4741 and BY4742 strains were transformed with the resulting fragments to generate strains CCY071 and CCY072, respectively. The transformants were incubated on YPD plates at 30°C for 1 day, and then selected to YPD plates with G418 (100 μg·ml<sup>-1</sup>). The correct insertion of GFP<sup>S65T</sup>-kanMX6 was verified by PCR using genomic DNA of candidates as the template. Diploid strain (CCY073) was generated by mating CCY071 with CCY072. The growth rate of resulting strain (CCY073) was confirmed to be identical to that of BY4743 strain by a spotting assay.

### Analysis of budding patterns

Cells grown to late log phase were harvested by centrifugation at 3000 × *g* for 5 min, washed twice, and resuspended in water. Cells were fixed with a paraformaldehyde solution to a final concentration of 3.7% with gentle agitation for 30 min at room temperature. The fixed cells were washed twice and resuspended in water and then incubated with a final concentration of 5 μg·ml<sup>-1</sup> calcofluor white for 20 min to visualize bud and birth scars. After washing twice with water, the fluorescent image of chitin rings was observed with an ECLIPSE 80i fluorescence microscope (Nikon, Tokyo) using a UV filter set.

Budding patterns were analyzed according to Chen *et al.* (2000). All analyses were performed in homozygous diploids. The bud-scar positions of daughter and mother cells were separately counted and analyzed. The daughter cells with the first bud scar were classified in relation to the bud position to the birth scar as “proximal” when one bud scar was at the proximal pole, as “distal” when one bud scar was at the distal pole, or as “random” when one bud scar was around equatorial region. The mother cells that had more than three bud scars were used (Chant and Pringle 1995) and classified on the basis of the relation of the bud positions to the birth scar as “bipolar” when one or more bud scars were at both poles, as “proximal” when all bud scars were at the proximal pole, as “distal” when all bud scars were at the distal pole, as “axial-like” when all bud scars were connected in a chain with at least one touching the birth scar, or as “random” when at least one or more bud scars were in the



**Figure 1** Budding pattern of the deletion mutants of each EKC/KEOPS component in diploid cells. (A) Representative image of budding patterns in both diploid daughter and mother cells. Daughter cells showed three budding patterns in the wild type and mutants. The positions of the first buds on diploid daughter cells were scored as being at the pole proximal to the birth scar (a), at the pole distal to the birth scar (b), or at a random site (c) on the cell. Mother cells showed five budding patterns in the wild type and mutants. Each diploid mother cell budding pattern was determined by counting cells with more than three bud scars. Bud scars were concentrated at the two poles (d, as bipolar), only at the pole proximal [e and g (as chain-like bud)] or distal (f) to the birth scar, and at the random sites (h). (B) Budding patterns of the diploid deletion mutants. Strains used were diploid wild type (BY4743) and *bud32Δ* (CCY003), *cgi121Δ* (CCY006), *gon7Δ* (CCY009), *bud32Δcgi121Δ* (CCY017), *bud32Δgon7Δ* (CCY018), and *cgi121Δgon7Δ* (CCY019). Budding positions are classified as in the budding patterns of diploid cells in Figure 1A. At least 150 cells were scored for each bud scar pattern from both daughter and mother cells (the percentages are indicated). In daughter cells, the solid, open, and dotted boxes indicate cells with buds at only the proximal pole, at only the distal pole, and at a random site, respectively. In mother cells, the shaded, solid, open, slashed, and dotted boxes indicate cells with buds at two poles, at only the proximal pole, at only the distal pole, at the axial-like, and at random sites, respectively.

equatorial region. A representative image of each pattern in both daughter and mother cells is shown in Figure 1A. Over 150 cells of each mother and daughter cell were counted for the classification. For each strain, the average value from two independent experiments is shown.

### Fluorescence microscopy

Cells expressing GFP-fused protein were grown for 12–16 h at 25° and then harvested by centrifugation at 3000 × g for 5 min, stained with calcofluor white, washed twice, and resuspended in water. The fluorescent images of the GFP-fused proteins and bud scars in cells were observed under a fluorescence microscope (BZ-9000; Keyence, Osaka, Japan) using GFP and UV filter sets, respectively.

### Western blotting

Yeast cell lysates were prepared as previously reported (Krappmann *et al.* 2007). Briefly, the same OD<sub>600</sub> units of cells were resuspended in an 80- $\mu$ l extraction buffer [50 mM Tris-HCl (pH 7.5), 50 mM DTT, 1 mM EDTA, and inhibitor cocktail; Roche, Basel, Switzerland], vortexed with glass beads for 10 min at 4° followed by an additional 20  $\mu$ l of membrane extraction buffer [50 mM Tris-HCl (pH 7.5), 50 mM DTT, 1 mM EDTA, 10% SDS, and 5% Triton X-100] and then vortexed at 4° for 1 min. The protein concentration of the resulting extracts was determined by the Bradford method (Kruger 1994). Proteins were subjected to SDS-PAGE with 15% low-bis gels (Hirano 1989). The separated

proteins were electroblotted onto polyvinylidene fluoride membranes as described in Hirano (1989). Membrane blocking was performed using the method of Kawasaki *et al.* (2008). Monoclonal mouse anti-FLAG (Sigma-Aldrich, St. Louis) and peroxidase-coupled goat anti-mouse immunoglobulin G (Santa Cruz Biotechnology, Santa Cruz, CA) were used as primary and secondary antibodies, respectively.

## Results

### Complex made with *Bud32p*, *Cgi121p*, and *Gon7p* regulates bipolar bud-site selection

The EKC/KEOPS complex consists of *Kae1p*, *Bud32p*, *Cgi121p*, *Gon7p*, and *Pcc1p* in haploid cells (Downey *et al.* 2006; Kisseleva-Romanova *et al.* 2006). We investigated whether the EKC/KEOPS complex is involved in bipolar bud-site selection. By using tandem affinity purification (TAP), we confirmed that the components of the complex pulled down with *Bud32p*-TAP or *Gon7p*-TAP from diploid cells are the same as those of the haploid EKC/KEOPS complex (Y. Kato, H. Kawasaki, and H. Hirano, unpublished results). The budding patterns of the deletion mutants of each EKC/KEOPS component were analyzed, except those of *KAE1* and *PCC1*, whose deletion showed a severe growth defect. The budding patterns of diploid cells were classified as shown in Figure 1A. The first buds in daughter diploid cells of these mutants were at the distal pole, as observed in

**Table 1 Budding patterns of homozygous diploid strains**

	Budding = 1			Budding > 3				
	Proximal	Distal	Random	Bipolar	Proximal	Distal	Random	Axial-like
BY4743	1	99	0	85	0	11	4	0
<i>bud32Δ</i>	6	94	0	26	1	10	63	0
<i>cgi121Δ</i>	6	94	0	21	2	16	61	0
<i>gon7Δ</i>	3	94	3	22	0	34	44	0
<i>bud32Δ cgi121Δ</i>	2	98	0	33	0	7	60	0
<i>bud32Δ gon7Δ</i>	1	98	1	21	1	10	68	0
<i>cgi121Δ gon7Δ</i>	4	96	0	24	0	13	63	0
<i>bud8Δ</i>	95	0	5	5	95	0	0	0
<i>bud9Δ</i>	0	99	1	0	2	98	0	0
<i>bud32Δbud8Δ</i>	67	18	15	29	28	2	41	0
<i>cgi121Δbud8Δ</i>	53	30	17	28	15	0	57	0
<i>gon7Δbud8Δ</i>	96	2	2	13	86	0	1	0
<i>bud32Δbud9Δ</i>	5	95	0	20	2	69	9	0
<i>cgi121Δbud9Δ</i>	<1	99	<1	23	1	66	10	0
<i>gon7Δbud9Δ</i>	<1	99	<1	16	0	81	3	0
<i>rax2Δ</i>	52	33	15	17	2	1	65	15
<i>bud32Δ rax2Δ</i>	51	35	14	18	1	0	70	11
<i>cgi121Δ rax2Δ</i>	50	37	13	22	0	1	66	11
<i>gon7Δ rax2Δ</i>	50	36	14	20	0	0	68	12
<i>K52A</i>	2	96	2	36	0	11	53	0
<i>K52Abud8Δ</i>	76	6	18	29	14	1	56	0
<i>K52Abud9Δ</i>	8	92	0	14	1	74	11	0
<i>sch9Δ</i>	3	97	0	91	1	4	4	0
<i>S258A</i>	3	96	1	85	0	11	4	0

Each value indicates the percentage of total cells.

the wild-type diploid cells (Figure 1B). The deletion of each component of the EKC/KEOPS complex did not affect the budding pattern of daughter diploid cells, while the mother diploid cells of these mutants showed a severe budding defect (Figure 1B and Table 1). About half of the mother cells showed a random budding pattern in these mutants, although the *gon7Δ* mutant displayed a slightly different phenotype with a higher distal budding ratio. We also examined the phenotype of several double mutants, in which two genes of the complex component were disrupted. The phenotypes of *bud32Δcgi121Δ*, *bud32Δgon7Δ*, and *cgi121Δgon7Δ* double mutants were very similar to those of the single mutants (Figure 1B and Table 1). These results suggest that *BUD32*, *CGI121*, and *GON7* act in the same process as a complex.


#### **Random budding of *bud32Δ*, *cgi121Δ*, and *gon7Δ* is suppressed by deletion of *BUD9* but not *BUD8***

*Bud8p* and *Bud9p* are cortical landmarks in diploid cells for the distal and proximal poles, respectively (Harkins *et al.* 2001), because each deletion of these landmarks displays unipolar budding at the distal or the proximal pole, respectively. The first buds of daughter cells in *bud32Δ*, *cgi121Δ*, or *gon7Δ* mutants formed at the distal pole as in wild-type cells (Figure 1B and Table 1). This indicates that *Bud32p*, *Cgi121p*, and *Gon7p* may not be related to marking of the distal pole with *Bud8p*. We observed the effect of the deletion of *BUD8* or *BUD9* in the *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutant cells. In the daughter cells of the *bud32Δbud8Δ* double mutant, the buds were observed largely at the proximal pole (~60%), but some were at the distal pole (20%)

and at the equatorial sites (~20%) (Figure 2A and Table 1). The budding pattern of daughter cells in the *cgi121Δbud8Δ* double mutant was similar to that of the *bud32Δbud8Δ* double mutant. In contrast, that of daughter cells in the *gon7Δbud8Δ* double mutant was similar to that of the *bud8Δ* mutant in that the first bud of >90% of cells shows the proximal budding. Because the proximal budding observed in the *bud8Δ* mutant would be dependent on *Bud9p*, *Bud9p* seems to act as the proximal marker in daughter cells of these double mutants. However, ~30–50% of cells in *bud32Δbud8Δ* and *cgi121Δbud8Δ* mutants budded at the distal pole or at the equatorial site. Thus, the proximal marker may be somewhat impaired in these mutants.

In the *bud32Δbud9Δ* double mutant, the first and subsequent buds were formed at the distal pole, as in the *bud9Δ* mutant (Figure 2A and Table 1). Similar results were obtained for the *cgi121Δbud9Δ* and *gon7Δbud9Δ* double mutants. Because distal budding of the *bud9Δ* mutant would be dependent on *Bud8p*, these observations suggest that *Bud8p* in daughter cells of these mutants still acts as the distal landmark.

As shown in Figure 1B, ~45–60% of *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutant mother cells showed a random budding pattern. Similarly, ~40–60% of *bud32Δbud8Δ* and *cgi121Δbud8Δ* double-mutant mother cells showed a random budding pattern (Figure 2B). A distal budding pattern, which was observed in ~15% of *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutant cells, was almost completely suppressed and replaced with a proximal budding pattern (Figure 1B and Figure 2B). However, the majority of the double mutants showed a random budding pattern, although the ratio of the proximal budding pattern increased slightly.



**Figure 2** Genetic interaction between *BUD8* or *BUD9* and *EKC/KEOPS* genes. (A and B) Budding pattern of daughter (A) and mother (B) cells with *BUD8* or *BUD9* deletion in the *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutant backgrounds. Strains used were the diploid wild type (BY4743), *bud8Δ* (CCY011), *bud9Δ* (CCY013), *bud32Δ* (CCY003), *bud32Δbud8Δ* (CCY026), *cgi121Δbud8Δ* (CCY027), *gon7Δbud8Δ* (CCY028), *bud32Δbud9Δ* (CCY029), *cgi121Δbud9Δ* (CCY030), and *gon7Δbud9Δ* (CCY031). Budding positions are classified as in the budding pattern of diploid cells in Figure 1A. At least 150 cells were scored for each bud scar pattern on both daughter and mother cells (the percentages are indicated). The solid, open, and dotted boxes in A indicate daughter cells budded at proximal, distal, and equatorial sites, respectively. The shaded, solid, open, and dotted boxes in B indicate mother cells with buds at two poles, at only the proximal pole, at only the distal pole, and at the random site, respectively.

In mother cells of *bud32Δbud9Δ* double mutants, the random budding pattern was replaced by a distal budding pattern, although bipolar budding observed in the *bud32Δ* mutant was still seen in the double mutant (Figure 2B and Table 1). The random budding of mother cells of *cgi121Δ* and *gon7Δ* mutants was also suppressed by *BUD9* deletion. In the absence of *Bud9p*, a random budding pattern clearly replaced the distal budding pattern in which *Bud8p* was marked at the distal pole. These results suggest that the proximal marking by *Bud9p* is regulated by *Bud32p* and other components of the EKC/KEOPS complex.


#### Localization of the bipolar landmarks *Bud8p* and *Bud9p*

We showed that *BUD9*, but not *BUD8*, is required for random budding of *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutants. Therefore, we observed the localization of these bipolar landmarks in the wild type, and these mutant cells in GFP-tagged *Bud8p* or *Bud9p* expressed with a high-copy vector with fluorescence microscopy. The expression of GFP-*Bud8p* or GFP-*Bud9p* in diploid *bud8Δ* or *bud9Δ* mutant cells complemented the budding defects, suggesting that these tagged proteins provide a normal function. In the wild-type cells, GFP-*Bud8p* was localized at the distal pole in 36% of unbudded cells ( $n = 439$ ) and at the proximal pole in 17% of such cells (Figure 3A, cells 1 and 2, and Table 2). The GFP-*Bud8p* signal in mother cells appeared to be weaker than that in daughter cells. It was localized at the bud tip

in 17% of small-/large-budded cells ( $n = 529$ ) and at the bud tip/neck in 22% of such cells (Figure 3A, cells 3–7, and Table 2). Localization only to the bud neck was also observed in 7% of such cells (Figure 3A, cell 8, and Table 2). GFP-*Bud9p* was localized at the proximal pole in 17% of unbudded cells ( $n = 200$ ), and the signal at the distal pole was observed in only 5% of such wild-type mother cells (Figure 3B, cells 1 and 2, and Table 3). In addition, GFP-*Bud9p* was also localized at the proximal pole (10%), bud neck (7%), and bud neck/proximal pole (9%) in small-/large-budded cells ( $n = 254$ ; Figure 3B, cells 3–8, and Table 3).

In the *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutants, the localization patterns of GFP-*Bud8p* were essentially identical to that of the wild type, but its GFP signal appeared to be very diffused around the poles, tips, and bud necks (Figure 3C, cells 1–5, and Table 2; GFP-*Bud8p* localization patterns in the *cgi121Δ* mutant are those of the representative mutant). In addition, GFP-*Bud8p* was also observed at the bud tip and bud neck in cells that had bud scars at the equatorial position (Figure 3C, cells 5–8).

However, the localization to both poles and the bud neck of the GFP-*Bud9p* signal dramatically decreased compared to those of the wild-type cells (~3% of all cells) (Table 3). No GFP signal was observed in most cells (Figure 3D, cells 1–4; localization patterns of GFP-*Bud9p* in the *cgi121Δ* mutant are those of the representative mutant), but a few cells were also present with the GFP signal in which normal



**Figure 3** Effect on the localization of GFP-Bud8p and GFP-Bud9p due to loss of EKC/KEOPS components. (A and B) Localization patterns of GFP-Bud8p and GFP-Bud9p in the wild type. Wild type (BY4743) expressing the full-length GFP-Bud8p or GFP-Bud9p from its own promoter in pYC14 or pYC06, respectively, was grown for 12–16 h at 25° in synthetic complete (SC)–Leu or –Ura liquid medium, stained with calcofluor white, and then suspended in water for observation. GFP-Bud8p or GFP-Bud9p and bud scars were observed with a fluorescence microscope with GFP and UV filter sets, respectively. (C and D) Localization of GFP-Bud8p and GFP-Bud9p in the *cgi121Δ* mutant. The *cgi121Δ* (CCY006) expressing full-length GFP-Bud8p or GFP-Bud9p from their own promoters in pYC14 or pYC06 was grown for 12–16 h at 25° in SC–Leu or –Ura liquid medium, stained with calcofluor white, and then suspended in water for observation. GFP-Bud8p or GFP-Bud9p and bud scars were observed with a fluorescence microscope with GFP and UV filter sets, respectively. (E) Expression level of Bud9p in the absence of Cgi121p. Diploid wild type (BY4743) and *cgi121Δ* mutant (CCY006), both expressing the FLAG<sub>6</sub>-Bud9p in pYC10, were analyzed by Western blotting. The wild type (BY4743) was considered the control, which carried the empty pRS426. Cells were grown overnight to log phase in SC medium lacking uracil. The total cell extract from the same OD<sub>600</sub> units of cells was analyzed by immunoblotting using an anti-FLAG antibody (top) and stained with Ponceau-S as a loading control (bottom).

*Bud9p* localization was observed (Figure 3D, cells 5–8, and Table 3). Another visible GFP signal was observed primarily as a dot-like structure along the cell periphery of both daughter and mother cells (Figure 3D, cells 9–15). Because no obvious difference was seen at the *BUD9* expression levels of mRNA and protein between the wild-type and mutant cells, we concluded that mislocalization of the GFP signal in these mutants was not due to a difference in *Bud9p* expression (Figure 3E and real-time PCR; Y. Kato, H. Kawasaki, and H. Hirano, unpublished data). These results also suggest that *Bud32p*, *Cgi121p*, and *Gon7p* are specifically related in the localization of *Bud9p*, but not *Bud8p*.

#### ***Bud32p* kinase activity is required for *Bud9p* localization**

We generated a kinase dead mutant (*bud32-K52A*) by replacing the 52nd lysine residue with alanine in *Bud32p*, which is the binding site for the  $\alpha$ - and  $\beta$ -phosphates of ATP (Mao *et al.* 2008), to investigate whether kinase activity is required for the regulation of bipolar bud-site selection. The mutant displayed a random budding pattern, as observed in the *bud32Δ* mutant (Figure S1A and Table 1). These results suggest that kinase activity is also required for regulation of bipolar bud-site selection. We observed the effect of deleting *BUD8* or *BUD9* in the kinase dead mutant (*bud32-K52A*); the deletion of *BUD9* suppressed random budding in the kinase dead mutant, but deletion of *BUD8* did not (Figure S1, B and C, and Table 1). Finally, we examined the localization of GFP-Bud8p and GFP-Bud9p in the mutant (Figure S1D). In the kinase dead mutant, the GFP-Bud8p localization pattern was similar to that of wild-type cells, whereas the GFP-Bud9p signal was not observed in the kinase dead mutant, as in the *bud32Δ* mutant (Figure S1D). These results indicate that kinase activity is required for proper localization of *Bud9p*, but not *Bud8p*.

#### **Mutation at Ser258 of *Bud32p* did not affect bipolar bud-site selection**

A recent report has suggested that *Bud32p* is involved in the *Sch9p* signaling pathway (Peggion *et al.* 2008). We investigated whether this signaling pathway is involved in bipolar bud-site selection. *Sch9p*, a homolog of yeast Akt, phosphorylates *Bud32p* at Ser258. This phosphorylation of *Bud32p* positively regulates its ability to interact with *Grx4p* and to phosphorylate it. We constructed a mutant *bud32-S258A* in which Ser258 on *Bud32p* was replaced with Ala. We observed the budding patterns of *sch9Δ* and *bud32-S258A*. The budding patterns of these two mutants were similar to those of wild-type daughter and mother cells (Figure S2 and Table 1). This result suggests that the *Sch9p* signaling cascade is not related to bipolar bud-site selection.

#### **Genetic interaction between *RAX2* and *EKC/KEOPS* genes**

*Rax1p* and *Rax2p* are interdependently localized and also involved in *Bud9p* localization via interactions (Kang *et al.* 2004). In daughter cells of the *rax2Δ* mutant, about half of the first buds were at the distal pole, and the rest were at the



**Table 2 Localization patterns of GFP-Bud8p**

	Unbudded cells			Budded cells				
	Distal	Both poles	No signal	Bud tip	Bud neck	Bud tip/neck	No signal	Others
BY4743	19	17	64	17	7	22	59	5
<i>bud32Δ</i>	20	10	70	20	6	20	51	3
<i>cgi121Δ</i>	15	14	71	15	6	29	46	4
<i>gon7Δ</i>	17	19	64	19	10	25	49	7

Each value indicates the percentage of total cells.

proximal pole. Mother cells of *rax2Δ* mutants largely showed a random budding pattern, as in the *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutants. However, the phenotype differed between *rax2Δ* and EKC/KEOPS mutants although the ratio of random budding in both mutants was almost the same. A small number of *rax2Δ* mutant cells showed axial-like budding, while a few *bud32Δ* mutant cells showed distal budding (Figure 4A). We deleted *RAX2* in the *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutant cells and observed the budding pattern of these mutants (Figure 4A). In both daughter and mother cells of the double mutants, the phenotype was changed to one similar to the *rax2Δ* mutant (Figure 4A and Table 1). The distal budding pattern of the *bud32Δ* mutant disappeared following *RAX2* deletion and was replaced with an axial-like budding pattern. The axial budding pattern was never observed in the single mutants *bud32Δ*, *cgi121Δ*, and *gon7Δ*.

We observed *Rax2p*-GFP in wild-type and mutant cells (Figure 4, B and C, and Figure S3). Cells chromosomally expressing *Rax2p*-GFP also displayed a bipolar budding pattern, as in wild-type cells, suggesting that *Rax2p*-GFP is functional. In wild-type cells, *Rax2p*-GFP localized mainly to bud scars and birth scars in unbudded cells and sometimes to bud tips in budded cells. Similar results were obtained from *bud32Δ*, *cgi121Δ*, and *gon7Δ* single mutants (Figure 4B and Figure S3A). In addition, the localization of *Rax2p* at bud scars could be observed even in equatorial bud scars, suggesting that the regulation for localization of *Rax2p* is normal in the mutants (Figure 4C and Figure S3B).

## Discussion

### *EKC/KEOPS complex is required for bipolar bud-site selection*

*BUD32* was originally found as a gene encoding an atypical protein kinase found in virtually all eukaryotic and archaeal organisms (Stocchetto *et al.* 1997). It was also identified as a gene that regulates bud-site selection of diploid cells

through a genome-wide analysis (Ni and Snyder 2001). *Bud32p* is involved in the signaling pathway of *Sch9p* kinase (Peggion *et al.* 2008) and is also involved in telomere or transcriptional regulation as a component of the EKC/KEOPS complex (Downey *et al.* 2006; Kisseleva-Romanova *et al.* 2006). In an analysis with gel-filtration chromatography in haploid cells, the subcomplex containing only *Kae1p* and *Bud32p* was observed in yeast cells although it could have been a preparation artifact (Kisseleva-Romanova *et al.* 2006). No other subcomplex has been reported. We purified the EKC/KEOPS complex from diploid cells using a TAP pull-down. The diploid EKC/KEOPS complex was composed of *Bud32p*, *Cgi121p*, *Kae1p*, *Gon7p*, and *Pcc1p*, which is identical to the haploid complex. Of the genes encoding these components, *CGI121* and *GON7* were identified as new genes required for bipolar bud-site selection (Figure 1B). The EKC/KEOPS complex regulates bipolar bud-site selection, as deletion of the components showed a very similar phenotype with random budding. We also revealed that the kinase activity of *Bud32p* is required for bipolar bud-site selection (Figure S1). This activity is essential for the functions of the EKC/KEOPS complex, such as transcriptional regulation and telomere maintenance, and it is also contributed to, but was not absolutely essential for, the *Sch9p* signaling cascade (Peggion *et al.* 2008). In addition, we observed that phosphorylation at Ser258 on *Bud32p* by *Sch9p*, the modification of which did not alter its kinase activity, was not necessary for bipolar bud-site selection (Figure S2). Therefore, we concluded that the EKC/KEOPS complex is involved in bipolar bud-site selection with a signaling pathway different from the *Sch9p* kinase.


### *EKC/KEOPS complex is involved in the regulation of Bud9p localization*

We showed that *BUD9* deletion in the *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutants suppressed random budding and increased budding at the distal pole. In concordance with these results,

**Table 3 Localization patterns of GFP-Bud9p**

	Unbudded cells			Budded cells				
	Proximal	Both poles	No signal	Proximal	Bud neck	Proximal/neck	No signal	Others
BY4743	12	5	83	10	7	9	71	3
<i>bud32Δ</i>	3	0	97	<1	3	<1	96	0
<i>cgi121Δ</i>	2	0	98	2	0	3	95	0
<i>gon7Δ</i>	1	0	99	3	1	4	92	0

Each value indicates the percentage of total cells.




**Figure 4** Genetic interaction between *RAX2* and *EKC/KEOPS* genes. (A) Budding pattern of cells with deletion of *RAX2* in the *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutant backgrounds. Strains used were the diploid wild type (BY4743), *rax2Δ* (CCY044), *bud32Δrax2Δ* (CCY046), *cgi121Δrax2Δ* (CCY048), *gon7Δrax2Δ* (CCY051), and *bud32Δ* (CCY003). Budding positions are classified as in the budding pattern of diploid cells in Figure 1A. At least 150 cells were scored for each bud scar pattern from both daughter and mother cells (the percentages are indicated). In daughter cells, the solid, open, and dotted boxes indicate cells with buds at only the proximal pole, at only the distal pole, and at a random site, respectively. In mother cells, the shaded, solid, open, slashed, and dotted boxes indicate cells with buds at two poles, only the proximal pole, only the distal pole, axial-like and at random sites, respectively. (B) Diploid wild type (CCY034) and *cgi121Δ*

(CCY038) expressing Rax2p-GFP from their own promoter at their chromosomal locus were grown overnight to log phase in YPD liquid medium, stained with calcofluor white, and then suspended in water for observation. Rax2p-GFP and bud scars were observed with a fluorescence microscope with GFP and UV filter sets, respectively. Bar, 5  $\mu$ m. (C) Bud scars at the equatorial position of random budding sites were also marked by Rax2p-GFP. The arrows indicate Rax2p-GFP localization at random positions. Bar, 5  $\mu$ m.

**Bud9p** localization was affected by deletion of *BUD32*, *CGI121*, and *GON7* (Figure 3D and Table 3). The deletion of *BUD8* did not suppress random budding of *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutants although proximal budding increased in the double mutants in place of the distal budding observed in the single mutants (Figure 2B). The proximal budding in the mutants might be due to the accidental localization of **Bud9p** at the proximal pole because **Bud9p** is expressed in a normal amount in these cells (Figure 3E).

Our data suggest that mislocalization of **Bud9p** by loss of the EKC/KEOPS components caused random budding.

Proper localization of **Bud9p** is also dependent on the timing of **Bud9** expression that was observed in the G1 phase (Schenkman *et al.* 2002). A possibility exists that the EKC/KEOPS complex might be related to the timing of **Bud9** expression because the complex controls transcription of several genes (Kisseleva-Romanova *et al.* 2006). In a promoter swapping assay, GFP-**Bud9p** expression from the *BUD8* or *CLB2* promoter specifically at the G2/M phase failed to rescue unipolar proximal budding in a *bud9Δ-1* mutant, indicating inefficient delivery or poor stability of **Bud9p**, or both (Schenkman *et al.* 2002). Furthermore,



**Figure 5** Regulation of the localization of the bipolar markers **Bud8p** and **Bud9p** by the EKC/KEOPS complex and Rax1p/Rax2p. The schematic illustration is based on figure 3 in Casamayor and Snyder (2002). For details, see the Discussion.

virtually no random budding was seen in these cases. Localization of GFP-Bud9p in *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutants was observed in a dot-like structure at cell periphery, as shown in Figure 3D. We confirmed that the stability and expression level of Bud9p were the same as that of the wild type (Figure 3E). GFP-Bud9p from the *BUD8* promoter in the *bud9Δ-1* mutant appeared to localize in internal vesicles, and the mutant showed mainly unipolar distal budding (Schenkman *et al.* 2002). If the time lag for *BUD9* expression occurs in the *bud32Δ*, *cgi121Δ*, and *gon7Δ*, it should largely be unipolar distal budding not random budding. Therefore, we conclude that random budding in the mutants is caused by mislocalization at the cell periphery after delivery of Bud9p to the plasma membrane, but not through a time lag in *BUD9* expression. Support for this conclusion was also provided by our observation that GFP-Bud8p localization patterns in the mutant deleted of each EKC/KEOPS component were similar to those of the wild type because the delivery of both Bud8p and Bud9p to the presumptive bud site is dependent on actin (Schenkman *et al.* 2002).

The deletion of *RAX2* in diploid cells induced random budding and axial-like budding, although ~20% of cells showed a bipolar budding pattern. Axial budding was suppressed in diploid cells, as *Axl1p*, on which axial budding is dependent, is transcriptionally repressed in diploid cells (Fujita *et al.* 1994). However, the allele mutant of both *BUD8* and *BUD9* uses axial budding cues (Zahner *et al.* 1996). Several components of the axial budding system, such as *Axl2p* and *Bud3p*, are also expressed in diploid cells. Indeed, *BUD3* deletion in the *bud8Δbud9Δ* double mutant essentially induces a random budding pattern (Harkins *et al.* 2001). Axial budding cues can be recognized with less efficiency in diploid cells when both bipolar markers are impaired. However, we never observed an axial-like budding pattern in the deletion mutant of EKC/KEOPS components (Figure 1 and Figure 2). This indicates that the axial budding system is essentially inactive in these cells although a very small part of the bud at the proximal pole might use such a cue.

We observed that deletion of *RAX2* in *bud32Δ*, *cgi121Δ*, and *gon7Δ* mutants changed the phenotype to one identical with *rax2Δ* (Figure 4A). In concordance with this observation, *Rax2p* distribution did not differ between the wild type and *bud32Δ*, *cgi121Δ*, or *gon7Δ* (Figure 4, B and C, and Figure S3). We also confirmed that regulation of Bud9p localization by the EKC/KEOPS complex is independent of septin by observing the localization of *Cdc11p*-GFP, one of the septin subunits, in the deletion mutant of the EKC/KEOPS component (Figure S4). These results suggest that the EKC/KEOPS complex acts to regulate Bud9p localization as a downstream factor of *Rax2p* and septin.

#### **Regulation of bipolar bud-site selection by the EKC/KEOPS complex**

We concluded that the EKC/KEOPS complex in diploid cells acts as a *Rax2p* downstream factor and that the complex maintains bipolar budding through regulation of Bud9p lo-

calization (Figure 5). The localization of the cortical markers Bud8p and Bud9p is regulated by several proteins as shown in Figure 5. The EKC/KEOPS complex is the factor regulating the localization of only Bud9p, downstream of *Rax2p* and septins. Recent reports suggest that the complex functions in telomere maintenance, transcriptional regulation, and t<sup>6</sup>A modification (Downey *et al.* 2006; Kisseleva-Romanova *et al.* 2006; Daugeron *et al.* 2011; Srinivasan *et al.* 2011), although direct target of the complex is still elusive. The complex localized mainly to the nucleus, and the minor portion of the EKC/KEOPS complex localized to cytosol (Y. Kato, H. Kawasaki, and H. Hirano, unpublished data), whereas Bud9p localized to the cell surface. Although the complex in the cytosol might be directly linked to Bud9p, we could not observe a physical interaction between these proteins. These results suggest that the regulation of Bud9p localization by the complex may not occur at the cell surface and that the localization of Bud9p might be regulated by factor(s) under the transcriptional control by the EKC/KEOPS complex. Further experiments, such as suppressor screening that can rescue Bud9p localization in the deletion mutants of each EKC/KEOPS component, might be able to identify the factors that are important for establishing and maintaining spatial cues in cells.

Our results suggest that the correct localization of Bud9p is essential for bipolar budding. Although many of the genes that specifically caused random budding when deleted are associated with the localization of Bud8p, so far no genes have been reported that specifically regulate only Bud9p, except for the septins (Ni and Snyder 2001; Schenkman *et al.* 2002). Our findings revealed that the EKC/KEOPS complex is specifically involved in the regulation of Bud9p localization. Further analysis of this regulation system will be an important step toward understanding asymmetric development from spatial cues in diploid cells.

#### **Acknowledgments**

We thank Nobuyuki Endo (Keyence Corporation) for technical advice on the BIOREVO BZ-9000 fluorescence microscope. We are also extremely grateful to Katsuki Asano for his assistance with the TAP purification. This work was partly supported by the 21st Century Centers-of-Excellence Program at Yokohama City University and by Grants-in-Aid for Scientific Research from KAKENHI (C).

#### **Literature Cited**

- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, *et al.*, eds. 1995 *Current Protocols in Molecular Biology*. John Wiley & Sons, New York.
- Bedinger, P. A., K. J. Hardeman, and C. A. Loukides, 1994 Travelling in style: the cell biology of pollen. *Trends Cell Biol.* 4: 132–138.
- Casamayor, A., and M. Snyder, 2002 Bud-site selection and cell polarity in budding yeast. *Curr. Opin. Microbiol.* 5: 179–186.
- Chant, J., and J. R. Pringle, 1995 Patterns of bud-site selection in the yeast *Saccharomyces cerevisiae*. *J. Cell Biol.* 129: 751–765.

- Chant, J., M. Mischke, E. Mitchell, I. Herskowitz, and J. R. Pringle, 1995 Role of Bud3p in producing the axial budding pattern of yeast. *J. Cell Biol.* 129: 767–778.
- Chen, T., T. Hiroko, A. Chaudhuri, F. Inose, M. Lord, *et al.*, 2000 Multigenerational cortical inheritance of the Rax2 protein in orienting polarity and division in yeast. *Science* 290: 1975–1978.
- Christianson, T. W., R. S. Sikorski, M. Dante, J. H. Shero, and P. Hieter, 1992 Multifunctional yeast high-copy-number shuttle vectors. *Gene* 110: 119–122.
- Daugeron, M. C., T. L. Lenstra, M. Frizzarin, B. EL Yacoubi, X. Liu, *et al.*, 2011 Gcn4 misregulation reveals a direct role for the evolutionary conserved EKC/KEOPS in the t6A modification of tRNAs. *Nucleic Acids Res.* (in press).
- Downey, M., R. Houlsworth, L. Maringele, A. Rollie, M. Brehme, *et al.*, 2006 A genome-wide screen identifies the evolutionarily conserved KEOPS complex as a telomere regulator. *Cell* 124: 1155–1168.
- Drubin, D. G., and W. J. Nelson, 1996 Origins of cell polarity. *Cell* 84: 335–344.
- Freifelder, D., 1960 Bud position in *Saccharomyces cerevisiae*. *J. Bacteriol.* 80: 567–568.
- Fujita, A., C. Oka, Y. Arikawa, T. Katagai, A. Tonouchi, *et al.*, 1994 A yeast gene necessary for bud-site selection encodes a protein similar to insulin-degrading enzymes. *Nature* 372: 567–570.
- Gietz, D., A. St. Jean, R. A. Woods, and R. H. Schiestl, 1992 Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Res.* 20: 1425.
- Halme, A., M. Michelitch, E. L. Mitchell, and J. Chant, 1996 Bud10p directs axial cell polarization in budding yeast and resembles a transmembrane receptor. *Curr. Biol.* 6: 570–579.
- Harkins, H. A., N. Page, L. R. Schenkman, C. De Virgilio, S. Shaw, *et al.*, 2001 Bud8p and Bud9p, proteins that may mark the sites for bipolar budding in yeast. *Mol. Biol. Cell* 12: 2497–2518.
- Hirano, H., 1989 Microsequence analysis of winged bean seed proteins electroblotted from two-dimensional gel. *J. Protein Chem.* 8: 115–130.
- Hyman, A. A., and J. G. White, 1987 Determination of cell division axes in the early embryogenesis of *Caenorhabditis elegans*. *J. Cell Biol.* 105: 2123–2135.
- Kang, P. J., A. Sanson, B. Lee, and H. O. Park, 2001 A GDP/GTP exchange factor involved in linking a spatial landmark to cell polarity. *Science* 292: 1376–1378.
- Kang, P. J., E. Angerman, K. Nakashima, J. R. Pringle, and H. O. Park, 2004 Interactions among Rax1p, Rax2p, Bud8p, and Bud9p in marking cortical sites for bipolar bud-site selection in yeast. *Mol. Biol. Cell* 15: 5145–5157.
- Kato, Y., N. Arakawa, Y. Masuishi, H. Kawasaki, and H. Hirano, 2009 Mutagenesis of longer inserts by the ligation of two PCR fragments amplified with a mutation primer. *J. Biosci. Bioeng.* 107: 95–97.
- Kawasaki, H., A. Okayama, Y. Iwafune, S. Yahagi, N. Arakawa, *et al.*, 2008 Multiplex detection and identification of proteins on a PVDF membrane blocked with a synthetic polymer-based reagent. *Electrophoresis* 29: 4377–4380.
- Kisseleva-Romanova, E., R. Lopreiato, A. Baudin-Baillieu, J. C. Rousselle, L. Ilan, *et al.*, 2006 Yeast homolog of a cancer-testis antigen defines a new transcription complex. *EMBO J.* 25: 3576–3585.
- Krappmann, A. B., N. Taheri, M. Heinrich, and H. U. Mosch, 2007 Distinct domains of yeast cortical tag proteins Bud8p and Bud9p confer polar localization and functionality. *Mol. Biol. Cell* 18: 3323–3339.
- Kraut, R., W. Chia, L. Y. Jan, Y. N. Jan, and J. A. Knoblich, 1996 Role of inscuteable in orienting asymmetric cell divisions in *Drosophila*. *Nature* 383: 50–55.
- Kruger, N. J., 1994 The Bradford method for protein quantitation. *Methods Mol. Biol.* 32: 9–15.
- Longtine, M. S., and A. McKenzie III, D. J. Demarini, N. G. Shah, A. Wach, *et al.*, 1998 Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* 14: 953–961.
- Mao, D. Y., D. Neculai, M. Downey, S. Orlicky, Y. Z. Haffani, *et al.*, 2008 Atomic structure of the KEOPS complex: an ancient protein kinase-containing molecular machine. *Mol. Cell* 32: 259–275.
- Mooseker, M. S., 1985 Organization, chemistry, and assembly of the cytoskeletal apparatus of the intestinal brush border. *Annu. Rev. Cell Biol.* 1: 209–241.
- Ni, L., and M. Snyder, 2001 A genomic study of the bipolar bud site selection pattern in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 12: 2147–2170.
- Park, H. O., and E. Bi, 2007 Central roles of small GTPases in the development of cell polarity in yeast and beyond. *Microbiol. Mol. Biol. Rev.* 71: 48–96.
- Peggion, C., R. Lopreiato, E. Casanova, M. Ruzzene, S. Facchin, *et al.*, 2008 Phosphorylation of the *Saccharomyces cerevisiae* Grx4p glutaredoxin by the Bud32p kinase unveils a novel signaling pathway involving Sch9p, a yeast member of the Akt/PKB subfamily. *FEBS J.* 275: 5919–5933.
- Roosen, J., K. Engelen, K. Marchal, J. Mathys, G. Griffioen, *et al.*, 2005 PKA and Sch9 control a molecular switch important for the proper adaptation to nutrient availability. *Mol. Microbiol.* 55: 862–880.
- Sanders, S. L., and I. Herskowitz, 1996 The BUD4 protein of yeast, required for axial budding, is localized to the mother/BUD neck in a cell cycle-dependent manner. *J. Cell Biol.* 134: 413–427.
- Schenkman, L. R., C. Caruso, N. Page, and J. R. Pringle, 2002 The role of cell cycle-regulated expression in the localization of spatial landmark proteins in yeast. *J. Cell Biol.* 156: 829–841.
- Srinivasan, M., P. Mehta, Y. Yu, E. Prugar, E. V. Koonin, *et al.*, 2011 The highly conserved KEOPS/EKC complex is essential for a universal tRNA modification, t6A. *EMBO J.* 30: 873–881.
- Stocchetto, S., O. Marin, G. Carignani, and L. A. Pinna, 1997 Biochemical evidence that *Saccharomyces cerevisiae* YGR262c gene, required for normal growth, encodes a novel Ser/Thr-specific protein kinase. *FEBS Lett.* 414: 171–175.
- Taheri, N., T. Kohler, G. H. Braus, and H. U. Mosch, 2000 Asymmetrically localized Bud8p and Bud9p proteins control yeast cell polarity and development. *EMBO J.* 19: 6686–6696.
- Zahner, J. E., H. A. Harkins, and J. R. Pringle, 1996 Genetic analysis of the bipolar pattern of bud site selection in the yeast *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 16: 1857–1870.

Communicating editor: M. D. Rose


# GENETICS

Supporting Information

<http://www.genetics.org/content/suppl/2011/05/30/genetics.111.128231.DC1>


## **Cell Polarity in *Saccharomyces cerevisiae* Depends on Proper Localization of the Bud9 Landmark Protein by the EKC/KEOPS Complex**

**Yu Kato, Hiroshi Kawasaki, Yoshifumi Ohyama, Takashi Morishita, Hiroshi Iwasaki, Tetsuro Kokubo,  
and Hisashi Hirano**




**Figure S1** Phenotype of the kinase dead mutant (*bud32-K52*). (A) Budding patterns of the kinase dead mutant. Strains used were diploid BY4743, the *bud32Δ* (CCY003), the *bud32-K52A* (CCY056). (B) Budding pattern of cells with *BUD8* deletion in the kinase dead mutant backgrounds. Diploid strains used were the *bud32Δ* (CCY003), *bud32Δbud8Δ* (CCY026), *K52Abud8Δ* (CCY058) and *bud8Δ* (CCY011). (C) Budding pattern of cells with *BUD9* deletion in the kinase dead mutant backgrounds. Diploid strains used were the *bud32Δ* (CCY003), *bud32Δbud9Δ* (CCY029), *K52Abud9Δ* (CCY060) and *bud9Δ* (CCY013). Budding positions are classified as in the budding pattern of diploid cells

in Figure 1A. At least 150 cells were scored for each bud scar pattern from both daughter and mother cells; the percentages are indicated. The black, white, and dotted boxes indicate daughter cells with bud at only the proximal pole, only the distal pole, or the random site, respectively. The gray, black, white, and dotted boxes indicate mother cells with buds at two poles, only the proximal pole, only the distal pole, or the random site, respectively. (D) Localization of GFP-Bud8p and GFP-Bud9p in the kinase dead mutant. The *bud32-K52A* (CCY065) mutant expressing full-length GFP-Bud8p and GFP-Bud9p from its own promoter in pYC14 or pYC06, respectively, was grown for 12-16h at 25°C in synthetic complete (SC)-Leu or -Ura liquid medium, stained with calcofluor white, and then suspended in water for observation. GFP-Bud8p/GFP-Bud9p and bud scars were observed with a fluorescence microscope with GFP and UV filter sets, respectively.




**Figure S2** Budding patterns of *bud32-S258A* and *sch9D* mutant. Strains used were diploid BY4743, *sch9Δ* (CCY070) and *bud32-S258A* (CCY068). Budding positions are classified as in the budding pattern of diploid cells in Figure 1A. At least 150 cells were scored for each bud scar pattern from both daughter and mother cells; the percentages are indicated. The black, white, and dotted boxes indicate daughter cells with bud at only the proximal pole, only the distal pole, or the random site, respectively. The gray, black, white, and dotted boxes indicate mother cells with buds at two poles, only the proximal pole, only the distal pole, or the random site, respectively.






B.



**Figure S3** Localization of Rax2p-GFP in *bud32Δ* and *gon7Δ* mutants. (A) Diploid *bud32Δ* (CCY036) and *gon7Δ* (CCY041) expressing Rax2p-GFP from their own promoter at their chromosomal locus were grown overnight to log phase in YPD liquid medium, stained with calcofluor white, and then suspended in water for observation. Rax2p-GFP and bud scars were observed with a fluorescence microscope using GFP and UV filter sets, respectively. Scale bar indicates 5  $\mu$ m. (B) Bud scars at the equatorial position of random budding sites were also marked by Rax2p-GFP in diploid *bud32Δ* (CCY036) and *gon7Δ* (CCY041) mutants. The arrows indicate Rax2p-GFP localization at random positions. Scale bar indicates 5  $\mu$ m.



**Figure S4** Effect on the localization of a septin component, Cdc11p-GFP. (A) Localization of Cdc11p-GFP in *cgi121*Δ mutants and wild type cells. Diploid wild-type (CCY073) and *cgi121*Δ (CCY075) mutant cells expressing Cdc11p-GFP at its chromosomal loci were grown overnight to log phase in YPD liquid medium and then suspended in water for observation. Cdc11p-GFP and bud scars were observed with fluorescence microscope using a GFP and UV filter sets, respectively. Scale bar indicates 5  $\mu$ m. (B) The septin ring is formed in the mutant cells at the equatorial position. Scale bar indicates 5  $\mu$ m.

**Table S1 Strains used in this study**

Strains	Relevant genotype <sup>a</sup>	Source
BY4741	a his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1	our stock
BY4742	α his3- Δ 1 leu2- Δ 1 ura3- Δ 0 lys- Δ 0	our stock
BY4743	a/α his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/lys- Δ 0	our stock
CCY001	a his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud32 Δ ::kanMX6	EUROSCARF
CCY002	α his3- Δ 1 leu2- Δ 1 ura3- Δ 0 lys- Δ 0 bud32 Δ ::kanMX6	Open Biosystems, Inc.
CCY003	a/α his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud32 Δ ::kanMX6/bud32 Δ ::kanMX6	Open Biosystems, Inc.
CCY004	a his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 cgi121 Δ ::kanMX6	EUROSCARF
CCY005	α his3- Δ 1 leu2- Δ 1 ura3- Δ 0 lys- Δ 0 cgi121 Δ ::kanMX6	Open Biosystems, Inc.
CCY006	a/α his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 cgi121 Δ ::kanMX6/cgi121 Δ ::kanMX	Open Biosystems, Inc.
CCY007	a his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 gon7 Δ ::kanMX6	EUROSCARF
CCY008	α his3- Δ 1 leu2- Δ 1 ura3- Δ 0 lys- Δ 0 gon7 Δ ::kanMX6	Open Biosystems, Inc.
CCY009	a/α his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 gon7 Δ ::kanMX6/gon7 Δ ::kanMX6	Open Biosystems, Inc.
CCY010	a his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud8 Δ ::kanMX6	EUROSCARF
CCY011	a/α his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud8 Δ ::kanMX6/bud8 Δ ::kanMX6	a/α conversion from CCY010
CCY012	a his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud9 Δ ::kanMX6	EUROSCARF
CCY013	a/α his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud9 Δ ::kanMX6/bud9 Δ ::kanMX6	a/α conversion from CCY012
CCY014	a his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud32 Δ ::kanMX6 cgi121 Δ ::kanMX6	segregant from CCY001xCCY005
CCY015	a his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud32 Δ ::kanMX6 gon7 Δ ::URA3	see textb

CCY016	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 cgi121 Δ ::kanMX6 gon7 Δ ::URA3	see textb
CCY017	a/ α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud32 Δ ::kanMX6/bud32 Δ ::kanMX6 cgi121 Δ ::kanMX6/cgi121 Δ ::kanMX6	a/ α conversion from CCY014
CCY018	a/ α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud32 Δ ::kanMX6/bud32 Δ ::kanMX6 gon7 Δ ::URA3/gon7 Δ ::URA3	a/ α conversion from CCY015
CCY019	a/ α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 cgi121 Δ ::kanMX6/cgi121 Δ ::kanMX6 gon7 Δ ::URA3/gon7 Δ ::URA3	a/ α conversion from CCY016
CCY020	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud8 Δ ::kanMX6 bud32 Δ ::kanMX6	segregant from CCY002xCCY010
CCY021	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud8 Δ ::kanMX6 cgi121 Δ ::kanMX6	segregant from CCY005xCCY010
CCY022	a	his3- Δ 1 leu2-D1 ura3- Δ 0 met15- Δ 1 bud8 Δ ::kanMX6 gon7 Δ ::URA3	see textb
CCY023	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud9 Δ ::kanMX6 bud32 Δ ::kanMX6	segregant from CCY002xCCY012
CCY024	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud9 Δ ::kanMX6 cgi121 Δ ::kanMX6	segregant from CCY005xCCY012
CCY025	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 bud9 Δ ::kanMX6 gon7 Δ ::URA3	see textb
CCY026	a/ α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud8 Δ ::kanMX6/bud8 Δ ::kanMX6 bud32 Δ ::kanMX6/bud32 Δ ::kanMX6	a/ α conversion from CCY020
CCY027	a/ α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud8 Δ ::kanMX6/bud8 Δ ::kanMX6 cgi121 Δ ::kanMX6/cgi121 Δ ::kanMX6	a/ α conversion from CCY021
CCY028	a/ α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud8 Δ ::kanMX6/bud8 Δ ::kanMX6 gon7 Δ ::URA3/gon7 Δ ::URA3	a/ α conversion from CCY022
CCY029	a/ α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud9 Δ ::kanMX6/bud9 Δ ::kanMX6 bud32 Δ ::kanMX6/bud32 Δ ::kanMX6	a/ α conversion from CCY023
CCY030	a/ α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud9 Δ ::kanMX6/bud9 Δ ::kanMX6 cgi121 Δ ::kanMX6/cgi121 Δ ::kanMX6	a/ α conversion from CCY024
CCY031	a/ α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 bud9 Δ ::kanMX6/bud9 Δ ::kanMX6 gon7 Δ ::URA3/gon7 Δ ::URA3	a/ α conversion from CCY025

CCY032	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 RAX2-GFP-kanMX6::RAX2	see text
CCY033	α	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 lys- Δ 0 RAX2-GFPkanMX6::RAX2	see text
CCY034	a/α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/lys- Δ 0 RAX2-GFP-kanMX6::RAX2/RAX2-GFP-kanMX6::RAX2	mating of CCY032 and CCY033
CCY035	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 RAX2-GFP-kanMX6::RAX2 bud32 Δ ::kanMX6	segregant from CCY002xCCY032
CCY036	a/α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 RAX2-GFP-kanMX6::RAX2/RAX2-GFP-kanMX6::RAX2bud32 Δ ::kanMX6/bud32 Δ ::kanMX6	a/α conversion from CCY035
CCY037	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 RAX2-GFP-kanMX6::RAX2 cgi121 Δ ::kanMX6	segregant from CCY005xCCY032
CCY038	a/α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met15- Δ 1 RAX2-GFP-kanMX6::RAX2/RAX2-GFP-kanMX6::RAX2 cgi121 Δ ::kanMX6/cgi121 Δ ::kanMX6	a/α conversion from CCY037
CCY039	a/α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/lys- Δ 0 RAX2-GFP-kanMX6::RAX2/RAX2-GFP-kanMX6::RAX2 gon7 Δ ::hisMX6/GON7	see text
CCY040	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 RAX2-GFP-kanMX6::RAX2 gon7 Δ ::hisMX6	segregant from CCY039
CCY041	a/α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/met1 RAX2-GFP-kanMX6::RAX2/ RAX2-GFP-kanMX6::RAX2 gon7 Δ ::hisMX6/gon7 Δ ::hisMX6	a/α conversion from CCY040
CCY042	a	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 met15- Δ 1 rax2 Δ ::kanMX6	EUROSCARF
CCY043	α	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 lys- Δ 0 rax2 Δ ::kanMX6	Open Biosystems, Inc.
CCY044	a/α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 met15- Δ 1/lys- Δ 0 rax2 Δ ::kanMX6/rax2 Δ ::kanMX6	mating of CCY042 and CCY043
CCY045	α	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 lys- Δ 0 rax2 Δ ::kanMX6 bud32 Δ ::kanMX6	segregant from CCY002xCCY042
CCY046	a/α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 lys- Δ 0/lys- Δ 0 rax2 Δ ::kanMX6/rax2 Δ ::kanMX6 bud32 Δ ::kanMX6/bud32 Δ ::kanMX6	a/α conversion from CCY045
CCY047	α	his3- Δ 1 leu2- Δ 1 ura3- Δ 0 lys- Δ 0 rax2 Δ ::kanMX6 cgi121 Δ ::kanMX6	segregant from CCY005xCCY042
CCY048	a/α	his3- Δ 1/his3- Δ 1 leu2- Δ 1/leu2- Δ 1 ura3- Δ 0/ura3- Δ 0 lys- Δ 0/lys- Δ 0 rax2 Δ ::kanMX6/rax2 Δ ::kanMX6 cgi121 Δ ::kanMX6/cgi121 Δ ::kanMX6	a/α conversion from CCY047

CCY049	a/α his3-Δ1/his3-Δ1 leu2-Δ1/leu2-Δ1 ura3-Δ0/ura3-Δ0 met15-Δ1/met15-Δ1 rax2Δ::kanMX6/rax2Δ::kanMX6 gon7Δ::hisMX6/GON7	see text
CCY050	a his3-Δ1 leu2-Δ1 ura3-Δ0 met15-Δ1 rax2Δ::kanMX6 gon7Δ::hisMX6	segregant from CCY049
CCY051	a/α his3-Δ1/his3-Δ1 leu2-Δ1/leu2-Δ1 ura3-Δ0/ura3-Δ0 met15-Δ1/met15-Δ1 rax2Δ::kanMX6/rax2Δ::kanMX6 gon7Δ::hisMX6/gon7Δ::hisMX6	a/α conversion from CCY050
CCY052	a his3-Δ1 leu2-Δ1 ura3-Δ0 met15-Δ1 BUD32-GFP::BUD32	see text
CCY053	a/α his3-Δ1/his3-Δ1 leu2-Δ1/leu2-Δ1 ura3-Δ0/ura3-Δ0 met15-Δ1/met15-Δ1 BUD32-GFP::BUD32/BUD32-GFP::BUD32	a/α conversion from CCY052
CCY054	a his3-Δ1 leu2-Δ1 ura3-Δ0 met15-Δ1 BUD32(K52A)-GFP::BUD32	see text
CCY055	α his3-Δ1 leu2-Δ1 ura3-Δ0 lys-Δ0 BUD32(K52A)-GFP::BUD32	see text
CCY056	a/α his3-Δ1/his3-Δ1 leu2-Δ1/leu2-Δ1 ura3-Δ0/ura3-Δ0 met15-Δ1/lys-Δ0 BUD32(K52A)-GFP::BUD32/BUD32(K52A)-GFP::BUD32	mating of CCY054 and CCY055
CCY057	a his3-Δ1 leu2-Δ1 ura3-Δ0 met15-Δ1 BUD32(K52A)-GFP::BUD32 bud8Δ::kanMX6	segregant from CCY10xCCY055
CCY058	a/α his3-Δ1/his3-Δ1 leu2-Δ1/leu2-Δ1 ura3-Δ0/ura3-Δ0 met15-Δ1/met15-Δ1 BUD32(K52A)-GFP::BUD32/ BUD32(K52A)-GFP::BUD32 bud8Δ::kanMX6/ bud8Δ::kanMX6	a/α conversion from CCY057
CCY059	a his3-Δ1 leu2-Δ1 ura3-Δ0 met15-Δ1 BUD32(K52A)-GFP::BUD32 bud9Δ::kanMX6	segregant from CCY12xCCY055
CCY060	a/α his3-Δ1/his3-Δ1 leu2-Δ1/leu2-Δ1 ura3-Δ0/ura3-Δ0 met15-Δ1/met15-Δ1 BUD32(K52A)-GFP::BUD32/ BUD32(K52A)-GFP::BUD32 bud9Δ::kanMX6/ bud9Δ::kanMX6	a/α conversion from CCY059
CCY062	a his3-Δ1 leu2-Δ1 ura3-Δ0 met15-Δ1 BUD32-13myc:BUD32	see text
CCY063	a/α his3-Δ1/his3-Δ1 leu2-Δ1/leu2-Δ1 ura3-Δ0/ura3-Δ0 met15-Δ1/met15-Δ1 BUD32-13myc:BUD32/ BUD32-13myc:BUD32	a/α conversion from CCY062
CCY064	a his3-Δ1 leu2-Δ1 ura3-Δ0 met15-Δ1 BUD32(K52A):BUD32	see text
CCY065	a his3-Δ1 leu2-Δ1 ura3-Δ0 met15-Δ1 BUD32(K52A)-13myc:BUD32(K52A)	see text
CCY066	a/α his3-Δ1/his3-Δ1 leu2-Δ1/leu2-Δ1 ura3-Δ0/ura3-Δ0 met15-Δ1/met15-Δ1 BUD32(K52A)-13myc:BUD32(K52A)/ BUD32(K52A)-13myc:BUD32(K52A)	a/α conversion from CCY065

CCY067	a	his3- $\Delta$ 1 leu2- $\Delta$ 1 ura3- $\Delta$ 0 met15- $\Delta$ 1 BUD32 (S258A)-GFP::BUD32	see text
CCY068	a/ $\alpha$	his3- $\Delta$ 1/his3- $\Delta$ 1 leu2- $\Delta$ 1/leu2- $\Delta$ 1 ura3- $\Delta$ 0/ura3- $\Delta$ 0 met15- $\Delta$ 1/met15- $\Delta$ 1 BUD32(S258A)-GFP:BUD32/BUD32(S258A)-GFP::BUD32	a/ $\alpha$ conversion from CCY067
CCY069	a	his3- $\Delta$ 1 leu2- $\Delta$ 1 ura3- $\Delta$ 0 met15- $\Delta$ 1 sch9 $\Delta$ ::kanMX6	Open Biosystems, Inc.
CCY070	a/ $\alpha$	his3- $\Delta$ 1/his3- $\Delta$ 1 leu2- $\Delta$ 1/leu2- $\Delta$ 1 ura3- $\Delta$ 0/ura3- $\Delta$ 0 met15- $\Delta$ 1/met15- $\Delta$ 1 sch9 $\Delta$ ::kanMX6/sch9 $\Delta$ ::kanMX6	a/ $\alpha$ conversion from CCY069
CCY071	a	his3- $\Delta$ 1 leu2- $\Delta$ 1 ura3- $\Delta$ 0 met15- $\Delta$ 1 CDC11-GFP-kanMX6::CDC11	see text
CCY072	$\alpha$	his3- $\Delta$ 1 leu2- $\Delta$ 1 ura3- $\Delta$ 0 lys- $\Delta$ 0 CDC11-GFP-kanMX6::CDC11	see text
CCY073	a/ $\alpha$	his3- $\Delta$ 1/his3- $\Delta$ 1 leu2- $\Delta$ 1/leu2- $\Delta$ 1 ura3- $\Delta$ 0/ura3- $\Delta$ 0 met15- $\Delta$ 1/lys- $\Delta$ 0 CDC11-GFP-kanMX6::CDC11/CDC11-GFP-kanMX6::CDC11	mating of CCY071 and CCY072
CCY074	a	his3- $\Delta$ 1 leu2- $\Delta$ 1 ura3- $\Delta$ 0 met15- $\Delta$ 1 CDC11-GFP-kanMX6::CDC11 cgi121 $\Delta$ ::kanMX6/cgi121 $\Delta$ ::kanMX6	segregant of CCY071xCCY005
CCY075	a/ $\alpha$	his3- $\Delta$ 1/his3- $\Delta$ 1 leu2- $\Delta$ 1/leu2- $\Delta$ 1 ura3- $\Delta$ 0/ura3- $\Delta$ 0 met15- $\Delta$ 1/met15- $\Delta$ 1 CDC11-GFP-kanMX6::CDC11/CDC11-GFP-kanMX6::CDC11 cgi121 $\Delta$ ::kanMX6/cgi121 $\Delta$ ::kanMX6	a/ $\alpha$ conversion from CCY074

---

a All strains are congeneric to BY4743.

b The strain was generated by a precise replacement of the GON7 open reading frame by URA3.

**Table S2 Plasmids used in this study**

Plasmid	Description	Source
pBluescript (SK-)	Ampicillin (cloning)	Stratagene
pFA6a-kanMX	Kanamycin (integrative)	Longtine <i>et al.</i> (1998)
pFA6a-GFP <sup>S65T</sup> -kanMX	Kanamycin (integrative)	Longtine <i>et al.</i> (1998)
pFA6a-13Myc-kanMX	Kanamycin (integrative)	Longtine <i>et al.</i> (1998)
pRS306	<i>URA3</i> (integrative)	Sikorski and Hieter (1989)
pRS425	<i>LEU2</i> (high copy)	Christianson <i>et al.</i> (1992)
pRS426	<i>URA3</i> (high copy)	Christianson <i>et al.</i> (1992)
pTM55	<i>XhoI/SalI</i> fragment containing triple FLAG tag	our stock
pYC01	3138 bps <i>XhoI/SpeI</i> fragment containing <i>BUD8</i> gene in pRS425	see text
pYC02	3245 bps <i>XhoI/SpeI</i> fragment containing <i>BUD9</i> gene in pRS426	see text
pYC03	<i>MluI</i> restriction site at just downstream of start codon on <i>BUD8</i> in pYC01	see text
pYC04	<i>MluI</i> restriction site at just downstream of start codon on <i>BUD9</i> in pYC02	see text
pYC05	<i>MluI</i> fragments containing <i>GFP</i> sequence in pBluescript (SK-)	see text
pYC06	<i>GFP-BUD9</i> in pRS426	Kato <i>et al.</i> (2009)
pYC07	1460 bps fragment containing <i>GON7</i> gene in pBluescript (SK-)	see text
pYC08	1089 bps fragment without <i>GON7</i> gene in pBluescript (SK-)	see text
pYC09	<i>GON7::URA3</i> in pBluescript (SK-)	see text
pYC10	<i>FLAG<sub>6</sub>-BUD9</i> in pRS426	see text
pYC11	2094 bps <i>SpeI/XhoI</i> fragment containing <i>BUD32</i> in pBluescript (SK-)	see text
pYC12	<i>MluI</i> restriction site at just upstream of stop codon on <i>BUD32</i> in pYC11	see text
pYC13	<i>BUD32 (K52A)</i> in pBluescript (SK-)	see text
pYC14	<i>GFP-BUD8</i> in pRS425	Kato <i>et al.</i> (2009)
pYC15	<i>BUD32 (S258A)</i> in pBluescript (SK-)	see text
pYC16	<i>BUD32 (K52A)</i> in pRS306	see text
pYC17	<i>BUD32-GFP</i> in pBluescript (SK-)	see text
pYC18	<i>BUD32 (K52A)-GFP</i> in pBluescript (SK-)	see text
pYC19	<i>BUD32 (S258A)-GFP</i> in pBluescript (SK-)	see text
pYC20	<i>BUD32-GFP</i> in pRS306	see text



pYC21	<i>BUD32 (K52A)-GFP</i> in pRS306	see text
pCY22	<i>BUD32 (S258A)-GFP</i> in pRS306	see text

---

<sup>a</sup>The GFP sequences encode *GFP* with the S65T substitution.

---

**Table S3 Primers used in this study**

Primer	sequences
T7 promoter	TAATACGACTCACTATAGGG
M13 reverse	CAGGAAACAGCTATGAC
BUD8a	AAAAAA <b>CTCGAG</b> ACTTCAACCACCTTCA
BUD8b	AAAAAA <b>ACTAGT</b> AGAAGTACGGTTACAAGACT
BUD8c	<b>ACGCGT</b> ATACAATCAGACGAAG
BUD8d	CATACTTCATGTAGAATCG
BUD9a	AAAAAA <b>CTCGAG</b> CACAAGAACATTTCTG
BUD9b	AAAAAA <b>ACTAGT</b> CTCTATCGTACTCCTC
BUD9c	<b>ACGCGT</b> ACGAAAATAACCAGAG
BUD9d	CATTCATAGGATGAAGAATG
BUD32a	AAAAAACTAGTAAACGTCGATTTTACCAG
BUD32b	AAAAAACTCGAGGATAGTCATTCTTTCC
BUD32c	AAGAGAAGTATGCTAGGAACGCGTTAATAAATGCTAGCG
BUD32d	GAGAAGTATGCTAGGAACGCGTTATAAATGCTAGCGTA
BUD32e	GCATATAGGCCACCAAAGCGTTATAG
BUD32f	GATAATATACTTTTGATGAGAATCCTTTGC
BUD32g	GCTATGCTAGGATAATAAATGCTAGCG
BUD32h	TCTCTTACGACCACGCAACCTG
BUD32i	AAAAGATTCTGAAGAGGTCAGGTTGCGTGGTCGTAAGAAGTATGCTAGGAGCAGGTCGA
BUD32j	TGTGCAGCGATATACAGGCAGTACGCTAGCATTTATATTAGAATTCGAGCTCGTTAAAC
GFPa	AAAAAAACGCGTGGGTTAATTAACAGTAAA
GFPb	<b>AACGCGT</b> TTTTGTATAGTTCATCCATGCC
GFPc	<b>AGATATC</b> GGGTTAATTAACAGTAAAGGAG
GFPd	<b>AGATATC</b> TTTTGTATAGTTCATCCATGCCATG
GON7a	AAAAGTAAATAGCGATTCTTCATCGTGAATG
GON7b	AAACTCGAGTAAACGGTCATTTGTTTTAACACAG
GON7c	GCAACGATATGTAGCGCTAGAG
GON7d	<b>ACGCGT</b> TGAGATATTAGGATAAGGTTC

GON7e	TATACAGCCGATAGTGCCTGGAACCTTATCCTAATATCTCACGGATCCCCGGTTAATT
GON7f	GACTCTTTTCTGTTTGTATATACTCTCTAGCGCTACATATCGT
URA3a	AAAA <b><i>ACGCGT</i></b> AAGCTTTGGCACATCAATG
URA3b	AAAA <b><i>ACGCGT</i></b> ACCGTTGTTATCAGAAATTC
RAX2a	AGAATGAAATGCTTGATACCGTCCACCCGAAAACTTATGAAGTTTGCCGGATCCCCGGTTAATTAA
RAX2b	AGTGTTCAATTTTTAAGTAGTTATATATTATATAATAACAACCCGATTAGAATTCGAGCTCGTTTAAAC
CDC11a	ATTGAAGCCAGGTTGGAAAAAGAGGCGAAAATCAAACAGGAAGAACGGATCCCCGGTTAATTAA
CDC11b	ATATAGAGAAAGAAGAAATAAGTGAGGAAGCCAAAAGCGGACGAATTCGAGCTCGTTTAAAC

---

The bold italics indicate restriction sites.

---