

Sir,
Visual impairment in children

Certification of visual impairment is used to register persons as visually impaired or severely visually impaired according to set criteria. This data is reported nationally and allows comparisons between regions and ages, and is essential in planning health, social, and educational services.¹

Given the size of the population of Bradford district, it would be expected that 1% of national visual impairment would be accounted for by the region. However, reported data suggests that Bradford district accounts for 13% of the national registered visual impairment and 16% of registered severe visual impairment in 0–4 year olds. This figure drops to 3.5% and 3% of visual impairment and severe visual impairment, respectively, in 5–17 year olds.

The age-specific rates of registered severe visual impairment and visual impairment in Bradford district are compared with the national average in Tables 1 and 2. These rates are significantly higher in Bradford district compared with the national average (*Z*-test, *P* < 0.001).

There are a number of possible explanations for these differences. Bradford district has a higher prevalence than the national average of children born with some forms of disability that may be associated with visual impairment,² thought to be particularly prevalent in Black and minority ethnic (BME) groups. This may be due, in part, to a high prevalence of autosomal recessive and inherited disorders.³ Areas with similar BME populations, such as Calderdale, Walsall, and Oldham, have none or very few cases of registered visual impairment. Therefore, this suggests that other causes may be contributing to the higher rate.

Another explanation may be differences in registering visual impairment between regions such as differing levels of access to an ophthalmologist. As Bradford

district has a high prevalence of visual impairment, clinicians may be more confident in registering children at an early age than areas with lower prevalences.

Registration rates of visual impairment in children is higher in Bradford district than the rest of England. The cause of this is unclear and is likely to be because of the combination of a genuinely higher prevalence of visual impairment locally and underreporting of visual impairment nationally.

References

- 1 NHS Information Centre. Registered Blind and Partially Sighted People Year Ending 31 March 2008 England. http://www.ic.nhs.uk/webfiles/publications/Social%20Care/Blind%202008/Blind%20Commentary%202008_final%20v5.pdf. (Accessed on 4 August 2010).
- 2 Corry PC. Intellectual disability and cerebral palsy in the UK community. *Community Genet* 2002; 5: 201–204.
- 3 Schwarz K, Yeung S, Symons N, Bradbury J. Survey of school children with visual impairment in Bradford. *Eye* 2002; 16(5): 530–534.

M Kliner¹, G Fell¹, RF Pilling² and JA Bradbury²

¹Department of Public Health, NHS Bradford and Airedale, Bradford, West Yorkshire, UK

²Department of Ophthalmology, Bradford Royal Infirmary, Bradford, West Yorkshire, UK
E-mail: meravkliner@gmail.com

Eye (2011) 25, 1097; doi:10.1038/eye.2011.85;
published online 15 April 2011

Table 1 Rates of registered severe visual impairment per 1000 population in England and Bradford district

	Rate of severe visual impairment in 0–4 years per 1000 population (95% confidence intervals)	Rate of severe visual impairment in 5–17 years per 1000 population (95% confidence intervals)
England	0.22 (0.20, 0.23)	0.44 (0.43, 0.45)
Bradford and Airedale TPCT	3.30 (2.73, 3.87)	1.43 (1.17, 1.68)

Table 2 Rates of visual impairment per 1000 population in England and Bradford district

	Rate of visual impairment in 0–4 years per 1000 population (95% confidence intervals)	Rate of visual impairment in 5–17 years per 1000 population (95% confidence intervals)
England	0.19 (0.18, 0.20)	0.57 (0.55, 0.58)
Bradford and Airedale TPCT	2.28 (2.28, 2.76)	2.26 (1.94, 2.58)

Sir,
Avulsed retinal vessels accompanying posterior vitreous detachment: a late complication of retinopathy of prematurity

Abnormal retinal vasculature and vitreous, in combination with aberrant vitreoretinal traction are factors in the development of late complications in adults with retinopathy of prematurity (ROP).¹ A case of intact arcades of retinal blood vessels separated from the retina (see Supplementary movie file), secondary to posterior vitreous detachment (PVD) in an adult with ROP is presented. We wish to highlight this rare late complication of ROP, which was described in adults by Tasman² in 1970 and in neonates by Kingham³ in 1982.

Case report

A 59-year-old woman attended eye casualty 6 months after uncomplicated cataract surgery with a 1-day history of floaters in her right eye. She was a high myope with a history of ROP (twin birth, 10 weeks premature). In the past, she had suffered from recurrent vitreous haemorrhage in this eye. The left eye was phthisical as a result of endophthalmitis after cataract surgery.

On examination, the right eye was pseudophakic with visual acuity of 6/9. Fundoscopy showed the optic disc and retinal blood vessels dragged nasally (Figure 1a). In the peripheral fundus, there was chorioretinal atrophy and a falciform fold. A PVD accompanied by intact