

CLINICAL UTILITY GENE CARD

Clinical utility gene card for: CHARGE syndrome

Kim Blake^{*,1}, Conny MA van Ravenswaaij-Arts², Lies Hoefsloot³ and Alain Verloes⁴

European Journal of Human Genetics (2011) **19**, doi:10.1038/ejhg.2011.45; published online 16 March 2011

1. DISEASE CHARACTERISTICS

1.1 Name of the disease (synonyms)

CHARGE syndrome (CHARGE association, Hall–Hittner syndrome).

1.2 OMIM# of the disease

#214800.

1.3 Name of the analysed genes or DNA/chromosome segments

CHD7.

1.4 OMIM# of the gene(s)

#608892.

1.5 Mutational spectrum

Predominantly heterozygous point mutations (72% nonsense or frame shift, 13% splice site and 10% missense). Less than 5% of the whole-exon deletions or microdeletions of 8q12.1, including CHD7.^{1–3}

1.6 Analytical methods

Sequencing of all coding exons, including their boundaries of CDH7, MLPA covering most coding exons, including the 5'UTR and the first non-coding exon of CHD7. Array CGH in selected cases.^{4,5}

Conventional cytogenetics is usually normal. Translocations with breakpoint through CHD7 have been reported incidentally (Jongmans¹).

1.7 Analytical validation

Sequence analysis detects >99% of the (point) mutations present in the area that has been investigated, MLPA has an estimated sensitivity of >90% for individual exons, and >95% for deletions covering more probes.

1.8 Estimated frequency of the disease

(incidence at birth ('birth prevalence') or population prevalence)

Prevalence at birth: 1:10 000 (ranges from 1:8500 to 1:15 000 in the literature).⁶

1.9 If applicable, prevalence in the ethnic group of investigated person

There is no evidence at present for a different prevalence in various ethnic groups.

1.10 Diagnostic setting

	Yes	No
A. (Differential) diagnostics	<input checked="" type="checkbox"/>	<input type="checkbox"/>
B. Predictive testing	<input checked="" type="checkbox"/>	<input type="checkbox"/>
C. Risk assessment in relatives	<input checked="" type="checkbox"/>	<input type="checkbox"/>
D. Prenatal	<input checked="" type="checkbox"/>	<input type="checkbox"/>

Comment:

2. TEST CHARACTERISTICS

Test	Genotype or disease		A: True positives	C: False negatives
	Present	Absent	B: False positives	D: True negatives
Positive	A	B	Sensitivity:	A/(A+C)
			Specificity:	D/(D+B)
Negative	C	D	Positive predictive value:	A/(A+B)
			Negative predictive value:	D/(C+D)

2.1 Analytical sensitivity

(proportion of positive tests if the genotype is present)

Depends on the method used. If only CHD7 sequencing is performed, deletions are missed less than 5% due to whole-exon or whole-gene deletions.^{5,7} If sequencing is combined with MLPA, 100%.

2.2 Analytical specificity

(proportion of negative tests if the genotype is not present)

Almost 100% (some variants may erroneously be interpreted as pathogenic).

2.3 Clinical sensitivity

(proportion of positive tests if the disease is present)

The clinical sensitivity can be dependent on variable factors such as age or family history. In such cases a general statement should be given, even if a quantification can only be made case by case.

Depends on the clinical criteria used. In over 95% of the patients who fulfil the criteria of Blake or Verloes^{8,9} a mutation is found.¹ In those who are suspected for CHARGE syndrome in 60–70% a mutation is found

¹Department of Pediatrics, Dalhousie University, Halifax, Nova Scotia, Canada; ²Department of Genetics, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands; ³Department of Human Genetics, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands; ⁴AP-HP, Groupe Hospitalier Pitié-Salpêtrière, UF de Génétique Clinique, Paris, France

*Correspondence: Professor K Blake, Department of Pediatrics, Dalhousie University, 5850/5980 University Avenue, Halifax, Nova Scotia, Canada. Tel: +1 902 470 6499; Fax: +1 902 470 6913; E-mail: kblake@dal.ca

(CHARGE syndrome sometimes can be excluded if a patient does not fulfil the clinical criteria and does not carry a mutation or deletion of CHD7).

Some conditions can mimic CHARGE syndrome: 22q11 deletion syndrome, VACTERL association, chromosomal disorders (eg, deletions 3p12p21.2¹⁰), disorders caused by teratogens (eg, maternal diabetes, Accutane), and Kallmann syndrome.

2.4 Clinical specificity (proportion of negative tests if the disease is not present)

The clinical specificity can be dependent on variable factors such as age or family history. In such cases a general statement should be given, even if a quantification can only be made case by case.

The clinical variability of the syndrome is considerable. If the diagnosis is based on the Blake or Verloes criteria, some people with CHARGE will be missed. The clinical specificity is over 95%, since less than 5% of the patients with a CHD7 mutation do not completely fulfil these criteria. However, it should be taken into account that the mild end of the phenotypic spectrum is not completely known yet. For example, CHD7 mutations are also found in patients diagnosed with Kallmann syndrome or hypogonadotropic hypogonadism with minimal additional features of CHARGE syndrome.^{11,12}

2.5 Positive clinical predictive value (life-time risk to develop the disease if the test is positive)

100%, but high clinical variability (see also 2.4).

2.6 Negative clinical predictive value (probability of not developing the disease if the test is negative)

Assume an increased risk based on family history for a non-affected person. Allelic and locus heterogeneity may need to be considered.

Index case in that family had been tested:

100%.

Index case in that family had not been tested:

This depends on the *a priori* chance of the index to find a mutation, which varies between 60–90%. There is always a residual risk, but complete analysis (sequencing and MLPA) will reduce this by 90–95%.

3. CLINICAL UTILITY

3.1 (Differential) diagnosis: The tested person is clinically affected (To be answered if in 1.10 'A' was marked)

3.1.1 Can a diagnosis be made other than through a genetic test?

No	<input type="checkbox"/> (continue with 3.1.4)	
Yes	<input checked="" type="checkbox"/>	
	Clinically	<input checked="" type="checkbox"/>
	Imaging	<input checked="" type="checkbox"/>
	Endoscopy	<input type="checkbox"/>
	Biochemistry	<input type="checkbox"/>
	Electrophysiology	<input type="checkbox"/>
	Other (please describe)	

3.1.2 Describe the burden of alternative diagnostic methods to the patient.

CHARGE syndrome can be diagnosed clinically but not by solely using the CHARGE acronym (C=coloboma, H=heart defect, A=choanal Atresia, R=retardation of growth and development, and E=ear abnormalities) or the major criteria (coloboma of the eyes, choanal atresia, characteristic ear malformations including deafness, and cranial nerve abnormalities). Scanning of the temporal bones often elicits abnormalities in the semi circular canals, which brings more specificity to the diagnosis.¹³

3.1.3 How is the cost effectiveness of alternative diagnostic methods to be judged?

Gene testing is still expensive, and for that reason many parents, particularly those with older children, have not had their child tested.¹⁴ Either gene testing or clinical criteria can miss some individuals with CHARGE syndrome. However, gene testing may be important in patients who do not have the classical CHARGE characteristics and may be at risk for the long-term complications of CHARGE syndrome.

3.1.4 Will disease management be influenced by the result of a genetic test?

No	<input type="checkbox"/>	
Yes	<input checked="" type="checkbox"/>	
	Therapy (please describe)	Depends on clinical manifestations: severe gastroesophageal reflux resulting in tube feeding, problems with swallowing and aspiration, and secretions are a cluster of the hidden problems in CHARGE Syndrome. Earlier identification of hearing and visual loss with multidisciplinary educational team, physiotherapy, occupational therapy, speech therapy, psychology. Some patients require growth hormone and many require hormones to enter puberty. Problems with bone mineral density means nutritional therapy and physiotherapy, early in life and into adolescence (increased levels of vitamin D and calcium may be required).
	Prognosis (please describe)	A positive genetic test will alert professionals to a diagnosis of CHARGE syndrome. An early diagnosis and management of the sensory impairments may improve prognosis. Parents are also required to be highly proactive and organized with this syndrome involving such complexity. Parents can also become involved with the international CHARGE Syndrome Support Group.
	Management (please describe)	This can be divided up into early middle and late. Early management may involve repair of choanal atresia, cardiac abnormalities, and tracheoesophageal fistula. Attention to gastroesophageal reflux and the swallowing mechanism is an important early management problem that is often forgotten. ¹⁵ Botox injections into the salivary glands have recently proven effective. ¹⁶ Middle – attention to physical therapy for balance problems. Occupational therapy for fine motor and feeding problems. Speech and language therapy for swallowing, feeding problems, and speech. In the last 8 years, children with CHARGE syndrome are less likely to have tracheostomies, but are more likely to have G or J tube feeding. Watching for hidden anomalies such as renal disease, t-cell dysfunction, continued assessment of hearing, and vision with a view to baha or cochlear implants. Late – hormone issues are common, growth hormone deficiency is less prevalent than abnormalities of sex hormones, especially in boys. Vitamin D and calcium intake throughout childhood and into adolescent hood needs to be maximized. We recommend increase calcium and vitamin D intake to protect bone development. ¹⁷

3.2 Predictive setting: The tested person is clinically unaffected but carries an increased risk based on family history

(To be answered if in 1.10 'B' was marked)

3.2.1 Will the result of a genetic test influence lifestyle and prevention?

If the test result is positive (please describe):

If the test result is positive, yes; see 3.1.4.

If the test result is negative (please describe):

If the test result is negative, depends on clinical manifestations.

3.2.2 Which options in view of lifestyle and prevention does a person at risk have if no genetic test has been done (please describe)?

No special options.

3.3 Genetic risk assessment in family members of a diseased person (To be answered if in 1.10 'C' was marked)

3.3.1 Does the result of a genetic test resolve the genetic situation in that family?

Not necessarily, but if a mutation is found in the patient and not in the parents, recurrence risk can be given with more certainty and prenatal diagnosis can be offered. Moreover, in that situation sibs of the patient do not have an increased risk for having children with CHARGE syndrome. If the mutation is found in mosaic or non-mosaic form in one of the parents, recurrence risk is increased (up to 50%) and prenatal or pre-implantation diagnosis should be discussed.

3.3.2 Can a genetic test in the index patient save genetic or other tests in family members?

Family members may have very mild characteristics of CHARGE syndrome and therefore should be tested. Somatic mosaicism has been described in parents. So in case of child wish, parents should be tested to obtain a more accurate recurrence risk.

3.3.3 Does a positive genetic test result in the index patient enable a predictive test in a family member?

Yes. If an index patient has CHARGE syndrome other family members can be genetically tested.

3.4 Prenatal diagnosis

(To be answered if in 1.10 'D' was marked)

3.4.1 Does a positive genetic test result in the index patient enable a prenatal diagnosis?

Yes. If an index patient has CHARGE syndrome, then subsequent pregnancies can be screened genetically and by ultrasound.

4. IF APPLICABLE, FURTHER CONSEQUENCES OF TESTING

Please assume that the result of a genetic test has no immediate medical consequences. Is there any evidence that a genetic test is nevertheless useful for the patient or his/her relatives? There is a lot of

discussion in Hartshorne *et al*¹² that might be helpful, including reasons for testing or not testing given by parents and a discussion of the ethical issues involved.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by EuroGentest, an EU-FP6-supported NoE, contract number 512148 (EuroGentest Unit 3: 'Clinical genetics, community genetics and public health', Workpackage 3.2).

- Jongmans MC, Admiraal RJ, van der Donk KP *et al*: CHARGE syndrome: the phenotypic spectrum of mutations in the CHD7 gene. *J Med Genet* 2006; **43**: 306–314.
- Lalani SR, Safiullah AM, Fernbach SD *et al*: Spectrum of CHD7 mutations in 110 individuals with CHARGE syndrome and genotype-phenotype correlation. *Am J Hum Genet* 2006; **78**: 303–314.
- Zentner GE, Layman WS, MArtin DM, Scacheri PC: Molecular and phenotypic aspects of CHD7 mutation in CHARGE syndrome. *Am J Med Genet Part A* 2010; **152A**: 674–686.
- Vissers LE, van Ravenswaaij CMA, Admiraal R *et al*: Mutations in a new member of the chromodomain gene family cause CHARGE syndrome. *Nat Genet* 2004; **36**: 955–957.
- Bergman JEH, de Wijs I, Hoefsloot LH, Jongmans MCJ, van Ravenswaaij CMA: Exon copy number alterations of the CHD7 gene are not a major cause of CHARGE syndrome. *Eur J Med Genet* 2008; **51**: 417–425.
- Issekutz K, Graham Jr J, Prasad C, Smith IM, Blake KD: Behavioural profiles and symptoms of autism in CHARGE syndrome: preliminary results from a Canadian study. *Am J Med Genet* 2005; **133A**: 309–317.
- Wincent J, Schulze A, Schoumans J: Detection of CHD7 deletions by MLPA in CHARGE syndrome patients with a less typical phenotype. *Eur J Med Genet* 2009; **52**: 271–272.
- Blake KD, Davenport SLH, Hall BD *et al*: CHARGE association: an update and review for the primary pediatrician. *Clin Genet* 1998; **37**: 159–173.
- Verloes A: Updated diagnostic criteria for CHARGE syndrome; a proposal. *Am J Med Genet* 2005; **133A**: 306–308.
- Wieczorek D, Bolt J, Schwachheimer K, Gillissen-Kaesbach G: A patient with interstitial deletion of the short arm of chromosome 3 (pterrrrp21.2::p12rarrqter) and a CHARGE-like phenotype. *Am J Med Genet* 1997; **69**: 413–417.
- Kim HG, Kurth I, Lan F *et al*: Mutations in CHD7, encoding a chromatin-remodelling protein, cause idiopathic hypogonadotropic hypogonadism and Kallmann syndrome. *Am J Hum Genet* 2008; **83**: 511–519.
- Jongmans MCJ, van Ravenswaaij CMA, Pitteloud N *et al*: CHD7 mutations in patients initially diagnosed with Kallmann Syndrome - the clinical overlap with CHARGE syndrome. *Clin Genet* 2009; **75**: 65–71.
- Sanlaville D, Verloes A: CHARGE syndrome: an update. *Eur J Hum Genet* 2007; **15**: 389–399.
- Hartshorne TS, Stratton KK, van Ravenswaaij-Arts CM: Prevalence of genetic testing in CHARGE syndrome. *J Genet Couns* 2011; **20**: 49–57.
- Dobbelsteyn C, Peacocke SD, Blake K, Crist W, Rashid M: Feeding difficulties in children with CHARGE syndrome: prevalence, risk factors, and prognosis. *Dysphagia* 2008; **23**: 127.
- MacCuspie J, Blake K, Corsten G: Botulinum toxin injections into salivary glands to decrease oral secretions in CHARGE syndrome: prospective case study. *Am J Med Genet* 2010.
- Forward K, Cummings EA, Blake KD: Risk factors for poor bone health in adolescent and adults with CHARGE syndrome. *Am J Med Genet* 2007; **143A**: 839–845.