

Review Article

Molecular aspects of renal cell carcinoma: a review

Hari Koul¹, Jung-Sik Huh¹, Kyle O. Rove², Luiza Crompton², Sweaty Koul¹, Randall B. Meacham², Fernando J. Kim³

¹Signal Transduction and Molecular Urology Laboratory, Program in Urosciences, Division of Urology, University of Colorado Comprehensive Cancer Center, Aurora, CO 80045, USA; ²University of Colorado Denver, Anschutz Medical Campus School of Medicine, Division of Urology, Department of Surgery, Aurora, CO 80045, USA; ³Denver Health Medical Center, 777 Bannock Street, Denver, CO 80204, USA.

Received December 13, 2010; accepted December 17, 2010; Epub December 18, 2010; Published January 1, 2011

Abstract: Renal cell carcinoma (RCC) is a disease in which cancer cells form in the tubules of the kidney. RCC, the incidence of which is increasing annually, represents five percent of adult epithelial cancers. Clear cell carcinoma represents the most frequent histological subtype. RCC is characterized by a lack of early warning signs, diverse clinical manifestations. Incidentally detected tumors in asymptomatic individuals have been steadily increasing owing to the increased usage of various imaging technologies. Currently there are no recommendations for screening to detect and make an early diagnosis of renal cancer. But in recent years, the discovery of new molecular and cytogenetic markers has led to the recognition and classification of several novel subtypes of RCC, and the introduction of molecular-targeted therapy for advanced-stage RCC. We performed a literature review using PubMed and discuss current knowledge of epidemiology, pathophysiology, evaluation, treatment, and future research directions of RCC.

Keywords: Renal cell carcinoma, von hippel lindau, chemotherapy

Introduction

Renal cell carcinoma (RCC), the incidence of which is increasing annually, will account for approximately 3.8% of adult malignancies and 90-95% of neoplasms arising from the kidney in 2010 [1]. Up to 30% of RCC patients present at advanced stages, and approximately 40% of patients who undergo curative surgical resection experience recurrence during subsequent follow-up [1,2]. RCC is characterized by a lack of early warning signs, diverse and variable clinical manifestations, resistance to radiation and chemotherapy, and infrequent but reproducible responses to immunotherapeutic agents such as interferon alpha (INF- α) and interleukin (IL-2). Newer agents, such as sorafenib and sunitinib, which are orally available, are multi-targeted tyrosine kinase inhibitors which specifically interfere with platelet-derived growth factor receptor (PDGF-R) and vascular endothelial growth factor (VEGF) [3,4]. Temsirolimus, a mammalian target of rapamycin (mTOR) inhibitor, was also recently approved by the FDA for the treatment

of RCC [5]. This article focuses on the epidemiology, pathophysiology, histology, staging, diagnostic evaluation, therapeutic options, and follow-up of this disease.

Epidemiology

There will be an estimated 58,240 new cases of and 13,040 deaths from renal cancer in 2010 in the United States, accounting for 2.3% of all cancer deaths in the United States [1]. The incidence has steadily increased during the past 50 years in the United States and has occurred in 9.1 per 100,000 in 1997, with a mortality rate of 3.5 per 100,000 [6,7,8]. The American Cancer Society predicts that there will be about 54,000 cases of kidney cancer in the United States in 2008, and about 13,010 people will die from this disease [9]. Reported worldwide incidence rates range from 0.6 per 100,000 to 14.7 per 100,000 [10]. Most tumors present in the fifth to seventh decades of life, with a median age at diagnosis of 66 years and median age at death of 70 years. The incidence is two

to three times higher in men and is slightly more common in blacks than in whites [6]. At autopsy, the incidence of renal tumors is approximately 2% [11]. In general, the tumors are usually solitary but may be multifocal in 6-25% of patients, and bilateral disease is diagnosed in 4% of all RCC patients [12]. Certain genetic conditions are associated with an increased incidence of RCC, including von Hippel-Lindau disease, hereditary papillary renal cancer, and possibly tuberous sclerosis [13]. RCC occurs in von Hippel-Lindau disease in 35-40% of patients, arises at a younger age, and is frequently bilateral (75%) or multifocal (87%) [14]. In these patients, the disease has a 5:1 male predominance. Overall, other suggested risk factors include cigarette smoking, obesity, diuretic use, exposure to petroleum products, chlorinated solvents, cadmium, lead, asbestos, ionizing radiation, high-protein diets, hypertension, kidney transplantation, and HIV infection [6,12,15,16].

Pathophysiology

Loss of the Von Hippel-Lindau tumor suppressor gene & HIF-related events in RCC tumor biology

Identification of the Von Hippel-Lindau (VHL) gene and understanding the function of this pathway in RCC tumorigenesis has played a major role in the development of RCC therapeutics. VHL disease is a hereditary cancer syndrome that has proven to be highly informative with respect to the pathogenesis of clear cell RCC. In affected families, cancer risk is transmitted in an autosomal dominant manner on chromosome 3p, and the syndrome is manifested by retinal angiomas, central nervous system hemangiomas and clear cell RCC [17]. Individuals with VHL disease carry in their germline one wild-type VHL allele and one inactivated VHL allele. Pathologic changes ensue in VHL disease when the remaining wild-type allele is somatically inactivated in a susceptible cell type. Therefore, VHL is a classic two-hit tumor suppressor gene [17].

It has been shown that inactivation of the VHL gene is an early step in the development of clear cell RCC associated with VHL disease [17]. The VHL tumor suppressor gene is mutated in all hereditary RCC and approximately 50% of sporadic RCC, and thus the majority of clear cell RCCs appear to be linked to biallelic VHL inactivation (see **Table 1**) [18]. The majority of clear

cell RCC demonstrates either a mutation of the VHL gene or downregulation of its protein product. Translation of VHL mRNA gives rise to protein that is referred to generically as VHL protein (pVHL), which has an important role in the cellular response to hypoxia [19]. Among the many functions attributed to pVHL, the one most clearly linked to the development of RCC is inhibition of hypoxia-inducible factor (HIF) [20]. HIF is a heterodimeric transcription factor consisting of an unstable α -subunit (HIF1 α) and a stable β -subunit [21].

Under normal oxygen conditions, the pVHL complex polyubiquitinates HIF1 α , tagging it for destruction by the proteasome [22]. Under low oxygen conditions or in cells lacking pVHL, HIF1 α accumulates, binds to HIF1 β and transcriptionally activates hypoxia-inducible genes [21].

The consequence of mutated pVHL is similar to that of cellular hypoxia causing HIF dimerization and stabilization. During hypoxia, there is accumulation of hydroxyl-free HIF that no longer binds to VHL. HIF1 α is stabilized by dimerization with the constitutively expressed HIF1 β subunit and translocates to the nucleus. The HIF1 α and HIF1 β complex binds to HIF inducible gene promoter regions, including genes implicated in angiogenesis, pH regulation, glycolysis, glucose transport, cell cycle, chemotaxis, signaling, and apoptosis [23].

VEGF-R pathway

Biallelic loss of VHL leads to upregulated transcription of growth factors such as VEGF, PDGF and TGF- α . These factors bind to their respective receptors (VEGF-R, PDGF-R and EGF-R), which are each tyrosine kinase receptors. Consequent binding of ligands to these receptors leads to downstream signaling that results in increased cell proliferation, upregulated angiogenesis and decreased apoptosis. RCC-associated mutations of pVHL are invariably inactivating the process of HIF destruction, suggesting that HIF plays a critical role in RCC carcinogenesis. Numerous HIF responsive genes have been described, with a number of these genes encoding proteins that are growth factor receptors or their ligands, some of which were listed above [24]. A number of HIF responsive gene products are implicated in tumorigenesis. Uncontrolled production of these growth factors

Molecular aspects of renal cell carcinoma

Table 1. Genetic alterations in renal epithelial neoplasms

Tumor type	Chromosome	Gene	Mechanism and manifestation	Other genetic alterations
Clear cell renal cell carcinoma	3p14.2 3p21 3p25	FHIT RASSF1A VHL	Deletion, mutation, methylation	+5q22,-6q,-8p12,-9p21,-9q22,-10q,-14q
Papillary renal cell carcinoma	7 17 7q31.1 7q31 Y	? FRA7G c-MET ?	Trisomy Trisomy Gain Gain Loss	+3q,+8,-9p21,+12,-14q,+16,+17q21,+20
Chromophobe renal cell carcinoma	1,2,6,10,13,17,21,Y	?	Multiple chromosome loss	-5q22,-8p,-9p23,-18q22
Oncocytoma	1,14 11q13	?	Loss Translocation	-1p,-8p,-11q13,14q,-19q,-21q,-X/Y, der(13)t(13;16)(p11;p11)
Collecting duct carcinoma	1,2,6,10,13q,13,14,15,22	?	Deletion	-1q32,-6p,-8p,-9p,-13q,-19q32,-21q
Renal carcinoma associated with Xp11.2 translocation	1p34 1q21 17q23 17q25 3q23 Xq12	PSF-TFE3 PRCC-TFE3 CLTC-TFE3 ASPL-TFE3 ? Non O-TFE3	t(X;1)(p11.2;p34) t(X;1)(p11.2;q21) t(X;17)(p11.2;q23) t(X;17)(p11.2;q25) t(X;3)(p11;q23) inv(X)(p11.2;q12)	
Mucinous tubular and spindle cell carcinoma	1,4,6,8,9,11,13,14,1,18,22	?	Multiple chromosome loss	-8p,-9p,-11q,+12q,+16q,+17,+20q
Metanephric adenoma	2p13 2p	?	Deletion Partial monosomy	Inv(9)(p12q13),t(1,22)(q22;q13), t(15;16)(q21;p13)

provides a stimulus for the tumor and endothelial cell proliferation.

Angiogenic stimuli produced secondary to metabolic demands of host tissues initiate the angiogenic response in healthy individuals. Upon binding to membrane receptors in vascular endothelial cells, a five-step process is triggered. Initially, the vascular endothelial basement membrane of the parent vessel breaks down, allowing a route for the development of a new capillary sprout. This is followed by migration of endothelial cells and chemoattraction [12]. This leading front of migrating cells is driven by enhanced proliferation of endothelial cells, followed by formation of capillary tubes via organization of the endothelial cells, and a recruitment of pericytes and vascular smooth muscle cells for capillary stabilization [25].

During tumorigenesis, the angiogenic switch is activated directly via induction of angiogenic growth factors or indirectly by recruiting host immune cells that release mediators of angiogenesis [26].

Induction of the HIF pathway results in production of VEGF. VEGF a key regulator of angiogenesis and functions are mediated through two tyrosine kinase receptors, VEGF-R1 and VEGF-R2, in vascular endothelial cells [27,28]. VEGF initially interacts with VEGF-R2 to promote endothelial cell proliferation, migration and vascular permeability, and subsequently activates VEGF-R1 to assist in the organization of new capillaries. Several therapeutics targeting angiogenic pathways are currently being evaluated in clinical trials for their efficacy and long-term clinical benefits, while others are being mechanistically

exploited toward the development of novel therapeutic modalities for treating RCC.

mTOR pathway

Another regulator of HIF1 α levels in the cell is mTOR, whose signaling activity acts to increase the cellular levels of HIF1 α , accentuating the overall elevation in levels caused by the absence of adequate pVHL function [29]. mTOR inhibitors have been previously described, and laboratory experiments have shown that anti-proliferative effects of these inhibitors in RCC may result from the interruption of essential survival pathways and autophagy [30]. The effect of mTOR inhibitors on angiogenesis is likely to have an important function in RCC pathogenesis, a highly vascular tumor [30].

mTOR is a highly conserved serine/threonine kinase that forms quaternary complexes and has a key function in apoptosis, cell growth and tumor proliferation by controlling cellular catabolism and anabolism [30]. mTOR may complex with a regulatory-associated protein of mTOR to form mTORC1 and can also complex with a rapamycin-insensitive companion of mTOR, to form another multimolecular complex named mTORC2. (Rapamycin is an inhibitor of mTOR.) mTORC1 may eventually be activated by growth factors and also the VEGF-R, PDGF-R, EGF-R, IGF receptor, and phosphatidylinositol 3-kinase/Akt (PI3K/Akt) pathways [31]. At the molecular level, it is known that tumor angiogenesis depends on vascular growth factors such as VEGF, PDGF, bFGF and members of the TGF- β superfamily. As these factors have been shown to be able to activate the PI3K/Akt/mTOR in cancer cells, endothelial cells or pericytes as described, mTOR complexes are also implicated in tumor angiogenesis biology [32].

Once activated, mTORC1 acts through its downstream effectors to stimulate protein synthesis, entrance into the G1 phase of the cell cycle and asserts control over proteins that regulate apoptosis. The witnessed activity of mTOR inhibitors in RCC has raised the possibility that patients who respond to this therapy share a common molecular phenotype that renders these tumors dependent on mTOR for growth and survival. In RCC, the PTEN gene has been shown to be downregulated in the majority of cases, presumably by epigenetic silencing [33].

It has been previously demonstrated that PI3K/

Akt/mTOR signaling pathway inhibitors target tumor growth indirectly at the tumor level by interacting with the maintenance of endothelial cells and pericytes that are required for tumor angiogenesis [34].

A major stimulus of cancer angiogenesis is tissue hypoxia likely driven by tumorigenesis and growth initially lacking adequate blood supply. These conditions activate HIF1 α , and the mTOR pathway further enhances the translation of HIF1 α mRNA, thereby increasing the overall vasculogenic effect [35]. The observed clinical efficacy of mTOR inhibitors in RCC is mediated in part by dependence on efficient HIF translation in the mTOR pathway by interfering with the VEGF/VEGF-R and/or PDGF/PDGF-R signaling cascades. In summary, these data strongly suggest that the anticancer effects of mTOR inhibitors involve antiangiogenetic processes mediated by effects on endothelial cells and pericytes, rather than on RCC themselves [29].

NF- κ B pathway

NF- κ B is a family of transcription factors that has been associated with diverse cellular functions. NF- κ B activation is associated with increased proliferation, tissue invasion, angiogenesis, inhibition of apoptosis, and the development of drug resistance [36]. NF- κ B activation has also been associated with proliferative responses mediated by induction of expression of cyclin D1, which drives the transition from the G1 to the S phase of the cell cycle [37]. VHL loss ultimately drives NF- κ B activation by resulting in HIF α accumulation, which induces expression of transforming growth factor alpha (TGF- α), with consequent activation of an EGF-R/phosphatidylinositol-3-OH kinase/protein kinase B (AKT)/I κ B-kinase alpha/NF- κ B signaling cascade [38].

NOx pathway

Reactive oxygen species (ROS) regulate hypoxia-dependent and hypoxia-independent activation of HIF1 α . NAD(P)H oxidase systems are major sources of ROS. The NOx family of NAD(P)H oxidases have a core structure consisting of six transmembrane domains, including two heme-binding regions located at the N terminus and a cytoplasmic C terminus containing FAD- and NADPH-binding regions [39,40]. Reactive oxygen species, generated by NAD(P)H oxidases, are involved in signaling cascades of malignant

growth. In VHL-deficient cells, NOx4 protein levels and NAD(P)H-dependent superoxide generation are increased. Reintroduction of VHL into the VHL-deficient cells down-regulates NADPH-dependent superoxide generation [41].

Tyrosin kinase pathway

Receptor tyrosine kinases (RTKs) constitute a superfamily of transmembrane proteins that relays signals from extracellular growth factors into the cell [42,43]. The TAM subfamily of RTKs contains the receptors Axl, Tyro3, and Mer [44,45]. They have in common a unique extracellular domain composed of two N-terminal immunoglobulin-like domains and two fibronectin type III repeats similar to the structure of neural cell adhesion molecules (NCAMs). TAM receptors share the same ligand, Gas6, a product of the growth arrest-specific gene 6 [46,47]. Gas6, cloned from serum-starved fibroblasts, is a member of the vitamin K-dependent family of Gla proteins homologous to the blood coagulation protein S [48]. Axl has been shown to affect neovascularization in vitro, and loss of Axl expression in tumor cells blocks growth of human neoplasms [49]. Perhaps, Axl on its own, by homophilic interactions and by a kinase domain-dependent mechanism [50], contribute to the disease-specific angiogenic programming during VHL loss in tumor cells in parallel with angiogenic factors such as VEGF. Gas6 signaling via Axl, on the other hand, has been shown to have inhibitory effects on the VEGFR-driven angiogenic program [51]. Gas6-mediated activation of Axl in clear cell carcinoma cells results in Axl phosphorylation, receptor down-regulation, decreased cell-viability and migratory capacity [52].

Mitogen-activated protein kinase pathway

Mitogen-activated protein kinase (MAPK) kinases (MKK) are crucial enzymes at the intersection of several biological pathways that regulate cell differentiation, proliferation, and survival. In response to a variety of extracellular stimuli, MKKs become activated and then phosphorylate MAPKs, including extracellular signal-regulated protein kinase (ERK), c-Jun-NH2 kinase (JNK), and p38 MAPK (p38) [53,54]. Overexpression of MKK has been described in human RCC cases [55]. Sustained activation of ERK has been established as a requirement for angiogenesis as well [56,57].

HSP70 pathway

The major hsp70 are encoded by a duplicated locus (hsp70-1, hsp70-2) located in the MHC region, 92 kb telomeric to the C2 gene [58]. This segment of the MHC has been proposed to be termed the class IV region since it includes at least seven genes implicated to some degree in inflammation and in stress responses [59]. The two intronless genes (hsp70-1 and hsp70-2) encode an identical protein product of 641 amino acids. Hsp70-2 may have a potential role in cancer pathogenesis by participating in the regulation of antitumor immunity such as acting as a chaperone molecule for immunogenic tumor-associated peptides but also in regulatory processes such as the cell cycle. The possibility that a mutated hsp70-2 chaperone might have a dominant effect in tumor cells in triggering the G2/M phase transition during mitotic cell cycle cannot be excluded at the present time. Interestingly, among more than 100 RCC tumors studied in the laboratory, RCC-7 is the most aggressive one, with a rapid doubling interval in vitro and a high growth rate in SCID/nu mice. Preliminary experiments testing either the immortalizing or the transforming capacity of the mutated versus wild-type hsp70-2 cDNA in recipient cells have not led, however, to any direct evidence that the mutation plays a role in the oncogenic process [60].

Clinical course of RCC

Symptomatology and presentation

RCC can remain clinically occult for the majority of its course. The classic clinical presentation of flank pain, hematuria, and a palpable mass is relatively uncommon (only 5-10% of cases). In addition, clinical symptomatology, if ever present, is often nonspecific—for example, anorexia, fatigue, weight loss, or fever of unknown origin [61]. Other clinical manifestations include varicocele formation in men (from tumor thrombus in the left renal vein or the inferior vena cava or from extrinsic compression of these structures impairing return of blood from gonadal veins, more commonly left than right) and disseminated malignancy. RCC may also present with a variety of paraneoplastic syndromes, such as polycythemia secondary to excessive secretion of erythropoietin, hypercalcemia secondary to derangement of serum factors regulating calcium, and hepatic dysfunction

(Stauffer syndrome). Incidentally detected tumors in asymptomatic individuals have been steadily rising with the increase in use of imaging techniques, including computed tomography (CT), MRI, and ultrasonography. Incidental lesions accounted for approximately 60% of renal tumors during the 1990s, compared with only 10% in the early 1970s [62].

Pathology

On gross pathology, tumors most often appear encapsulated. Tumors may be solid, cystic, or mixed and can have calcification present [63]. As many as 10% of tumors have some cystic component [64], and such tumors may be more clinically aggressive [65]. Each of these subtypes of RCC has different cytogenetic and immunohistochemical profiles. Histopathologic grading of the nuclei of the tumor is made by dividing them into the four-tier Fuhrman nuclear classification [66], with grade I representing well-differentiated and grade IV the most anaplastic, poorly differentiated.

Histologic subtypes according to the Heidelberg classification (and relative incidence) include clear cell (conventional) adenocarcinoma (80%), papillary (15%), chromophobe (5%), collecting duct (1%), and unclassified (4%) [67-69].

Clear cell carcinoma displays large uniform cells with abundant clear cytoplasm rich in glycogen and lipid. Clear cell carcinoma is typically highly vascular. Papillary tumors are subdivided into type I tumors, which occur sporadically and metastasize somewhat late and are often composed of papillae covered by single-layered small cells with pale cytoplasm and round ovoid nuclei. Type II papillary RCC lesions, which are more likely inherited, may be multicentric, have pseudostratified large cells with relatively abundant eosinophilic cytoplasm, and often present with a higher Fuhrman grade and poorer prognosis. Collecting duct tumors arise from the medullary collecting duct and may show morphological features of both adenocarcinoma and urothelial carcinoma, often occur in younger patients and are associated with a poor overall prognosis. Renal medullary carcinoma is a rare subtype of collecting duct carcinoma and has a very poor prognosis. It is seen more commonly in young patients with sickle cell anemia or the sickle trait. Chromophobe tumors and oncocytomas, both of which arise from collecting duct epithelium, are similar on histologic examina-

tion but have differing immunohistochemical profiles. Chromophobe tumors have the best overall prognosis, and oncocytomas are benign [18,70].

Staging

The tumor, nodes, and metastases (TNM) classification is endorsed by the American Joint Committee on Cancer (AJCC). The major advantage of the TNM system is that it clearly differentiates individuals with tumor thrombi from those with local nodal disease. In the Robson system, stage III inferior vena caval involvement (IIIA) is the same stage as local lymph node metastases (IIIB) [71]. These entities can have highly differing clinical courses. The TNM classification system is presented in **Tables 2, 3** and **4**.

Diagnostic evaluation

Currently, there are no accepted methods for screening or early diagnosis of renal cancer. Although population-based screening by renal ultrasound would undoubtedly detect renal tumors in asymptomatic individuals, this is not considered cost effective, and the implications of population screening from a public health perspective are enormous. Cost, increased morbidity and mortality, and reduced quality of life from increased detection and intervention cannot be ignored when considering secondary prevention (screening) strategies. Neither the US Preventative Task Force nor American Cancer Society, for example, has official screening recommendations for or against screening for RCC. No serological tests are currently available for the early detection of renal cancer. Initial tests performed at the time of diagnosis include urinalysis, CBC count with differential, electrolytes, renal profile (serum creatinine and/or glomerular filtration rate), liver function tests, serum calcium, erythrocyte sedimentation rate, prothrombin time, activated partial thromboplastin time. Other tests may be indicated for symptomatic patients.

A large proportion of patients diagnosed with renal cancer have small tumors. A number of different diagnostic imaging modalities, such as excretory urography, CT scan, ultrasonography, arteriography, venography, magnetic resonance imaging (MRI), and positron emission tomography (PET), are used to evaluate and stage renal masses.

Molecular aspects of renal cell carcinoma

Table 2. TNM Staging of Renal Cell Carcinoma. Adapted from current version of American Joint Committee On Cancer (2002)

Stage	Description
TX	Primary tumor cannot be assessed
T0	No evidence of primary tumor
T1	Tumor < 7 cm in greatest dimension, limited to kidney
T1a	Tumor < 4 cm in greatest dimension, limited to kidney
T1b	Tumor > 4 cm but < 7 cm in greatest dimension, limited to kidney
T2	Tumor ≥7 cm in greatest dimension, limited to kidney
T3	Tumor extends into major veins or invades adrenal gland or perinephric tissues, but not beyond Gerota's fascia
T3a	Tumor invades adrenal gland or perinephric tissues but not beyond Gerota's fascia
T3b	Tumor grossly extends into renal vein(s) or vena cava below diaphragm
T3c	Tumor grossly extends into vena cava above diaphragm
T4	Tumor invades beyond Gerota's fascia
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in a single regional lymph node
N2	Metastasis in more than one regional lymph node
MX	Distant metastasis cannot be assessed
M0	No distant metastasis
M1	Distant metastasis

Table 3. TNM stage groupings. Adapted from current version of American Joint Committee On Cancer (2002)

Stage Grouping	T stage	N stage	M stage
I	T1	N0	M0
II	T2	N0	M0
III	T1	N1	M0
	T2	N1	M0
	T3	N0	M0
	T3	N1	M0
IV	T4	N0	M0
	T4	N1	M0
	Any T	N2	M0
	Any T	Any N	M1

The goals of radiologic imaging include detection and staging of the primary tumor. In most institutions, CT is the primary imaging technique used for the evaluation of renal tumors. In some instances, such as when a patient has an allergy to iodinated contrast medium, MRI or ultrasonography are suitable alternatives. With MRI and its common gadolinium-based contrast agents, there is a small risk of nephrogenic sys-

temic fibrosis in patients with impaired renal function and should be considered carefully when staging with MRI is deemed advisable [72].

For the evaluation of metastatic disease and local recurrence, CT has excellent sensitivity and is the standard imaging test. Nuclear medicine studies with PET have limited sensitivity for

Table 4. Survival by TNM stage. Adapted from current version of American Joint Committee On Cancer (2002)

Disease Extent	TNM Stage	5-year Survival Rate (%)
All organ-confined	T1-T2NOMO	70-90
<4cm	T1aNOMO	90-100
>4 but <7cm	T1bNOMO	80-90
>7cm	T2NOMO	70-80
Invasion of perinephric fat	T3aNOMO	60-80
Adrenal involvement	T3aNOMO	0-30
Venous involvement	T3b-T3cNOMO	40-65
Locally advanced	T4NOMO	0-20
Lymphatic involvement	Any T,N+,M0	0-20
Systemic metastases	Any T, any N,M1	1-10

evaluating metastatic RCC and particularly for small metastatic lesions. However, a positive PET scan should be considered strongly suspicious for local recurrence or metastasis, because of the high specificity and PPV of this test. A combined test (PET-CT) may be necessary if important management decisions are to be based on the test result. This would take advantage of the high sensitivity of CT and high specificity of PET in patients with metastatic RCC [73,74].

Chest CT should be performed if the primary tumor is large or locally aggressive because metastases are more common in these patients. Chest radiography without CT should be reserved for patients with a low risk of metastatic disease or for those in long-term follow-up [75]. Sonography can be useful for assessing the presence and extent of venous thrombus. It can also be helpful in distinguishing cysts from hypovascular solid tumors seen on CT (e.g., papillary RCC). Sonography can reveal septations within a lesion better because of complex interfaces and the ultrasound beam. Sonography has reported accuracies for T staging of 77-85% [76,77] and for detection of venous thrombus of 87% of the time [78]. However, it has limitations in visualizing the retroperitoneum and perinephric tissues [78,79], although some proponents argue otherwise [77].

MRI is generally only used when optimal CT imaging cannot be performed (iodinated contrast allergies, poor renal function, pregnancy). MRI has similar reported overall staging accuracies to those of CT [80]. Its multiplanar capability, however, is particularly useful for delineating the superior extent of tumor in the IVC [81]. Pre-

operative percutaneous biopsy of renal lesions is generally not undertaken because the results usually do not affect what therapy will be recommended except in patients with multiple tumors or occasionally in patients with an underlying predisposing condition. Percutaneous biopsies may be considered in selected cases—for example, when an abscess or metastatic disease from a known primary tumor is suspected, especially from lymphoma or melanoma [12].

Approximately 20% of patients have multiple renal arteries, and many surgeons find preoperative CT or MR angiograms to be valuable, particularly when partial nephrectomy or laparoscopic approaches are planned. Three-dimensional and multiplanar reformatted images, as well as angiographic displays, aid appreciation of the relationships of the tumor to the collecting system, adjacent normal parenchyma, and vascular supply at the renal pedicle [12,82,83].

The efficacy of PET in renal malignancy remains under investigation. It shows some potential in staging, the detection of unsuspected metastases, follow-up, and the evaluation of indeterminate renal masses, but CT remains the standard of care [84-87].

Treatment

Radical nephrectomy, which remains the most commonly performed standard surgical procedure today for treatment of localized renal carcinoma, involves complete removal of Gerota's fascia and its contents, including a resection of kidney, perirenal fat, and ipsilateral adrenal gland, with or without ipsilateral lymph node

dissection. Radical nephrectomy provides a better surgical margin than simple removal of the kidney, since perinephric fat may be involved in some patients. Twenty to thirty percent of patients with clinically localized disease develop metastatic disease after nephrectomy. Some surgeons believe that the adrenal gland should not be removed because of the low probability of ipsilateral adrenal metastasis and the morbidity associated with adrenalectomy. In the absence of distant metastatic disease with locally extensive and invasive tumors, adjacent structures such as bowel, spleen, or psoas muscle may be excised en bloc during radical nephrectomy [12].

Ipsilateral adrenalectomy is included in the classic radical nephrectomy. However, ipsilateral adrenal metastases occur in only 1-10% of patients. It occurs especially in large left-sided upper pole tumors, usually by direct extension (i.e., T3a tumors) [88]. One of the roles of imaging is to assist in allowing adrenal-sparing nephrectomies in order to reduce the risk of future adrenal insufficiency [8].

Laparoscopic nephrectomy is a less invasive procedure than open radical nephrectomy, incurs less morbidity, and is associated with shorter recovery time and less blood loss. The need for pain medications is reduced, but operating room time and costs are higher. Disadvantages include concerns about spillage and technical difficulties in defining surgical margins. Laparoscopic partial nephrectomy can be considered at centers with experience in this procedure for early stage renal cell cancers generally less than 4 cm in largest dimension [12].

Palliative nephrectomy should be considered in patients with metastatic disease for alleviation of symptoms such as pain, hemorrhage, malaise, hypercalcemia, erythrocytosis, or hypertension. Several randomized studies show improved overall survival in patients presenting with metastatic kidney cancer who have nephrectomy followed by either interferon or IL-2. If the patient has good physiological status, then nephrectomy should be performed prior to immunotherapy [89].

Nephron-sparing surgery (i.e., partial nephrectomy) has been shown to be equally as efficacious as radical nephrectomy, with reported local recurrence rates of < 2% and 5-year survival rates of 87-90%, which are comparable to

those from radical nephrectomy [59]. This more limited treatment of a presumed cancerous lesion is generally reserved for lesions less than 4 cm in size along the largest dimension, for patients with poor renal reserve or function, or solitary kidney. In the case of a solitary kidney, though, there is increased risk of developing proteinuria, focal segmental glomerulosclerosis, and progressive renal failure exists if more than 50% of the renal mass is removed [83].

Micrometastasis to lymph nodes may be present in 10-25% of patients. The 5-year survival rate in patients with regional node involvement is substantially lower than in patients with stage I or II disease. Regional lymphadenectomy adds little in terms of operative time or risk and should be included in conjunction with radical nephrectomy [12,83,90,91]. Cryoablation and radiofrequency ablation, which may be undertaken laparoscopically or percutaneously, are promising techniques for treating small tumors [83].

Chemotherapy

A recent phase 2 trial of weekly intravenous gemcitabine (600 mg/m² on days 1, 8, and 15) with continuous infusion fluorouracil (150 mg/m²/day for 21 days in 28-day cycle) in patients with metastatic renal cell cancer produced a partial response rate of 17%. No complete responses were noted. Eighty percent of patients had multiple metastases, and 83% had received previous treatment. The mean progression-free survival duration of 28.7 weeks was significantly longer than that of historic control [91].

Floxuridine (5-fluoro 2'-deoxyuridine [FUDR]), 5-fluorouracil (5-FU), and vinblastine, paclitaxel (Taxol), carboplatin, ifosfamide, gemcitabine, and anthracycline (doxorubicin) have each been studied and used in the treatment of metastatic RCC. Floxuridine infusion has a mean response rate of 12%, while vinblastine infusion yielded an overall response rate of 7%. 5-FU alone has a response rate of 10%, but when used in combination with interferon, it had a 19% response rate in some studies [92].

RCC is refractory to most chemotherapeutic agents because of multidrug resistance mediated by p-glycoprotein normally present in all cells. This protein is responsible for efflux of drugs and other compounds deemed foreign to

cells. Normal renal proximal tubules and renal cell carcinoma both express high levels of p-glycoprotein [93].

Follow-Up

For stage I and II disease, complete history, physical examination, chest radiographs, liver function tests, BUN and creatinine, and calcium are recommended every six months for two years, then annually for five years. Abdominal CT scan is recommended once at 4-6 months and then as indicated.

For stage III renal cell carcinoma, physical examination, chest radiographs, liver function tests, BUN and creatinine, and calcium are recommended every 4 months for 2 years, every 6 months for 3 years, and then annually for 5 years. Abdominal CT scan should be performed at 4-6 months, then annually or as indicated [94].

Prognosis

Investigators have attempted to identify pathological and morphological features within tumors that correlate with survival in patients with RCC. Tumor stage remains the most important factor predictive of survival in RCC [96-99]. In addition, tumor size [97], histological pattern [99], cell type [99], nuclear grade [97], DNA content [100], and nuclear morphometry [101] have been reported as prognostic surrogates for survival. Additionally, variables such as performance status [102], weight loss, time to progression, number and type of metastases [103], vascular invasion [104] and several laboratory values, e.g. haemoglobin level, ESR and alkaline phosphatase levels, have been studied in relation to prognosis [103]. Increasing knowledge of cytogenetic abnormalities and the role of oncogenes and tumor suppressor genes in RCC is critical, but the study of molecular mechanisms underlying RCC is still in its infancy; future studies will have to provide information on the implications for the prognosis of patients with RCC [105].

Immunotherapy and targeted therapies

The mainstay of systemic therapy for metastatic RCC has historically been immunotherapy (or cytokine therapy) with interleukin-2 (IL-2) and interferon- α (IFN- α). High-dose IL-2 has consis-

tently produced a 15-20% response rate, 6-8% complete remission rate, and approximately 5% cure rate, however, it is a fairly toxic regimen [106,107]. IFN- α has provided modest survival benefit, has a more favorable toxicity profile, and is more easily administered than IL-2. As a result, IFN- α has been adopted as the control arm in many clinical trials evaluating the merits and characteristics of novel agents.

Novel therapies for metastatic RCC have targeted downstream effects of von Hippel-Lindau (VHL) gene inactivation and the resulting up-regulation of HIF target genes, notably VEGF and PDGF. Five targeted agents are presently in clinical use for metastatic RCC: sunitinib, sorafenib, temsirolimus, everolimus, and bevacizumab. Sunitinib (Sutent, Pfizer), interacts with an adenosine triphosphate binding site in multiple tyrosine kinase domains and prevents autophosphorylation; this multiple tyrosine kinase inhibitor (MKI) has shown an objective response rate of 31% and a median progression-free survival of 11 months for sunitinib-treated patients [108,109]. It is considered the standard-of-care treatment for patients with advanced-, good- and intermediate-risk conventional clear cell RCC. Sorafenib (Nexavar, Bayer HealthCare), another MKI, is considered second-line therapy after cytokine failure [4]. Temsirolimus (Torisel, Wyeth), an inhibitor of mammalian target of rapamycin (mTOR), is considered appropriate for patients with poor-risk metastatic RCC, irrespective of histology [110,111]. Bevacizumab (Avastin, Genentech), a humanized antibody to VEGF, has shown promise in patients with good- and intermediate-risk metastatic RCC [112,113] but has not yet been approved for metastatic RCC by the U.S. Food and Drug Administration (**Table 5**).

Bevacizumab (+ interferon-alpha), sunitinib, and temsirolimus (in poor-risk groups) have proven to be effective as first-line palliative treatments. Sorafenib has demonstrated benefits in patients that have failed prior therapy, as has everolimus after failure of sorafenib and/or sunitinib [114].

Conclusion

Renal cell carcinomas comprise a heterogenous group of epithelial tumors with variable clinical outcomes. Hopefully, genetic hallmarks will be identified in the various histological subtypes

Molecular aspects of renal cell carcinoma

Table 5. Clinical activity of selected vascular endothelia growth factor-targeted agents in advanced renal cell carcinoma

Reference	Agent	No	Trial design	Prior Therapy	Overall response rate
Yang et al	Bevacizumab	116	Randomized phase II high-dose bevacizumab versus low dose bevacizumab versus placebo	Cytokines	10% vs 0%
Motzer et al	Sunitinib	63, 106	Single-arm phase II single agent sunitinib	Cytokines	40%,34%
Motzer et al	Sunitinib	750	Randomized phase III sunitinib versus IFN-a	None	31 vs 6%
Escudier et al	Sorafenib	903	Randomized phase III sorafenib versus placebo	Various	2% vs 0%
Hudes G et al	Temsirolimus	626	Randomized phase III temsirolimus versus IFN-a versus temsirolimus+IFN-a	None	9% vs 7% vs 11%

and improve understanding of the antiapoptotic and immune escape mechanisms that allow for oncogenic growth. These mechanisms may then be specifically targeted with novel agents in the treatment of advanced disease and will result in novel approaches allowing rational combinations of drugs with immunostimulating effect. Much work still needs to understand, treat, and ultimately control RCC.

Please address correspondence to: Hari K. Koul, PhD, Signal transduction and Molecular Urology Laboratory, Division of Urology, Department of Surgery, University of Colorado School of Medicine, Denver Veteran's Medical Center and The University of Colorado Cancer Center, Building RC2, 12700 E 19th Avenue, Room 6001, Aurora, CO 80045, USA. Tel: (303)724-6300, E-mail: hari.koul@ucdenver.edu

References

- [1] Jemal A, Siegel R, Xu J, Ward E. Cancer Statistics, 2010. *CA Cancer J Clin* 2010; 60: 277-300.
- [2] Janzen NK, Kim HL, Figlin RA, Belldegrun AS. Surveillance after radical or partial nephrectomy for localized renal cell carcinoma and management of recurrent disease. *Urol Clin North Am* 2003; 30: 843-52.
- [3] Abrams TJ, Lee LB, Murray LJ, Pryer NK, Cherrington JM. SU11248 Inhibits KIT and platelet-derived growth factor receptor β in preclinical models of human small cell lung cancer. *Mol Cancer Ther* 2003; 2: 471-8.
- [4] Escudier B, Eisen T, Stadler WM, Szczylik C, Oudard S, Siebels M, Negrier S, Chevreau C, Solska E, Desai AA, Rolland F, Demkow T, Hutson TE, Gore M, Freeman S, Schwartz B, Shan M, Simantov R, Bukowski RM; TARGET

Study Group. Sorafenib in advanced clear-cell renal-cell carcinoma. *N Engl J Med* 2007; 356: 125-34.

- [5] Oya M. Renal cell carcinoma: Biological features and rationale for molecular-targeted therapy. *Keio J Med* 2009; 58: 1-11.
- [6] Chow WH, Devesa SS, Warren JL, Fraumeni JF Jr. Rising incidence of renal cell cancer in the United States. *JAMA* 1999; 281: 1628-31.
- [7] Katner HP, Baynham SA. Increasing incidence of renal cell cancer. *JAMA* 1999; 282: 2119-20.
- [8] Pantuck AJ, Zisman A, Belldegrun AS. The changing natural history of renal cell carcinoma. *J Urol* 2001; 166: 1611-23.
- [9] Arsanoious A, Bjarnason GA, Yousef GM. From bench to bedside: current and future applications of molecular profiling in renal cell carcinoma. *Molecular Cancer* 2009; 8: 20-31.
- [10] Parkin DM, Pisani P, Ferlay J. Estimates of the worldwide incidence of 25 major cancers in 1990. *Int J Cancer* 1999; 80: 827-41.
- [11] Choyke PL, Amis ES Jr, Bigongiari LR, Bluth EI, Bush WH Jr, Fritzsche P, Holder L, Newhouse JH, Sandler CM, Segal AJ, Resnick MI, Rutsky EA. Renal cell carcinoma staging. American College of Radiology. ACR appropriateness criteria. *Radiology* 2000; 215: 721-5.
- [12] Russo P. Renal cell carcinoma: presentation, staging, and surgical treatment. *Semin Oncol* 2000; 27: 160-76.
- [13] Bjornsson J, Short MP, Kwiatkowski DJ, Henske EP. Tuberous sclerosis-associated renal cell carcinoma: clinical, pathological, and genetic features. *Am J Pathol* 1996; 149: 1201-8.
- [14] Levine E, Weigel JW, Collins DL. Diagnosis and management of asymptomatic renal cell carcinomas in von Hippel-Lindau syndrome. *Urology* 1983; 21: 146-50.
- [15] Figlin RA. Renal cell carcinoma: management of advanced disease. *J Urol* 1999; 161: 381-7.

Molecular aspects of renal cell carcinoma

- [16] Godley P, Kim SW. Renal cell carcinoma. *Curr Opin Oncol* 2002; 14: 280-5.
- [17] Kim WY, Kaelin WG. Molecular pathways in renal cell carcinoma - rationale for targeted treatment. *Semin Oncol* 2006; 33: 588-95.
- [18] Cheug L, Zhang S, MacLennan GT, Lopez-Beltran A, Montironi R. Molecular and cytogenetic insights into the pathogenesis, classification, differential diagnosis, and prognosis of renal epithelial neoplasms. *Human Pathology* 2009; 40: 10-29.
- [19] Blakenship C, Naglich JG, Whalet JM, Seizinger B, Kley N. Alternate choice of initiation codon procedures a biologically active product of the von Hippel Lindau gene with tumor suppressor activity. *Oncogene* 1999; 18: 1529-35.
- [20] Kaelin WG. Molecular basis of the VHL hereditary cancer syndrome. *Nat Rev Cancer* 2002; 2: 673-82.
- [21] Semenza GL. HIF-1 and mechanisms of hypoxia sensing. *Curr Opin Cell Biol* 2001; 13: 167-71.
- [22] Jaakkola P, Mole DR, Tian YM, Wilson MI, Gielbert J, Gaskell SJ, et al. Targeting of HIF- α targeted to the von Hippel Lindau ubiquitylation complex by O₂ regulated prolyl hydroxylation. *Science* 2001; 292: 468-72.
- [23] Harris AL. Hypoxia - a key regulatory factor in tumor growth. *Nat Rev Cancer* 2002; 2: 38-47.
- [24] Caldwell MC, Hough C, Fürer S, Linehan WM, Morin PJ, Gorospe M. Serial analysis of gene expression in renal carcinoma cells reveals VHL-dependent sensitivity to TNF- α cytotoxicity. *Oncogene* 2002; 21: 929-36.
- [25] Sakamoto S, Ryan AJ, Kyprianu N. Targeting vasculature in Urologic tumors: mechanistic and therapeutic significance. *J Cell Biochem* 2008; 103: 691-708.
- [26] Folkman J. Clinical applications of research on angiogenesis. *N Engl J Med* 1995; 333: 1757-63.
- [27] de Vries C, Escobedo JS, Ueno H, Houck K, Ferrara N, Williams LT. The fms-like tyrosine kinase, a receptor for vascular endothelial growth factor. *Science* 1992; 255: 989-91.
- [28] Millauer B, Wizigmann-Voos S, Schnürch H, et al. High affinity VEGF binding and developmental expression suggests flk-1 as a major regulator of vasculogenesis and angiogenesis. *Cell* 1993; 72: 835-46.
- [29] Cho D, Signoretti S, Regan M, Mier JW, Atkins MB. The role of mammalian target of rapamycin inhibitors in the treatment of advanced renal cancer. *Clin Cancer Res* 2007; 13: S758-S763.
- [30] Le Tourneau C, Faivre S, Raymond E. mTORC1 inhibitors: is temsirolimus in renal cancer telling us how they really work? *Br J Cancer* 2008; 99: 1197-203.
- [31] Faivre S, Kraeme G, Ramond E. Current development of mTOR inhibitors as anticancer agents. *Nat. Rev. Drug Discov* 2006; 5: 671-88.
- [32] Guba M, von Breitenbuch P, Steinbauer M, Koehl G, Flegel S, Hornung M, Bruns CJ, Zuelke C, Farkas S, Anthuber M, Jauch KW, Geissler EK. Rapamycin inhibits primary and metastatic tumor growth by anti-angiogenesis involvement of vascular endothelial growth factor. *Nat Med* 2002; 8: 128-35.
- [33] Brenner W, Farber G, Herget T, Lehr HA, Hengstler JG, Thuroff W. Loss of tumor suppressor protein PTEN during renal angiogenesis. *Int J Cancer* 2002; 99: 53-7.
- [34] Hamada K, Sasaki T, Koni PA, et al. The PTEN/PI3K pathway governs normal vascular development and tumor angiogenesis. *Genes Dev* 2005; 19: 2054-65.
- [35] Arsham AM, Plas DR, Thompson CB, Simon MC. Phosphatidylinositol 3-kinase/Akt signaling is neither required for hypoxic stabilization of HIF-1 α nor sufficient for HIF-1 dependent target gene transcription. *J Biol Chem* 2002; 277: 15162-70.
- [36] Baldwin AS. Control of oncogenesis and cancer therapy resistance by the transcription factor NF- κ B. *J Clin Invest* 2001; 107: 241-6.
- [37] Guttridge DC, Albanese JY, Pestell PG, Baldwin AS Jr. NF- κ B controls cell growth and differentiation through transcriptional regulation of cyclin D1. *Mol Cell Biol* 1999; 19: 5785-99.
- [38] An J, Rettig MB. Mechanism of von Hippel-Lindau Protein-Mediated Suppression of Nuclear Factor κ B Activity. *Mol Cell Biol* 2005; 25: 7546-56.
- [39] Srinivas V, Leshchinsky I, Sang N, King MP, Minchenko A, Caro J. *J Biol Chem* 2001; 276: 21995-8.
- [40] Chandel NS, McClintock DS, Feliciano CE, et al. *J Biol Chem* 2000; 275: 25130-8.
- [41] Block K, Gorin Y, Hoover P, et al. NAD(P)H oxidases regulate HIF-2 α protein expression. *J Biol Chem* 2007; 282: 8019-26.
- [42] Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell* 2000; 103: 211-25.
- [43] Robinson DR, Wu YM, Lin SF. The protein tyrosine kinase family of the human genome. *Oncogene* 2000; 19: 5548-57.
- [44] Hafizi S, Dahlback B. Signalling and functional diversity within the Axl subfamily of receptor tyrosine kinases. *Cytokine Growth Factor Rev* 2006; 17: 295-304.
- [45] Linger RM, Keating AK, Earp HS, Graham DK. TAM receptor tyrosine kinases: biologic functions, signaling, and potential therapeutic targeting in human cancer. *Adv Cancer Res* 2008; 100: 35-83.
- [46] Hafizi S, Dahlback B. Gas6 and protein S. Vitamin K-dependent ligands for the Axl receptor tyrosine kinase subfamily. *Febs J* 2006; 273: 5231-44.
- [47] Bellido-Martin L, de Frutos PG. Vitamin K-dependent actions of Gas6. *Vitam Horm* 2008; 78: 185-209.

Molecular aspects of renal cell carcinoma

- [48] Manfioletti G, Brancolini C, Avanzi G, Schneider C. The protein encoded by a growth arrest-specific gene (*gas6*) is a new member of the vitamin K-dependent proteins related to protein S, a negative coregulator in the blood coagulation cascade. *Mol Cell Biol* 1993; 13: 4976-85.
- [49] Holland SJ, Powell MJ, Franci C, Chan EW, et al. Multiple roles for the receptor tyrosine kinase *axl* in tumor formation. *Cancer Res* 2005; 65: 9294-303.
- [50] Bellosta P, Costa M, Lin DA, Basilico C. The receptor tyrosine kinase *ARK* mediates cell aggregation by homophilic binding. *Mol Cell Biol* 1995; 15: 614-25.
- [51] Gallicchio M, Mitola S, Valdembrì D, Fantozzi R, Varnum B, Avanzi GC, Bussolino F. Inhibition of vascular endothelial growth factor receptor 2-mediated endothelial cell activation by *Axl* tyrosine kinase receptor. *Blood* 2005; 105: 1970-6.
- [52] Gustafsson A, Boström AK, Ljungberg B, Axelsson H, Dahlbäck B. *Gas6* and the receptor tyrosine kinase *Axl* in clear cell renal cell carcinoma. *PLoS One* 2009 Oct 30; 4: e7575.
- [53] Chang L, Karin M. Mammalian MAP kinase signalling cascades. *Nature* 2001; 410: 37-40.
- [54] Johnson GL, Lapadat R. Mitogen-activated protein kinase pathways mediated by ERK, JNK, and p38 protein kinases. *Science* 2002; 298: 1911-2.
- [55] Oka H, Chatani Y, Hoshino R, Ogawa O, Kakehi Y, Terachi T, Okada Y, Kawaichi M, Kohno M, Yoshida O. Constitutive activation of mitogen-activated protein (MAP) kinases in human renal cell carcinoma. *Cancer Res* 1995; 55: 4182-7.
- [56] Dong G, Chen Z, Li ZY, Yeh NT, Bancroft CC, Van Waes C. Hepatocyte growth factor/scatter factor-induced activation of MEK and PI3K signal pathways contributes to expression of proangiogenic cytokines interleukin-8 and vascular endothelial growth factor in head and neck squamous cell carcinoma. *Cancer Res* 2001; 61: 5911-8.
- [57] Eliceiri BP, Klemke R, Stromblad S, Cheresh DA. Integrin $\alpha v \beta 3$ requirement for sustained mitogen-activated protein kinase activity during angiogenesis. *J Cell Biol* 1998; 140: 1255-63.
- [58] Milner, C. M., R. D. Campbell. Structure and expression of the three MHC-linked HSP70 genes. *Immunogenetics* 1990; 32: 242-51.
- [59] Gruen JR, Weissman SM. Evolving views of the major histocompatibility complex. *Blood* 1997; 90: 4252.
- [60] Gaudin C, Kremer F, Angevin E, Scott V, Triebel F. A *hsp70-2* mutation recognized by CTL on a human renal cell carcinoma. *J Immunol* 1999; 162: 1730-8.
- [61] Luciani LG, Cestari R, Tallarigo C. Incidental renal cell carcinoma: age and stage characterization and clinical implications, a study of 1092 patients (1982-1997). *Urology* 2000; 56: 58-62.
- [62] Jayson M, Sanders H. Increased incidence of serendipitously discovered renal cell carcinoma. *Urology* 1998; 51: 203-5.
- [63] Hilton S. Imaging of renal cell carcinoma. *Semin Oncol* 2000; 27: 150-9.
- [64] Parianty RA, Pradel J, Parianty I. Cystic renal cancers: CT characteristics. *Radiology* 1985; 157: 741-4.
- [65] Brinker DA, Amin MB, de Peralta-Venturina M, et al. Extensively necrotic cystic renal cell carcinoma: a clinicopathologic study with comparison to other cystic and necrotic renal cancers. *Am J Surg Pathol* 2000; 24: 988-95.
- [66] Fuhrman SA, Lasky LC, Limas C. Prognostic significance of morphologic parameters in renal cell carcinoma. *Am J Surg Pathol* 1982; 6: 655-63.
- [67] Kovacs G, Akhtar M, Beckwith BJ, Bugert P, Cooper CS, Delahunt B, Eble JN, Fleming S, Ljungberg B, Medeiros LJ, Moch H, Reuter VE, Ritz E, Roos G, Schmidt D, Srigley JR, Störkel S, van den Berg E, Zbar B. The Heidelberg classification of renal cell tumors. *J Pathol* 1997; 183: 131-3.
- [68] Shanks JH. Pathology of renal cell carcinoma: recent developments. *Clin Oncol (R Coll Radiol)* 1999; 11: 263-8.
- [69] Bostwick DG, Eble JN. Diagnosis and classification of renal cell carcinoma. *Urol Clin North Am* 1999; 26: 627-35.
- [70] Ng CS, Wood CG, Silverman PM, et al. Renal cell carcinoma: diagnosis, staging, and surveillance. *Am J Roentgenol* 2008; 191: 1220-32.
- [71] American Joint Committee on Cancer. *AJCC cancer staging manual*. New York, NY: Springer-Verlag, 2002.
- [72] Broome DR, Girguis MS, Baron PW, Cottrell AC, Kjellin I, Kirk GA. Gadodiamide-associated nephrogenic systemic fibrosis: why radiologists should be concerned. *Am J Roentgenol* 2007; 188: 586-92.
- [73] Schöder H, Larson S. Positron emission tomography for prostate, bladder, and renal cancer. *Semin Nucl Med* 2004; 34: 274-92.
- [74] Coll DM, Uzzo RG, Herts BR, Davros WJ, Wirth SL, Novick AC. Three-dimensional volume rendered computerized tomography for preoperative evaluation and intraoperative treatment of patients undergoing nephron sparing surgery. *J Urol* 1999; 161: 1097-102.
- [75] Chin AI, Lam JS, Figlin RA, Belldegrün AS. Surveillance strategies for renal cell carcinoma patients following nephrectomy. *Rev Urol* 2006; 8: 1-7.
- [76] Frohmüller HG, Grups JW, Heller V. Comparative value of ultrasonography, computerized tomography, angiography and excretory urography in the staging of renal cell carcinoma. *J Urol* 1987; 138: 482-4.
- [77] Bos SD, Mensink HJ. Can duplex Doppler ultrasound replace computerized tomography in

Molecular aspects of renal cell carcinoma

- staging patients with renal cell carcinoma? *Scand J Urol Nephrol* 1998; 32: 87-91.
- [78] Habboub HK, Abu-Yousef MM, Williams RD, See WA, Schweiger GD. Accuracy of color Doppler sonography in assessing venous thrombus extension in renal cell carcinoma. *Am J Roentgenol* 1997; 168: 267-71.
- [79] Fritzsche PJ, Millar C. Multimodality approach to staging renal cell carcinoma. *Urol Radiol* 1992; 14: 3-7.
- [80] Beer AJ, Dobritz M, Zantl N, Weirich G, Stollfuss J, Rummeny EJ. Comparison of 16-MDCT and MRI for characterization of kidney lesions. *Am J Roentgenol* 2006; 186: 1639-1650.
- [81] Pretorius ES, Wickstrom ML, Siegelman ES. MR imaging of renal neoplasms. *Magn Reson Imaging Clin N Am* 2000; 8: 813-6
- [82] Fielding JR, Aliabadi N, Renshaw AA, Silverman SG. Staging of 119 patients with renal cell carcinoma: the yield and cost-effectiveness of pelvic CT. *Am J Roentgenol* 1999; 172: 23-5.
- [83] Novick AC. Advances in the management of localized renal cell cancer. *Can J Urol* 2000; 7: 960-6.
- [84] Montravers F, Grahek D, Kerrou K, Younsi N, Doublet JD, Gattegno B, Rossert J, Costa de Beauregard MA, Thibault P, Talbot JN. Evaluation of FDG uptake by renal malignancies (primary tumor or metastases) using a coincidence detection gamma camera. *J Nucl Med* 2000; 41: 78-84.
- [85] Ramdave S, Thomas GW, Berlangieri SU, Bolton DM, Davis I, Danguy HT, Macgregor D, Scott AM. Clinical role of F-18 fluorodeoxyglucose positron emission tomography for detection and management of renal cell carcinoma. *J Urol* 2001; 166: 825-30.
- [86] Shvarts O, Han KR, Seltzer M, Pantuck AJ, Beldegren AS. Positron emission tomography in urologic oncology. *Cancer Control* 2002; 9: 335-42.
- [87] Majhail NS, Urbain JL, Albani JM, Kanvinde MH, Rice TW, Novick AC, Mekhail TM, Olencki TE, Elson P, Bukowski RM. F-18 fluorodeoxyglucose positron emission tomography in the evaluation of distant metastases from renal cell carcinoma. *J Clin Oncol* 2003; 21: 3995-4000.
- [88] Sandock DS, Seftel AD, Resnick MI. Adrenal metastases from renal cell carcinoma: role of ipsilateral adrenalectomy and definition of stage. *Urology* 1997; 49: 28-31.
- [89] Novick AC, Stroom S, Montie JE, Pontes JE, Siegel S, Montague DK, Goormastic M. Conservative surgery for renal cell carcinoma: a single-center experience with 100 patients. *J Urol* 1989; 141: 835-9.
- [90] Rini BI, Vogelzang NJ. Prognostic factors in renal carcinoma. *Semin Oncol* 2000; 27: 213-20.
- [91] Blom JH, van Poppel H, Marechal JM, Jacqmin D, Sylvester R, Schröder FH, de Prijck L. EORTC Genitourinary Group. Radical nephrectomy with and without lymph node dissection: preliminary results of the EORTC randomized phase III protocol 30881. *Eur Urol* 1999; 36: 570-5.
- [92] Rini BI, Vogelzang NJ, Dumas MC, Wade JL 3rd, Taber DA, Stadler WM. Phase II trial of weekly intravenous gemcitabine with continuous infusion fluorouracil in patients with metastatic renal cell cancer. *J Clin Oncol* 2000; 18: 2419-26.
- [93] Yagoda A, Abi-Rached B, Petrylak D. Chemotherapy for advanced renal-cell carcinoma: 1983-1993. *Semin Oncol* 1995; 22: 42-60.
- [94] Saidi JA, Newhouse JH, Sawczuk IS. Radiologic follow-up of patients with T1-3a,b,c or T4N+M0 renal cell carcinoma after radical nephrectomy. *Urology* 1998; 52: 1000-3.
- [95] Robson CJ, Churchill BM, Anderson W. The results of radical nephrectomy for renal cell carcinoma. *J Urol* 1969; 101: 297-301.
- [96] Skinner DG, Colvin RB, Vermillion CD, Pfister RC, Leadbetter WF. Diagnosis and management of renal cell carcinoma. A clinical and pathologic study of 309 cases. *Cancer* 1971; 28: 1165-77.
- [97] Medeiros LJ, Gelb AB, Weiss LM. Renal cell carcinoma. Prognostic significance of morphologic parameters in 121 cases. *Cancer* 1988; 61: 1639-51.
- [98] Giberti C, Oneto F, Martorana G, Rovida S, Carmignani G. Radical nephrectomy for renal cell carcinoma: long-term results and prognostic factors on a series of 328 cases. *Eur Urol* 1997; 31: 40-8.
- [99] Golimbu M, Joshi P, Sperber A, Tessler A, Al-Askari S, Morales P. Renal cell carcinoma: survival and prognostic factors. *Urology* 1986; 27: 291-301.
- [100] Ljungberg B, Forsslund G, Stenling R, Zetterberg A. Prognostic significance of the DNA content in renal cell carcinoma. *J Urol* 1986; 135: 422-6.
- [101] Murphy WM, Chandler RW, Trafford RM. Flow cytometry of deparaffinized nuclei compared to histological grading for the pathological evaluation of transitional cell carcinomas. *J Urol* 1986; 135: 694-7.
- [102] Thrasher JB & Paulson DF. Prognostic factors in renal cancer. *Urol Clin North Am* 1993; 20: 247-62.
- [103] Citterio G, Bertuzzi A, Tresoldi M, Galli L, Di Lucca G, Scaglietti U, Rugarli C. Prognostic factors for survival in metastatic renal cell carcinoma: retrospective analysis from 109 consecutive patients. *Eur Urol* 1997; 31: 286-91.
- [104] Van Poppel H, Vandendriessche H, Boel K, et al. Microscopic vascular invasion is the most relevant prognosticator after radical nephrectomy for clinically nonmetastatic renal cell carcinoma. *J Urol* 1997; 158: 45-9.
- [105] Van Brussel JP, Mickisch GH. Prognostic factors in renal cell and bladder cancer. *BJU Int* 1999; 83: 902-8.
- [106] Mignogna C, Staibano S, Altieri V, et al. Prognostic significance of multidrug-resistance pro-

Molecular aspects of renal cell carcinoma

- tein (MDR-1) in renal clear cell carcinomas: a five year follow-up analysis. *BMC Cancer* 2006; 6: 293-303.
- [107] Yang JC, Sherry RM, Steinberg SM, Topalian SL, Schwartzentruber DJ, Hwu P, Seipp CA, Rogers-Freezer L, Morton KE, White DE, Liewehr DJ, Merino MJ, Rosenberg SA. Randomized study of high-dose and low-dose interleukin-2 in patients with metastatic renal cancer. *J Clin Oncol* 2003; 21: 3127-32.
- [108] Motzer RJ, Hutson TE, Tomczak P, Michaelson MD, Bukowski RM, Rixe O, Oudard S, Negrier S, Szczylik C, Kim ST, Chen I, Bycott PW, Baum CM, Figlin RA. Sunitinib versus interferon-alfa in metastatic renal-cell carcinoma. *N Engl J Med* 2007; 356: 115-24.
- [109] Motzer RJ, Hutson TE, Tomczak P, Michaelson MD, Bukowski RM, Rixe O, Oudard S, Negrier S, Szczylik C, Kim ST, Chen I, Bycott PW, Baum CM, Figlin RA. Sunitinib versus interferon-alfa (IFN- α) as first-line treatment of metastatic renal cell carcinoma (mRCC): updated results and analysis of prognostic factors. *J Clin Oncol* 2007; 25 [meeting abstracts suppl]: 5024.
- [110] Hudes G, Carducci M, Tomczak P, Dutcher J, Figlin R, Kapoor A, Staroslawska E, Sosman J, McDermott D, Bodrogi I, Kovacevic Z, Lesovoy V, Schmidt-Wolf IG, Barbarash O, Gokmen E, O'Toole T, Lustgarten S, Moore L, Motzer RJ; Global ARCC Trial. Temsirolimus, interferon-alfa, or both for advanced renal-cell carcinoma. *N Engl J Med* 2007; 356: 2271-81.
- [111] Figlin RA, de Souza P, McDermott D, Dutcher JP, Berkenblit A, Thiele A, Krygowski M, Strahs A, Feingold J, Boni J, Hudes G. Analysis of PTEN and HIF-1 α and correlation with efficacy in patients with advanced renal cell carcinoma treated with temsirolimus versus interferon- α . *Cancer* 2009; 115: 3651-60.
- [112] Yang JC, Haworth L, Sherry RM, et al. A randomized trial of bevacizumab, an anti-vascular endothelial growth factor antibody, for metastatic renal cancer. *N Engl J Med* 2003; 349: 427-34.
- [113] Melichar B, Koralewski P, Ravaud A, Pluzanska A, Bracarda S, Szczylik C, Chevreau C, Filipek M, Delva R, Sevin E, Négrier S, McKendrick J, Santoro A, Pisa P, Escudier B. First-line bevacizumab combined with reduced dose interferon- α 2a is active in patients with metastatic renal cell carcinoma. *Ann Oncol* 2008; 19: 1470-6.
- [114] Seront E, Machiels JP. Targeted therapies in the treatment of advanced renal cell carcinoma. *Recent Pat Anticancer Drug Discov* 2009; 4: 146-56.