

Treatment of seasonal affective disorders

Nicole Praschak-Rieder, MD; Matthäus Willeit, MD



Seasonal affective disorder (SAD) is a subform of major depressive disorder, recurrent, or bipolar disorder with a regular onset of depressive episodes at a certain time of year, usually the winter. The treatment of SAD is similar to that of other forms of affective disorder, except that bright light therapy is recommended as the first-line option. Light therapy conventionally involves exposure to visible light of at least 2500 lux intensity at eye level. The effects of light therapy are thought to be mediated exclusively by the eyes, not the skin, although this assumption has not yet been verified. Morning light therapy has proven to be superior to treatment regimens in the evening. Response rates to light therapy are about 80% in selected patient populations, with atypical depressive symptoms being the best predictor of a favorable treatment outcome. Data from randomized, controlled trials suggest that antidepressants are effective in the treatment of SAD. Three double-blind, placebo-controlled trials have been conducted showing promising results for the selective serotonin reuptake inhibitors (SSRIs) sertraline and fluoxetine, as well as for moclobemide, a reversible inhibitor of monoamine oxidase A.

© 2003, LLS SAS

Dialogues Clin Neurosci. 2003;5:389-398.

Keywords: seasonal affective disorder; depression; light therapy; pharmacotherapy; treatment guidelines

Author affiliations: Centre for Addiction and Mental Health, PET Centre, Toronto, ON, Canada

Seasonal affective disorder (SAD), as originally described in 1984,¹ is a condition characterized by the annual recurrence of depressive episodes in fall and winter followed by remission of depressive symptoms in spring and summer.¹ Patients with SAD have to meet diagnostic criteria for major depression, recurrent, or bipolar disorder. In the latest version of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, SAD is listed as a specifier of either bipolar or recurrent major depressive disorder, with a seasonal pattern of major depressive episodes.² Subsyndromal SAD is a disorder with similar but milder symptoms that do not grossly disrupt patients social and occupational functioning.³ The four central features characterizing SAD are listed in *Table I*. Patients with SAD have the usual symptoms of depression, including low mood, lack of drive, decreased concentration, and reduced interest. Typically, many SAD patients also tend to have a specific symptom cluster consisting of the so-called reverse vegetative or atypical depressive symptoms. These symptoms include increased sleep (70%-90% of SAD patients), increased appetite (70%-80%), carbohydrate craving (80%-90%), and weight gain (70%-80%).⁴

Pathophysiology

The etiology of SAD remains unclear. It is thought that the decreasing daylight period as winter approaches triggers depressive episodes in individuals vulnerable to SAD. However, although bright light exposure is used in the treatment of SAD, no causal relation can be drawn between the occurrence of SAD and the shortage of light in fall and winter. Patients with SAD may be sensitive to factors that are common to various forms of recurrent

Address for correspondence: Nicole Praschak-Rieder, MD, Centre for Addiction and Mental Health, PET Centre, 250 College Street, Toronto, Ontario, Canada M5T 1R8
(e-mail: nicole@camhpet.on.ca)

Clinical research

affective disorder, and SAD can be seen as a disorder driven by endogenous annual rhythms and characterized by an imbalance of indoleamines, serotonin, and melatonin, as well as catecholamines, over the year.

Light therapy

Bright light therapy (BLT) has become a first-line clinical standard for treatment of SAD (*Table II*). The use of BLT as a therapy for SAD evolves directly out of neuroscience. In the early 1980s, knowledge that light could shift circadian and seasonal rhythms in animals, together with the heuristic idea of extending daylight during the winter months, led to the first clinical study on BLT in SAD.¹ Since then, numerous studies have not only proven its efficacy, but also greatly refined our knowledge on treatment strategies, thereby allowing for astonishingly high response rates of 80% in selected patient populations.⁵ BLT is safe, effective, and it has few and benign side effects. It is generally well accepted by the patients,⁶ and indications other than SAD, eg, disturbances of circadian rhythm due to jet lag or shift work,^{7,8} circadian-phase-related disturbances in dementia,^{9,10} sleep disorders,^{11,12} and nonseasonal affective disorders,¹³⁻¹⁵ are expanding research fields.

Efficacy

The first controlled clinical trial¹ already showed the beneficial effects of light against symptoms of SAD. Since then, more than 60 controlled studies and two meta-analyses^{16,17} have shown the efficacy of this treatment. Using stringent criteria, the meta-analysis of Terman et al¹⁶ found remission rates of up to 67% of patients with milder depression and up to 40% of more severely depressed patients. These benefits were seen as early as 1 week after beginning treatment. However, it is now

Recurrent major depressive episodes that start around the same time each year (eg, fall and winter) and end around the same time each year (eg, spring and summer)
Full remission of symptoms during the unaffected period of the year (eg, summer)
Over the lifetime course of the illness, there are relatively more seasonal depressive episodes than nonseasonal episodes
Seasonal depressive episodes occur in at least 2 consecutive years

Table I. Features of seasonal affective disorder (SAD).

known that improvement may sometimes be seen as late as 2 to 4 weeks after beginning BLT.¹⁸

The obvious obstacles in the search for plausible placebos and “blinded” designs for studying the efficacy of BLT have led to concerns about the adequacy of control conditions. Early studies mostly used dim-light of an intensity of 300 lux or less, delivered through a light source otherwise identical to that used in the active condition (mostly light of 2500 lux intensity). As it was still possible to distinguish between the bright and dim-light conditions, these studies have been questioned as to whether they measured real biological effects and whether they could reliably separate a true antidepressant effect from placebo effects. However, fantasy and creativity of researchers in the field has led to further studies, which now unequivocally prove that light is an active biological agent with antidepressant effects in SAD. Some of these studies used (deactivated) negative ion generators as placebo condition.¹⁸⁻²⁰ Like a light box, the negative ion generator is a device, has a plausible mechanism of antidepressant action, and requires the subject to sit beside it. Light had the better antidepressant effect and produced significantly more remissions, although expectations were equal for both conditions. Interestingly enough, high doses of negative ions do seem to have an antidepressant effect as well.²⁰

Light therapy is an effective first-line treatment for seasonal affective disorder
A fluorescent light box with light intensities greater than 2500 lux is the preferred device for light therapy
An optimal starting dose for light therapy is 10 000 lux for 30 min/day
Light boxes emitting 2500 lux are effective with a treatment duration of 2 h/day
Morning light is more effective than evening light. Patients should be encouraged to undergo light therapy as early as possible (eg, before/during breakfast). However, evening light may be effective for some patients
Many patients will respond as early as after 1 week. However, in some cases a response may occur after only 2 to 4 weeks
If there is no sufficient response after two treatment weeks, the dose should be doubled to 30 min in the morning plus 30 min in the evening
If this regimen does not lead to sufficient improvement, consider adding a pharmacological treatment (best evidence available for serotonin reuptake inhibitors)

Table II. Guidelines for bright light therapy (BLT).

Another challenge for efficacy studies on light therapy is the “light-noise” inevitably encountered in a normal living environment. Depending on weather conditions, outdoor light intensities can easily surmount the intensities delivered by a light box, even in the winter. This has been compared to a study trying to prove the superiority of an antidepressant drug over placebo, with antidepressants occasionally delivered through the drinking water to all study subjects.²¹ Nevertheless, today there is sound evidence showing that antidepressant effects of light in SAD are real biological treatment effects.^{16,20,22}

Dose

Earlier studies have mostly used light intensities of 2500 lux. This is much more than the typical indoor illumination, ranging from 100 lux in average rooms to about 500 lux in brightly illuminated ones. Outdoor light intensities greatly vary with weather conditions ranging from about 2000 lux on a rainy winter day to 10 000 lux or more (usually 50 000 to 300 000 lux) in direct sunshine. Today, light treatment with intensity of 10 000 lux has become clinical standard. One great advantage of higher intensity light is that it allows for shorter exposure times. Current clinical guidelines recommend beginning treatment with 10 000 lux for 30 min in the morning.²³ Nevertheless, intensities of 2500 lux have shown to have antidepressant effects when applied for 2 h daily.

Timing

A further finding that emerged from BLT studies is the superiority of morning light over light administered in the evening.^{16,20,22} By further refining timing of light administration in relationship to the position of the circadian phase, Terman and coworkers achieved remission rates up to 80% in selected patient populations.⁵ They were able to show that response to BLT critically depends on time of delivery relative to the position of the circadian phase as determined by the onset of melatonin secretion in the evening (dim-light melatonin onset). The study suggests that the ideal therapy time is around 8.5 h after melatonin onset.

Although the superiority of morning light in the majority of patients with SAD had been demonstrated before,^{16,20,22} Terman and colleagues showed that, for clinical purposes, circadian phase position can simply and reliably be determined by administration of a modified version Horne-Öst-

berg morningness-eveningness scale.²⁴ The authors also provide an online-questionnaire²⁵ together with recommendations for individually optimized light therapy timing as based on morningness-eveningness or individual circadian phase position. In summary, now there is sufficient evidence that light administered in the early morning is superior to evening light.

Dawn stimulation

Many patients with SAD experience markedly increased duration of sleep during the winter months.^{1,4,26} Usually, most of these patients have to force themselves out of bed during the weekdays despite feeling excessively drowsy. Dawn stimulation is a form of light therapy involving gradually increasing bedside light in the morning before awakening.²⁷ Dawn stimulation has shown to improve symptoms of SAD compared with placebo light signals.^{28,29} In addition, dawn stimulation appears to be effective in ameliorating the difficulty awakening and morning drowsiness in SAD.³⁰ In a comparison study, dawn stimulation using 100 to 300 lux for 60 to 90 min every morning improved symptoms of SAD similarly to light therapy using 1500 to 2500 lux for 2 h every morning.³¹

Adverse effects

The adverse effects of light therapy include headache, eye-strain, nausea, and agitation.^{32,33} Usually, adverse effects are mild and subside spontaneously or with dose reduction. Bright light in the evening may be associated with sleep disturbances, and, occasionally, hypomania may arise during BLT.³³ However, subjective benefits of light consistently outweigh its adverse effects.^{32,33} Altogether, it remains questionable whether the frequency of these symptoms under BLT significantly exceeds the frequency of side effects seen under placebo conditions.

Risks

There are no absolute contraindications for light therapy.²³ Animal studies suggest increased risk for retinal damage with lithium, β -blockers, tricyclic antidepressants, and tryptophan. However, no such interactions have been reported in humans, and there is no evidence that light therapy is associated with ocular or retinal damage in humans. Patients with severe ophthalmological conditions or patients taking photosensitizing medication should have

Clinical research

an ophthalmological examination before starting light therapy. However, it is important that the UV spectrum is filtered out of the therapeutic light source.

Although suicidality is commonly regarded as being rather infrequent in SAD, our own group has reported severe suicidal ideation and suicide attempts in three patients after the initiation of light therapy.³⁴ All three patients had suicidal thoughts before light therapy was started. As always when dealing with depressed patients, patients with SAD should be carefully assessed for suicidality before light therapy, and therapy outcome should frequently and regularly be evaluated by health care professionals.

Treatment predictors

Atypical depressive symptoms, specifically hyperphagia, hypersomnia, and carbohydrate craving, seem to be associated with favorable response to BLT.^{35,36} Younger age also seems to predict a good response,³⁷ while comorbid personality disorders seem to compromise the response to BLT.^{38,39}

Mechanism of action

Theories on the mechanism of action of BLT are closely connected to what is known about the pathogenesis of SAD.⁴⁰ Two main—mutually not exclusive—theories have been raised by researchers in the field: one concentrates on the evidence for reduced serotonin neurotransmission in SAD, the other theory relates light therapy-induced improvement to corrections of altered circadian rhythms during depression in SAD.

Serotonin

Several lines of evidence suggest an alteration in serotonin neurotransmission in SAD.⁴⁰⁻⁴² A keystone of the serotonin hypothesis on the mechanism of action of light therapy is the finding that lowering brain serotonin by tryptophan depletion leads to a transient depressive relapse in patients with SAD who are in light therapy-induced remission.⁴³ In line with this theory is the beneficial effect of selective serotonin reuptake inhibitors (SSRIs) in SAD. Although there is evidence for a seasonal variation in serotonin neurotransmission,⁴⁴ and although there seems to be a close relationship between brain serotonin and atypical depressive symptoms,⁴¹ serotonergic alterations are not specific for the pathogenesis of SAD or the antidepres-

sant action of light. They rather seem to constitute a pathway common to depressive syndromes and their treatment in general.

Circadian phase shifts

More specific for SAD are the theories concerning alterations in circadian and circannual rhythms. Neurons of the hypothalamic suprachiasmatic nucleus (SCN) act as the main “zeitgeber” in the mammalian organism. Having an intrinsic near to 24-h rhythm, they are also known as the “internal clock.” These neurons are reset by environmental light, and they are believed to be the main determinant for the position of the circadian phase. Several body functions, such as hormonal rhythms, including nocturnal melatonin secretion, sleep, or eating behavior, are subjected to a specific circadian rhythm. The best studied marker for the position of the circadian phase is the onset of melatonin secretion by the pineal gland.⁴⁵ In humans, the beginning of melatonin secretion occurs in the evening, usually between 1 and 2 h before falling asleep. Light can shift the position of the circadian phase, and amount and direction of that shift greatly depend on the time of light exposure: light in the evening leads to a phase delay (eg, melatonin onset occurs later), morning light advances the circadian phase.^{20,22}

Early theories on the pathogenesis of SAD held that a delay in the circadian phase was responsible for the appearance of SAD symptoms.⁴⁶ Although the phase-delay hypothesis on the pathogenesis of SAD did not hold up, as there does not seem a consistent pattern of phase alterations in SAD, recent work has shown the circannual variation in circadian phase to be altered in patients with SAD when compared with healthy control subjects.⁴⁷ Recent work by Terman and colleagues showed a correlation between light-induced changes in the “phase angle” (the relationship between the circadian phase as measured by melatonin onset and, eg, sleep onset) and antidepressant response to light in SAD.⁵

Practical issues

Sufficient and clear instructions to patients are critical for a satisfactory treatment response. Patients should be informed that the beneficial effects of light are not enduring, ie, that a relapse is to be expected after a few days when treatment is discontinued. Although one study⁴⁸ suggested a transcutaneous effect of light on melatonin secre-

tion, these results have not been replicated.^{49,50} It is so far safe to state that the antidepressant effect of light is mediated by the eye. The patient therefore needs to make sure that light of sufficient intensity meets the eye. Light projection does not need to be foveal, ie, it is not necessary to look directly into the light source. The ideal is an angle of about 30° to 60° to let enough light meet the eye and allow for the patient to, for example, read or eat during light therapy. The intensity of light critically depends upon the distance from the light source. Light boxes should be powerful enough to deliver an intensity of about 10 000 lux at a distance of 60 to 90 cm. If the light box is less powerful, treatment time should be expanded (see above). Patients should be encouraged to seek exposure to environmental light on sunny days. Sunlight has much higher intensity than light delivered by a light therapy device (see above). Despite the fact that light therapy is now recommended as a treatment of choice for SAD, only in Switzerland has the economic argument that in the long run, light is cheaper than drugs, attained government endorsement and mandatory reimbursement by medical insurance.⁵¹ The fact that there is no reimbursement for light therapy has been widely criticized by patients with SAD, their relatives, and experts in the field of SAD.⁵² Case reports on SAD patients resistant to several antidepressants, but finally responsive to light therapy illustrate that, although depressive symptoms may often be only moderate, SAD can lead to severe impairment in occupational and social functioning and can precipitate catastrophic life events.^{52,53}

Pharmacotherapy

Although light therapy is recommended as the first-line option for SAD, some patients do not experience sufficient relief of depressive symptoms with light. BLT can then be supplemented with antidepressant drugs. Other patients with SAD feel unable to integrate light therapy into their daily routine, or other logistical difficulties in administering light therapy are present. The evidence of SAD being associated with a dysfunction in brain serotonin systems has guided the search for promising pharmacological treatments of SAD. Data emerging from multicenter placebo-controlled trials has led to the recommendation of the SSRIs sertraline and fluoxetine as first-line treatments of SAD. Other antidepressant compounds, like monoamine oxidase inhibitors, dopaminergic and noradrenergic agents, melatonin, β -blockers as melatonin antagonists, herbs, and nutritional supplements

like L-tryptophan and vitamin D have been investigated in small studies. The efficacy of these medications has not yet been proven in SAD. Open trials, controlled studies, and placebo-controlled studies in SAD are listed in *Tables III to V*.⁵⁴⁻⁷⁵ New pharmacological agents are of potential value in the treatment of SAD, for example, agomelatine (Valdoxan). This new dual melatonergic and specific serotonergic antidepressant has been shown to be efficient in the treatment of major depression⁷⁶: it exhibits a specific core action on circadian rhythms, and therefore could be of particular value in the treatment of SAD. More specific studies are underway to more obtain information about its activity in SAD.

Open studies

A survey of open studies in SAD is given in *Table III*.⁵⁴⁻⁶¹ There is some suggestion from pilot data with small sample sizes that serotonergic agents like fluoxetine, citalopram, and trazodone may be treatment options for SAD.^{54,57} Tranylcypromine, a nonselective monoamine oxidase inhibitor was effective in the treatment of 14 patients leading to an average 91% reduction in depressive symptoms within 4 weeks of initiation of treatment.⁵⁵ A study in 20 patients indicates that St John's wort (*Hypericum perforatum*) may be helpful in treating SAD. An add-on therapy with bright light in 10 of these patients treated with hypericum did not lead to a significantly better treatment outcome.⁵⁹ Two studies in 6 patients each report beneficial effects of the benzodiazepine alprazolam.^{56,60} A 6-week open trial investigating efficacy and tolerability of reboxetine, a selective noradrenaline reuptake inhibitor, led to rapid full remission of depressive symptoms in 11 out of 16 patients.⁶¹ A rapid relief of preexistent severe atypical symptoms was observed in 9 patients within the

Authors	Number of patients	Medication
Jacobsen et al, ⁵⁴ 1989	n=3	Fluoxetine, trazodone
Dilsaver and Jaekle, ⁵⁵ 1990	n=14	Tranylcypromine
Teicher and Glod, ⁵⁶ 1990	n=6	Alprazolam
Wirz-Justice et al, ⁵⁷ 1992	n=1	Citalopram
Lingjaerde et al, ⁵⁸ 1993	n=5	Moclobemide
Martinez et al, ⁵⁹ 1994	n=20	Hypericum
Yamadera et al, ⁶⁰ 2001	n=6	Alprazolam
Hilger et al, ⁶¹ 2001	n=16	Reboxetine

Table III. Open studies of pharmacotherapy of seasonal affective disorder (SAD).⁵⁴⁻⁶¹

Clinical research

first week of treatment. This finding is of pathophysiological interest since, so far, atypical depressive symptoms like increased appetite, carbohydrate craving, and hypersomnia have been strongly associated with a dysfunction in brain serotonin systems.

Controlled studies

Controlled studies of pharmacotherapy in SAD are presented in *Table IV*.⁶²⁻⁶⁶ A study by Ruhrmann et al comparing the SSRI fluoxetine and light therapy in 40 patients with SAD found no significant difference in treatment outcome between the groups, but a faster onset of antidepressant action in the light therapy group.⁶³

Because SAD is prevalent in winter when vitamin D stores are typically low, and because light therapy includes wavelengths that allow the skin to produce vitamin D, the potential role of vitamin D in SAD has been investigated in a small pilot study.⁶⁵ Vitamin D was reported to lead to a greater improvement of depressive symptoms than light therapy. However, no difference in vitamin D levels has been observed between patients with SAD and healthy subjects,^{70,77} and the antidepressant effect of light therapy has been shown to be independent of changes in vitamin D levels.⁷⁷ So far, any benefits of vitamin D on SAD remain unproven. In two small, preliminary trials, 4 to 6 g daily doses of the amino acid L-tryptophan, the precursor of serotonin, were as effective as light therapy.^{62,64} In a postal survey using an 11-item rating scale, 301 patients with SAD treated with hypericum at 300 mg three times daily for 8 weeks were asked to report changes in their symptoms.⁶⁶ Of these patients,

133 used additional light therapy. Significant overall improvement was reported in both treatment groups. Improvement in sleep was greater in the hypericum and light therapy group. However, double-blind research is needed to confirm the usefulness of hypericum (St John's wort) for treating SAD.

Placebo-controlled studies

Table V^{58,67-75} presents placebo-controlled studies of pharmacotherapy in SAD. The best evidence for efficacy of antidepressants in SAD comes from studies of SSRIs. Multicenter, double-blind, randomized studies of fluoxetine and sertraline confirm that these medications are effective in the treatment of SAD. In the fluoxetine study (68 patients), significant improvement in mood was present in both fluoxetine and placebo-treated patients at termination of the study. However, there was significant superiority of fluoxetine over placebo in the clinical response rates (59% versus 34%, respectively).⁷¹ In the sertraline study (187 patients), a significant superiority to placebo in both clinical response rates (62% versus 46%, respectively) and depression scores was found. Although they have been widely cited, the data from the sertraline study have only been published as an abstract so far.⁷⁸ A double-blind study by Lingjaerde et al⁵⁸ investigating the efficacy of moclobemide, a reversible inhibitor of monoamine oxidase A, versus placebo over 14 weeks found no significant difference in depression scores between groups at study termination. However, within the first week of treatment, patients in the moclobemide group, but not in the placebo group, had a significant reduction in atypical depression symptoms. Testing the

Authors	Number of patients	Medication and study design	Outcome
McGrath et al, ⁶² 1990	n=13	L-Tryptophan and LT versus L-tryptophan alone	No significant difference
Ruhrmann et al, ⁶³ 1998	n=40	Fluoxetine versus LT	No significant difference, but faster onset of antidepressant action in LT group
Ghadirian et al, ⁶⁴ 1998	n=13	L-Tryptophan versus LT, crossover study	No significant difference
Gloth et al, ⁶⁵ 1999	n=15	Vitamin D versus LT	Vitamin D superior to LT
Wheatley, ⁶⁶ 1999	n=168	Hypericum and LT versus hypericum alone	Significant improvement in both groups, improvement in sleep greater in hypericum and LT group

Table IV. Controlled studies of pharmacotherapy of seasonal affective disorder (SAD).⁶²⁻⁶⁶ LT, light therapy.

hypothesis that a dopaminergic deficiency plays a role in the pathophysiology of SAD, Oren et al conducted a small study investigating the efficacy of levodopa plus carbidopa as a treatment for SAD.⁶⁸ No differences to placebo were found in the rates of response. The melatonin hypothesis of SAD was tested in two studies using the β -blockers atenolol⁶⁷ and propranolol⁶⁹ to suppress melatonin secretion. No difference in antidepressant efficacy was found between atenolol and placebo. Propranolol was superior to placebo in preventing a depressive relapse in patients with SAD who had previously responded to an open treatment with propranolol. Supplementation with melatonin has shown to be ineffective in patients with SAD when taken at night or in the morning.⁷⁹ Melatonin has also been reported to even reverse the benefits of light therapy.⁸⁰ However, a small pilot study with low doses of melatonin in the afternoon showed a significant decrease in depression ratings compared to placebo.⁷² The authors argue that a replication of this finding in an adequate sample with documentation of expected phase shifts would substantially support the phase shift hypothesis of SAD. A recent 1-year pilot study⁷³ aimed at investigating possible advantages of combining light therapy with the SSRI citalopram. No significant group difference

was found during the initial 10-day light therapy period. However, during the follow-up period depression ratings were significantly lower in the citalopram group compared with the placebo group. The authors conclude that light therapy with continued SSRI treatment may be a useful strategy to achieve beneficial long-term effects in patients with SAD. A study by Lingjaerde et al⁷⁴ testing the hypothesis that *Ginkgo biloba* extract may prevent the symptoms of winter depression in patients with SAD yielded negative results. In a recent study metergoline, a nonspecific serotonin antagonist, did not demonstrate a sustained significant effect on mood compared with placebo.⁷⁵

General management issues

Since there is little evidence comparing light therapy with antidepressant medication, the choice between these alternatives relies on individual assessment of risks and benefits.²³ Generally, light therapy is very well accepted by patients. Availability and costs of a light therapy device are sometimes limiting factors, as is the time patients need to commit for daily light therapy. However, BLT should be considered first-line treatment for moderately depressed

Authors	Number of patients	Medication and study design	Outcome
Rosenthal et al, ⁶⁷ 1988	n=19	Atenolol versus placebo	No significant difference
Lingjaerde et al, ⁵⁸ 1993	n=34	Moclobemide versus placebo	No significant difference, but significant reduction of atypical symptoms in moclobemide group
Oren et al, ⁶⁸ 1994	n=25	Levodopa + carbidopa versus placebo	No significant difference
Schlager, ⁶⁹ 1994	n=23	Propranolol versus placebo	Significantly higher relapse rates in placebo group
Oren et al, ⁷⁰ 1994	n=27	Vitamin B ₁₂ (cyanocobalamine) versus placebo	No significant difference
Lam et al, ⁷¹ 1995	n=78	Fluoxetine versus placebo	No significant difference, higher response rate in fluoxetine group
Lewy et al, ⁷² 1998	n=10	Melatonin versus placebo	Melatonin superior to placebo
Thorell et al, ⁷³ 1999	n=8	LT and citalopram versus LT and placebo	No significant difference between citalopram and placebo during LT; after LT citalopram superior to placebo
Lingjaerde et al, ⁷⁴ 1999	n=27	<i>Ginkgo biloba</i> versus placebo	No significant difference; ginkgo significant more effective in preventing new depressive episodes
Turner et al, ⁷⁵ 2002	n=16	Metergoline versus placebo	No significant difference

Table V. Placebo-controlled studies of pharmacotherapy of seasonal affective disorder (SAD).^{58,67-75} LT, light therapy.

Clinical research

patients and patients with prominent atypical depressive symptoms. Generally, light therapy alone or in combination with antidepressants should be given for the duration of the dark time of year, ie, until April or May in the northern hemisphere. A trial of light therapy should last at least 2 to 4 weeks. A trial of antidepressants should last 4 to 6 weeks. Light therapy and medication trials should be applied sequentially, as combining them from the beginning on will lead to a loss of information about which treatment is beneficial, or which treatment is causing side effects. A combination of both treatments should be con-

sidered if there is insufficient response to either pharmacological or light treatment. If a depressive episode is resistant to the combination of BLT and an antidepressant, options are lengthening light treatment time, raising the dose of the antidepressant, or switching to a drug of a different class. Although there are no specific data available for SAD, treatment should follow algorithms for treatment-resistant depression if a sufficient response still cannot be achieved. Pharmacological augmentation strategies, electroconvulsive therapy, or sleep deprivation procedures⁸¹ should then be considered. □

Tratamiento de los trastornos afectivos estacionales

El trastorno afectivo estacional (TAE) es un subtipo del trastorno depresivo mayor recurrente o del trastorno bipolar, con una aparición regular de episodios depresivos en cierta época del año, especialmente en invierno. El tratamiento del TAE es similar al de otros tipos de trastorno afectivo, excepto que la terapia con luz brillante se recomienda como opción de primera línea. Convencionalmente la terapia lumínica se refiere a la exposición a luz visible de al menos 2500 lux de intensidad a nivel del ojo. Se piensa que los efectos de la fototerapia son mediados exclusivamente por los ojos, no por la piel, aunque esta hipótesis no ha sido aun verificada. La terapia lumínica matinal ha probado ser superior a las aplicaciones vespertinas. Los porcentajes de respuesta a la fototerapia son cercanos al 80% en poblaciones seleccionadas de pacientes y los síntomas depresivos atípicos son el mejor predictor de una evolución favorable al tratamiento. Datos de ensayos controlados, randomizados sugieren que los antidepressivos son efectivos en el tratamiento del TAE. Se han realizado tres estudios doble ciego, controlados con placebo que han mostrado resultados promisorios para los inhibidores selectivos de la recaptación de serotonina (ISRS) sertralina y fluoxetina, y también para moclobemida, un inhibidor reversible de la monoamino-oxidasa A.

Traitement du trouble affectif saisonnier

Le trouble affectif saisonnier (TAS) représente un sous-groupe des troubles dépressif majeur récurrent ou bipolaire et se caractérise par l'installation régulière d'épisodes dépressifs à certaines périodes de l'année, généralement en hiver. Le traitement du TAS est semblable à celui des autres formes de troubles affectifs, si ce n'est que la photothérapie à lumière vive est recommandée en première intention. La photothérapie conventionnelle implique l'exposition à une lumière du spectre visible d'une intensité supérieure à 2 500 lux au niveau de l'œil. Les effets de la photothérapie sont considérés comme médiés exclusivement par les yeux et non la peau, bien que cette hypothèse n'ait pas encore été vérifiée. La photothérapie matinale s'est avérée supérieure au schéma thérapeutique vespéral. Les taux de réponse à la photothérapie sont d'environ 80 % dans des populations sélectionnées de patients, les symptômes dépressifs atypiques étant le meilleur facteur prédictif d'évolution thérapeutique favorable. Des données issues d'études randomisées, contrôlées, suggèrent que les antidépresseurs sont efficaces dans le traitement du TAS. Trois études en double aveugle, contre placebo, ont été réalisées, montrant des résultats prometteurs pour les inhibiteurs sélectifs de la recapture de la sérotonine (ISRS), la sertraline et la fluoxétine, ainsi que pour un inhibiteur de la monoamine oxydase sélectif A (IMAO sélectif A), le moclobémide.

REFERENCES

1. Rosenthal NE, Sack DA, Gillin JC, et al. Seasonal affective disorder. A description of the syndrome and preliminary findings with light therapy. *Arch Gen Psychiatry*. 1984;41:72-80.
2. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed. Washington, DC: American Psychiatric Association; 1994.
3. Kasper S, Wehr TA, Bartko JJ, Gaist PA, Rosenthal NE. Epidemiological findings of seasonal changes in mood and behavior. A telephone survey of Montgomery County, Maryland. *Arch Gen Psychiatry*. 1989;46:823-833.
4. Winkler D, Willeit M, Praschak-Rieder N, et al. Changes of clinical pattern in seasonal affective disorder (SAD) over time in a German-speaking sample. *Eur Arch Psychiatry Clin Neurosci*. 2002;252:54-62.
5. Terman JS, Terman M, Lo ES, Cooper TB. Circadian time of morning light administration and therapeutic response in winter depression. *Arch Gen Psychiatry*. 2001;58:69-75.
6. Michalak EE, Hayes S, Wilkinson C, Hood K, Dowrick C. Treatment compliance in light therapy. Do patients do as they say they do? *J Affect Disord*. 2002;68:341-342.
7. Horowitz TS, Tanigawa T. Circadian-based new technologies for night workers. *Ind Health*. 2002;40:223-236.
8. Burgess HJ, Sharkey KM, Eastman CI. Bright light, dark and melatonin can promote circadian adaptation in night shift workers. *Sleep Med Rev*. 2002;6:407-420.
9. Graf A, Wallner C, Schubert V, et al. The effects of light therapy on minimal state examination scores in demented patients. *Biol Psychiatry*. 2001;50:725-727.
10. Ancoli-Israel S, Martin JL, Kripke DF, Marler M, Klauber MR. Effect of light treatment on sleep and circadian rhythms in demented nursing home patients. *J Am Geriatr Soc*. 2002;50:282-299.
11. Terman M, Lewy AJ, Dijk DJ, Boulos Z, Eastman CI, Campbell SS. Light treatment for sleep disorders: consensus report. IV. Sleep phase and duration disturbances. *J Biol Rhythms*. 1995;10:135-147.
12. Montgomery P, Dennis J. Bright light therapy for sleep problems in adults aged 60+. *Cochrane Database Syst Rev*. 2002;CD003403.
13. Loving RT, Kripke DF, Shuchter SR. Bright light augments antidepressant effects of medication and wake therapy. *Depress Anxiety*. 2002;16:1-3.
14. Oren DA, Wisner KL, Spinelli M, et al. An open trial of morning light therapy for treatment of antepartum depression. *Am J Psychiatry*. 2002;159:666-669.
15. Benedetti F, Colombo C, Pontiggia A, Bernasconi A, Florita M, Smeraldi E. Morning light treatment hastens the antidepressant effect of citalopram: a placebo-controlled trial. *J Clin Psychiatry*. 2003;64:648-653.
16. Terman M, Terman JS, Quitkin FM, McGrath PJ, Stewart JW, Rafferty B. Light therapy for seasonal affective disorder. A review of efficacy. *Neuropsychopharmacology*. 1989;2:1-22.
17. Thompson C, Rodin I, Birtwhistle J. Light therapy for seasonal and non-seasonal affective disorder: a Cochrane meta-analysis. *Soc Light Treatment Biol Rhythms Abstr*. 1999;11. Abstract.
18. Eastman CI, Young MA, Fogg LF, Liu L, Meaden PM. Bright light treatment of winter depression: a placebo-controlled trial. *Arch Gen Psychiatry*. 1998;55:883-889.
19. Eastman CI, Lahmeyer HW, Watell LG, Good GD, Young MA. A placebo-controlled trial of light treatment for winter depression. *J Affect Disord*. 1992;26:211-221.
20. Terman M, Terman JS, Ross DC. A controlled trial of timed bright light and negative air ionization for treatment of winter depression. *Arch Gen Psychiatry*. 1998;55:875-882.
21. Avery DH. A turning point for seasonal affective disorder and light therapy research? *Arch Gen Psychiatry*. 1998;55:863-864.
22. Lewy AJ, Bauer VK, Cutler NL, et al. Morning vs evening light treatment of patients with winter depression. *Arch Gen Psychiatry*. 1998;55:890-896.
23. Lam RW, Levitt AJ. *Canadian Consensus Guidelines for the Treatment of SAD. A Summary of the Report of the Canadian Consensus Group on SAD*. Vancouver, Canada: Clinical & Academic Publishing; 1999.
24. Horne JA, Östberg O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int J Chronobiol*. 1976;4:97-110.
25. The Automated Morningness-Eveningness Questionnaire. Available at: <http://www.cet.org>. Norwood, NJ: Center for Environmental Therapeutics. Accessed 22 September 2003.
26. Thalen BE, Kjellman BF, Morkrid L, Wetterberg L. Seasonal and non-seasonal depression. A comparison of clinical characteristics in Swedish patients. *Eur Arch Psychiatry Clin Neurosci*. 1995;245:101-108.
27. Terman M, Schlager D, Fairhurst S, Perlman B. Dawn and dusk simulation as a therapeutic intervention. *Biol Psychiatry*. 1989;25:966-970.
28. Avery DH, Bolte MA, Dager SR, et al. Dawn simulation treatment of winter depression: a controlled study. *Am J Psychiatry*. 1993;150:113-117.
29. Avery DH, Bolte MA, Wolfson JK, Kazaras AL. Dawn simulation compared with a dim red signal in the treatment of winter depression. *Biol Psychiatry*. 1994;36:180-188.
30. Avery DH, Kouri ME, Monaghan K, Bolte MA, Hellekson C, Eder D. Is dawn simulation effective in ameliorating the difficulty awakening in seasonal affective disorder associated with hypersomnia? *J Affect Disord*. 2002;69:231-236.
31. Lingjaerde O, Foreland AR, Dankertsen J. Dawn simulation vs lightbox treatment in winter depression: a comparative study. *Acta Psychiatr Scand*. 1998;98:73-80.
32. Terman M, Terman JS. Bright light therapy: side effects and benefits across the symptom spectrum. *J Clin Psychiatry*. 1999;60:799-808. Quiz 809.
33. Labbate LA, Lafer B, Thibault A, Sachs GS. Side effects induced by bright light treatment for seasonal affective disorder. *J Clin Psychiatry*. 1994;55:189-191.
34. Praschak-Rieder N, Neumeister A, Hesselmann B, Willeit M, Barnas C, Kasper S. Suicidal tendencies as a complication of light therapy for seasonal affective disorder: a report of three cases. *J Clin Psychiatry*. 1997;58:389-392.
35. Oren DA, Jacobsen FM, Wehr TA, Cameron CL, Rosenthal NE. Predictors of response to phototherapy in seasonal affective disorder. *Compr Psychiatry*. 1992;33:111-114.
36. Terman M, Amira L, Terman JS, Ross DC. Predictors of response and nonresponse to light treatment for winter depression. *Am J Psychiatry*. 1996;153:1423-1429.
37. Lam RW. Morning light therapy for winter depression: predictors of response. *Acta Psychiatr Scand*. 1994;89:97-101.
38. Reichborn-Kjennerud T, Lingjaerde O. Response to light therapy in seasonal affective disorder: personality disorders and temperament as predictors of outcome. *J Affect Disord*. 1996;41:101-110.
39. Levitt AJ, Joffe RT, Brecher D, MacDonald C. Anxiety disorders and anxiety symptoms in a clinic sample of seasonal and non-seasonal depressives. *J Affect Disord*. 1993;28:51-56.
40. Lam RW, Levitan RD. Pathophysiology of seasonal affective disorder: a review. *J Psychiatry Neurosci*. 2000;25:469-480.
41. Neumeister A, Konstantinidis A, Praschak-Rieder N, et al. Monoaminergic function in the pathogenesis of seasonal affective disorder. *Int J Neuropsychopharmacol*. 2001;4:409-420.
42. Willeit M, Praschak-Rieder N, Neumeister A, et al. [¹²⁵I]-Beta-CIT SPECT imaging shows reduced brain serotonin transporter availability in drug-free depressed patients with seasonal affective disorder. *Biol Psychiatry*. 2000;47:482-489.
43. Neumeister A, Praschak-Rieder N, Besselmann B, Rao ML, Gluck J, Kasper S. Effects of tryptophan depletion on drug-free patients with seasonal affective disorder during a stable response to bright light therapy. *Arch Gen Psychiatry*. 1997;54:133-138.
44. Neumeister A, Pirker W, Willeit M, et al. Seasonal variation of availability of serotonin transporter binding sites in healthy female subjects as measured by [¹²⁵I]-2 beta-carbomethoxy-3 beta-(4-iodophenyl)tropane and single photon emission computed tomography. *Biol Psychiatry*. 2000;47:158-160.
45. Lewy AJ, Cutler NL, Sack RL. The endogenous melatonin profile as a marker for circadian phase position. *J Biol Rhythms*. 1999;14:227-236.
46. Lewy AJ, Sack RL, Miller LS, Hoban TM. Antidepressant and circadian phase-shifting effects of light. *Science*. 1987;235:352-354.
47. Wehr TA, Duncan WC, Jr, Sher L, et al. A circadian signal of change of season in patients with seasonal affective disorder. *Arch Gen Psychiatry*. 2001;58:1108-1114.

Clinical research

48. Campbell SS, Murphy PJ. Extraocular circadian phototransduction in humans. *Science*. 1998;279:396-399.
49. Lockley SW, Skene DJ, Thapan K, et al. Extraocular light exposure does not suppress plasma melatonin in humans. *J Clin Endocrinol Metab*. 1998;83:3369-3372.
50. Eastman CI, Martin SK, Hebert M. Failure of extraocular light to facilitate circadian rhythm reentrainment in humans. *Chronobiol Int*. 2000;17:807-826.
51. Wirz-Justice A. Beginning to see the light. *Arch Gen Psychiatry*. 1998;55:861-862.
52. Kanofsky JD, Aspengren KL, Watts GR. Medicaid reimbursement for light therapy. *Am J Psychiatry*. 2003;160:796-797.
53. Ibatoullina E, Praschak-Rieder N, Kasper S. Severe atypical symptoms without depression in SAD: effects of bright light therapy. *J Clin Psychiatry*. 1997;58:495.
54. Jacobsen FM, Murphy DL, Rosenthal NE. The role of serotonin in seasonal affective disorder and the antidepressant response to phototherapy. In: Rosenthal NE, Blehar MC, eds. *Seasonal Affective Disorder and Phototherapy*. New York, NY: Guilford; 1989:333-341.
55. Dilsaver SC, Jaekle RS. Winter depression responds to an open trial of tranlycypromine. *J Clin Psychiatry*. 1990;51:326-329.
56. Teicher MH, Glod CA. Seasonal affective disorder: rapid resolution by low-dose alprazolam. *Psychopharmacol Bull*. 1990;26:197-202.
57. Wirz-Justice A, van der Velde P, Bucher A, Nil R. Comparison of light treatment with citalopram in winter depression: a longitudinal single case study. *Int Clin Psychopharmacol*. 1992;7:109-116.
58. Lingjaerde O, Reichborn-Kjennerud T, Haggag A, Gartner I, Narud K, Berg EM. Treatment of winter depression in Norway. II. A comparison of the selective monoamine oxidase A inhibitor moclobemide and placebo. *Acta Psychiatr Scand*. 1993;88:372-380.
59. Martinez B, Kasper S, Ruhrmann S, Moller HJ. Hypericum in the treatment of seasonal affective disorders. *J Geriatr Psychiatry Neurol*. 1994;7(suppl 1):S29-S33.
60. Yamadera H, Okawa M, Takahashi K. Open study of effects of alprazolam on seasonal affective disorder. *Psychiatry Clin Neurosci*. 2001;55:27-30.
61. Hilger E, Willeit M, Praschak-Rieder N, Stastny J, Neumeister A, Kasper S. Reboxetine in seasonal affective disorder: an open trial. *Eur Neuropsychopharmacol*. 2001;11:1-5.
62. McGrath RE, Buckwald B, Resnick EV. The effect of L-tryptophan on seasonal affective disorder. *J Clin Psychiatry*. 1990;51:162-163.
63. Ruhrmann S, Kasper S, Havelle B, et al. Effects of fluoxetine versus bright light in the treatment of seasonal affective disorder. *Psychol Med*. 1998;28:923-933.
64. Ghadirian AM, Murphy BE, Gendron MJ. Efficacy of light versus tryptophan therapy in seasonal affective disorder. *J Affect Disord*. 1998;50:23-27.
65. Gloth FM, 3rd, Alam W, Hollis B. Vitamin D vs broad spectrum phototherapy in the treatment of seasonal affective disorder. *J Nutr Health Aging*. 1999;3:5-7.
66. Wheatley D. Hypericum in seasonal affective disorder (SAD). *Curr Med Res Opin*. 1999;15:33-37.
67. Rosenthal NE, Jacobsen FM, Sack DA, et al. Atenolol in seasonal affective disorder: a test of the melatonin hypothesis. *Am J Psychiatry*. 1988;145:52-56.
68. Oren DA, Moul DE, Schwartz PJ, Wehr TA, Rosenthal NE. A controlled trial of levodopa plus carbidopa in the treatment of winter seasonal affective disorder: a test of the dopamine hypothesis. *J Clin Psychopharmacol*. 1994;14:196-200.
69. Schlager DS. Early-morning administration of short-acting β -blockers for treatment of winter depression. *Am J Psychiatry*. 1994;151:1383-1385.
70. Oren DA, Schulkin J, Rosenthal NE. 1,25-(OH)₂-Vitamin D₃ levels in seasonal affective disorder: effects of light. *Psychopharmacology (Berl)*. 1994;116:515-516.
71. Lam RW, Gorman CP, Michalon M, et al. Multicenter, placebo-controlled study of fluoxetine in seasonal affective disorder. *Am J Psychiatry*. 1995;152:1765-1770.
72. Lewy AJ, Bauer VK, Cutler NL, Sack RL. Melatonin treatment of winter depression: a pilot study. *Psychiatry Res*. 1998;77:57-61.
73. Thorell LH, Kjellman B, Arned M, Lindwall-Sundel K, Walinder J, Wetterberg L. Light treatment of seasonal affective disorder in combination with citalopram or placebo with 1-year follow-up. *Int Clin Psychopharmacol*. 1999;14(suppl 2):S7-S11.
74. Lingjaerde O, Foreland AR, Magnusson A. Can winter depression be prevented by *Ginkgo biloba* extract? A placebo-controlled trial. *Acta Psychiatr Scand*. 1999;100:62-66.
75. Turner EH, Schwartz PJ, Lowe CH, et al. Double-blind, placebo-controlled study of single-dose metergoline in depressed patients with seasonal affective disorder. *J Clin Psychopharmacol*. 2002;22:216-220.
76. L o H, Hale A, D'haenen H. Determination of the dose of agomelatine, a melatonergic agonist and selective 5-HT_{2c} antagonist, in the treatment of major depressive disorder: a placebo-controlled dose range study. *Int Clin Psychopharmacol*. 2002;17:239-247.
76. Partonen T, Vakkuri O, Lamberg-Allardt C, Lonnqvist J. Effects of bright light on sleepiness, melatonin, and 25-hydroxyvitamin D(3) in winter seasonal affective disorder. *Biol Psychiatry*. 1996;39:865-872.
77. Moscovitch A, Blashko C, Wiseman R. A double-blind, placebo-controlled study of sertraline in patients with seasonal affective disorder. New Research Abstracts, 151st Meeting of the American Psychiatric Association. Toronto, Canada. 1995. Abstract.
78. Wirz-Justice A, Graw P, Krauchi K, et al. Morning or night-time melatonin is ineffective in seasonal affective disorder. *J Psychiatr Res*. 1990;24:129-137.
79. Rosenthal NE, Sack DA, Jacobsen FM, et al. Melatonin in seasonal affective disorder and phototherapy. *J Neural Transm Suppl*. 1986;21:257-267.
80. Graw P, Haug HJ, Leonhardt G, Wirz-Justice A. Sleep deprivation response in seasonal affective disorder during a 40-h constant routine. *J Affect Disord*. 1998;48:69-74.