

Published in final edited form as:

*J Am Chem Soc.* 2011 August 17; 133(32): 12584–12589. doi:10.1021/ja206013j.

## Synthetic Versatility in C—H Oxidation: A Rapid Approach to Differentiated Diols and Pyrans from Simple Olefins

Paul E. Gormisky and M. Christina White\*

Roger Adams Laboratory, Department of Chemistry, University of Illinois, Urbana, Illinois 61801

### Abstract

Conventionally, C—H oxidation reactions are used to install functional groups. The use of C—H oxidation to transform simple starting materials into highly versatile intermediates, which enable rapid access to a range of complex target structures is a new area with tremendous potential in synthesis. Herein we report a Pd(II)/sulfoxide-catalyzed allylic C—H oxidation to form *anti*-1,4-dioxan-2-ones from homoallylic oxygenates. These versatile building blocks are rapidly elaborated to differentiated *syn*-1,2-diols, stereodefined amino-polyols, and *syn*-pyrans, structures ubiquitous in medicinally important complex molecules found in Nature. We also demonstrate that a C—H oxidation approach to the synthesis of these motifs is orthogonal and complementary to other state-of-the-art methods.

### Introduction

Selective C—H oxidation reactions have emerged as powerful methods for rapidly introducing functional groups onto preformed carbon skeletons,<sup>1</sup> often enabling a significant streamlining of the synthesis of complex molecules.<sup>2</sup> Tremendous untapped opportunity also exists to use C—H oxidation as a means of transforming simple starting materials into versatile synthetic intermediates en route to more complex products.<sup>3</sup> Ideally, these pluripotent intermediates could rapidly be transformed into a range of structurally and functionally diverse products of high synthetic value. Herein we describe the realization of this concept for the straightforward synthesis of *anti*-1,4-dioxan-2-ones from simple olefins using Pd(II)/sulfoxide catalysis. These highly versatile synthetic intermediates are rapidly elaborated to varied motifs prevalent in biologically active, oxygenated natural products: differentiated *syn*-1,2-diol functionality, polyfunctionalized acyclic structures, as well as *syn*-pyran skeletons.

### Design Principles

In addition to being present in some classes of natural products,<sup>4</sup> dioxanones are a particularly valuable target because of their potential for high synthetic versatility. Enolization of *anti*-dioxanones is known to promote skeletal rearrangement with relay of stereochemistry to furnish *syn*-pyrans.<sup>5</sup> Furthermore, the development of a method for selective cleavage of the dioxanone C—O bonds (etheral vs. acyl) would enable efficient installation of differentially protected *syn*-1,2-diol functionality.<sup>6</sup> In considering C—H activation platforms to access dioxanones, Pd(II)/sulfoxide catalysis presents unique opportunities for efficiency and selectivity.<sup>7–9</sup> Given our success in using tethered

Corresponding Author: white@scs.uiuc.edu.


Supporting Information. Complete experimental procedures, compound characterization data and NMR spectra for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

nucleophiles for diastereoselective allylic C—H aminations,<sup>8a, d</sup> we postulated that an intramolecular process would provide rapid access to stereochemically-defined 1,4-dioxan-2-ones via allylic C—H oxidation of readily available chiral homoallylic alcohols with a carboxylic acid tether. Moreover, given the proven levels of broad functional group tolerance and high chemoselectivity for  $\alpha$ -olefins versus more electron rich  $\pi$ -systems, a palladium(II)/sulfoxide catalyzed allylic C—H oxidation reaction would provide complementary reactivity to other olefin oxidation methods. The terminal olefin in these products may also be utilized for iterative C—H oxidation sequences, providing a new approach for retrosynthetic simplification of polyfunctionalized targets.<sup>10</sup> The ability to selectively transform simple homoallylic oxygenates to differentiated diols, polyfunctionalized acyclic structures, or stereochemically defined pyrans presents a unique opportunity to use C—H oxidation as a means of rapidly constructing diverse and biologically privileged oxygenated motifs.

## Results and Discussion

### Reaction Development

Despite the success of using carboxylic acid nucleophiles for intermolecular allylic C—H esterifications with a wide range of substrates,<sup>7</sup> we were surprised to find that substrates having oxygenation in the homoallylic position afforded only poor yields of 1,2-dioxygenation products with poor regioselectivities (Table 1, linear(L):branched(B) ratio). This low reactivity persists even when *p*-nitrobenzoic acid (previously shown to be the most effective nucleophile in intermolecular C—H esterifications) is used as a functionalization reagent and is unaffected by changing the nature of the homoallylic ether substituent (Table 1, R = CH<sub>2</sub>CO<sub>2</sub>Me or Bn). This result stands in stark contrast to the same reaction with alkyl homoallylic substitution, which proceeds in good yield (73%) and outstanding regioselectivity (>20:1 B:L) favoring the branched product, albeit with no diastereoselectivity.<sup>2a</sup> Intermolecular C—H esterification is thought to proceed via a serial ligand catalysis mechanism involving Pd(II)/sulfoxide promoted allylic C—H cleavage to generate a  $\pi$ -allylPd-carboxylate followed by a benzoquinone (BQ) mediated C—O bond forming event from within the sphere of the metal (eq. 1).<sup>7b, 11</sup> We hypothesize that homoallylic oxygen substituents may chelate to the palladium and occupy a requisite site for inner-sphere functionalization thereby forcing functionalization to proceed via an outer-sphere pathway. In support of this, the large amount of linear regioisomer formed (Table 1) is consistent with an outer-sphere, non-BQ dependent functionalization pathway.<sup>8b, c, 12</sup>



We hypothesized that by tethering the carboxylic acid nucleophile to the homoallylic oxygen substituent, a productive Pd-carboxylate chelate would be formed, which upon BQ activation, would lead to the desired C—O bond formation. In addition to overcoming reactivity and selectivity issues inherent to the intermolecular process, we considered that such an intramolecular process would also expand the synthetic utility of this method beyond formation of 1,2-dioxygenation patterns. We therefore examined the reaction of 2-((2-methylhex-5-en-3-yl)oxy)acetic acid **5a**, containing a carboxylic tether, which could

fulfill these goals (Table 2). Under standard allylic C—H oxidation conditions,<sup>7b</sup> dioxanone **6a** was produced in improved yield (38%), compared to intermolecular oxidation, and good diastereoselectivity favoring the *anti*-stereoisomer (Table 2, entry 1). We next examined diisopropylethylamine (DIPEA) and Cr(salen)Cl as catalytic additives due to their known ability to promote functionalization of carboxylic acid and carbamate nucleophiles without attenuating C—H cleavage.<sup>8b, c, 13</sup> While addition of 10 mol % DIPEA provided only a modest increase in yield (entry 2), 10 mol % Cr(salen)Cl markedly improved the yield to 83% with identical stereoselectivity (entry 3).<sup>14</sup> Interestingly, Cr(salen)Cl did not improve reactivity in the intermolecular C—H esterification system (Table 1, entries 2 and 4). We also performed a series of experiments to probe if this reaction, like the intermolecular version,<sup>7b</sup> proceeds via a serial ligand catalysis mechanism. Use of 2,6-dimethyl-1,4-benzoquinone (2,6-Me<sub>2</sub>BQ), a sterically encumbered quinone, which does not coordinate well to the  $\pi$ -allylPd intermediate, failed to produce the dioxanone (Table 2, entry 4). Moreover, Pd(OAc)<sub>2</sub> in the absence of bis-sulfoxide ligand also failed to furnish significant quantities of the desired product (entry 5). Finally, Cr and oxidant alone fail to promote the reaction (entry 6).<sup>15</sup> These experimental findings are consistent with the restoration of a serial ligand catalysis mechanism in the intramolecular C—H esterification reaction where both sulfoxide and quinone ligands working sequentially with the palladium are required to furnish the C—H oxidation product.

## Scope

We next evaluated a series of substrates to examine the effects of neighboring substituents (R group, Table 3) on reactivity and selectivity. Proximal aryl moieties are well tolerated (entry 2). Reactions with unbranched substrates proceeded in good yield, but diminished stereoselectivity (entry 3); whereas a substrate with a bulky quaternary alkyl substituent gave poor yields, but excellent diastereoselectivity (entry 4). Despite the apparent sensitivity to sterics, tertiary centers are well tolerated (entries 5 and 6). Notably, competing methods for forming these diol products would require setting the geometry of a tri-substituted olefin.<sup>16</sup> Additionally,  $\alpha$ -stereocenters influence the magnitude of the diastereoselectivity (entries 7 and 8), but the overall diastereomeric outcome of the reaction favoring the *anti*-dioxanone is relayed exclusively from the stereocenter bearing the carboxylic acid tether. Significantly, these reactions feature a high level of operational simplicity. The reactions are run with no precautions taken to exclude O<sub>2</sub> or water and the analytical scales directly translate to preparative scales with no diminishment in yield or selectivity (10 mmol scale, 80% yield, entry 1). Moreover, the major *anti*-dioxanone diastereomer can generally be isolated in pure form using standard chromatography techniques.

## Chemoselectivity

The C—H oxidation approach to these 1,2-dioxygenated motifs also provides orthogonal and complementary selectivity to the Sharpless asymmetric dihydroxylation (SAD).<sup>17</sup> When multiple olefins are present in a molecule, SAD generally selects for the more electron rich olefin with variable chemoselectivity. For example, SAD of a terminal diene to generate the vinylic diol products produced from C—H oxidation gives mixtures of regioisomeric diols.<sup>17b</sup> Consequently, SAD is often not used to install diols when multiple olefins are present and alternate, lengthy routes must be devised, reducing overall synthetic efficiency.<sup>18</sup> In contrast, the allylic C—H oxidation reaction gives complete selectivity for the terminal olefin allowing a diol to be installed in the presence of tetra-substituted, tri-substituted, and *trans*- and *cis*-di-substituted olefins (Table 4).

## Differentiated 1,2-Diols

We first sought to use this intramolecular C—H esterification reaction to develop a streamlined route to differentiated, chiral *syn*-1,2-diols (Scheme 1); compounds, which

generally require extensive functional group manipulations to access. In order to achieve this goal, we needed to: 1) evaluate the feasibility of generating optically enriched dioxanones using this method, and 2) develop a novel deprotection sequence for converting the chemically inequivalent oxygens in the dioxanone products to differentiated, chiral *syn*-1,2-diols. Starting from a simple aldehyde, asymmetric allylation affords an enantioenriched homoallylic alcohol. Alkylation with bromoacetic acid followed by C—H oxidation catalyzed by Pd(II)/bis-sulfoxide **1** gives *anti*-dioxanone (–)-**6a** with no erosion in enantioenrichment. The *anti*-dioxanone is rapidly unmasked via base-promoted lactone opening to generate a hydroxyacid, which can be protected to afford (–)-**9**. The  $\alpha$ -alkoxy ester moiety in (–)-**9**, is then cleaved under mild reducing conditions using SmI<sub>2</sub><sup>19</sup> to furnish a free alcohol, which may be protected as desired. Benzyloxymethyl protection gave differentiated diol (–)-**10**, a compound that is challenging to obtain via other oxidation methods, like the Sharpless asymmetric dihydroxylation (SAD), which directly affords unprotected *syn*-1,2-diol motifs.

### Polyoxidized Motifs via Iterative C—H Oxidation

The  $\alpha$ -olefin moiety of the differentiated *syn*-1,2-diol products (e.g. (–)-**10**) can be used as a functional handle to iterate this and other allylic C—H oxidation processes for the purpose of rapidly constructing densely functionalized chains from simple starting materials (Scheme 2).<sup>10</sup> A hydroboration-oxidation sequence affords an intermediate aldehyde that can undergo asymmetric allylation to generate a homoallylic alcohol as one diastereomer. After carbamate formation to yield (–)-**11**, a masked *syn*-1,2-amino alcohol moiety can be installed in good yields via Pd(II)/bis-sulfoxide **1** catalyzed allylic C—H amination.<sup>8a</sup> Amino-polyols like (–)-**12**, which are generated in optically pure form via this sequence, are found in several classes of natural products,<sup>20</sup> such as bengazole A (a potent antifungal agent) and AAL Toxin T<sub>A</sub>. Traditional approaches to these polyoxidized motifs often rely on chiral relay strategies, making the synthesis of multiple stereoisomeric compounds challenging. This sequence highlights the power of diastereoselective C—H oxidation and amination reactions, when used in combination with powerful reagent controlled alkylation methodology, to facilitate the synthesis of a wide assortment of diastereomers with varied oxygen and nitrogen motifs.

### Syn-Pyrans

Efficient generation of dioxanones via C—H oxidation additionally allows for rapid structural diversification from topologically simple starting materials. 1,4-dioxan-2-ones undergo stereoselective Ireland-Claisen<sup>5d</sup> rearrangements to form *syn*-pyran skeletons.<sup>5,21</sup> We present two case studies illustrating the utility and advantages of C—H oxidation for the synthesis of *syn*-pyrans. First, bifunctional pyrans like (–)-**16** are powerful building blocks used in a number of natural product syntheses. For example, dioxanone **14**, readily formed via C—H oxidation, undergoes facile rearrangement to give dihydropyran **15** (Scheme 3). Hydrogenation, reduction of the ester, benzylation and desilylation gives (–)-**16**, an intermediate in the synthesis of SCH 351448.<sup>22</sup> Our intramolecular C—H oxidation not only sets a stereocenter selectively, which is relayed to the pyran 2-position, but the tether can also be incorporated into the final product, eliminating extraneous masking or directing groups.

Particularly in the context of complex molecule synthesis, C—H oxidation methods must be effective in the presence of dense functionality. Scheme 4 shows the rapid synthesis of highly oxygenated dioxanone (+)-**18** and subsequent rearrangement to dihydropyran (+)-**19**. Reduction of the ester, benzyl protection and hydrolysis of the acetonide gives diol (+)-**20**, an intermediate in the synthesis of *ent*-goniodomin A, a potent antifungal agent.<sup>23</sup> This route compares favorably to the reported synthesis of this pyran (7 versus 10 steps from

commercial material) and demonstrates the ability of C—H oxidation to enable the rapid construction of densely functionalized pyran skeletons.

## Conclusion

We have developed a novel approach to differentiated polyoxygenated motifs and bifunctional *syn*-pyrans from homoallylic alcohols using Pd(II)/bis-sulfoxide C—H oxidation catalysis. This work underscores the power of selective C—H oxidation reactions for installing versatile functionality that enables rapid access to functionally and topologically diverse structures from simple starting materials.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgments

Financial support was provided by NIH/NIGMS (GM07615). P.E.G. is the recipient of the University of Illinois Synthetic Organic and NSF Graduate Research Fellowships. We thank TCI and Aldrich Chemical Company for generous gifts of commercial bis-sulfoxide/Pd(OAc) catalyst (**1**). We also thank Mr. Marinus Bigi for checking the experimental procedure in Table 3, entry 1.


## References

1. For examples of C—H activation reactions to introduce oxygen see: (a) Chen MS, White MC. *Science*. 2010; 327:566. [PubMed: 20110502] (b) Chen MS, White MC. *Science*. 2007; 318:783. [PubMed: 17975062] (c) Groves JT, Nemo TE. *J Am Chem Soc*. 1982; 105:6243. (d) Chen K, Que L Jr. *J Am Chem Soc*. 2001; 123:6327. [PubMed: 11427057] (e) Mello R, Fiorentino M, Fusco C, Curci R. *J Am Chem Soc*. 1989; 111:6749. (f) Brodsky BH, Du Bois J. *J Am Chem Soc*. 2005; 127:15391. [PubMed: 16262401] (g) Zhou M, Schley ND, Crabtree RH. *J Am Chem Soc*. 2010; 132:12550. [PubMed: 20731395] (h) Dick AR, Hull KL, Sanford MS. *J Am Chem Soc*. 2004; 126:2300. [PubMed: 14982422] (i) Zhang YH, Yu JQ. *J Am Chem Soc*. 2009; 131:14654. [PubMed: 19788192]
2. For examples of late-state C—H oxidations in total synthesis see: (a) Stang EM, White MC. *Nat Chem*. 2009; 1:547. [PubMed: 21378935] (b) Hinmann A, DuBois J. *J Am Chem Soc*. 2003; 125:11510. [PubMed: 13129349] (c) Davies HML, Dai X, Long MS. *J Am Chem Soc*. 2006; 128:2485. [PubMed: 16478205] (d) Kim J, Ashenurst JA, Movassaghi M. *Science*. 2009; 324:238. [PubMed: 19359584] (e) Nicolaou KC, Yang Z, Liu JJ, Ueno H, Nantermet PG, Guy RK, Claibourne CF, Reynaud J, Couladouros EA, Paulvannan K, Sorenson EJ. *Nature*. 1994; 367:630. [PubMed: 7906395] (f) Wender PA, Jesudason CD, Nakhira H, Tamura N, Tebe AL, Ueno YJ. *J Am Chem Soc*. 1997; 119:12976.
3. For an example of C—H activation/Cope rearrangement for forming carbocyclic skeletons see: (a) Davies HML, Jin Q. *Proc Nat Acad Sci*. 2004; 101:5472. [PubMed: 15024094] (b) Hansen J, Gregg TM, Ovalles SR, Lian Y, Autschbach J, Davies HML. *J Am Chem Soc*. 2011; 133:5076. [PubMed: 21384883]
4. Washida K, Abe N, Sugiyama Y, Hirota A. *Biosci Biotechnol Biochem*. 2009; 73:1355. [PubMed: 19502726]
5. (a) Burke SD, Armistead DM, Schoenen FJ, Fevig JM. *Tetrahedron*. 1986; 42:2787. For the dioxanone to pyran approach previously applied to total synthesis see: (b) Burke SD, Sametz GM. *Org Lett*. 1999; 1:71. [PubMed: 10822536] (c) Burke SD, Piscopio AD, Kort ME, Matulenko MA, Parker MH, Armistead DM, Shankaran K. *J Org Chem*. 1994; 59:332. For the seminal report of the Ireland-Claisen rearrangement see: (d) Ireland RE, Mueller RH. *J Am Chem Soc*. 1972; 94:5897.
6. For an example of the synthesis of differentiated cyclic 1,2-diols see: Lim SM, Hill N, Myers AG. *J Am Chem Soc*. 2009; 131:5763.
7. For C—H esterification see: (a) Chen MC, White MC. *J Am Chem Soc*. 2004; 126:1346. [PubMed: 14759185] (b) Chen MS, Prabakaran N, Labenz NA, White MC. *J Am Chem Soc*. 2005; 127:6970.

- [PubMed: 15884938] (c) Fraunhofer KF, Prabakaran N, Sirois LE, White MC. *J Am Chem Soc.* 2006; 128:9032. [PubMed: 16834366]
8. For C—H amination see: (a) Fraunhofer KJ, White MC. *J Am Chem Soc.* 2007; 129:7274. [PubMed: 17516648] (b) Reed SA, White MC. *J Am Chem Soc.* 2008; 130:3316. [PubMed: 18302379] (c) Reed SA, Mazzotti AR, White MC. *J Am Chem Soc.* 2009; 131:11701. [PubMed: 19645492] (d) Rice GT, White MC. *J Am Chem Soc.* 2009; 131:11707. [PubMed: 19645489]
  9. For C—H alkylation see: (a) Young AJ, White MC. *J Am Chem Soc.* 2008; 130:14090. [PubMed: 18831588] (b) Young AJ, White MC. *Angew Chem Int Ed.* 2011; 50:1002/anie.201101654(c) Lin S, Song CX, Cai GX, Wang WH, Shi ZJ. *J Am Chem Soc.* 2008; 130:12901. [PubMed: 18778061]
  10. For examples of an iterative approach to polyol synthesis see: (a) Zacuto MJ, Leighton JL. *J Am Chem Soc.* 2000; 122:8587.(b) Iwata M, Yazaki R, Suzuki Y, Kumagai N, Shibasaki M. *J Am Chem Soc.* 2009; 131:18244. [PubMed: 19994862] (c) Menz H, Krisch SF. *Org Lett.* 2009; 11:5634. [PubMed: 19919037] (d) Hassan A, Lu Y, Krische MJ. *Org Lett.* 2009; 11:3112. [PubMed: 19586067]
  11. An analogous intermediate has been proposed for the Pd(OAc)<sub>2</sub>-catalyzed 1,4-diacetoxylation of 1,3-dienes and stereochemical evidence supports an inner-sphere C—O bond forming process. See: Bäckvall JE, Byström SE, Nordberg RE. *J Org Chem.* 1984; 49:4619.
  12. For example, intermolecular C—H amination, thought to proceed via outer-sphere C—N bond formation, forms predominantly the linear regioisomer and is tolerant of homoallylic oxygen functionality. See ref. 8b.
  13. For a detailed exploration of the role of Cr(salen)X in allylic C—H oxidations see: (a) Covell DJ, White MC. *Angew Chem, Int Ed.* 2008; 47:1. For other applications of Cr(salen)Cl in C—H oxidations see references 8b and 8c and: (b) Qi X, Rice GT, Lall MS, Plummer MS, White MC. *Tetrahedron.* 2010; 66:4816. [PubMed: 21379377] (c) Wu L, Qiu S, Liu G. *Org Lett.* 2009; 11:2707. [PubMed: 19456146]
  14. The use of lower palladium loadings for C—H dioxanone formation led to significantly diminished yields. For a discussion on the need for high catalyst loadings in Pd(II)-catalyzed oxidations see: (a) Steinhoff BA, King AE, Stahl SS. *J Org Chem.* 2006; 71:1861. [PubMed: 16496970] For recent C—H oxidation processes used in drug discovery that operate at 10 mol% Pd(II) catalyst loadings see: ref. 13b and (b) Dai HX, Stepan AF, Plummer MS, Zhang YH, Yu JQ. *J Am Chem Soc.* 2011; 133:7222. [PubMed: 21488638]
  15. For PhI(OAc)<sub>2</sub>-mediated benzylic C—H lactonizations see: Dohi T, Takenaga N, Goto A, Maruyama A, Kita Y. *Org Lett.* 2007; 9:3129. [PubMed: 17616204] In contrast, BQ and PhI(OAc)<sub>2</sub> alone failed to promote this reaction: see Supporting Information S8–9.
  16. For an example in total synthesis see: Taillier C, Gille B, Bellosta V, Cossy J. *J Org Chem.* 2005; 70:2097. [PubMed: 15760193]
  17. (a) Kolb HC, VanNieuwenhze MS, Sharpless BK. *Chem Rev.* 1994; 94:2483.(b) Xu D, Crispino GA, Sharpless KB. *J Am Chem Soc.* 1992; 114:7570.
  18. For an example of installation of a diol motif in the presence of unsaturated functionality see: Sabitha G, Gopal P, Reddy N, Yadav JS. *Tetrahedron Lett.* 2009; 50:6298.
  19. For general applications of SmI<sub>2</sub> see: (a) Molander GA, Harris CR. *Chem Rev.* 1996; 96:307. [PubMed: 11848755] For cleavage of  $\alpha$ -alkoxy esters see: (b) Kasuda K, Inanaga J, Yamaguchi M. *Tetrahedron Lett.* 1989; 30:2945.
  20. (a) Chandrasekhar S, Sudhakar A. *Org Lett.* 2010; 12:236. [PubMed: 20017536] (b) Shier WT, Abbas HK, Badria FA. *Tetrahedron Lett.* 1995; 36:1571.
  21. For selected examples of pyran synthesis in complex molecules see: (a) Smith AB III, Verhoest PR, Minbiole KP, Schelhaas M. *J Am Chem Soc.* 2001; 123:4834. [PubMed: 11457294] (b) Simpson GL, Heffron TP, Merino E, Jamison TF. *J Am Chem Soc.* 2006; 128:1056. [PubMed: 16433504] (c) Keck GE, Kraft MB, Truong AP, Li W, Sanchez CC, Kedei N, Lewin NE, Blumberg PM. *J Am Chem Soc.* 2008; 130:6660. [PubMed: 18452293]
  22. Cheung LL, Marumoto S, Anderson CD, Rychnovsky SD. *Org Lett.* 2008; 10:3101. [PubMed: 18543942]
  23. Saito T, Fuwa H, Sasaki M. *Org Lett.* 2009; 11:5274. [PubMed: 19905029]




**Figure 1.**  
Rapid Access to Complex Structures via Allylic C—H Oxidation



**Scheme 1. Access to Differentiated *syn*-1,2-Diols**

<sup>a</sup>Conditions: (a) 10% **1**, 10% Cr(salen)Cl, BQ (2.0 equiv), dioxane, 45 °C (72% of >20:1 *anti*-diastereomer; 83%, 9:1 crude dr); (b)(1) LiOH (2.0 equiv), 3:1 THF:H<sub>2</sub>O, 0 °C, (2) TBSOTf (3.0 equiv), 2,6-lutidine (6.0 equiv), CH<sub>2</sub>Cl<sub>2</sub> 0 °C, (3) MeI (3.0 equiv), K<sub>2</sub>CO<sub>3</sub> (3.0 equiv, DMF, RT (89%, 3 steps); (c) SmI<sub>2</sub> (3.0 equiv), ethylene glycol (1.2 equiv), THF/HMPA, RT (61%, 12% rsm); (d) BOMCl (1.5 equiv), *i*Pr<sub>2</sub>NEt (1.75 equiv, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to RT (79%).




**A. Generalized Approach**


\* =reagent controlled asymmetric allylation or alkylation

□ =diastereoselective allylic C–H oxidation or amination

X=O,  $n = 0$ ; X=N,  $n = 0, 1$


**B. Successful Implementation****Scheme 2. Iterative C–H Oxidation for the Synthesis of Polyoxidized Motifs**

<sup>a</sup>Conditions: (a) (1)  $\text{BH}_3\text{-Me}_2\text{S}$  (2.4 equiv), THF; 2-methyl-2-butene (4.7 equiv); **7**, 0 to 45 °C; 3.0 M NaOH, 30% wt.  $\text{H}_2\text{O}_2$ , (2) PCC (1.3 equiv),  $\text{CH}_2\text{Cl}_2$ , RT (74%, 2 steps); (b) (+)-Ipc<sub>2</sub>B-allyl,  $\text{Et}_2\text{O}$ , –78 °C; 3.0 M NaOH, 30% wt.  $\text{H}_2\text{O}_2$  (95%); (c) NsNCO (1.5 equiv), THF, RT (90%); (d) 10% **1**, 5% 1,2-bis(phenylsulfonyl)ethane, PhBQ (1.05 equiv), THF, 45 °C (65%, 1.6:1 crude dr).



### Scheme 3. Synthesis of a Useful Bifunctional Pyran

<sup>a</sup>Conditions: (a) NaH (3.0 equiv), BrAcOH (1.1 equiv), THF/DMF 0 °C to RT (70%); (b) 10% **1**, 10% Cr(salen)Cl, BQ (2.0 equiv), dioxane, 65 °C (83%), 3:1 crude dr, mixture of diastereomers taken forward); (c) (1) LiHMDS (2.0 equiv), 1:1 v:v TMSCl:Et<sub>3</sub>N, THF, -78 °C then reflux in toluene, (2) MeI (3.0 equiv), K<sub>2</sub>CO<sub>3</sub> (3.0 equiv), DMF, RT (83%); (d) 10% wt. of 5% Pd/C, H<sub>2</sub> (1 atm), EtOAc, RT (68% of >20:1 *syn*-diastereomer, 3:1 crude dr); (e) LiAlH<sub>4</sub> (2.0 equiv), THF, 0 °C; (f) BnBr (2.0 equiv), NaH (2.0 equiv), DMF, 0 °C to RT; (g) 3 M HCl, EtOH, RT (74%, 3 steps).



#### Scheme 4. Synthesis of a Densely Functionalized Pyran

<sup>a</sup>Conditions: (a) NaH (3.0 equiv), BrAcOH (1.1 equiv), THF/DMF 0 °C to RT (54%); (b) 10% **1**, 10% Cr(salen)Cl, BQ (2.0 equiv), dioxane, 65 °C (56% of >20:1 dr *anti*-diastereomer; 73%, 3:1 crude dr); (c) (1) LiHMDS (2.0 equiv), 1:1 v:v TMSCl:Et<sub>3</sub>N, THF, -78 °C then reflux in toluene, (2) MeI (3.0 equiv), K<sub>2</sub>CO<sub>3</sub> (3.0 equiv), DMF, RT (82%); (d) LiAlH<sub>4</sub> (2.0 equiv), THF, 0 °C; (e) BnBr (2.0 equiv), NaH (2.0 equiv), DMF, 0 °C to RT; (f) 1N HCl, THF, RT (60%, 3 steps).

Table 1


Intermolecular C—H Esterification with Homoallylic Oxygen Substitution

entry	R	R <sup>a</sup>	additive	isolated yield <sup>b</sup>	L:B <sup>c</sup> 3/4	E:Z <sup>d</sup> (3) <sup>d</sup>	dr (4) <sup>e</sup>
1	CH <sub>2</sub> CO <sub>2</sub> Me	Ac	none	11	1:2	>20:1	16:1
2	CH <sub>2</sub> CO <sub>2</sub> Me	Ac	Cr(salen)Cl	10	>20:1	>20:1	N/A
3	CH <sub>2</sub> CO <sub>2</sub> Me	<i>p</i> -NO <sub>2</sub> Bz	none	16	1:2	nd	>20:1
4	CH <sub>2</sub> CO <sub>2</sub> Me	<i>p</i> -NO <sub>2</sub> Bz	Cr(salen)Cl	>5	nd	nd	nd
5	Bn	<i>p</i> -NO <sub>2</sub> Bz	none	>5	nd	nd	nd

<sup>a</sup> AcOH (4.0 equiv) or *p*-NO<sub>2</sub>BzOH (1.5 equiv) used as the acid nucleophile.<sup>b</sup> Average of 2 runs at 0.5 mmol.<sup>c</sup> Linear(L):Branched(B) ratio determined by GC of the crude reaction after workup.<sup>d</sup> E:Z ratio determined by GC of the crude reaction after workup.<sup>e</sup> Diastereomeric ratio determined by GC of the crude reaction after workup. BQ = *p*-benzoquinone, salen = 1,2-cyclohexanediamine-*N,N'*-bis(3,5-di-*tert*-butylsalicylidine).

Table 2

## Optimization of the Intramolecular C—H Oxidation








entry	additive	quinone	Pd catalyst	isolated yield	dr (anti:syn) <sup>d</sup>
1	none	BQ	<b>1</b>	38	9:1
2	DIPEA	BQ	<b>1</b>	46	7:1
3	Cr(salen)Cl	BQ	<b>1</b>	83	9:1
4	none	2,6-Me <sub>2</sub> BQ	<b>1</b>	0	nd
5	none	BQ	Pd(OAc) <sub>2</sub>	>5	nd
6	Cr(salen)Cl	BQ	none	0	nd

<sup>d</sup>Diastereomeric ratio determined by <sup>1</sup>H NMR of the crude reaction after workup. DIPEA = diisopropylethylamine.

Table 3

Scope of the Intramolecular C—H Oxidation




entry	R	product	isolated yield <sup>a</sup>	dr (anti:syn) <sup>b</sup>
1	<i>i</i> -Pr	(-)- <b>6a</b>	83 (80) <sup>c</sup>	9:1 <sup>d</sup>
2	4-BrPh	<b>6b</b>	57	3:1
3	<i>n</i> -Pent	<b>6c</b>	82	2:1
4	<i>t</i> -Bu	<b>6d</b>	22	11:1
5		<b>6e</b> R=Ph	57	3:1
6		<b>6f</b> R= <i>n</i> -Bu	60	3:1
7		(-)- <b>6g</b>	62	8:1
8		(+)- <b>6h</b>	62	4:1

<sup>a</sup> Average of 2 runs at 0.3 mmol.<sup>b</sup> Diastereomeric ratio determined by GC of the crude reaction after workup.<sup>c</sup> Average of 2 runs at 10 mmol.<sup>d</sup> Diastereomeric ratio determined by <sup>1</sup>H NMR of the crude reaction after workup.

Table 4

## Chemoselectivity of the Intramolecular C—H Oxidation



entry	product	isolated yield <sup>a</sup>	dr (anti:syn) <sup>b</sup>
1	 <b>8a</b>	74	4:1
2	 <b>8b</b>	77	6:1
3	 <b>8c</b>	53	7:1
4	 <b>8d</b>	72	3:1
5	 <b>8e</b>	52	2:1

<sup>a</sup> Average of 2 runs at 0.3 mmol.

<sup>b</sup> Diastereomeric ratio determined by GC of the crude reaction after workup.