

REVIEW

Open Access

Manual therapy with and without vestibular rehabilitation for cervicogenic dizziness: a systematic review

Reidar P Lystad*, Gregory Bell, Martin Bonnevie-Svensen and Catherine V Carter

Abstract

Background: Manual therapy is an intervention commonly advocated in the management of dizziness of a suspected cervical origin. Vestibular rehabilitation exercises have been shown to be effective in the treatment of unilateral peripheral vestibular disorders, and have also been suggested in the literature as an adjunct in the treatment of cervicogenic dizziness. The purpose of this systematic review is to evaluate the evidence for manual therapy, in conjunction with or without vestibular rehabilitation, in the management of cervicogenic dizziness.

Methods: A comprehensive search was conducted in the databases Scopus, Mantis, CINAHL and the Cochrane Library for terms related to manual therapy, vestibular rehabilitation and cervicogenic dizziness. Included studies were assessed using the Maastricht-Amsterdam criteria.

Results: A total of fifteen articles reporting findings from thirteen unique investigations, including five randomised controlled trials and eight prospective, non-controlled cohort studies were included in this review. The methodological quality of the included studies was generally poor to moderate. All but one study reported improvement in dizziness following either unimodal or multimodal manual therapy interventions. Some studies reported improvements in postural stability, joint positioning, range of motion, muscle tenderness, neck pain and vertebralbasilar artery blood flow velocity.

Discussion: Although it has been argued that manual therapy combined with vestibular rehabilitation may be superior in the treatment of cervicogenic dizziness, there are currently no observational and experimental studies demonstrating such effects. A rationale for combining manual therapy and vestibular rehabilitation in the management of cervicogenic dizziness is presented.

Conclusion: There is moderate evidence to support the use of manual therapy, in particular spinal mobilisation and manipulation, for cervicogenic dizziness. The evidence for combining manual therapy and vestibular rehabilitation in the management of cervicogenic dizziness is lacking. Further research to elucidate potential synergistic effects of manual therapy and vestibular rehabilitation is strongly recommended.

Keywords: Cervicogenic dizziness, Vertigo, Manual therapy, Vestibular rehabilitation, Spinal manipulation, mobilisation

Background

Dizziness is a non-specific symptom that is commonly encountered by primary health care practitioners [1], and the prevalence has been reported to be between 11.1% and 28.9% [2-5]. It can be experienced as faintness, unsteadiness, perception of spinning and

disorientation [6-8]. The mechanisms producing these symptoms are multiple and can involve several different organ systems. Ardc, Topuz and Kara [9] reported the most frequent diagnosis of patients suffering from dizziness to be benign paroxysmal positional vertigo, endolymphatic hydrops, migraine, central decompensation, acute vestibulopathy and autonomic dysfunction. Furthermore, it is not uncommon for patients experiencing dizziness to have more than one diagnosis.

* Correspondence: reidar.lystad@mq.edu.au
Department of Chiropractic, Macquarie University, Sydney, Australia

Dizziness is commonly seen in whiplash patients, affecting 20-58% of individuals with flexion-extension injuries [10].

One specific type of dizziness is cervicogenic dizziness. The disorder was first described as "cervical vertigo" by Ryan and Cope in 1955 [11]. Other terms used for the same disorder are proprioceptive vertigo, cervicogenic vertigo and cervical dizziness [12-14]. Although the diagnosis has remained controversial since its introduction, several observations have led to the proposal of a plausible pathophysiological mechanism. The deep intervertebral muscles in the cervical spine possess a high density of muscle spindles and are assumed to play an important role in postural control [15-18]. Cervical afferents are known to be involved in the cervico-collic reflex, the cervico-ocular reflex and the tonic neck reflex, which work in conjunction with other reflexes influenced by visual and vestibular systems to stabilise the head, the eyes and posture [19]. Vestibular and proprioceptive input is linearly combined for computing egocentric, body-centred coordinates [20].

Several authors have demonstrated that anaesthetic injections to the upper cervical dorsal nerve roots can produce dizziness and nystagmus [21-23]. Electrical stimulation to cervical muscles has also been shown to induce a sensation of tilting or falling [24]. Brandt and Bronstein [25] proposed a mechanism where changed firing characteristics of cervical somatosensory receptors due to neck pain lead to a sensory mismatch between vestibular and cervical input, resulting in cervical vertigo.

Several authors have proposed manual therapy interventions for the treatment of dizziness of a cervical origin [26-28]. Indeed, it has been suggested that the management of cervicogenic dizziness should be the same as for cervical pain [25]. In a systematic review of the literature, Reid and Rivett [29] concluded that there is limited evidence to support manual therapy treatment of cervicogenic dizziness. Moreover, it was recommended that further research be conducted, especially randomised controlled trials (RCTs), to provide more conclusive evidence of the role of manual therapy for cervicogenic dizziness.

Another treatment modality that is advocated for cervical pain is sensorimotor rehabilitation exercises [19,30]. These exercises fall under the scope of exercises included in vestibular rehabilitation therapy. Vestibular rehabilitation emerged as a group of exercises for peripheral vestibular disorders, aiming to maximise central nervous system compensation to vestibular pathology [31,32]. These exercises are usually movement based, and can be further subcategorised according to different physiological rationales: (i) compensatory responses using motion to habituate activity in the vestibular

nuclei; (ii) adaptation for visual-vestibular interaction and possibly eye/hand coordination, using repetitive and provocative movements of the head and/or eyes; (iii) substitution which promotes the use of individual or combinations of sensory inputs to bias use away from dysfunctional vestibular input; (iv) postural control exercises, falls prevention, relaxation training, reconditioning activities and functional/occupational retraining, which are based on motor learning principles [33,34].

Hillier and Hollohan [34] concluded that there was moderate to strong evidence that vestibular rehabilitation is safe and effective in the management of unilateral peripheral vestibular disorders. Moreover, several authors encourage the implementation of vestibular rehabilitation in treatment of dizziness of a cervical origin [10,32,35], and published case studies have reported positive outcomes when combining manual therapy and vestibular rehabilitation [36,37].

To the authors' knowledge, the evidence of implementing vestibular rehabilitation with manual therapy in the management of cervicogenic dizziness has not been systematically reviewed. Thus, the purpose of this systematic review was: (i) to provide an updated systematic review of manual therapy for cervicogenic dizziness by including higher level evidence published since the previous review by Reid and Rivett [29], and (ii) to compare the evidence of (a) manual therapy with vestibular rehabilitation for cervicogenic dizziness with (b) manual therapy without vestibular rehabilitation for cervicogenic dizziness.

Methods

This systematic review adhered to the guidelines outlined in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Statement [38].

Eligibility criteria

This systematic review was limited to prospective, controlled or non-controlled intervention studies published in peer-reviewed journals. Retrospective study designs, case reports, case series, commentaries, letters to the editor, and expert opinions were excluded from this review. No language restrictions were applied in this review.

Eligible studies had to investigate a cohort of patients diagnosed with cervicogenic dizziness. Cervicogenic dizziness was defined as the presence of dizziness, imbalance or unsteadiness related to movements or position of the cervical spine, or occurring with a stiff or painful neck [29]. Studies investigating populations diagnosed with cardiovascular disorders, central nervous system disorder (e.g. cerebellar ataxia, stroke, demyelination), Mal de Debarquement syndrome, migraine-associated vertigo, psychogenic dizziness, vestibular disorders (e.g.

benign paroxysmal positional vertigo, Meniere's disease, peripheral vestibulopathy), were not included in this review. Studies were also excluded if the study population was comprised of patients with a history of active inflammatory joint disease, spinal cord pathology, cervical spine cancer or infection, bony disease or marked osteoporosis, marked cervical spine disc protrusion, acute cervical nerve root symptoms, fracture or dislocation of the neck, or previous surgery to the upper cervical spine.

This review considered two possible interventions, namely manual therapy alone and manual therapy in conjunction with vestibular rehabilitation. For the purposes of this review, manual therapy was defined as spinal manipulation (high velocity, low amplitude techniques) or mobilisation (low-velocity, small or large amplitude techniques) [29]. Vestibular rehabilitation was defined as an exercise-based group of approaches with the aim of maximising the central nervous system compensation for vestibular pathology [39]. Vestibular rehabilitation techniques included habituation (movement provoking) with gaze stabilising (adaptation), sensory substitution, and balance and gait/activity training [34].

Search Strategy

A comprehensive search of the literature was conducted, including electronic searches of the Scopus, Mantis, and CINAHL databases from January 1955 to June 2010. In addition, the Cochrane Library was searched from inception (1993) to June 2010 to identify any relevant Cochrane Reviews. Keywords used in the literature search included "cervicogenic dizziness" and "manual therapy". Alternative spellings, synonyms and related terms, and truncated versions of both the condition and the intervention were included. In addition, bibliographies of included studies and relevant review articles were hand searched to identify potentially eligible studies not captured by the electronic searches.

Study selection

Citations from the electronic searches were combined in a single list and duplicate records were discarded. Two reviewers screened all titles and abstracts to identify and remove obviously irrelevant citations. Full text versions of all potentially eligible articles were retrieved and evaluated by two independent reviewers to determine eligibility for inclusion in this review. Any differences were resolved by mutual consensus with a third independent reviewer.

Data extraction process

Data from eligible studies were extracted and compiled in a spreadsheet. For the purposes of this systematic review the following data were extracted: (i) study

population (e.g. age, gender, diagnosis, and sample size); (ii) study design; (iii) intervention; (iv) outcome measures; and (v) main findings.

Data analysis

Owing to the clinically heterogeneous nature of the included studies (i.e. varying study designs, interventions, outcome measures, and quality of data), a meta-analysis was deemed unfeasible. Thus, in this review only a qualitative analysis of included studies was undertaken. As per the previous review by Reid and Rivett [29], qualitative analysis was achieved by attributing levels that rate the scientific evidence, i.e. Level 1: Strong evidence (provided by generally consistent findings in multiple higher quality RCTs); Level 2: Moderate evidence (provided by generally consistent findings in one higher quality RCT and one or more lower quality RCTs); Level 3: Limited evidence (provided by generally consistent findings in one or more lower quality RCTs); and Level 4: No evidence (if there were no RCTs or if the results were conflicting).

Assessment of methodological quality

The methodological quality of the included studies was assessed using the Maastricht-Amsterdam criteria [40]. The Maastricht-Amsterdam criteria list, which consists of 19 items assessing patient selection, interventions, outcome measures and statistics, is included in Additional file 1. Two independent reviewers assessed methodological quality and any differences were resolved by mutual consensus with a third independent reviewer. Each item was answered "yes", "no", or "don't know", and one point was assigned for each "yes" (fulfilled item). The assessed studies were categorised as either poor, moderate or good based on the percentage of fulfilled items from the Maastricht-Amsterdam criteria list. In accordance with other authors using similar quality assessment methods, the cut-off percentage values were arbitrarily set at < 50% (poor), 50-80% (moderate), and > 80% (good) [41-43].

Results

The electronic searches returned 658 hits, which included 335 duplicate records and 323 unique citations. After removing duplicate records and screening titles and abstracts to discard obviously irrelevant citations, a total of 42 potentially eligible studies were identified. A hand search revealed four additional studies that were not captured by the electronic search. Thus, a total of 46 potentially eligible studies were evaluated for inclusion in this systematic review. Thirty-one studies [29,36,44-72] did not meet the inclusion criteria and were excluded from this review. See Additional file 2 for a list of excluded studies including reasons for

exclusion. Figure 1 contains a flow diagram of the study selection process. Two articles [26,73] reported data from the same RCT, and the results from one cohort study were published in two separate articles [74,75] Thus, this review included reports from a total of thirteen unique investigations. See Table 1 for a description of included studies.

The included studies comprised five RCTs [26,76-79] and eight prospective, non-controlled, cohort studies [75,80-86], with samples sizes ranging from 12 to 168. One study [81] did not report on the gender distribution of recruited participants, however all but one of the

remaining studies included more females, ranging from 52% to 88%.

Six studies [75-77,80-82], including two RCTs [76,77], used only spinal manipulation or mobilisation, or both, as the intervention. The remaining seven investigations [26,78,79,83-86], including three RCTs [26,78,79] utilised a multimodal approach consisting of several different interventions (e.g. spinal manipulation and mobilisation, soft tissue therapy, electrotherapy, and medications) and home exercise programs. However, none of the included studies used manual therapy in conjunction with vestibular rehabilitation.




Figure 1 PRISMA flow diagram.

Table 1 Included studies

Study	Participants	Interventions		Outcome measures	Main findings	Quality
		Manual therapy	Vestibular rehabilitation			
<i>Randomised controlled trials:</i>						
Karlberg et al. 1996 [26]; Malmström et al. 2007 [44]	n = 34 (88% females) Mean age: 37 Age range: 25-55 Country: Sweden Setting: primary care centers and a tertiary referral center	Mobilisation; Soft tissue therapy (relaxation techniques, stabilisation techniques); Home training program; Ergonomic changes at work	Nil	Dizziness frequency; Dizziness intensity; Posturography	- Significantly improved dizziness intensity and neck pain post-treatment ($p < 0.05$). - Significantly improved postural sway post-treatment ($p < 0.2$). - 14 patients (82%) reported improvements at 6 months post-treatment. [†] - 11 patients (65%) reported improvements in dizziness at 2 year post-treatment. [†]	Moderate
Reid et al. 2008 [47]	n = 34 (62% females), 1 drop-out Mean age: 63.5 Age range: not reported Country: Australia Setting: University of Newcastle	Mobilisation (SNAGs)	Nil	DHI; Dizziness severity (VAS); Dizziness frequency; Neck pain (VAS); Posturography	- Significantly reduced DHI, dizziness severity, dizziness frequency and neck pain in the treatment group at 6 and 12 weeks post-treatment ($p < 0.05$). - No difference in dizziness severity at 12 weeks post-treatment. - No difference in dizziness frequency at either 6 or 12 weeks post-treatment.	Good
Kang, Wang and Ye 2008 [48]	n = 76 (49% females) Mean age: 32.4 Age range: 18-45 Country: China Setting: hospital	<i>Group A:</i> Spinal manipulation <i>Group B:</i> Acupressure	Nil	TCD-US; TCM syndrome diagnostic criteria	- Significantly reduced VBA blood flow velocity post-treatment in both groups ($p < 0.01$). - Significantly larger reduction in left and right vertebral artery blood flow velocity in Group B compared with Group A ($p < 0.01$). - Group differences remained statistically significant at a 6-month follow-up.	Moderate
Fang 2010 [49]	n = 168 (73% females) Mean age: 37.5 Age range: not reported Country: China Setting: hospital	<i>Treatment group:</i> Spinal manipulation; Soft tissue therapy <i>Control group:</i> TCM medication	Nil	CVSFAS; Colour Doppler ultrasonography	- Significant improvements in dizziness ($p < 0.01$), shoulder/neck pain ($p < 0.05$), and headache ($p < 0.01$) post-treatment. - Significant reduction of cervical artery spasm index and atlantoaxial displacement index in the treatment group post-treatment ($p < 0.05$).	Moderate
Du et al. 2010 [50]	n = 70 (54% females) Mean age: 37.6 Age range: 21-45 Country: China Setting: hospital	<i>Treatment group:</i> Spinal manipulation; Soft tissue therapy <i>Control group:</i> Traction; Medication	Nil	CVSFAS; Radiography; TCD-US; TCM syndrome diagnostic criteria	- Significant improvements in dizziness scores, vertebral displacement post-treatment ($p < 0.01$). - Significantly reduced left and right vertebral artery blood flow velocity post-treatment ($p < 0.01$). - Significantly improved clinical outcomes six months post-treatment ($p < 0.01$).	Moderate
<i>Prospective cohort studies:</i>						

Table 1 Included studies (Continued)

Konrad and Gerencser 1990 [51]	n = 54 (74% females) Mean age: 34.7 Age range: not reported Country: Hungary Setting: hospital	Mobilisation; Manipulation	Nil	Dizziness (instrument not specifically stated); Electronystagmography	- 40 patients (74%) experienced improvement of dizziness post-treatment. [†]	Poor
Mahlstedt, Westhofen and König 1992 [52]	n = 28 (gender distribution not reported) Mean age: not reported Age range: not reported Country: Germany Setting: not reported	Spinal manipulation	Nil	No information provided	- 19 patients (68%) reported reduced dizziness post-treatment. [†]	Poor
Uhlemann et al. 1993 [53]	n = 12* (gender distribution not reported) Mean age: 40.7 Age range: not reported Country: Germany Setting: not reported	Mobilisation; Spinal manipulation (traction)	Nil	Cervical turn test	- 5 out of 9 patients went from testing positive to testing negative on the cervical turn test post-treatment.	Poor
Bracher et al. 2000 [54]	n = 15 (80% females), 3 drop-outs Mean age: 41 Age range: 27-82 Country: Brazil Setting: chiropractic clinic	Spinal manipulation; Soft tissue therapy; Electrotherapy; Labyrinth sedation medication; sEMG biofeedback; Exercise program	Nil	Dizziness (instrument not specifically stated; "improvement of symptoms was based on patient's reports")	- 9 patients (60%) reported complete remission of dizziness, 3 patients (20%) reported consistent improvement with rare recurrence of episodes of mild intensity, and 3 patients (20%) reported no change. [†]	Poor
Hülse and Hölzl 2000 [55]	n = 67 (52% females) Mean age: 49 Age range: 18-66 Country: Germany Setting: not reported	Soft tissue therapy (traction massage, PIR, occipital-base-release technique, atlas-impulse-therapy)	Nil	Craniocorpography; Posturography	- Significant improvements in pathological vestibulospinal reactions found post-treatment (p < 0.001).	Poor
Chen and Zhan 2003 [56]	n = 16 (38% females) Mean age: 42.4 Age range: 38-58 Country: China Setting: hospital	Spinal manipulation; Soft tissue therapy	Nil	TCD-US; Radiography; TCM syndrome diagnostic criteria	- 14 patients (87.5%) reported marked improvement or complete remission of symptoms. - Significantly decreased vertebral artery mean blood flow velocity post-treatment (p < 0.05). - Significantly reduced vertebral displacement post-treatment (p < 0.05).	Poor

Table 1 Included studies (Continued)

Wu et al. 2006 [45]; Wu et al. 2008 [46]	n = 121 (73% females) Mean age: not reported Age range: 20-71 Country: China Setting: hospital	Tuina manipulation therapy (pressing-kneading manipulation applied continuously to bilateral vertebrae for 5 minutes)	Nil	Custom-made instrument to measure skull 3D motion and head repositioning.	- Significant improvements in skull spatial offset repositioning ability post-manipulation ($p < 0.01$).	Poor
Strunk and Hawk 2009 [57]	n = 21 (63% females), 2 drop-outs Mean age: 70 Age range: 44-85 Country: USA (California) Setting: Cleveland Chiropractic College	Spinal manipulation; Soft tissue therapy (myofascial release, PIR, and heat or cold therapy)	Nil	DHI SF-BBS NDI	- Improved DHI and SF-BBS scores. [†] - Improved balance. [†] - Decreased dizziness and neck pain. [†]	Moderate

CVSFAS: cervical vertigo, symptoms and functional assessment scale; DHI: Dizziness Handicap Inventory; NDI: Neck Disability Index; HVLA: high-velocity, low amplitude; PIR: post-isometric relaxation; RCT: randomised, controlled trial; ROM: range of motion; SF-BBS: Berg Balance Scale (short form); sEMG: surface electromyography; SNAGs: sustained natural apophyseal glides; TCD: transcranial Doppler ultrasonography; TCM: traditional Chinese medicine; VAS: Visual Analogue Scale.

* Of the 42 patients that were recruited for this study only 12 patients were included in the manual therapy group, of which only 9 patients actually tested positive on the cervical turn test pre-treatment.

† No inferential statistics reported

Twelve studies, including all five RCTs, reported improvements in dizziness and associated symptoms (e.g. neck pain) following manual therapy intervention. The remaining study measured skull spatial offset repositioning ability, and found a significant improvement following soft tissue manipulation [75]. In addition to reduction in dizziness and associated symptoms, two RCTs [77,79] reported significant changes in vertebral artery blood flow velocity post-treatment, and a further two RCTs [26,76] found improvement in balance performance measured with posturography.

The methodological quality of the included studies was generally poor [75,80-85] to moderate [26,77-79,86]. However, one study [76] was found to be of good methodological quality. Not surprisingly, there was a trend towards more robust study designs (i.e. RCTs) and more recently published studies attaining higher quality scores. Overall, common methodological weaknesses included: lack of control group; failure to provide information allocation concealment and participant, provider, and assessor blinding; omitting performing appropriate statistical analysis; omitting reporting on patient compliance and drop-outs; and including long-term follow-up measurements. A tabulated overview of methodological quality scores is provided in Additional file 3.

Only three studies commented on adverse reactions. Two RCTs [26,76] reported no adverse reactions, and one prospective cohort study [86] found minor adverse

reactions associated with the interventions in eight of nineteen participants.

Discussion

In a previous review of the literature, Reid and Rivett [29] concluded there was limited (Level 3) evidence for manual therapy in the treatment of cervicogenic dizziness. The current systematic review has identified additional studies published since the previous review, including: four RCTs [76-79], three prospective cohort studies [75,85,86], and a long-term follow up [73] of the intervention group from the RCT published by Karlberg et al. [26].

The RCT by Reid et al. [76], which was deemed to be of good methodological quality, assessed the effectiveness of a specific type of spinal mobilisation known as sustained natural apophyseal glides (SNAGs). Reid et al. [76] found significant improvement in dizziness severity and frequency, lower scores on the Dizziness Handicap Inventory (DHI), and decreased neck pain in the treatment group at both six and twelve weeks post-treatment. In comparison the placebo group had significant changes only at the 12-week follow-up in three outcome measures (dizziness severity, DHI, and neck pain). The remaining four RCTs [26,77-79] were deemed to be of moderate methodological quality. The findings from the RCT by Karlberg et al. [26] (including the long-term follow-up by Malmstrom et al. [44]) appear to corroborate

the evidence provided by Reid et al. [76]. The RCTs by Kang, Wang and Ye [77], Fang [78], and Du et al. [79] all utilised spinal manipulation in the intervention group and reported improvements in clinical outcomes.

In addition to five RCTs the current systematic review identified eight prospective cohort studies, of which seven [80-86] reported improvements in dizziness following manual therapy. Although these were generally of poor methodological quality they also reported improvements in additional outcome measures, including: neck pain [86], reduction of pathological vestibulo-spinal activity [84], balance [86], and reduced vertebral displacement and vertebrobasilar artery blood flow velocity [85]. The remaining cohort study [75] reported improvements in skull spatial offset repositioning ability post treatment. Collectively, these findings provide further rationale for the use of manual therapy in the treatment of cervicogenic dizziness. Overall, the evidence evaluated in the current systematic review suggests that there is moderate (Level 2) evidence in a favourable direction to support the use of manual therapy for cervicogenic dizziness.

Although positive clinical outcomes have been demonstrated, the underlying biological mechanism remains a controversial subject. It has been theorised that disturbances to the afferent input from cervical spine mechanoreceptors may lead to a sensory mismatch between vestibular and cervical input subsequently resulting in symptoms such as dizziness, unsteadiness, and visual disturbances [25]. There is an experimental body of evidence indicating that the biomechanical forces of spinal manipulation and mobilisation impacts primary afferent neurons in paraspinal tissues, which in turn leads to physiological consequences such as gating of nociception at the spinal cord and spinal reflex activity to alter muscle activity [87,88]. Thus it is believed that manual therapy serves to normalise disturbances to the afferent input from deep neck proprioceptors and their subsequent reflex arcs (e.g. cervico-colic, cervico-ocular, and tonic neck), which in turn restores the ability to utilise internal vestibular orienting information to resolve inaccurate information from the somatosensory and visual subsystems (i.e. reducing sensory mismatch) [89].

Alas, no experimental or observational studies reporting the effect of combining manual therapy and vestibular rehabilitation in the management of cervicogenic dizziness could be identified. Collins and Misukanis [36] and Schenk et al. [90] have published case studies in which they argue that manual therapy combined with vestibular rehabilitation may be superior in the treatment of cervicogenic dizziness. Notwithstanding the paucity of such investigations, consideration of vestibular dysfunction is paramount in patients with dizziness. Unilateral peripheral vestibular dysfunction can be

characterised by complaints of dizziness, visual or gaze disturbances and balance impairment [34]. In a recent meta-analysis of vestibular rehabilitation for unilateral peripheral vestibular dysfunction it was concluded that vestibular rehabilitation is a safe and effective therapy [34].

The original vestibular rehabilitation protocols were developed by Cooksey [91] in 1946. These included: mental exercise, occupational therapy, physical exercise with the aim of restoring balance and joint position sense, and training of the eyes, to compensate for permanent vestibular dysfunction [91]. More recently, Hillier and Hollohan [34] stated vestibular rehabilitation may include: learning to coordinate eye and head movements, improving balance and walking skills, learning to bring on the symptoms to desensitize the vestibular system, patient education, coping strategies, and physical activity. There are four mechanisms of vestibular rehabilitation techniques that may contribute to its benefits, namely: (i) the compensatory response, (ii) adaptation, (iii) substitution, and (iv) postural control exercises. The compensatory responses are applied using motion to minimise the responsiveness to repetitive stimuli and to rebalance tonic activity within the vestibular nuclei. Adaptation for visual-vestibular interaction uses repetitive and provocative movements of the head and/or eyes to minimise error and restore vestibulo-ocular reflex gain. Substitution encourages the use of other sensory inputs to compensate for dysfunctional afferent systems. Postural control exercises and functional retraining are applied for movement behaviour and fitness.

The four mechanisms canvas a rationale for the inclusion of vestibular rehabilitation in the management of patients with cervicogenic dizziness. Stability and posture of the cervical spine is achieved by a combination of reflexes mediated by vestibular, visual and cervical sensory input [19]. The cerebellum plays an important role in integrating this sensory information [92]. It can be hypothesised that a well-integrated vestibulo-cerebellar system would be more capable of compensating for the altered cervical sensory input in cases of cervicogenic dizziness. Thus, one can argue that when normal cervical afferent input is compromised, vestibular rehabilitation may strengthen the vestibulo-cerebellar system to improve the ability to adapt to the situation. Further research to elucidate the effectiveness of manual therapy in conjunction with vestibular rehabilitation for cervicogenic dizziness is strongly recommended.

There are insufficient data to provide guidelines on dosage and frequency of manual therapy in general, and spinal manipulation in particular, especially in the context of management of cervicogenic dizziness. With this in mind, it is recommended that caution is taken when delivering any sensory stimulation in the form of

manual therapy or vestibular rehabilitation, or both, to affect dysfunctions in the afferent system in patients with cervicogenic dizziness. Further research is necessary to determine appropriate treatment dosage, scheduling of interventions, and which manual therapy and vestibular rehabilitation techniques are most effective in managing patients with cervicogenic dizziness.

Methodological limitations of this systematic review included lack of blinding during the quality assessment and the quality and utility of the quality assessment tool itself. Meta-analysis of the finding was precluded by the lack of robust research methodologies and heterogeneity of outcome measures in the studies included in this systematic review.

Conclusion

This systematic review has found that there is moderate (Level 2) evidence in a favourable direction to support the use of manual therapy (spinal mobilisation and/or manipulation) for cervicogenic dizziness. The evidence for combining manual therapy and vestibular rehabilitation in the management of cervicogenic dizziness remains inconclusive due to no observational and experimental studies investigating manual therapy in conjunction with vestibular rehabilitation. However, there is a reasonable rationale for utilising manual therapy in conjunction with vestibular rehabilitation for cervicogenic dizziness, and further research to elucidate the potential synergistic effects is strongly recommended.

Additional material

Additional file 1: Maastricht-Amsterdam criteria list. The Maastricht-Amsterdam criteria list is an instrument developed by van Tulder et al. [40] to assess methodological quality clinical trials. It consists of nineteen items that can be rated individually using one of three options: yes, no, or *don't know*. The overall methodological quality score is determined by adding up all of the 'yes' ratings, with a maximum score of nineteen.

Additional file 2: Excluded studies. Alphabetic list of excluded studies, including the reasons for exclusion.

Additional file 3: Methodological quality assessment scores of included studies. Methodological quality assessment scores of included studies.

Acknowledgements

We sincerely thank Lee-Lian Yeo who translated the Chinese language articles to English. We also thank Marius Monsveen for his contributions to the conception and design of the study in its early stages.

Authors' contributions

MBS, CVC and GB conceived of the study, participated in the design of the study, and helped to draft and edit the manuscript. RPL participated in the design and coordination of the study, helped to draft, edit and revise the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 25 April 2011 Accepted: 18 September 2011

Published: 18 September 2011

References

1. Sloane PD: Dizziness in primary care: Results from the National Ambulatory Medical Care Survey. *Journal of Family Practice* 1989, **29**(1):33-38.
2. Yardley L, Owen N, Nazareth I, Luxon L: Prevalence and presentation of dizziness in a general practice community sample of working age people. *British Journal of General Practice* 1998, **48**(429):1131-1135.
3. Johansson M, Andersson G: Prevalence of dizziness in relation to psychological factors and general health in older adults. *Audiological Medicine* 2006, **4**(3):144-150.
4. Stevens KN, Lang IA, Guralnik JM, Melzer D: Epidemiology of balance and dizziness in a national population: findings from the English Longitudinal Study of Ageing. *Age and Ageing* 2008, **37**(3):300-305.
5. Neuhauser HK: Epidemiology of dizziness and vertigo. *Der Nervenarzt* 2009, **80**(8):887-894.
6. Drachman D, Hart C: An approach to the dizzy patient. *Neurology* 1972, **22**(4):323-334.
7. Froehling D, Silverstein M, Mohr D, Beatty C: Does this dizzy patient have a serious form of vertigo. *Journal of American Medical Association* 1994, **271**(5):385-388.
8. Enloe LJ, Shields RK: Evaluation of health-related quality of life in individuals with vestibular disease using disease-specific and general outcome measures. *Physical Therapy* 1994, **77**(9):890-903.
9. Ardç FN, Topuz B, Kara CO: Impact of multiple etiology on dizziness handicap. *Otology and Neurology* 2006, **27**(5):676-680.
10. Wrisley DM, Sparto PJ, Whitney SL, Furman JM: Cervicogenic dizziness: a review of diagnosis and treatment. *Journal of Orthopaedic & Sports Physical Therapy* 2000, **30**(12):755-766.
11. Ryan G, Cope S: Cervical vertigo. *Lancet* 1955, **31**:1355-1358.
12. Lafon H: Les vertiges d'origine proprioceptive par lesion mecanique vertebro-cervicale. *Journal Francais d'Oto-Rhino-Laryngologie* 1990, **39**(5):269-278.
13. Fitz-Ritson D: Assessment of cervicogenic vertigo. *Journal of Manipulative and Physiological Therapeutics* 1991, **14**(3):193-198.
14. Tjell C, Rosenhall U: Smooth pursuit neck torsion test: A specific test for cervical dizziness. *American Journal of Otolaryngology* 1998, **19**(1):76-81.
15. Abrahams VC: The physiology of neck muscles; their role in head movement and maintenance of posture. *Canadian Journal of Physiology and Pharmacology* 1977, **55**(3):332-338.
16. Kulkarni V, Chandy MJ, Babu KS: Quantitative study of muscle spindles in suboccipital muscles of human foetuses. *Neurology India* 2001, **49**(4):355-359.
17. Boyd-Clark LC, Briggs CA, Galea MP: Muscle spindle distribution, morphology, and density in longus colli and multifidus muscles of the cervical spine. *Spine* 2002, **27**(7):694-701.
18. Liu J-X, Thornell L-E, Pedrosa-Domellöf F: Muscle spindles in the deep muscles of the human neck: A morphological and immunocytochemical study. *Journal of Histochemistry and Cytochemistry* 2003, **51**(2):175-186.
19. Treleaven J: Sensorimotor disturbances in neck disorders affecting postural stability, head and eye movement control. *Manual Therapy* 2008, **13**(1):2-11.
20. Karnath H-O: Subjective body orientation in neglect and the interactive contribution of neck muscle proprioception and vestibular stimulation. *Brain* 1994, **117**(5):1001-1012.
21. Cohen LA: Role of eye and neck proprioceptive mechanisms in body orientation and motor coordination. *Journal of Neurophysiology* 1961, **24**:1-11.
22. Biemond A, de Jong JMBV: On cervical nystagmus and related disorders. *Brain* 1969, **92**(2):437-458.
23. de Jong PTVM, de Jong JMBV, Cohen B, Jongkees LBW: Ataxia and nystagmus induced by injection of local anesthetics in the neck. *Annals of Neurology* 1977, **1**(3):240-246.
24. Wapner S, Werner H, Chandler KA: Experiments on the sensory-tonic field theory of perception: 1. Effect of extraneous stimulation of the visual perception of verticality. *Journal of Experimental Psychology* 1951, **42**(5):351-357.
25. Brandt T, Bronstein AM: Cervical vertigo. *Journal of Neurology Neurosurgery and Psychiatry* 2001, **71**(1):8-12.

26. Karlberg M, Magnusson M, Malmström E-M, Melander A, Moritz U: **Postural and symptomatic improvement after physiotherapy in patients with dizziness of suspected cervical origin.** *Archives of Physical Medicine and Rehabilitation* 1996, **77**(9):874-882.
27. Galm R, Rittmeister M, Schmitt E: **Vertigo in patients with cervical spine dysfunction.** *European Spine Journal* 1998, **7**(1):55-58.
28. Zhou W, Jiang W, Li X, Zhang Y, Wu Z: **Clinical study on manipulative treatment of derangement of the atlantoaxial joint.** *Journal of Traditional Chinese Medicine* 1999, **19**(4):273-278.
29. Reid SA, Rivett DA: **Manual therapy treatment of cervicogenic dizziness: A systematic review.** *Manual Therapy* 2005, **10**(1):4-13.
30. Sjölander P, Michaelson P, Jaric S, Djupsjöbacka M: **Sensorimotor disturbances in chronic neck pain-Range of motion, peak velocity, smoothness of movement, and repositioning acuity.** *Manual Therapy* 2008, **13**(2):122-131.
31. Denham T, Wolf A: **Vestibular rehabilitation.** *Rehabilitation Management* 1997, **10**(3):93-94, 144.
32. Hansson EE: **Vestibular rehabilitation - For whom and how? A systematic review.** *Advances in Physiotherapy* 2007, **9**(3):106-116.
33. Gans RE: **Vestibular rehabilitation: Critical decision analysis.** *Seminars in Hearing* 2002, **23**(2):149-159.
34. Hillier SL, McDonnell M: **Vestibular rehabilitation for unilateral peripheral vestibular dysfunction.** *Cochrane Database of Systematic Reviews* 2011, : 2: art. no. CD005397.
35. Hansson EE, Månsson N-O, Ringsberg KAM, Håkansson A: **Dizziness among patients with whiplash-associated disorder: A randomized controlled trial.** *Journal of Rehabilitation Medicine* 2006, **38**(6):387-390.
36. Collins ME, Misukanis TM: **Chiropractic management of a patient with post traumatic vertigo of complex origin.** *Journal of Chiropractic Medicine* 2005, **4**(1):32-38.
37. Schenk R, Coons LB, Bennett SE, Huijbregts PA: **Cervicogenic dizziness: A case report illustrating orthopaedic manual and vestibular physical therapy comanagement.** *The Journal of Manual & Manipulative Therapy* 2006, **14**(3):56-68.
38. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group: **Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement.** *Annals of Internal Medicine* 2009, **151**(4):264-269.
39. Denham T, McKinnon WA: **Vestibular Rehabilitation.** *Rehabilitation Management* 1997, **10**(14):93-94.
40. van Tulder MW, Assendelft WJ, Koes BW, Bouter LM: **Method guidelines for systematic reviews in the Cochrane Collaboration Back Review Group for spinal disorders.** *Spine* 1997, **22**(20):2323-2330.
41. Olmos M, Antelo M, Vazquez H, Smecuol E, Maurino E, Bai JC: **Systematic review and meta-analysis of observational studies on the prevalence of fractures in coeliac disease.** *Digestive and Liver Disease* 2008, **40**(1):46-53.
42. Lystad RP, Pollard H, Graham PL: **Epidemiology of injuries in competition taekwondo: A meta-analysis of observational studies.** *Journal of Science and Medicine in Sport* 2009, **12**(6):614-621.
43. Swain MS, Lystad RP, Pollard H, Bonello R: **Incidence and severity of neck injury in Rugby Union: A systematic review.** *Journal of Science and Medicine in Sport* 2011, **14**(5):383-389.
44. Becker VF: **Dizziness complaints from the viewpoint of manual therapy.** *Manuelle Medizin* 1978, **16**(5):95-104.
45. Biesinger E: **Diagnosis and therapy of vertebrogenic vertigo.** *Laryngologie, Rhinologie, Otologie* 1987, **66**(1):32-36.
46. Borg-Stein J, Rauch SD, Krabak B: **Evaluation and management of cervicogenic dizziness.** *Critical Reviews in Physical and Rehabilitation Medicine* 2001, **13**(4):255-264.
47. Bronfort G, Haas M, Evans R, Leininger B, Triano J: **Effectiveness of manual therapies: the UK evidence report.** *Chiropractic & Osteopathy* 2010, **18**:3.
48. Eber AM: **Reeducation of patients with vertigo.** *La Revue du Praticien* 1994, **44**(3):367-71.
49. El-Kahky AM, Kingma H, Dolmans M, de Jong I: **Balance control near the limit of stability in various sensory conditions in healthy subjects and patients suffering from vertigo or balance disorders: impact of sensory input on balance control.** *Acta Oto-Laryngologica* 2000, **120**(4):508-516.
50. Falkenau HA: **The pathogenesis and management of cervical vertigo.** *HNO* 1976, **24**(10):339-341.
51. Garcia FV: **Disequilibrium and its management in elderly patients.** *International Tinnitus Journal* 2009, **15**(1):83-90.
52. Grgic V: **Cervicogenic proprioceptive vertigo: ethiopathogenesis, clinical manifestations, diagnosis and therapy with special emphasis on manual therapy.** *Liječnicki Vjesnik* 2006, **128**(9-10):288-295.
53. Grod JP: **Effect of neck pain on verticality perception: a cohort study.** *Archives of Physical Medicine and Rehabilitation* 2002, **83**(3):412-415.
54. Hansson EE, Håkansson A: **Physical therapy of vertigo.** *Lakartidningen* 2009, **106**(35):2147-2149.
55. Hansson EE, Månsson N-O, Håkansson A: **Balance performance and self-perceived handicap among dizzy patients in primary health care.** *Scandinavian Journal of Primary Health Care* 2005, **23**(4):215-220.
56. Hawk C, Cambron J: **Chiropractic care for older adults: effects on balance, dizziness, and chronic pain.** *Journal of Manipulative and Physiological Therapeutics* 2009, **32**(6):431-437.
57. Hawk C, Khorsan R, Lisi AJ, Ferrance RJ, Evans MW: **Chiropractic care for nonmusculoskeletal conditions: a systematic review with implications for whole systems research.** *The Journal of Alternative and Complementary Medicine* 2007, **13**(5):491-512.
58. Heikkilä H, Johansson M, Wenngren B-I: **Effects of acupuncture, cervical manipulation and NSAID therapy on dizziness and impaired head repositioning of suspected cervical origin: a pilot study.** *Manual Therapy* 2000, **5**(3):151-157.
59. Jäger S: **Cervical vertigo in manual therapy.** *Zeitschrift für Physiotherapeuten* 2004, **56**(8):1398-1411.
60. Jepsen O: **Vertigo corrected from the cervical spine.** *Nordisk Medicin* 1963, **69**(23):675-676.
61. Karlberg M, Persson L, Magnusson M: **Impaired postural control in patients with cervico-brachial pain.** *Acta Oto-Laryngologica Supplementum* 1995, **520**:440-442.
62. Maffei G: **Vertigo in the pathology of the cervical spine.** *Acta Bio-Medica de l'Ateneo Parmense* 1983, **54**(Supplement 1):21-26, no. 1.
63. Persson L, Karlberg M, Magnusson M: **Effects of different treatments on postural performance in patients with cervical root compression: a randomized prospective study assessing the importance of the neck in postural control.** *Journal of Vestibular Research* 1996, **6**(6):439-453.
64. Rapaccini A, Pascucci W: **Rehabilitative therapy of tinnitus and vertigo.** *Recenti Progressi in Medicina* 2003, **94**(7-8):323.
65. Rohmer F, Collard M: **Vertigo of cervical origin.** *La Revue du Praticien* 1974, **24**(1):95-107.
66. Rong G: **General rehabilitation of the cervical syndrome in 500 cases.** *Chinese Journal of Clinical Rehabilitation* 2003, **7**(6):1036.
67. Scherer H: **Neck-induced vertigo.** *Archives of Oto-Rhino-Laryngology Supplement* 1985, **2**:107-124.
68. Seifert K: **Peripheral vestibular vertigo and functional disorders of the craniovertebral joint.** *HNO* 1987, **35**(9):363-371.
69. Takizawa H, Graille R, Dumolard P, Gignoux B: **Vertigo and the cervical column.** *Le Journal de Medecine de Lyon* 1967, **48**(131):1639-1654.
70. Teixeira LJ, Prado GF: **Impact of physical therapy in vertigo treatment.** *Revista Neurociencias* 2009, **17**(2):112-118.
71. Thomas D: **Dizziness in osteopathic practice.** *Osteopathische Medizin* 2009, **10**(2):29-31.
72. Yardley L, Beech S, Zander L, Evans T, Weinman J: **A randomized controlled trial of exercise therapy for dizziness and vertigo in primary care.** *British Journal of General Practice* 1998, **48**(429):1136-1140.
73. Malmström E-M, Karlberg M, Melander A, Magnusson M, Moritz U: **Cervicogenic dizziness - musculoskeletal findings before and after treatment and long-term outcome.** *Disability and Rehabilitation* 2007, **29**(15):1193-1205.
74. Wu J-R, Fang M, Hu J, Shen G-Q, Jiang S-Y: **Effects of manipulation on head repositioning skill in patients with cervical vertigo.** *Journal of Chinese Integrative Medicine* 2006, **4**(1):76-78.
75. Wu J, Fang M, Hu J, Shen G, Jiang S: **Action of tuina on retro-positioning of skull spatial offset in patients with cervical vertigo.** *Journal of Acupuncture and Tuina Science* 2008, **6**(2):83-86.
76. Reid SA, Rivett DA, Katekar MG, Callister R: **Sustained natural apophyseal glides (SNAGs) are an effective treatment for cervicogenic dizziness.** *Manual Therapy* 2008, **13**(4):357-366.
77. Kang F, Wang Q-C, Ye Y-G: **A randomized controlled trial of rotatory reduction manipulation and acupoint massage in the treatment of younger cervical vertigo.** *Chinese Journal of Orthopedics & Trauma* 2008, **21**(4):270-272.

78. Fang J: **Observation of curative effect on fixed-point spin reduction of spinal manipulation therapy for cervical vertigo.** *Chinese Journal of Orthopedics & Trauma* 2010, **23**(2):99-101.
79. Du H, Wei H, Huang M-Z, Jiang Z, Ye S-L, Song H-Q, Yu J-W, Ning X-T: **Randomized controlled trial on manipulation for the treatment of cervical vertigo of high flow velocity type.** *Chinese Journal of Orthopedics & Trauma* 2010, **23**(3):212-215.
80. Konrad K, Gerencser F: **Manual treatment in patients with vertigo.** *Manuelle Medizin* 1990, **28**(4):62-64.
81. Mahlstedt K, Westhofen M, König K: **Therapy of functional disorders of the craniovertebral joints in vestibular diseases.** *Laryngorhinootologie* 1992, **71**(5):246-250.
82. Uhlemann C, Gramowski K-H, Endres U, Callies R: **Manual diagnosis and therapy in cervical giddiness.** *Manuelle Medizin* 1993, **31**(4):77-81.
83. Bracher ESB, Almeida CIR, Almeida RR, Duprat AC, Bracher CBB: **A combined approach for the treatment of cervical vertigo.** *Journal of Manipulative and Physiological Therapeutics* 2000, **23**(2):96-100.
84. Hülse M, Hölzl M: **Vestibulospinal reflexes in patients with cervical disequilibrium. Cervicogenic imbalance.** *HNO* 2000, **48**(4):295-301.
85. Chen I, Zhan H-S: **A transcranial Doppler ultrasonography and X-ray study of cervical vertigo patients treated by manipulation in supine position.** *Journal of Chinese Integrative Medicine* 2003, **1**(4):262-264.
86. Strunk RG, Hawk C: **Effects of chiropractic care on dizziness, neck pain, and balance: a single-group, preexperimental, feasibility study.** *Journal of Chiropractic Medicine* 2009, **8**:156-164.
87. Pickar JG: **Neurophysiological effects of spinal manipulation.** *Spine Journal* 2002, **2**(5):357-371.
88. Schmid A, Brunner F, Wright A, Bachmann LM: **Paradigm shift in manual therapy? Evidence for a central nervous system component in the response to passive cervical joint mobilisation.** *Manual Therapy* 2008, **13**(5):387-396.
89. Kristjansson E, Treleaven J: **Sensorimotor function and dizziness in neck pain: implications for assessment and management.** *Journal of Orthopaedic & Sports Physical Therapy* 2009, **39**(5):364-377.
90. Schenk R, Coons LB, Bennett SE, Huijbregts PA: **Cervicogenic dizziness: A case report illustrating manual and vestibular physical therapy comanagement.** *The Journal of Manual & Manipulative Therapy* 2006, **14**(3): E56-E68.
91. Cooksey FS: **Rehabilitation in vestibular injuries.** *Proceedings of the Royal Society of Medicine* 1946, **39**(5):273-278.
92. Manzoni D: **The cerebellum may implement the appropriate coupling of sensory inputs and motor responses: Evidence from vestibular physiology.** *The Cerebellum* 2005, **4**:178-188.

doi:10.1186/2045-709X-19-21

Cite this article as: Lystad et al.: Manual therapy with and without vestibular rehabilitation for cervicogenic dizziness: a systematic review. *Chiropractic & Manual Therapies* 2011 **19**:21.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

