

ORIGINAL ARTICLE

Detecting population structure in a high gene-flow species, Atlantic herring (*Clupea harengus*): direct, simultaneous evaluation of neutral vs putatively selected lociC André¹, LC Larsson^{2,9}, L Laikre², D Bekkevold³, J Brigham⁴, GR Carvalho⁵, TG Dahlgren^{1,10}, WF Hutchinson⁴, S Mariani⁶, K Mudde⁷, DE Ruzzante⁸ and N Ryman²

¹Department of Marine Ecology-Tjärnö, University of Gothenburg, Strömstad, Sweden; ²Division of Population Genetics, Department of Zoology, Stockholm University, Stockholm, Sweden; ³National Institute of Aquatic Resources, Technical University of Denmark, Silkeborg, Denmark; ⁴Department of Biological Sciences, University of Hull, Humberside, UK; ⁵School of Biological Sciences, Bangor University, Environment Centre Wales, Gwynedd, UK; ⁶Marine Biodiversity Ecology & Evolution, UCD School of Biology & Environmental Science, University College Dublin, Dublin, Ireland; ⁷Wageningen Institute of Animal Sciences, Wageningen University, Wageningen, The Netherlands and ⁸Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada

In many marine fish species, genetic population structure is typically weak because populations are large, evolutionarily young and have a high potential for gene flow. We tested whether genetic markers influenced by natural selection are more efficient than the presumed neutral genetic markers to detect population structure in Atlantic herring (*Clupea harengus*), a migratory pelagic species with large effective population sizes. We compared the spatial and temporal patterns of divergence and statistical power of three traditional genetic marker types, microsatellites, allozymes and mitochondrial DNA, with one microsatellite locus, *Cpa112*, previously shown to be influenced by divergent selection associated with salinity, and one locus located in the major histocompatibility complex class IIA (*MHC-IIA*) gene, using the same individuals across analyses. Samples were collected in 2002 and 2003 at two locations in the North Sea, one location in the Skagerrak

and one location in the low-saline Baltic Sea. Levels of divergence for putatively neutral markers were generally low, with the exception of single outlier locus/sample combinations; microsatellites were the most statistically powerful markers under neutral expectations. We found no evidence of selection acting on the *MHC* locus. *Cpa112*, however, was highly divergent in the Baltic samples. Simulations addressing the statistical power for detecting population divergence showed that when using *Cpa112* alone, compared with using eight presumed neutral microsatellite loci, sample sizes could be reduced by up to a tenth while still retaining high statistical power. Our results show that the loci influenced by selection can serve as powerful markers for detecting population structure in high gene-flow marine fish species.

Heredity (2011) **106**, 270–280; doi:10.1038/hdy.2010.71; published online 16 June 2010

Keywords: genetic markers; hitchhiking selection; statistical power; pelagic fish; MHC; Baltic

Introduction

Population genetic principles and methods are increasingly used in the management of harvested species and the conservation of natural populations (Allendorf and Luikart, 2007; Allendorf *et al.*, 2008). Population subdivision results from genome-wide processes, such as reduced migration and genetic drift, that on an average affect all genes in the same way, as well as locus-specific processes such as recombination, selection and mutation that are limited to specific genomic regions. Population

structure is typically assessed by estimating divergence in the allele frequencies at neutral marker loci, whereas coding gene regions are usually not used for testing demographic independence or for estimating migration rates because selection biases parameter estimation (for example, Whitlock and McCauley, 1999). However, in some cases, such as mixed stock analysis in harvested species, gene loci influenced by selection can be valuable as population markers on ecological time scales where isolated populations have not yet diverged at neutral loci (O'Malley *et al.*, 2007; Westgaard and Fevolden, 2007; Hauser and Carvalho, 2008; Nielsen *et al.*, 2007, 2009a). Moreover, coding loci will help in identifying locally adapted populations (Hemmer-Hansen *et al.*, 2007; Andersen *et al.*, 2009; Gebremedhin *et al.*, 2009; Nielsen *et al.*, 2009a; Arnason *et al.*, 2009).

In marine fishes, population differentiation through drift is typically weak because populations have relatively shallow histories and are often very large (Hauser and Carvalho, 2008). Although some marine fish

Correspondence: Dr C André, Department of Marine Ecology-Tjärnö, University of Gothenburg, Strömstad SE-452 96, Sweden.

E-mail: carl.andre@marecol.gu.se

⁹Current address: GM Sutton Avian Research Center, University of Oklahoma, PO Box 2007, Bartlesville, OK 74005, USA

¹⁰Current address: Uni Research, Thormøhlensgt. 49, Bergen 5006, Norway

Received 5 November 2009; revised 14 April 2010; accepted 5 May 2010; published online 16 June 2010

species show population structure even at small spatial scales (Bekkevold *et al.*, 2005; Jorde *et al.*, 2007; Galarza *et al.*, 2009), a high number of neutral loci and large sample sizes are needed to resolve structure. Genetic differentiation at coding loci requires heterogeneous selection patterns and limited migration, and is expected to be more pronounced in large populations, in which even weak selection may override genetic drift. Temporal stability of divergence, whether selected or neutral, increases the likelihood that the detected structuring is biologically significant, and thus that the loci are useful as population markers (Waples, 1998; Nielsen *et al.*, 2007).

In this study, we compare the performance of three putatively neutral genetic marker types (microsatellites, allozymes and mitochondrial DNA (mtDNA)) with two genetic markers allegedly influenced by selection (the major histocompatibility complex (MHC)-linked locus *Clha-DAA-INTR3* and a microsatellite locus *Cpa112*) to detect structuring in Atlantic herring (*Clupea harengus*) populations within the North Sea/Baltic Sea region. Although there are several cases in which each of these three classes of neutral markers have shown to be under the influence of selection (Karl and Avise, 1992; Grant *et al.*, 2006; Nielsen *et al.*, 2006), the expectation is, in the absence of evidence to the contrary, that their evolutionary characteristics follow neutral expectations (Baer, 1999). It is, however, important to apply appropriate tests of neutrality beyond traditional measures of population structure (Rand, 1996).

Atlantic herring has traditionally been divided into different geographical 'stocks' based on spawning behavior and meristic or morphological characters. However, there has been little concordance between these characters and genetic differentiation (Ryman *et al.*, 1984; but see Clausen *et al.*, 2007 and Jørgensen *et al.*, 2008). Recent genetic investigations using microsatellite DNA and allozyme markers have found little differentiation among spatially discrete populations in the North Sea, but concordant divergence between the Baltic and North Sea populations (Bekkevold *et al.*, 2005; Mariani *et al.*, 2005; Larsson *et al.*, 2007, 2010). Genetic mixed-stock analyses show that coastal Skagerrak is used as a nursery both for the Baltic and North Sea spawning populations (Ruzzante *et al.*, 2006). Earlier studies of population structure in the NE Atlantic using mtDNA have generally indicated little divergence among the populations (Dahle and Eriksen, 1990).

Using two different methods, Larsson *et al.* (2007) and Gaggiotti *et al.* (2009) showed that a microsatellite locus, *Cpa112*, displayed highly divergent allele frequencies among populations, suggesting that this locus may be affected by hitchhiking selection. The identity and function of associated gene(s) is unknown, but the divergence pattern was stable over a 25-year time span (Larsson *et al.*, 2010).

MHC genes encode proteins in the vertebrate immune response, and molecular variation among alleles has been linked directly to fitness and survival (Bos *et al.*, 2008). For example, disease challenge trials on Atlantic salmon confirm that polymorphism at the MHC genes are associated with increased disease resistance (Grimholt *et al.*, 2003). Wild populations show evidence for local adaptation (de Eyto *et al.*, 2007), and several studies indicate that MHC genes, as well as a number of

linked loci, can be more powerful than neutral microsatellites in detecting population differentiation in spatially structured fish species such as salmonids (Bernatchez and Landry, 2003; Beacham *et al.*, 2005; Hansen *et al.*, 2007). However, with the exception of a precursory characterization of the MHC class IIA gene and an embedded microsatellite locus in Atlantic herring (Stet *et al.*, 2008), no studies exist to date that describe the MHC variation in an abundant and fully marine fish. It is worth noting, however, that divergence in genes under selection is not always expected; if fish populations share similar parasite environments, balancing selection can result in lower levels of differentiation in MHC-linked microsatellites than neutral microsatellites (Muirhead, 2001; Bernatchez and Landry, 2003).

Different genetic markers reflect different demographic histories (Avise, 2004) and also have varying statistical properties that affect the extent to which they can resolve the patterns of divergence (Ryman *et al.*, 2006). For example, it has been shown that organelle markers such as mtDNA may be more powerful for detecting differentiation in recently diverged populations and especially when migration among populations is relatively high (Larsson *et al.*, 2009).

In this study, we present an investigation of spatial and temporal patterns of divergence and statistical power obtained using three traditional genetic markers, allozymes, mtDNA and microsatellites, and two markers potentially influenced by selection, MHC and the microsatellite locus *Cpa112*. Using the same individuals for all analyses, we compare fish collected in 2002, and again in 2003, from four spawning grounds spanning the environmental gradient from the Baltic Sea to the North Sea.

Materials and methods

Sampling

A total of eight samples of Atlantic herring were collected in 2002 and 2003 at four locations in the North Sea/Baltic Sea region: autumn-spawning herring from (1) Flamborough and (2) Berwick in the North Sea, (3) a spring-spawning population from Tjøme in the Skagerrak and (4) a second spring-spawning population from Kalix in the inner Baltic Sea (Figure 1; Table 1). These samples cover a salinity gradient from the fully marine environment in the North Sea (35‰) to almost fresh water in the Bothnian Bay of the Baltic Sea (3‰). Each sample consisted of ca 100 fish in full spawning condition. Otolith (sagitta) winter rings were counted as a proxy for age following standard procedures (ICES, 2003). The hatching season of individual fish (spring, autumn or winter) was estimated from otolith central area microstructure according to Clausen *et al.* (2007).

Genetic analyses

The fish were genotyped using four genetic marker types: microsatellites, allozymes, mtDNA and MHC. One of the microsatellite loci, *Cpa112*, has previously been demonstrated to be influenced by selection (Gaggiotti *et al.*, 2009; Larsson *et al.*, 2007, 2010). Parts of the microsatellite, allozyme and MHC data have been presented previously (see Table 1), although here we provide full data sets to allow a direct comparison.




Figure 1 Sampling locations for Atlantic herring collected in 2002 and 2003. Sea surface salinities are indicated.

Table 1 Sample descriptions including locations, coordinates, dates, spawning seasons, percentage of fish in spawning condition and sample sizes (*n*) of genetic markers used

Location	Sampling coordinates	Date	Spawning season	Spawning (%)	Fish (n)	Microsatellites (including <i>Cpa112</i>) (n)	MHC (n)	Allozymes (n)	mtDNA (n)	All markers in the same fish (n)
Flamborough	54.60N/00.59W	14-09-2002	Autumn	100	80	80 ^{a,b}	73	55	80	48
Flamborough	54.57N/00.45W	17-09-2003	Autumn	99	93	93 ^{a,b,c,d}	80	93 ^c	91	79
Berwick	56.30N/00.97E	20-08-2002	Autumn	98	90	90 ^{a,b}	83 ^e	85	85	73
Berwick	57.70N/01.45W	22-08-2003	Autumn	88	96	96 ^{a,b,c}	90	96 ^c	80	72
Tjøme	59.35N/10.55E	13-03-2002	Spring	98	95	95 ^{b,f}	95 ^e	95	84	84
Tjøme	59.35N/10.55E	04-03-2003	Spring	100	104	104 ^{b,c,d,f}	94	104 ^c	86	75
Kalix	65.80N/22.72E	12-06-2002	Spring	44	97	97 ^g	90	97 ^g	85	78
Kalix	65.80N/22.72E	19-06-2003	Spring	94	88	88 ^{c,g}	79	88 ^{c,g}	69	64
				Sum	743	743	684	713	660	573

Abbreviations: MHC, major histocompatibility complex; mtDNA, mitochondrial DNA.

Only fish that were successfully analyzed for at least three different genetic marker types were used in subsequent statistical analyses. Genotypes previously used by ^aMariani *et al.* (2005), ^bRuzzante *et al.* (2006), ^cLarsson *et al.* (2007), ^dGaggiotti *et al.* (2009), ^eStet *et al.* (2008), ^fBekkevold *et al.* (2005) and ^gLarsson *et al.* (2010).

DNA for microsatellite, mitochondrial and MHC analyses was extracted from fin clip or muscle tissue using standard kits or the HotSHOT method (Truett *et al.*, 2000). Nine microsatellite loci, *Cha1017*, *Cha1020*, *Cha1027*, *Cha1202* (McPherson *et al.*, 2001), *Cpa101*, *Cpa111*, *Cpa113*, *Cpa114* and *Cpa112* (Olsen *et al.*, 2002), were amplified and genotyped using a BaseStation (MJ Research, MJ Research, Skovlunde, Denmark, Denmark) and Pharmacia ALF express (Amersham Pharmacia, Uppsala, Sweden) automated sequencers. Scoring consistency among runs and platforms was ensured by analyzing two heterozygote control individuals spanning the anticipated allelic ranges on all gels, in addition to internal and external size ladders.

MtDNA variation was examined by analysis of restriction fragment length polymorphism in the

mitochondrial NADH dehydrogenase ND3/4 subunit. A 2.4 Kb sequence was PCR amplified using an existing universal primer (Hauser *et al.*, 2001) and a new primer designed from a sardine (*Sardinops melanostictus*) mtDNA sequence (Inoue *et al.*, 2000) published on GenBank (accession number NC 002616):

Universal vertebrate primer: 5'-TTTTGGTTCCTAAGACCAA(C/T)GGAT-3'

Newly designed primer: 5'-AAGACAGTACAGGTGGCTTCAA-3'.

The PCR products were digested using six restriction enzymes: *AluI*, *Hae III*, *Hinf I*, *Mbo I*, *Msp I* and *Rsa I*, which produced 26 cut sites in Atlantic herring. Restricted fragments were separated on 6% polyacrylamide gels, together with a pGEM size marker (Promega, Southampton, UK), and stained using a standard silver

nitrate protocol. Fragments were sized using NEBCUTTER (<http://tools.neb.com/NEBcutter2/index.php>), and fragment patterns were expected to add up to a total of ~2.4Kb if the PCR product had been fully digested; partial digests were discarded.

MHC polymorphism was investigated using the linked microsatellite *Clha-DAA-INTR3*, located in the third intron of the MHC class II alpha gene *Clha-DAA* (Stet *et al.*, 2008). PCR amplification of *Clha-DAA* was performed using standard cycling conditions with annealing at 50 °C. PCR products were analyzed with an ABI 377 automated sequencer and fragment sizes assessed by the GENOTYPER software (Applied Biosystems, Foster City, CA, USA).

Allozyme starch gel electrophoresis was performed using procedures described in Andersson *et al.* (1981). The following eleven polymorphic loci were scored, (locus abbreviations used in early studies, enzyme names and EC numbers are given in parentheses): *mAAT** (AAT-2, aspartate aminotransferase, EC 2.6.1.1), *G3PDH** (*GPD-1*, glycerol-3-phosphate dehydrogenase, EC 1.1.1.8), *GPI** (glucosephosphate isomerase, EC 5.3.1.9), *mIDHP** (*IDH-2*, isocitrate dehydrogenase, EC 1.1.1.42), *LDH-1**, *LDH-2** (lactate dehydrogenase, EC 1.1.1.27), *MDH-4** (malate dehydrogenase, EC 1.1.1.37), *sMEP** (*ME-1*, malic enzyme, EC 1.1.1.40), *mMEP** (*ME-2*, malic enzyme, EC 1.1.1.40), *PGM-1** (phosphoglucumutase, EC 5.4.2.2, formerly EC 2.7.5.1) and *SOD** (superoxide dismutase, EC 1.15.1.1). These loci have previously been shown to be variable in herring from this region (Andersson *et al.*, 1981; Ryman *et al.*, 1984; Larsson *et al.*, 2007).

Statistical analyses

It was not possible to genotype all fish for all marker loci, and the original data set with ca 100 fish from each sample was cropped to include individual fish that were successfully analyzed for at least three of the four genetic marker types (microsatellites incl. *Cpa112*, the MHC locus *Clha-DAA-INTR3*, allozymes and mtDNA; Table 1). In total, 743 fish were analyzed, out of which 573 were genotyped for all marker types.

Amounts of genetic variation within herring samples were estimated as allelic richness (*AR*) and expected heterozygosity (gene diversity, *H_E*). *F*-statistics were calculated using FSTAT (Goudet, 2001). Statistical significance of *F_{ST}* for allozymes, MHC and microsatellites were examined using Fisher's exact test implemented in GENEPOP 3.4 (Raymond and Rousset, 1995), the χ^2 -test implemented in CHIFISH 1.3 (Ryman, 2006) and permutation tests using FSTAT (Goudet, 2001); the results were very similar and only *P*-values obtained using FSTAT (with 10 000 permutations and not assuming Hardy–Weinberg equilibrium) are presented here for brevity. Departure from Hardy–Weinberg equilibrium was tested using 10 000 permutations in FSTAT. We used GDA version 1.1 (Lewis and Zaykin, 2001) to examine linkage disequilibria. ARLEQUIN 3.1 (Excoffier *et al.*, 2005) was used to estimate gene diversity and pairwise *F_{ST}* between samples for the mtDNA haplotype data; statistical significance was obtained from 10 000 permutations. The presence of outlier loci was investigated with the test of Beaumont and Nichols (1996), in which the null distribution was created using FDIST2 (Beaumont, 2002).

STATISTICA 6.0 (StatSoft Inc., Uppsala, Sweden) was used to visualize genetic relationships among samples for the different marker types with multidimensional scaling using pairwise *F_{ST}* estimates. Concordance in *F_{ST}* estimates between markers was investigated using correlation of *F_{ST}* and Mantel tests between all sample pairs.

Temporal stability in differentiation for the different marker types was estimated using hierarchical analysis of molecular variance, with sampling years nested within locations. We used the locus-by-locus option in ARLEQUIN 3.1; statistical significance was obtained from 10 000 permutations. Further, we divided all fish into year classes, based on otolith ageing, and tested for genetic heterogeneity among cohorts with $n \geq 20$, within and among locations.

The statistical power for detecting genetic differentiation at various levels of *F_{ST}* was evaluated for the separate markers using the software POWSIM (Ryman and Palm, 2006; Ryman *et al.*, 2006). This analysis simulates sampling from a specified number of populations that have diverged to predefined levels of divergence. We used sample sizes, number of loci and allele frequencies obtained from the eight samples in this study. Finally, we used a modified version of POWSIM to obtain sampling distributions of *F_{ST}* for the different markers; using the observed allele frequencies over all samples, we simulated a drift process in which two large populations diverge until an expected *F_{ST}* of 0.002 is reached (see Table 2). We then sampled these simulated populations at $n=100$ and calculated *F_{ST}*. This procedure was repeated 10 000 times.

In all statistical analyses that involve multiple tests, we provide both uncorrected and sequential Bonferroni-corrected significance levels for comparison.

Results

Sampling

The hatching date determined from otolith microstructure verified that the collected samples represented the presumed seasonal spawning stocks: autumn spawners in the North Sea and spring spawners in the Skagerrak and the Baltic. Between 48 and 84 individuals from each sample were genotyped for all five genetic markers: the three putatively neutral genetic markers (8 microsatellite loci, 11 allozyme loci, 1 mtDNA locus) and the two genetic markers allegedly influenced by selection (the microsatellite *Cpa112* and the MHC-embedded *Clha-DAA-INTR3*).

Within-population descriptive analyses

Allelic richness and expected heterozygosity, *H_E*, for the MHC-embedded microsatellite locus ranged between 8.5–11.6 and 0.70–0.80 among samples, respectively (Supplementary Appendix 1a). These values are similar to those obtained for the other microsatellite loci, including *Cpa112*, in the same populations. For allozymes the average number of alleles per locus and *H_E* were considerably lower, ranging between 2.1–2.7 and 0.12–0.14 among samples (Supplementary Appendix 1b). In total, we found 237 mtDNA haplotypes among the 660 analyzed fish. Accordingly, gene (haplotype) diversity within samples was high, ranging between 0.93 and 0.98 among samples (Supplementary Appendix 1a).

Table 2 F -statistics (F_{IS} , F_{ST} , F_{IT}) for allozyme, microsatellite, MHC and mtDNA loci for unpooled ($s = 8$) and pooled ($s = 4$) temporal herring samples

Locus	$s = 8$			$s = 4$, temporal samples pooled		
	F_{IS}	F_{ST}	F_{IT}	F_{IS}	F_{ST}	F_{IT}
<i>Allozymes</i>						
<i>mAAT*</i>	-0.0115	0.0020	-0.0095	-0.0113	0.0020	-0.0092
<i>G3PDH*</i>	-0.0032	-0.0008	-0.0040	-0.0041	0.0003	-0.0038
<i>GPI*</i>	0.0100	-0.0006	0.0095	0.0104	-0.0011	0.0092
<i>mIDHP</i>	-0.0155	0.0011	-0.0143	-0.0164	0.0025	-0.0139
<i>LDH-1*</i>	0.0092	0.0031*	0.0123	0.0125	-0.0007	0.0118
<i>LDH-2*</i>	-0.0317	0.0029*	-0.0287	-0.0322	0.0041*	-0.0280
<i>MDH-4*</i>	0.0422	0.0016	0.0438	0.0440	-0.0007	0.0434
<i>sMEP*</i>	0.0000	-0.0003	-0.0003	-0.0007	0.0008	0.0001
<i>mMEP*</i>	-0.0072	-0.0026	-0.0098	-0.0084	-0.0014	-0.0098
<i>PGM-1*</i>	-0.0101	0.0020	-0.0080	-0.0123	0.0052	-0.0069
<i>SOD*</i>	0.0754	0.0162**	0.0904**	0.0781	0.0150**	0.0919**
Allozymes 11 loci	0.0184	0.0011*	0.0195	0.0192	0.0002**	0.0194
<i>Microsatellites</i>						
<i>Cha1017</i>	0.0679***	-0.0012	0.0668***	0.0669***	0.0001	0.0669***
<i>Cha1020</i>	0.0020	0.0056***	0.0075	0.0026	0.0057***	0.0083
<i>Cha1027</i>	0.0364**	0.0004*	0.0368***	0.0365***	0.0002*	0.0368***
<i>Cha1202</i>	0.0355*	0.0051***	0.0405**	0.0385*	0.0019***	0.0403**
<i>Cpa101</i>	0.0109	0.0008	0.0117	0.0109	0.0010	0.0118
<i>Cpa111</i>	-0.0008	0.0062***	0.0054	-0.0007	0.0071***	0.0064
<i>Cpa113</i>	0.0172	0.0025**	0.0197*	0.0169	0.0034***	0.0202*
<i>Cpa114</i>	0.0334**	0.0017*	0.0350**	0.0348**	0.0000	0.0348**
Microsatellites 8 loci	0.0268***	0.0024***	0.0291***	0.0273***	0.0021***	0.0294***
<i>Cpa112</i>	0.0132	0.0360***	0.0487***	0.0135	0.0415***	0.0544***
MHC, <i>Clha-DAA-INTR3</i>	0.0020	0.0027	0.0047	0.0012	0.0043	0.0055
MtDNA	—	0.0145***	—	—	0.0065***	—
All loci, excluding mtDNA	0.0225***	0.0052***	0.0276***	0.0230***	0.0055***	0.0283***
All, excluding mtDNA, <i>Cpa112</i>	0.0234***	0.0022***	0.0256***	0.0238***	0.0020***	0.0258***
All, excluding mtDNA <i>Cpa112</i> , <i>SOD</i>	0.0233***	0.0021***	0.0254***	0.0237***	0.0019***	0.0256***

Abbreviations: MHC, major histocompatibility complex; mtDNA, mitochondrial DNA.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Sample-specific heterozygosities and F_{IS} estimates for microsatellite loci and MHC are presented in Supplementary Appendix 1.

Deficiency of heterozygotes relative to Hardy–Weinberg expectations was found in one locus-sample combination for the allozymes, and in 12 out of 64 for the presumed neutral microsatellites; there was no sample- or locus-specific pattern, and only one persisted after Bonferroni correction. One sample showed heterozygote deficiency at *Cpa112*, whereas no deviations were found at the MHC locus (Supplementary Appendix 1a). Combining the information from all localities and/or samples show that microsatellites displayed significantly high overall F_{IS} values at four loci (Table 2); however, separate statistically significant locus-sample combinations appeared evenly distributed throughout the locus-sample table (Supplementary Appendix 1a). There was little indication of linkage disequilibria between loci. Out of a total of 1848 pairwise locus combinations (22 loci) within samples, 57 showed significant deviation from linkage equilibrium (ca 3% of the tests). Only one significance remained after Bonferroni correction.

Outlier test

The microsatellite locus *Cpa112* was identified as an outlier (Figure 2), indicating that this locus is influenced by directional selection. While within the 95% confidence limits of the simulated F_{ST} distribution, the allozyme locus *SOD** had a conspicuously large F_{ST} value compared with the other allozyme loci (Table 2; Figure 2).

There was no indication of selection affecting the MHC locus.

Statistical power and expected F_{ST}

Before comparing statistical power, we investigated the expected distribution of F_{ST} for all markers in this case under the assumption of selective neutrality. Two simulated populations differentiated at $F_{ST} = 0.002$ were sampled repeatedly at $n = 100$. The various markers had similar mean F_{ST} values, close to 0.002, but the medians and variances differed (Figure 3). The F_{ST} distributions were markedly skewed to the right, except for the eight microsatellites, in which the distribution was more symmetric and had a smaller variance (Figure 3). This pattern is also seen in our empirical data in which the average and variance were 0.0022 and 24.5×10^{-6} for the 11 allozyme loci and 0.0026 and 7.4×10^{-6} for the eight microsatellite loci (see Table 2; also see Larsson *et al.*, 2007).

Analysis of statistical power indicated that all marker sets will detect $F_{ST} = 0.0050$ or larger with a probability close to one, for present sample sizes (Table 3). At lower levels of divergence, eight microsatellites are more powerful for detecting structuring than eleven allozymes, mtDNA, MHC and *Cpa112*. The individual microsatellite loci have, however, statistical power comparable to the latter markers (Table 3).




Figure 2 F_{ST} for individual microsatellite, allozyme and MHC loci plotted against heterozygosity. Lines denote the 2.5, 50 and 97.5 percentiles of simulation-based expected distributions of F_{ST} assuming an infinite allele or stepwise mutation model, respectively. The simulations were based on the combined overall $F_{ST}=0.002$ for all three marker types.




Figure 3 Distributions of F_{ST} for various marker types based on 10,000 simulated samples of 100 individuals from two populations with a true F_{ST} of 0.002. The sample parameters mean, median and variance ($\times 10^{-6}$) are for microsatellites (eight loci): 0.0020, 0.0019, 1.4; allozymes: 0.0019, 0.0009, 19; mtDNA: 0.0020, 0.0014, 9.3; MHC: 0.0021, 0.0008, 25; *Cpa112*: 0.0020, 0.0014, 10.

The locus *Cpa112* showed strong divergence and had an F_{ST} more than ten times higher than the other loci (Table 2). This raises the question: How much smaller could sample sizes be using *Cpa112* and still have the same power as several neutral loci to detect divergence? Using POWSIM, we estimated power at *Cpa112* for different sample sizes, and we show that a sample size reduced to $n \approx 20$ would be sufficient to detect $F_{ST}=0.025$ (Table 4).

Population differentiation

The patterns of differentiation among the herring populations using the different marker types were

Table 3 Statistical power of different marker types for detecting various true levels of population differentiation (F_{ST}) by means of Fisher's exact test when using present sample sizes and allele frequencies obtained from the eight samples in this study

F_{ST}	Micro-satellites 8 loci	Allozymes 11 loci	mtDNA	MHC	<i>Cpa112</i>	Median of individual microsatellite loci
0.0000	0.063	0.057	0.063	0.064	0.049	0.054
0.0001	0.125	0.056	0.072	0.068	0.070	0.065
0.0010	0.958	0.342	0.455	0.240	0.386	0.366
0.0020	1.000	0.774	0.866	0.572	0.799	0.770
0.0050	1.000	0.998	1.000	0.982	0.998	1.000
0.0100	1.000	1.000	1.000	1.000	1.000	1.000

Abbreviations: MHC, major histocompatibility complex; mtDNA, mitochondrial DNA.

Table 4 Statistical power of *Cpa112* for detecting various true levels of population differentiation (F_{ST}) by means of Fisher's exact test when using allele frequencies obtained from the eight samples in this study and different sample sizes (n)

F_{ST}	$n=100$	$n=50$	$n=30$	$n=20$	$n=10$
0.000	0.056	0.049	0.049	0.057	0.051
0.001	0.512	0.216	0.145	0.088	0.064
0.002	0.898	0.462	0.220	0.163	0.083
0.005	1.000	0.956	0.669	0.397	0.171
0.010	1.000	1.000	0.971	0.778	0.325
0.025	1.000	1.000	1.000	1.000	0.844
0.050	1.000	1.000	1.000	1.000	0.996

explored in three different ways: using individual samples ($s=8$), pooling temporal samples from the same locality ($s=4$), and using year-class cohorts at sampling locations as the basic unit ($s=10$; Supplementary




Figure 4 Multidimensional scaling plots of herring samples based on pairwise F_{ST} estimated from different genetic markers.

Appendix 3). Overall F_{ST} ($s = 8$) was 0.0024 for microsatellites, 0.0011 for allozymes, 0.0145 for mtDNA, 0.0027 for MHC and 0.0360 for *Cpa112* (Table 2). F_{ST} over all nuclear marker types (that is, excluding mtDNA) was 0.0052. Pooling the temporal samples within locations gave similar estimate of F_{ST} for microsatellites, whereas the estimates decreased for mtDNA and allozymes, and increased for MHC and *Cpa112* (Table 2).

Spatial and temporal patterns

The microsatellite markers showed significant differentiation between the three regions, the North Sea, the Skagerrak and the Baltic (Figure 4, Table 2, Supplementary Appendix 2). At both Kalix and Tjøme, samples were genetically similar between years. In contrast, the four North Sea samples clustered by year of sampling. The cohort analysis indicates that this could be a year-class effect; younger fish from Berwick and Flamborough are more similar genetically than they are to older fish from the same locations (Supplementary Appendices 3, 4). The locus *Cpa112* showed a similar pattern, but with a more pronounced divergence for the Baltic samples. Pairwise comparisons involving the Baltic had one magnitude higher F_{ST} values than comparisons outside the Baltic. Larsson *et al.* (2007) showed that the differentiation in *Cpa112* is due to frequency differences especially for one allele, *Cpa112*(306). In this study, this allele

had average frequencies of 38.6% at Kalix, 10.5% at Tjøme and 3.4% among the North Sea samples (see Figure 5). This pattern corroborates that *Cpa112* is likely influenced by selection. The allozymes showed little divergence: 4 out of 11 loci had negative overall F_{ST} , and 16 out of 28 pairwise comparisons also had a negative F_{ST} estimate. As an exception, the herring sample collected in Tjøme in 2002 was highly divergent from other samples at the *SOD** locus (Figure 4, Supplementary Appendix 2). Owing to this rare pattern, we repeated the genotyping of this specific sample, with an identical result. Notwithstanding this specific sample, and despite the many negative F_{ST} values, the North Sea samples clearly grouped together (Figure 4). Similar to the allozymes, the mtDNA data included a single highly divergent sample (Figure 4, Supplementary Appendix 2): the Kalix 2003 sample had one haplotype occurring at a frequency of 20% compared with around 2% in all other samples. In addition, mtDNA displayed a number of statistically significant pairwise differences. Finally, MHC showed a more complex pattern. Some pairwise comparisons were negative and some had relatively high point estimates of F_{ST} of around 1%, without any obvious geographical pattern (Figure 4, Supplementary Appendix 2).

The hierarchical analysis of molecular variance analysis (Table 5) indicated spatial differentiation among basins for microsatellites and *Cpa112*. MtDNA, on the

other hand, showed a strong temporal variation, presumably masking any spatial pattern (see Table 2, $s = 4$).

Marker correlations

Across samples, there was strikingly little concordance in sample pairwise F_{ST} between marker types ($R = 0.000-0.005$; $P > 0.05$), except between *Cpa112* and microsatellites ($R = 0.70$; $P < 0.001$), and *Cpa112* and mtDNA ($R = 0.44$; $P = 0.021$).

Discussion

Using the same fish for all analyses, thus excluding year-class effects, we show that levels of population differentiation in herring were low for the presumed neutral genetic markers (allozymes, mtDNA and microsatellites), in agreement with earlier studies from the same region (Ryman *et al.*, 1984; Hauser *et al.*, 2001; Bekkevold *et al.* 2005; Jørgensen *et al.*, 2005; Mariani *et al.*, 2005; Larsson *et al.*, 2007). For the putatively selected markers, the microsatellite locus *Clha-DAA-INTR3* embedded in the *MHC* gene showed characteristics similar to those of the neutral microsatellite loci, and there was no indication of selection influencing this locus. *Cpa112* on the other hand was highly divergent for Baltic samples, indicating strong influence by selection as reported in previous studies (Gaggiotti *et al.*, 2009; Larsson *et al.*, 2007, 2010). The higher level of divergence in *Cpa112* makes this locus efficient at detecting even weak overall population structure in Atlantic herring.




Figure 5 Frequency of the allele 306 in the locus *Cpa112* in samples of herring collected in 2002 and 2003 along the salinity gradient from the North Sea to the inner part of the Baltic Sea. Genetic data are compiled from Mariani *et al.*, 2005; Bekkevold *et al.*, 2005; Larsson *et al.*, 2010 and André C (unpublished) (sample collected in 2002 in Idefjord, a sheltered Skagerrak fjord with low salinity, see Figure 1). Salinities were obtained as in Bekkevold *et al.*, 2005 or by the Swedish Meteorological and Hydrological Institute.

Statistical power of neutral vs selected markers

All genetic markers, including *Cpa112*, displayed high statistical power for detecting low levels of genetic differentiation in simulations; the applied set of eight neutral microsatellites could even detect F_{ST} as low as 0.0010 with a power of 95% at $n \approx 100$. Statistical power is related to the distribution of expected F_{ST} under neutrality, and the smaller variance of the eight microsatellites explains their higher power (Figure 3). In turn, this is likely an effect of the larger total number of alleles (Kalinowski, 2002; Ryman *et al.*, 2006; Larsson *et al.*, 2007). The eight microsatellite loci had 192 alleles in total, whereas the eleven allozyme loci had 38 alleles. Interestingly, the median microsatellite locus harboring 26 alleles had a power comparable to the allozymes (Table 3).

These evaluations were made under the assumption of selective neutrality; if selection is taken into account, expectations change. For Baltic vs North Sea herring, *Cpa112* displayed a 10–20 times higher overall F_{ST} compared with the eight neutral microsatellite loci (Table 2), and this higher level of differentiation increase the statistical power. Consequently, a substantial reduction in sample size is possible when using a selected locus vs neutral markers (Table 4), in agreement with predictions by Schmidt *et al.* (2008). However, it should be noted that these estimates relate to F_{ST} over all eight samples, and that power varied with specific comparisons, that is, North Sea vs Skagerrak comparisons were less powerful, and Baltic vs Skagerrak or North Sea comparisons were even more powerful than the estimates above (see Supplementary Appendix 2 for detailed pairwise comparisons of F_{ST}).

Several pairwise comparisons using mtDNA were statistically significant (Supplementary Appendix 2), indicating that mtDNA can be a powerful marker, even at low levels of nuclear divergence (Hoarau *et al.*, 2004; Larsson *et al.*, 2009).

Spatial and temporal patterns of differentiation

Neutral markers: The microsatellite results support the existence of a barrier to gene flow between Baltic and Skagerrak/North Sea populations, which has been shown earlier for herring (Bekkevold *et al.*, 2005; Larsson *et al.*, 2007), as well as for several other species (Johannesson and André, 2006; Limborg *et al.*, 2009). Using the microsatellite data for the same samples collected at Berwick and Tjøme as in this study, as well as several other samples, Ruzzante *et al.* (2006) showed that herring collected at Rügen in the southwestern Baltic is clearly distinct from the populations in the North Sea and Skagerrak. In this study, the differentiation estimated with neutral microsatellites was stable over

Table 5 Hierarchical analysis of genetic differences (AMOVA) among four sampling locations (see Figure 1) and between temporal samples (2002 and 2003) of Atlantic herring for different genetic markers

Source of variation	d.f.	Microsatellites		Allozymes		MtDNA		MHC		Cpa112	
		F-value	P-value	F-value	P-value	F-value	P-value	F-value	P-value	F-value	P-value
Among locations	3	0.0016	0.005	−0.0007	0.78	−0.0013	0.63	0.0052	0.067	0.0410	0.010
Between yrs within locations	4	0.0011	0.037	0.0018	0.25	0.016	0.001	−0.0017	0.81	0.0007	0.32

Abbreviations: AMOVA, analysis of molecular variance; MHC, major histocompatibility complex; mtDNA, mitochondrial DNA.

two subsequent years (Figure 4). The allozyme data for the 2003 herring showed a concordant spatial pattern of differentiation, as was also reported by Larsson *et al.* (2007). In 2002, however, the sample collected at Tjøme was highly divergent at the *SOD** locus. Owing to the low overall level of divergence, a single odd genotype at one locus had a disproportional impact on the F_{ST} estimate. A similar case is shown in the mtDNA data in which one specific haplotype had an elevated frequency in the Kalix 2003 sample (Figure 4). Owing to the low levels of differentiation typical for marine fish, a slightly different set of samples, or loci, may yield different patterns of population structure (*cf.* Waples, 1998; Allendorf and Seeb, 2000; Hauser and Carvalho, 2008). Such variance is particularly pronounced in genetic markers with low heterozygosities and a high expected range of F_{ST} , such as allozymes (Figure 3; also refer Fauvelot *et al.*, 2007). Therefore, careful sampling including temporal replication is warranted, especially in species that undertake ontogenetic shifts in habitat use, such as herring and cod, in which the inclusion of juvenile fish may bias patterns of spatial genetic differentiation (Nielsen *et al.*, 2009b).

Markers influenced by selection

The MHC-embedded microsatellite locus, *Clha-DAA-INTR3*, showed similar patterns of allelic richness and divergence as the presumed neutral microsatellite loci (Table 2; Supplementary Appendix 1a), and there was no indication of either directional or balancing selection acting on *Clha-DAA-INTR3* (Figure 2). The locus is situated in the third intron of the *MH IIA* gene, and the extent to which the variability in the microsatellite reflects exonic variability is unknown (Stet *et al.*, 2008). Previous attempts to estimate the variability in MHC genes using embedded micro- or minisatellites in salmonids have yielded varying results (Stet *et al.*, 2002; Grimholt *et al.*, 2002, 2003; Hansen *et al.*, 2007; de Eyto *et al.*, 2007). de Eyto *et al.* (2007) found a one-to-one relationship between minisatellite and MH class II allelic variability; however, for MH class I, a specific microsatellite allele could be found in several different MH I alleles. Wynne *et al.* (2007) showed that, although alleles of a microsatellite located in the untranslated region of the salmon *MH II* gene were linked to several different MH II sequences, the microsatellite diversity was correlated to disease susceptibility, and thus acted as a good proxy for actual MH class II polymorphism. In a recent study, Tonteri *et al.* (2010) demonstrated that 18 microsatellite loci linked to immune-relevant genes show stronger signal of selection than 76 loci without such association.

Baltic herring are highly divergent at the microsatellite locus *Cpa112*. Larsson *et al.* (2010) showed this for herring collected in the 1980s, and it is also evident for herring collected in 2002 and 2003 (Figure 4; Supplementary Appendix 2). Skagerrak and North Sea samples are also divergent at this locus, but to a substantially lesser degree. Presently, there is no information on any gene associated with *Cpa112*, but the lower salinity of the Baltic Sea is an obvious candidate as selective agent (Figure 5), although other environmental factors that covary with salinity may also have a role. This is also supported by preliminary data indicating a high frequency of the *Cpa112(306)* allele in the low-saline Idefjord in NE Skagerrak (Figure 5).

Conclusion

Neutral markers have traditionally been used to estimate population structure and also migration rates. A decade ago, Whitlock and McCauley (1999) pointed out that indirect estimates of migration rates using gene frequency data often tend to be flawed because underlying assumptions are violated. Recently, new ways of estimating migration, m , independently of F_{ST} that do not rely on equilibrium assumptions have been developed (Beerli and Felsenstein, 2001), but most investigations of population structure *per se* still use presumed neutral markers, most likely because it is believed that selection may bias F_{ST} . However, as long as selection pressure is temporally stable (decades–centuries), genetic markers influenced by selection, such as *Cpa112*, can be of use in detecting population structure on ecological time scales, where neutral loci have not yet diverged substantially. Selected loci may be particularly useful in genetic stock identification based on individual assignment and as markers for population origin in mixed-stock analysis (see Nielsen *et al.*, 2009a, p. 3136), for example, MHC in Pacific sockeye salmon (Beacham *et al.*, 2005), the pantophysin locus in mixed stocks of coastal and oceanic cod in Lofoten, northern Norway (Wennevik *et al.*, 2008) and *Cpa112* in mixed-stock analysis of herring in the Skagerrak (Ruzzante *et al.*, 2006). It may be preferable to use selection-influenced genetic markers with known function (Hoffmann and Willi, 2008), or with at least some knowledge of the nature of the selection pressure, for example, hemoglobin (Andersen *et al.*, 2009), genes involved in osmoregulation (Larsen *et al.*, 2008) and depth adaptation (Arnason *et al.*, 2009). With recent advances in molecular technologies, it is anticipated that such information will become increasingly available. Moreover, loci under selection may allow for an estimation of the adaptive value of subpopulations, useful for conservation priorities (Bonin *et al.*, 2007; Gebremedhin *et al.*, 2009).

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

This paper is dedicated to Rene Stet who completed the initial research on the herring MHC. We thank two anonymous reviewers for very valuable comments and suggestions. Sampling and fish ageing were undertaken by the Marine Scotland Marine Laboratory in the United Kingdom, at IMR in Norway, and by the Swedish National Board of Fisheries. This work is part of the research project HERGEN (<http://www.hull.ac.uk/hergen/>) funded by the 5th EU Frame work program, and the efforts of the consortium are acknowledged. Financial support was provided by the Swedish Research Council for Environment FORMAS (CA, LL, NR), the Swedish Research Council VR (CA, LL, NR), and Sven and Lilly Lawski's Foundation (LCL). Parts of the analyses were conducted within the framework of the BaltGene project funded by BONUS Baltic Organisations' Network for Funding Science EEIG, and the Linnaeus Centre for Marine Evolutionary Biology at University of Gothenburg (<http://www.cemeb.science.gu.se>).

References

- Allendorf FW, Seeb LW (2000). Concordance of genetic divergence among sockeye salmon populations at allozyme, nuclear DNA, and mitochondrial DNA markers. *Evolution* **54**: 640–651.
- Allendorf FW, Luikart G (2007). *Conservation and the Genetics of Populations*. Blackwell Publishing: Malden, MA.
- Allendorf FW, England PR, Luikart G, Ritchie PA, Ryman N (2008). Genetic effects of harvest on wild animal populations. *Trends Ecol Evol* **23**: 327–337.
- Andersen Ø, Wetten OF, De Rosa MC, André C, Alinovi CC, Colafranceschi M *et al.* (2009). Haemoglobin polymorphisms affect the oxygen-binding properties in Atlantic cod populations. *Proc R Soc B* **276**: 833–841.
- Andersson L, Ryman N, Rosenberg R, Ståhl G (1981). Genetic variability in Atlantic herring (*Clupea harengus harengus*): description of protein loci and population data. *Hereditas* **95**: 69–78.
- Árnason E, Hernandez UB, Kristinsson K (2009). Intense habitat-specific fisheries-induced selection at the molecular *Pan I* locus predicts imminent collapse of a major cod fishery. *PLoS ONE* **4**: e5529.
- Avise JC (2004). *Molecular Markers, Natural History, and Evolution* 2 edn, Sinauer: Sunderland, MA.
- Baer CF (1999). Among-locus variation in F_{ST} : fish, allozymes and the Lewontin-Krakauer test revisited. *Genetics* **152**: 653–659.
- Beacham TD, Candy JR, McIntosh B, MacConnachie C, Tabata A, Kaukinen K *et al.* (2005). Estimation of stock composition and individual identification of sockeye salmon on a Pacific rim basis using microsatellite and major histocompatibility complex variation. *Trans Am Fish Soc* **134**: 1146.
- Beaumont MA (2002). FDIST2. Available from <http://rubic.rdg.ac.uk/~mab/software.html>.
- Beaumont MA, Nichols RA (1996). Evaluating loci for use in the genetic analysis of population structure. *Proc R Soc B* **263**: 1619–1626.
- Beerli P, Felsenstein J (2001). Maximum likelihood estimation of a migration matrix and effective population sizes in n subpopulations using a coalescent approach. *Proc Natl Acad Sci USA* **98**: 4563–4568.
- Bekkevold D, André C, Dahlgren TG, Clausen LAW, Torstensen E, Mosegaard H *et al.* (2005). Environmental correlates of population differentiation in Atlantic herring. *Evolution* **59**: 2656–2668.
- Bernatchez L, Landry C (2003). MHC studies in nonmodel vertebrates: what have we learned about natural selection in 15 years? *J Evol Biol* **16**: 363–377.
- Bonin A, Nicole F, Pompanon F, Miaud C, Taberlet F (2007). Population adaptive index: a new method to help measure intraspecific genetic diversity and prioritize populations for conservation. *Conserv Biol* **21**: 697–708.
- Bos DH, Gopurenko D, Williams RN, DeWoody JA (2008). Inferring population history and demography using microsatellites, mitochondrial DNA, and major histocompatibility complex (MHC) genes. *Evolution* **62**: 1458–1468.
- Clausen LAW, Bekkevold D, Hatfield EMC, Mosegaard H (2007). Application and validation of otolith microstructure as a stock identification method in mixed Atlantic herring (*Clupea harengus*) stocks in the North Sea and western Baltic. *ICES J Mar Sci* **64**: 377–385.
- Dahle G, Eriksen AG (1990). Spring and autumn spawners of herring (*Clupea harengus*) in the North Sea, Skagerrak and Kattegat; population genetic analysis. *Fisheries Res* **9**: 131–141.
- de Eyto E, McGinnity P, Consuegra S, Coughlan J, Tufto J, Farrell K *et al.* (2007). Natural selection acts on Atlantic salmon major histocompatibility (MH) variability in the wild. *Proc R Soc B* **274**: 861–869.
- Excoffier L, Laval G, Schneider S (2005). ARLEQUIN (version 3.0): an integrated software package for population genetics data analysis. *Evol Bioinform Online* **1**: 47–50.
- Fauvelot C, Lemaire C, Planes S, Bonhomme F (2007). Inferring gene flow in coral reef fishes from different molecular markers: which loci to trust? *Hereditas* **99**: 331–339.
- Gaggiotti OE, Bekkevold D, Jørgensen HBH, Foll M, Carvalho GR, André C *et al.* (2009). Disentangling the effects of evolutionary, demographic, and environmental factors influencing genetic structure of natural populations: Atlantic herring as a case study. *Evolution* **63**: 2939–2951.
- Galarza JA, Carreras-Carbonell J, Macpherson E, Pascual M, Roques S, Turner GF, Rico C (2009). The influence of oceanographic fronts and early-life-history traits on connectivity among littoral fish species. *Proc Natl Acad Sci USA* **106**: 1473–1478.
- Gebremedhin B, Ficetola GF, Naderi S, Rezaei H-R, Maudet C, Rioux D *et al.* (2009). Combining genetic and ecological data to assess the conservation status of the endangered Ethiopian walia ibex. *Anim Conserv* **12**: 89–100.
- Goudet J (2001). FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3). Available from <http://www.unil.ch/izea/software/fstat.html>.
- Grant WS, Spies IB, Canino MF (2006). Biogeographic evidence for selection on mitochondrial DNA in North Pacific walleye pollock *Theragra chalcogramma*. *J Hered* **97**: 571–580.
- Grimholt U, Drablos F, Jørgensen SM, Hoyheim B, Stet RJM (2002). The major histocompatibility class I locus in Atlantic salmon (*Salmo salar* L. polymorphism, linkage analysis and protein modelling. *Immunogenetics* **54**: 570–581.
- Grimholt U, Larsen S, Nordmo R, Midtlyng P, Kjoeglum S, Storset A *et al.* (2003). MHC polymorphism and disease resistance in Atlantic salmon (*Salmo salar*); facing pathogens with single expressed major histocompatibility class I and class II loci. *Immunogenetics* **55**: 210–219.
- Hansen MM, Skaala O, Jensen LF, Bekkevold D, Mensberg KLD (2007). Gene flow, effective population size and selection at major histocompatibility complex genes: brown trout in the Hardanger Fjord, Norway. *Mol Ecol* **16**: 1413–1425.
- Hauser L, Turan C, Carvalho GR (2001). Haplotype frequency distribution and discriminatory power of two mtDNA fragments in a marine pelagic teleost (Atlantic herring, *Clupea harengus*). *Hereditas* **87**: 621–630.
- Hauser L, Carvalho GR (2008). Paradigm shifts in marine fisheries genetics: ugly hypotheses slain by beautiful facts. *Fish Fish* **9**: 333–362.
- Hemmer-Hansen J, Nielsen EE, Frydenberg J, Loeschcke V (2007). Adaptive divergence in a high gene flow environment: *Hsc70* variation in European flounder (*Platichthys flesus* L.). *Hereditas* **99**: 592–600.
- Hoarau G, Piquet AMT, Van der Veer HW, Rijnsdorp AD, Stam WT, Olsen JL (2004). Population structure of plaice (*Pleuronectes platessa* L.) in northern Europe: a comparison of resolving power between microsatellites and mitochondrial DNA data. *J Sea Res* **51**: 183–190.
- Hoffmann AA, Willi Y (2008). Detecting genetic responses to environmental change. *Nat Rev Genet* **9**: 421–432.
- ICES (2003). Report of the Herring Assessment Working Group for the area South of 62°N. ICES CM 2003 /ACFM: 17.
- Inoue JG, Miya M, Tsukamoto K, Nishida M (2000). Complete mitochondrial DNA sequence of the Japanese sardine *Sardinops melanostictus*. *Fish Sci* **66**: 924–932.
- Johannesson K, André C (2006). Life on the margin: genetic isolation and diversity loss in a peripheral marine ecosystem, the Baltic Sea. *Mol Ecol* **15**: 2013–2029.
- Jorde PE, Knutsen H, Espeland SH, Stenseth NC (2007). Spatial scale of genetic structuring in coastal cod *Gadus morhua* and the geographic extent of local populations. *Mar Ecol Prog Ser* **343**: 229–237.
- Jørgensen HBH, Hansen MM, Bekkevold D, Ruzzante DE, Loeschcke V (2005). Marine landscapes and population genetic structure of herring (*Clupea harengus* L.) in the Baltic Sea. *Mol Ecol* **14**: 3219–3234.
- Jørgensen HBH, Pertoldi C, Hansen MM, Ruzzante DE, Loeschcke V (2008). Genetic and environmental correlates

- of morphological variation in a marine fish: the case of Baltic Sea herring (*Clupea harengus*). *Can J Fish Aquat Sci* **65**: 389–400.
- Kalinowski ST (2002). How many alleles per locus should be used to estimate genetic distances? *Heredity* **88**: 62–65.
- Karl SA, Avise JC (1992). Balancing selection at allozyme loci in oysters: implications from nuclear RFLPs. *Science* **256**: 100–102.
- Larsen PF, Nielsen EE, Koed A, Thomsen DS, Olsvik PA, Loeschcke V (2008). Interpopulation differences in expression of candidate genes for salinity tolerance in winter migrating anadromous brown trout (*Salmo trutta* L.). *BMC Genetics* **9**: 12.
- Larsson LC, Laikre L, Palm S, André C, Carvalho GR, Ryman N (2007). Concordance of allozyme and microsatellite differentiation in a marine fish, but evidence of selection at a microsatellite locus. *Mol Ecol* **16**: 1135–1147.
- Larsson LC, Charlier J, Laikre L, Ryman N (2009). Statistical power for detecting genetic divergence—organelle versus nuclear markers. *Conserv Genet* **10**: 1255–1264.
- Larsson LC, Laikre L, André C, Dahlgren TG, Ryman N (2010). Temporally stable genetic structure of heavily exploited Atlantic herring (*Clupea harengus*) in Swedish waters. *Heredity* **104**: 40–51.
- Lewis PO, Zaykin D (2001). Genetic Data Analysis: Computer program for the analysis of allelic data. Version 1.0 (d16c). Free program distributed over the internet from <http://hydrodictyon.eeb.uconn.edu/people/plewis/software.php>.
- Limborg MT, Pedersen JS, Hemmer-Hansen J, Tomkiewicz J, Bekkevold D (2009). Genetic population structure of European sprat, *Sprattus sprattus*: differentiation across a steep environmental gradient in a small pelagic fish. *Mar Ecol Prog Ser* **379**: 213–224.
- Mariani S, Hutchinson WF, Hatfield EMC, Ruzzante DE, Simmonds EJ, Dahlgren TG *et al.* (2005). North Sea herring population structure revealed by microsatellite analysis. *Mar Ecol Prog Ser* **303**: 245–257.
- McPherson AA, O'Reilly PT, McParland TL, Jones MW, Bentzen P (2001). Isolation of nine novel tetranucleotide microsatellites in Atlantic herring (*Clupea harengus*). *Mol Ecol Notes* **1**: 31–32.
- Muirhead CA (2001). Consequences of population structure on genes under balancing selection. *Evolution* **55**: 1532–1541.
- Nielsen EE, Hansen MM, Meldrup D (2006). Evidence of microsatellite hitch-hiking selection in Atlantic cod (*Gadus morhua* L.) implications for inferring population structure in nonmodel organisms. *Mol Ecol* **15**: 3219–3229.
- Nielsen EE, MacKenzie BR, Magnussen E, Meldrup D (2007). Historical analysis of *Pan I* in Atlantic cod (*Gadus morhua*): temporal stability of allele frequencies in the southeastern part of the species distribution. *Can J Fish Aquat Sci* **64**: 1448–1455.
- Nielsen EE, Hemmer-Hansen J, Larsen PF, Bekkevold D (2009a). Population genomics of marine fishes: identifying adaptive variation in space and time. *Mol Ecol* **18**: 3128–3150.
- Nielsen EE, Wright PJ, Hemmer-Hansen J, Poulsen NA, Gibb IM, Meldrup D (2009b). Microgeographical population structure of cod *Gadus morhua* in the North Sea and west of Scotland: the role of sampling loci and individuals. *Mar Ecol Prog Ser* **376**: 213–225.
- O'Malley KG, Camara MD, Banks MA (2007). Candidate loci reveal genetic differentiation between temporally divergent migratory runs of Chinook salmon (*Oncorhynchus tshawytscha*). *Mol Ecol* **16**: 4930–4941.
- Olsen JB, Lewis CJ, Kretschmer EJ, Wilson SL, Seeb JE (2002). Characterization of 14 tetranucleotide microsatellite loci derived from Pacific herring. *Mol Ecol Notes* **2**: 101–103.
- Rand DM (1996). Neutrality tests of molecular markers and the connection between DNA polymorphism, demography and conservation biology. *Conserv Biol* **10**: 665–671.
- Raymond M, Rousset F (1995). GENEPOP (Version 1.2): population genetics software for exact tests and ecumenicism. *J Hered* **86**: 248–249.
- Ruzzante DE, Mariani S, Bekkevold D, André C, Mosegaard H, Clausen LAW *et al.* (2006). Biocomplexity in a highly migratory pelagic marine fish, Atlantic herring. *Proc R Soc B* **273**: 1459–1464.
- Ryman N, Lagercrantz U, Andersson L, Chakraborty R, Rosenberg R (1984). Lack of correspondence between genetic and morphologic variability patterns in Atlantic herring. *Heredity* **53**: 687–704.
- Ryman N (2006). CHIFISH: a computer program testing for genetic heterogeneity at multiple loci using chi-square and Fisher's exact test. *Mol Ecol Notes* **6**: 285–287.
- Ryman N, Palm S (2006). POWSIM: a computer program for assessing statistical power when testing for genetic differentiation. *Mol Ecol Notes* **6**: 600–602.
- Ryman N, Palm S, André C, Carvalho GR, Dahlgren TG, Jorde PE *et al.* (2006). Power for detecting genetic divergence: differences between statistical methods and marker loci. *Mol Ecol* **15**: 2031–2045.
- Schmidt PS, Serrão EA, Pearson GA, Riginos C, Rawson PD, Hillish TJ *et al.* (2008). Ecological genetics in the North Atlantic: environmental gradients and adaptation at specific loci. *Ecology* **89**: S91–S107.
- Stet RJM, de Vries B, Mudde K, Hermsen T, van Heerwaarden J, Shum BP *et al.* (2002). Unique haplotypes of co-segregating major histocompatibility class II A and class II B alleles in Atlantic salmon (*Salmo salar*) give rise to diverse class II genotypes. *Immunogenetics* **54**: 320–331.
- Stet RJM, Mudde K, Wynne JW, Nooijen A, Dahlgren TG, Ruzzante DE *et al.* (2008). Characterization of a major histocompatibility class II A gene (*Clha-DAA*) with an embedded microsatellite marker in Atlantic herring (*Clupea harengus* L.). *J Fish Biol* **73**: 367–381.
- Tonteri A, Vasemägi A, Lumme J, Primmer CR (2010). Beyond MHC: signals of elevated selection pressure on Atlantic salmon (*Salmo salar*) immune-relevant loci. *Mol Ecol* **19**: 1273–1282.
- Truett GE, Heeger P, Mynatt RL, Truett AA, Walker JA, Warman ML (2000). Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris (HotSHOT). *Biotechniques* **29**: 52–54.
- Waples RS (1998). Separating the wheat from the chaff: patterns of genetic differentiation in high gene flow species. *J Hered* **89**: 438–450.
- Wennevik V, Jørstad KE, Dahle G, Fevolden S-E (2008). Mixed stock analysis and the power of different classes of molecular markers in discriminating coastal and oceanic Atlantic cod (*Gadus morhua* L.) on the Lofoten spawning grounds, Northern Norway. *Hydrobiologia* **606**: 7–25.
- Westgaard J-I, Fevolden S-E (2007). Atlantic cod (*Gadus morhua* L.) in inner and outer coastal zones of northern Norway display divergent genetic signature at non-neutral loci. *Fisheries Res* **85**: 306–315.
- Whitlock MC, McCauley DE (1999). Indirect measures of gene flow and migration: $F_{ST} \neq 1/(4Nm+1)$. *Heredity* **82**: 117–125.
- Wynne JW, Cook MT, Nowak BE, Elliott NG (2007). Major histocompatibility polymorphism associated with resistance towards amoebic gill disease in Atlantic salmon (*Salmo salar* L.). *Fish Shellfish Immunol* **22**: 707–717.

Supplementary Information accompanies the paper on Heredity website (<http://www.nature.com/hdy>)