

Corrigendum

High efficiency expression of transfected genes in a *Drosophila melanogaster* haploid (1182) cell line

by Susan E.Saunders, John M.Rawls, Catherine J.Wardle and Julian F.Burke

*Nucleic Acids Research*, 17, 6205–6216 (this issue)

The legends for figures 1 and 2 of the above paper were incomplete. The correct legends are printed below.





Figure 1

Activity of promoters in various cell lines. The indicated promoters linked to *cat* were transfected by the calcium phosphate procedure into *D.hydei* DH33 [cross-hatched], and *D.melanogaster* S<sub>3</sub> [dotted], D<sub>1</sub> [solid black], 1182-6 [horizontal lines] and 1182-4 [vertical lines] cell lines. The cells were harvested after 72 hours and assayed for CAT activity as described in Materials and Methods. The histogram shows the average CAT activity (plotted on a log scale and expressed as arbitrary units) from two transfected plates. All data is from one experiment. All cell lines received an equivalent portion of a pooled calcium phosphate precipitate for each plasmid



**Figure 2**

Non facilitated transfection of D1 (hatched) and 1182-4 (white) *Drosophila* cells. Cells, either in suspension (s) or as a monolayer (m), were transfected with pCV1cat as described in Materials and Methods, either by adding DNA to cells in serum free media (A), in serum free media plus 10% PEG 1500 (B), or as a calcium phosphate precipitate (C). The CAT activity was measured as described in Materials and Methods and expressed in arbitrary units plotted on a log scale.