

Published in final edited form as:

*Angew Chem Int Ed Engl.* 2011 February 11; 50(7): 1673–1677. doi:10.1002/anie.201005352.

## $\alpha,\beta$ -Unsaturated Acyl Azoliums from N-Heterocyclic Carbene Catalyzed Reactions: Observation and Mechanistic Investigation\*\*

**Jessada Mahatthananchai,**

Laboratory for Organic Chemistry, Department of Chemistry and Applied Biosciences, Swiss Federal Institute of Technology (ETH) Zurich, Wolfgang-Pauli-Strasse 10, 8093 Zurich (Switzerland)

**Dr. Pinguan Zheng,** and

Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, PA 19104 (USA)

**Prof. Dr. Jeffrey W. Bode\***

Laboratory for Organic Chemistry, Department of Chemistry and Applied Biosciences, Swiss Federal Institute of Technology (ETH) Zurich, Wolfgang-Pauli-Strasse 10, 8093 Zurich (Switzerland)

### Abstract

Catalytically generated acyl azoliums **I** and their  $\alpha,\beta$ -unsaturated counterparts **II** are thought to be key reactive intermediates in a rapidly growing number of transformations promoted by N-heterocyclic carbene (NHC) catalysts.<sup>[1]</sup> Acyl azoliums are invoked in the postulated catalytic cycles of nearly all of the new NHC-catalyzed reactions of  $\alpha$ -functionalized aldehydes reported since 2004, in which they are generally assumed to possess the reactivity of an activated carboxylic acid, that is, analogous to an activated ester.<sup>[2]</sup> In NHC-catalyzed processes, they are most often obtained through internal redox reactions of functionalized aldehydes but have also been prepared by oxidations of the Breslow intermediates<sup>[3]</sup> or additions to ketenes.<sup>[4]</sup> Acyl azoliums **I** are important intermediates in thiamine pyrophosphate (ThPP) dependent enzymatic reactions.<sup>[5]</sup> Townsend et al. have recently proposed that unsaturated acyl azolium **III** is the key intermediate in clavulanic acid biosynthesis;<sup>[6]</sup> despite careful efforts, **III** or its analogues **II** have never been characterized or independently synthesized. Here, we document the observation and characterization of  $\alpha,\beta$ -unsaturated acyl azoliums **1** and **2** (Scheme 1) and demonstrate that their corresponding hemiacetals (**1'** and **2''**) are the kinetically important intermediates in both their acylation and annulation reactions.

### Keywords

acyl azolium; N-heterocyclic carbenes; organocatalysis; reaction mechanisms; redox reactions

\*\*We are grateful to NIGMS (National Institutes of Health, GM-079339), the National Science Foundation (CHE-0449587), Bristol Myers Squibb, and ETH Zurich for support of this research. We thank Darko Santrac for a preliminary study, and NMR and MS services (ETHZ) for spectroscopic data.

© 2011 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

\*Fax: (+41)44-633-1235, bode@org.chem.ethz.ch, Homepage: <http://www.bodegroup.org>.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201005352>.

Simple acyl azoliums prepared under stoichiometric conditions were extensively studied in a seminal work by Breslow<sup>[7]</sup> and continued by Daigo,<sup>[8]</sup> White and Ingraham,<sup>[9]</sup> Bruice,<sup>[10]</sup> Lienhard,<sup>[11]</sup> and Owen.<sup>[12]</sup> These investigations revealed the unique and rich chemistry of acyl azoliums, including their remarkable reluctance to acylate amines<sup>[12, 13]</sup> and high preference for reactions with water or alcohols. Despite the more than 120 publications since 2004 that feature acyl azoliums, including the rediscovery of their unusual chemoselectivity, there have been no reports of the isolation, detection, or properties of novel acyl azoliums thought to be generated under catalytic conditions.<sup>[14]</sup>

Even at high catalyst loadings, most NHC-catalyzed reactions do not give any detectable intermediates that can be observed with conventional techniques such as UV, IR, NMR spectroscopy,<sup>[15]</sup> or MS methods.<sup>[16]</sup> For example, no intermediates could be observed during the redox esterification of cinnamaldehyde, even when the reaction was run with high catalyst loadings or in the absence of base to ensure slow reaction. This is consistent with NMR studies of the redox esterification of ynals by Zeitler, in which the postulated  $\alpha,\beta$ -unsaturated acyl azolium **II** was not observed.<sup>[17]</sup>

Our recent studies of reactions of ynals catalyzed by azolium salts revealed that the rate-limiting step of the catalytic cycle occurs after the formation of the acyl azolium.<sup>[18]</sup> These findings strongly suggested that generation of substantial amounts of the  $\alpha,\beta$ -unsaturated acyl azolium intermediate should be possible in the absence of a nucleophile. Careful preparation of a mixture of triazolium **3** and *para*-chlorophenyl ynal **4** in anhydrous CDCl<sub>3</sub> gives a clear solution (Figure 1). Upon addition of NaOAc, the solution rapidly becomes yellow, and turns deep red within 20 minutes. The color change was monitored by UV/Vis spectroscopy, and we observed a maximum absorption at 355 nm, with a significantly smaller absorption at 520 nm, for the solution containing acyl azolium **1** (Figure 2). Townsend et al. have speculated that the related protein-bound species **III** (Scheme 1) has a characteristic UV/Vis absorption at 310–320 nm.<sup>[6a]</sup> Our mixture was further investigated by electrospray ionization–high resolution mass spectrometry (ESI-HRMS), which verified the molecular formula of **1** (Figure 2).

To confirm the formation of **1**, the reaction was monitored by <sup>1</sup>H NMR spectroscopy. Figure 1 shows the <sup>1</sup>H NMR spectra of the region from  $\delta = 1.0$ – $13.0$  ppm before (Figure 1a) and after (Figure 1b) treatment of the mixture of azolium **3** and aldehyde **4** with NaOAc. A new product, characterized by doublets at  $\delta = 8.7$  ppm (<sup>1</sup>J=18.0 Hz) and at  $\delta = 7.6$  ppm (<sup>1</sup>J=18.0 Hz), which is consistent with the expected acyl azolium **1**, was detected and grew in intensity over time, reaching a maximum of 30% conversion after about 40 minutes. This new product immediately disappeared upon the addition of MeOH (Figure 1c), cleanly forming the corresponding unsaturated ester **5** and regenerating the protonated triazolium **3** (C-2 proton peak at  $\delta = 12.1$  ppm)<sup>[19]</sup>

Although this solution could be filtered away from undissolved NaOAc or stored for several hours in the NMR tube, acyl azolium **1** was exceedingly prone to hydrolysis to give *para*-chlorocinnamic acid; all attempts to isolate **1** in pure form were unsuccessful. However, this intermediate was successfully characterized by COSY, HSQC, and HMBC NMR experiments that were performed on the mixture (see the Supporting Information for spectra and details). Correlation spectroscopy (Figure 3) clearly showed the bond between the acyl group containing an  $\alpha,\beta$ -unsaturated moiety and the triazolium, and ruled out a triazole adduct, which we had previously observed in prior attempts at detecting catalytically generated acyl azoliums with non-anhydrous conditions.<sup>[20]</sup> <sup>1</sup>H NMR experiments were also repeated with *para*-methoxyphenyl ynal, which led to the formation of acyl azolium **2** with higher conversion (about 50%) relative to that of the aldehyde, but was much more susceptible to hydrolysis (see the Supporting Information).

The reactivity of **1**, generated in this pseudo-stoichiometric fashion, paralleled the chemistry observed under catalytic conditions (Scheme 2). It reacts instantaneously with water or MeOH to give acid **6** or ester **5**, but only a complex, unidentifiable mixture was obtained when **1** was treated with piperidine, an observation consistent with the known chemistry of saturated acyl azoliums and the reluctance of NHC-catalyzed acylations to deliver amide products in the absence of a suitable co-catalyst.<sup>[8, 12, 13]</sup> The addition of the kojic acid derivative **7** gives annulation product **8** through a mechanism that we believed to be a Coates–Claisen rearrangement from a hemiacetal intermediate (Scheme 3). Very recent related works by Lupton,<sup>[21]</sup> Xiao,<sup>[22]</sup> and Studer,<sup>[23]</sup> however, invoked a direct 1,4-addition of a nucleophile to the  $\alpha,\beta$ -unsaturated acyl azolium.

With the role of the acyl azolium **1** in both the annulation and acylation chemistry established, we sought to further clarify the mechanistic pathways and chemical properties of  $\alpha,\beta$ -unsaturated acyl azoliums by kinetic investigations. In NHC-catalyzed reactions, kinetic studies are often complicated by the complex reaction mechanism and catalyst inhibition events.<sup>[15c]</sup> Our recent finding that ynals are sufficiently reactive under NHC-catalyzed reactions in the absence of added base<sup>[24]</sup> simplified the reaction systems and allowed determination of the rate laws and activation parameters for both the esterification and annulation pathways.

This observed rate equation of the redox esterifications reaction is first order in aldehyde, partial order in the triazolium precatalyst, and partial negative order in MeOH (Scheme 3). The negative order in the nucleophile is initially surprising as we had expected that turnover of the acyl azolium to the ester and the NHC would be the rate-determining step. The negative order could be attributed to catalyst inhibition events. The inhibitory effects of the nucleophile were also observed even in the presence of base; the exact nature of this inhibition is not entirely clear.<sup>[25]</sup> Nonetheless, we were able to rationalize this observed partial rate orders with the derived rate law (see the Supporting Information). From the experimental rate law, we performed an Eyring analysis, which revealed an activation enthalpy ( $\Delta H^\ddagger$ ) of +23.60 kcal mol<sup>-1</sup> and an activation entropy ( $\Delta S^\ddagger$ ) of -2.93 calK<sup>-1</sup> mol<sup>-1</sup> for the redox esterification reaction.<sup>[26]</sup> These activation parameters can be contrasted to those of our recent studies on Coates–Claisen reactions<sup>[27]</sup> of ynals and enols, in which a large negative  $\Delta S^\ddagger$ , a lower  $\Delta H^\ddagger$ , and higher reaction rates with electron-deficient ynals were observed (Scheme 3).

With the role of  $\alpha,\beta$ -unsaturated acyl azolium in the catalytic cycle involving **3** and ynals established, we postulated that the mode of reactivity of NHC-catalyzed reactions of ynals and nucleophilic partner is governed by the pathways available to the corresponding hemiacetal (**1'** or **2'**) derived from the  $\alpha,\beta$ -unsaturated acyl azolium (**1** or **2**). To investigate this hypothesis, we undertook the measurement of the difference in the activation parameters ( $\Delta\Delta H^\ddagger$  and  $\Delta\Delta S^\ddagger$ )<sup>[28]</sup> between the Claisen and the redox reactions from the reaction of an ynal with sesamol—a reactant that could undergo both reactions (Scheme 4). We measured a  $\Delta\Delta H^\ddagger$  of 7.61 kcal mol<sup>-1</sup> and a  $\Delta\Delta S^\ddagger$  of 26.62 calK<sup>-1</sup> mol<sup>-1</sup> from an Eyring analysis of the product ratio ( $k_{\text{rel}} = \% \mathbf{10}/\% \mathbf{9}$ ) at five temperatures (see the Supporting Information). These values parallel the data obtained from the rate measurements of the independent processes (Scheme 3) and substantiates our postulate for the role of the kinetically important hemiacetal in the catalytic cycle of both the annulation and esterification reactions. The activation parameters of the competing processes arising for the same hemiacetal intermediate determine the reaction outcome.

Confirmation of the involvement of hemiacetal **V** as the final intermediate in the catalytic cycle of the esterification reaction was obtained by linear free-energy relationship analysis.<sup>[29]</sup> The Hammett plot revealed that electron-rich ynals underwent esterification

faster than their more electrophilic counterparts (Figure 4), an observation consistent with a mechanism featuring the breakdown of hemiacetal **V** as the rate-limiting step in the catalytic cycle (Scheme 5). This intermediate, which we cannot detect, must be in rapid equilibrium with unsaturated acyl azolium **IV** (i.e. the keto form). Although the equilibrium favors the  $\alpha,\beta$ -unsaturated acyl azolium, the reaction is governed by the kinetic profile of the pathways available to the hemiacetal. Electron-withdrawing groups on the acyl azolium stabilize the hemiacetal and decrease the rate of ester formation. This is consistent with the negative  $\rho$  value (albeit small) from the Hammett plot for redox esterification reactions (Figure 4), as well as the previous observation that electron-deficient ynals underwent faster Claisen reaction as these compounds have a higher proportion of the hemiacetal intermediate required for the Claisen rearrangement. This finding also corresponds with the studies of Breslow,<sup>[7]</sup> Daigo,<sup>[8]</sup> and Owen,<sup>[12]</sup> who have studied the reactions of simpler acyl azoliums generated under stoichiometric conditions and found the hemiacetal to be kinetically relevant. Our work demonstrates that the unusual reactivities and kinetics observed under catalytic conditions are governed by the unique properties of the catalytically generated  $\alpha,\beta$ -unsaturated acyl azoliums rather than by any of the prior C–C bond formations or protonation steps<sup>[30]</sup> involved in its formation.

In summary, we have successfully generated and characterized the acyl azolium intermediates key to the reactions of N-heterocyclic carbenes and  $\alpha$ -functionalized aldehydes. The confirmation that this intermediate is involved in both NHC-catalyzed acylation and annulation reactions has allowed us to demonstrate that a common intermediate, hemiacetal **V**, is the kinetically important species in both of these processes. These studies reveal the origin of the unique reactivities and selectivities observed in NHC-catalyzed reactions and confirm that NHC-catalyzed annulation reactions of  $\alpha,\beta$ -unsaturated acyl azoliums proceed through a Coates–Claisen reaction rather than a direct Michael addition. Our kinetics investigations provide a rate law (see the Supporting Information) that rationalizes the observed kinetics of both the esterifications and annulation pathways and considers the catalyst generation and inhibition processes that are the origin of the unexpected rate orders in the nucleophiles. These observations will serve as a guideline for future mechanistic investigations and identification of new reactions proceeding through catalytically generated acyl azoliums.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## References

1. a) Moore, J.L.; Rovis, T. *Topics in Current Chemistry*. List, B., editor. Vol. 291. Springer; Berlin: 2009. p. 77-144. b) Enders D, Niemeier O, Henseler A. *Chem Rev*. 2007; 107:5606–5655. [PubMed: 17956132] c) Zeitler K. *Angew Chem*. 2005; 117:7674–7678. *Angew Chem Int Ed*. 2005; 44:7506–7510. d) Enders D, Balensiefer T. *Acc Chem Res*. 2004; 37:534–541. [PubMed: 15311952] e) Chiang, P.C.; Bode, J.W. *N-Heterocyclic Carbenes: From Laboratory Curiosities to Efficient Synthetic Tools*. Díez-González, S., editor. Royal Society of Chemistry; Cambridge: 2010. p. 399-435.
2. a) Sohn SS, Rosen EL, Bode JW. *J Am Chem Soc*. 2004; 126:14370–14371. [PubMed: 15521753] b) Reynolds NT, Read de Alaniz J, Rovis T. *J Am Chem Soc*. 2004; 126:9518–9519. [PubMed: 15291537] c) Chow KYK, Bode JW. *J Am Chem Soc*. 2004; 126:8126–8127. [PubMed: 15225048] d) Burstein C, Glorius F. *Angew Chem*. 2004; 116:6331–6334. *Angew Chem Int Ed*. 2004; 43:6205–6208.
3. a) Guin J, De Sarkar S, Grimme S, Studer A. *Angew Chem*. 2008; 120:8855–8858. *Angew Chem Int Ed*. 2008; 47:8727–8730. b) Maki BE, Chan A, Phillips EM, Scheidt KA. *Org Lett*. 2007; 9:371–374. [PubMed: 17217307] c) Inoue H, Higashiura K. *J Chem Soc Chem Commun*. 1980:549–550.



4. a) Zhang YR, He L, Wu X, Shao PL, Ye S. *Org Lett.* 2008; 10:277–280. [PubMed: 18085789] b) Duguet N, Campbell CD, Slawin AMZ, Smith AD. *Org Biomol Chem.* 2008; 6:1108–1113. [PubMed: 18327339] c) Wang XN, Lv H, Huang XL, Ye S. *Org Biomol Chem.* 2009; 7:346–350. [PubMed: 19109680]
5. a) Kluger R, Tittmann K. *Chem Rev.* 2008; 108:1797–1833. [PubMed: 18491870] b) Frank R, Leeper F, Luisi B. *Cell Mol Life Sci.* 2007; 64:892–905. [PubMed: 17429582]
6. a) Merski M, Townsend CA. *J Am Chem Soc.* 2007; 129:15750–15751. [PubMed: 18052280] b) Khaleeli N, Li R, Townsend CA. *J Am Chem Soc.* 1999; 121:9223–9224.
7. Breslow R, McNelis E. *J Am Chem Soc.* 1960; 82:2394–2395.
8. Daigo K, Reed LJ. *J Am Chem Soc.* 1962; 84:659–662.
9. White FG, Ingraham LL. *J Am Chem Soc.* 1962; 84:3109–3111.
10. Bruice TC, Kundu NG. *J Am Chem Soc.* 1966; 88:4097–4098.
11. Lienhard GE. *J Am Chem Soc.* 1966; 88:5642–5649.
12. a) Owen TC, Richards A. *J Am Chem Soc.* 1987; 109:2520–2521. b) Owen TC, Harris JN. *J Am Chem Soc.* 1990; 112:6136–6137.
13. a) Movassaghi M, Schmidt MA. *Org Lett.* 2005; 7:2453–2456. [PubMed: 15932221] b) Bode JW, Sohn SS. *J Am Chem Soc.* 2007; 129:13798–13799. [PubMed: 17956104] c) Vora HU, Rovis T. *J Am Chem Soc.* 2007; 129:13796–13797. [PubMed: 17929821] d) De Sarkar S, Grimme S, Studer A. *J Am Chem Soc.* 2010; 132:1190–1191. [PubMed: 20055393]
14. a) Grasa GA, Kissling RM, Nolan SP. *Org Lett.* 2002; 4:3583–3586. [PubMed: 12375893] b) Sohn SS, Bode JW. *Org Lett.* 2005; 7:3873–3876. [PubMed: 16119920] c) He M, Bode JW. *Org Lett.* 2005; 7:3131–3134. [PubMed: 15987223] d) Sohn SS, Bode JW. *Angew Chem.* 2006; 118:6167–6170. *Angew Chem Int Ed.* 2006; 45:6021–6024. e) Suzuki Y, Muramatsu K, Yamauchi K, Morie Y, Sato M. *Tetrahedron.* 2006; 62:302–310. f) Burstein C, Tschan S, Xie X, Glorius F. *Synthesis.* 2006:2418–2439. g) Chiang PC, Kim Y, Bode JW. *Chem Commun.* 2009:4566–4568. h) Lathrop SP, Rovis T. *J Am Chem Soc.* 2009; 131:13628–13630. [PubMed: 19731910] i) Maki BE, Chan A, Phillips EM, Scheidt KA. *Tetrahedron.* 2009; 65:3102–3109. [PubMed: 20161230] j) De Sarkar S, Studer A. *Org Lett.* 2010; 12:1992–1995. [PubMed: 20359171]
15. Breslow R, Kim R. *Tetrahedron Lett.* 1994; 35:699–702. Chen YT, Barletta GL, Haghjoo K, Cheng JT, Jordan F. *J Org Chem.* 1994; 59:7714–7722. White MJ, Leeper FJ. *J Org Chem.* 2001; 66:5124–5131. [PubMed: 11463265] Pignataro L, Papalia T, Slawin AMZ, Goldup SM. *Org Lett.* 2009; 11:1643–1646. [PubMed: 19320507] e) a recent NMR study of the Breslow intermediate revealed a structure other than that expected and commonly formulated: Berkessel A, Elfert S, Etzenbach-Effers K, Teles JH. *Angew Chem.* 2010; 122:7063–7063. *Angew Chem Int Ed.* 2010; 49:7120–7124.
16. Schrader W, Handayani PP, Burstein C, Glorius F. *Chem Commun.* 2007:716–718.
17. Zeitler first invoked the intermediacy of  $\alpha,\beta$ -unsaturated acyl azoliums in the NHC-catalyzed redox esterification of ynals to give (*E*)- $\alpha,\beta$ -unsaturated esters. Zeitler K. *Org Lett.* 2006; 8:637–640. [PubMed: 16468730]
18. Our recently reported NHC-catalyzed Coates–Claisen reaction also features  $\alpha,\beta$ -unsaturated acyl azoliums as important intermediate in the catalytic cycle. Kaeobamrung J, Mahatthananchai J, Zheng P, Bode JW. *J Am Chem Soc.* 2010; 132:8810–8812. [PubMed: 20550127]
19. a) Harris TK, Washabaugh MW. *Biochemistry.* 1995; 34:14001–14011. [PubMed: 7577998] b) Harris TK, Washabaugh MW. *Biochemistry.* 1995; 34:13994–14000. [PubMed: 7577997] c) Washabaugh MW, Jencks WP. *Biochemistry.* 1988; 27:5044–5053. [PubMed: 2844248]
20. Carboxylic acids appear to form adducts with triazolium salt **3** through an addition to C-2. This is similar to the behavior of other triazolium salts, see: Enders D, Breuer K, Raabe G, Runsink J, Teles JH, Melder JP, Ebel K, Brode S. *Angew Chem.* 1995; 107:1119–1122. *Angew Chem Int Ed Engl.* 1995; 34:1021–1023. Vora HU, Rovis T. *J Am Chem Soc.* 2010; 132:2860–2861. [PubMed: 20151675]. The formation of the carboxylic acid is suppressed by employing activated molecular sieves
21. Ryan SJ, Candish L, Lupton DW. *J Am Chem Soc.* 2009; 131:14176–14177. [PubMed: 19757788]; b) this process was characterized as a rearrangement in a most recent report: Candish L, Lupton DW. *Org Lett.* 2010; 12:4836–4839. [PubMed: 20873828]

22. Zhu ZQ, Xiao JC. *Adv Synth Catal.* 2010; 352:2455–2458.
23. De Sarkar S, Studer A. *Angew Chem.* 2010; 122:9452–9455. *Angew Chem Int Ed.* 2010; 49:9266–9269.
24. The discussion regarding NHC-catalyzed reaction without added base has been reported in Ref. [18]. Our current understanding is that the counter anion serves the role of base in generating a trace amount of the active NHC species, which initiates the catalytic cycle.
25. For various discussions regarding the interaction of NHCs and protic nucleophiles, see: Kamber NE, Jeong W, Waymouth RM, Pratt RC, Lohmeijer BGG, Hedrick JL. *Chem Rev.* 2007; 107:5813–5840. [PubMed: 17988157] Schmidt MA, Müller P, Movassaghi M. *Tetrahedron Lett.* 2008; 49:4316–4318. Kuhn N, Mallah E, Maichle-Mößmer C, Steimann M. *Z Naturforsch B.* 2009; 64:835–839. Myles L, Gore R, Spulak M, Gathergood N, Connon SJ. *Green Chem.* 2010; 12:1157–1162.
26. The observed non-integral rate orders may be attributed to an inhibition event. The rationalization of the observed rate law is discussed in the rate law derivation (see the Supporting Information). From the derivation, the rate law may be reformulated as:  $\text{rate} = k_{\text{obs}}[\text{ynal}]^1[\text{precatalyst } \mathbf{3}]^1[\text{MeOH}]^0$ . This reformulation changes the activation entropy for the redox esterification reaction to + 3.95 calK<sup>-1</sup> mol<sup>-1</sup> while the activation enthalpy remains unchanged. In any case, the change in  $\Delta S^\ddagger$  does not affect the interpretation of the proposed mechanism.
27. Coates RM, Rogers BD, Hobbs SJ, Curran DP, Peck DR. *J Am Chem Soc.* 1987; 109:1160–1170.
28. For a similar approach to the measurements of the difference in the activation parameters between two competing pathways, see: Nigam M, Platz MS, Showalter BM, Toscano JP, Johnson R, Abbot SC, Kirchhoff MM. *J Am Chem Soc.* 1998; 120:8055–8059.
29. Hansch C, Leo A, Taft RW. *Chem Rev.* 1991; 91:165–195.
30. The redox reaction–protonation step is excluded from being rate limiting based on isotopic labeling experiments, which show no kinetic isotope effect for protonation (see the Supporting Information).






**Figure 1.** <sup>1</sup>H NMR investigation of  $\alpha,\beta$ -unsaturated acyl azolium **1** generated from **3** and **4**, and its methanolysis to **5** (the labeled H atoms correlate to the indicated signals in the spectra).




**Chemical Formula:**  $C_{30}H_{27}ClN_3O_2^+$   
***m/z*:** 496.1793 (found)  
***m/z*:** 496.1792 (calculated)

**Figure 2.** UV/Vis and ESI-HRMS characterizations of **1**. Black: pre-catalyst + aldehyde; gray: acyl azolium.







**Figure 3.**  
Key NMR correlation assignments of acyl azolium **1**.




**Figure 4.** Linear free-energy relationship (Hammett study) for NHC-catalyzed redox reaction with *para*-substituted ynals:  $\rho = -0.69$  ( $Z = \text{CF}_3, \text{Cl}, \text{H}, \text{Me}, \text{or OMe}$  as indicated in the plot).  $\gamma = -0.6901 x + 0.0326$ ,  $R^2 = 0.903$ .




**Scheme 1.**  
Various acyl azoliums and the hemiacetals.

**Scheme 2.**


Reactions of a catalytically generated acyl azolium with various nucleophiles.

**Scheme 3.**

Activation parameters, rate constants, and rate laws comparison between NHC-catalyzed redox and Claisen reactions.

**Scheme 4.**

a) Differential activation parameters measurement from a competition experiment and b) the mechanistic rationale.



**Scheme 5.** Breakdown of hemiacetal **V** is the rate-limiting step in NHC-catalyzed redox esterifications of ynals.