

**SHORT REPORT**

**Open Access**

# Retrotransposition of R2 elements in somatic nuclei during the early development of *Drosophila*

Michael T Eickbush<sup>1,2</sup> and Thomas H Eickbush<sup>1\*</sup>

## Abstract

**Background:** R2 retrotransposable elements exclusively insert in the 28S rRNA genes of their host. Their RNA transcripts are produced by self-processing from a 28S R2 cotranscript. Because full-length R2 transcripts are found in most tissues of R2-active animals, we tested whether new R2 insertions occurred in somatic tissues even though such events would be an evolutionary dead end.

**Findings:** PCR assays were used to identify somatic R2 insertions in isolated adult tissues and larval imaginal discs of *Drosophila simulans*. R2 somatic mosaics were detected encompassing cells from individual tissues as well as tissues from multiple body segments. The somatic insertions had 5' junction sequences characteristic of germline insertions suggesting they represented authentic retrotransposition events.

**Conclusions:** Body segments are specified early in *Drosophila* development, thus the detection of the same somatic insertion in cells from multiple tissues suggested that the R2 retrotransposition events had occurred before the blastoderm stage of *Drosophila* development. R2 activity at this stage, when embryonic nuclei are rapidly dividing in a common cytoplasm, suggests that some retrotransposition events appearing as germline events may correspond to germline mosaicism.

## Findings


Mobile element insertions during the development of somatic tissues provide no benefit to the element, as these insertions are not transferred to subsequent generations. Thus in animals, where the separation of somatic and germline tissues is established early, the ability of a mobile element to generate new insertions in somatic tissues would most likely be selected against. Consistent with this prediction, early studies in *Drosophila melanogaster* showed that P element transpositions were dependent upon a germline-specific RNA splicing component [1], and I elements were only transcribed in ovaries [2]. However, counter to this model, mobile elements in other animals have been shown to generate new insertions in somatic tissues (for example, Tc1 elements in *Caenorhabditis elegans* [3], L1 elements in mammals [4,5]).

Several explanations can be put forward for the somatic activity of mobile elements. First, somatic events are inconsequential to the host and thus there is little selective pressure for a mobile element to evolve specificity to the germline. Second, somatic events are harmful, however it is risky for a mobile element to become dependent on a germline-specific mechanism, as it provides another opportunity for the host to control the element. Third, on occasions somatic events provide a benefit to the host. This last fascinating possibility has been suggested to explain the ability of L1 to retrotranspose in nerve tissues [6].

R2 non-LTR retrotransposable elements specifically insert into the tandemly repeated rRNA genes of many animal genera, Figure 1A[7,8]. Each R2 insertion blocks the production of functional 28S rRNA from the inserted gene. Because animals contain many more rRNA genes than are needed for transcription [9,10], in most individuals inserted rRNA units are simply not transcribed. However, studies in *Drosophila simulans* indicate that in individuals where R2-inserted units are

\* Correspondence: eick@mail.rochester.edu

<sup>1</sup>Department of Biology, University of Rochester, Rochester, NY, USA  
Full list of author information is available at the end of the article



distributed throughout the rDNA locus, inserted rDNA units are transcribed [11-13]. The R2 transcripts are processed from the cotranscript [14], and new germline retrotransposition events can be observed in the progeny. Because rRNA transcription is essential in all tissues, and full-length R2 transcripts are readily detected in most larval and adult tissues of active lines ([12] and D. Eickbush and T. Eickbush, unpublished results), we

tested whether R2 retrotranspositions also occur in somatic tissues. The *D. simulans* stock selected for study, sim89, had high levels of R2 transcripts in many tissues and new insertions could be detected in progeny that had originated in either the male or female parent [12,15]. For each animal the screens for R2 somatic mosaics were conducted with either seven adult tissues (antenna, proboscis, the rest of the head, wing, haltere


and individual legs from different thoracic segments), or with four third instar larval tissues that are precursors to adult tissues (brain, and three pairs of imaginal discs). Larval-specific tissues were not used because they are composed of polytene cells, which under-replicate R2-inserted rDNA units [16]. New insertions were assayed using the same 5' junction PCR assays previously used to detect germline events [11,15,17]. These assays utilize the property that while all R2 insertions occur at the identical location in the 28S genes when monitored from the 3' end, over half of the R2 retrotransposition events result in a deletion of element sequences starting at its 5' end and extending to locations throughout its 3.6 kb length. These 5' truncated copies can also contain short deletions or duplications of upstream 28S gene sequences. As a result somatic R2 insertions containing 5' truncations will generate PCR bands of lengths that seldom match the lengths of the PCR bands derived from the germline inherited 5' truncated elements. The PCR assays were conducted using a single primer located 80 bp upstream of the R2 insertion site in combination with a series of primers to sequences spaced throughout the length of the R2 element (Figure 1B) [12].

Somatic mosaics were defined as the presence of unique PCR bands in only a subset of the tissues tested from a single animal. To be scored as a somatic insertion, each new PCR band also had to be reproducibly detected using two different PCR primer combinations. Examples of an R2 insertion in one tissue of the four tested from a third instar larva, and of another insertion detected in three of seven adult tissues are shown in Figure 1C, D. The PCR bands representing potential somatic events were less intense than the bands derived from the R2 elements inherited from the mother or father, as expected if not all cells of a tissue type contained the insertion. Generally new bands could be reproducibly observed if they corresponded to at least one-tenth the intensity of those bands derived from inherited R2 copies. In total, tissues from 29 individuals (14 females, 15 males) were scored. A total of 15 potential somatic insertions were detected in 7 animals (2 females with 4 total events, and 5 males with 11 total events). The detection of greater numbers of new insertions in males compared to females was likely due to the greater sensitivity of the PCR assay in males. The rRNA genes in *D. simulans* are located on the X chromosome [18]: thus males contain a single rDNA locus, compared to two copies of the rDNA locus in females. The somatic events were detected in essentially all tissues examined, although the numbers of events were not sufficient to make conclusions about relative frequencies.

To confirm that the PCR bands detected in only a subset of tissues corresponded to new R2 insertions

arising from retrotransposition mechanisms similar to that of germline events, PCR bands representing 12 events that were well separated from the germline bands were excised from the gel, reamplified and the product sequenced. As shown in Figure 2 the 5' ends of the somatic R2 insertions had the characteristics associated with germline R2 insertions [19]. First, the 5' junctions of the R2 sequences with the 28S gene occurred at a variety of positions near the R2 insertion site. As with germline insertions, most junctions were within a few base pairs of the insertion site, but a few were found at distances of approximately 25 and 50 nucleotides. Second, seven insertions had microidentities of from 1-3 nucleotides between the R2 element and the upstream 28S sequence (sequences highlighted in blue). These microidentities are suggested to arise by the R2 DNA polymerase (also known as reverse transcriptase) annealing the upstream target DNA of the 28S gene to the newly made cDNA strand to prime second strand DNA synthesis. Microidentities at the 5' junction of truncated copies is a common property of L1 and other non-LTR retrotransposons [20]. Third, in those cases with no microidentity, from 1-9 nucleotides were present at the junction that did not correspond to either the upstream 28S gene or the R2 element (sequences highlighted in orange). These bases are postulated to represent non-templated synthesis by the R2 reverse transcriptase on the second DNA strand cleavage site until a microidentity between these added nucleotides and the cDNA strand enables the polymerase to prime second strand DNA synthesis. In summary, the physical properties of the 5' junctions of the new PCR bands detected in somatic tissues suggest they represent authentic retrotransposition events.

Because the development of *Drosophila* has been intensively investigated, the timing of the retrotransposition events that generated the observed somatic mosaics can be estimated. By mid-embryogenesis (10-12 h), small clusters of cells (10-40 cells) are specified to become individual imaginal discs [21,22]. Each imaginal disc primordium divides during the 3 larval instars to form from 10,000 to 60,000 cells by late larval development [23]. Because the observed somatic events were present in a significant fraction of the cells present in a third instar larval disc or an adult tissue, the retrotransposition events probably occurred before or early in imaginal disc development. Those retrotransposition events detected in more than one disc or adult appendage probably occurred even earlier in development, before determination of body segments at the blastoderm stage (2-3 h). Of the 15 events we observed, 5 were detected in cells derived from more than 1 body segment. Because we surveyed only a fraction of all body segments in either the larvae or adult, it is likely



that a larger fraction of the somatic R2 insertion events we observed occurred before the blastoderm stage. This developmental period corresponds to rapid nuclear division in a common cytoplasm. During this period there is little RNA synthesis but active protein synthesis using the RNA synthesized by the nurse cells and deposited in the oocyte during oogenesis [24]. Because rRNA synthesis also does not occur in these first hours of development [25], R2 retrotransposition events occurring during this time probably use RNA templates synthesized by the nurse cells during oogenesis.

It should be noted that the observed somatic retrotransposition events likely occurred at a time when embryonic nuclei had not yet entered the pole plasma

of the egg to become the germline. Thus in addition to somatic mosaicism there is also likely to be germline mosaicism of R2 elements in *Drosophila*. As a result, re-evaluation of a previous study of retrotransposition in the germline of males and females appears warranted [15]. We have previously suggested that the rate of R2 insertion inherited through the male germline was one-third to one-quarter the rate of insertions through the female germline. Based on the findings in this report, it is possible that all of the insertions scored as inherited through the male germline (that is, during spermatogenesis), actually occurred during early embryogenesis.

Because preblastoderm development in male and female embryos are similar, we suggest the higher rate

of R2 insertions observed through the female germline represents this germline mosaicism as well as authentic germline events during oogenesis. Two separate periods of R2 activity in females was also consistent with experiments to monitor a large fraction of the offspring from individual females. In the most comprehensive study, new insertions were assayed in 213 progeny of a single female [15]. Of the 32 different R2 insertions detected in these progeny, 27 were found in only 1 individual and 4 were detected in 2 individuals. These insertions appeared to have occurred late in the development of the germline (that is, during oogenesis). The final R2 insertion was detected in 13 progeny, and could correspond to an insertion during early development. Additional evidence for germline mosaicism was found in the analysis of progeny from another female in which 6 of 17 individuals contained the same new R2 insertion.

In conclusion, we suggest that R2 elements are active early in *Drosophila* development, and as in the case with L1 elements in mouse and humans [4,5], can lead to both somatic and germline mosaicism. To determine if R2 elements are also active in somatic tissues later in development will require assaying many smaller samples from individual tissues or more sensitive approaches to detect insertions in smaller percentages of cells. Finally, R2 should serve as a reminder in the study of other mobile elements that events early in development can give rise to insertion mosaics that could be misinterpreted as germline events in the subsequent generation.

#### Acknowledgements

The research was supported by funds from the National Institutes of Health grant GM42790. The authors would like to thank B Burke for help with the DNA sequencing, D Eickbush for comments on the manuscript, and M Welte for discussions of early *Drosophila* development.

#### Author details

<sup>1</sup>Department of Biology, University of Rochester, Rochester, NY, USA. <sup>2</sup>Fred Hutchinson Cancer Research Center, Seattle, WA, USA.

#### Authors' contributions

MTE helped design the experiments, conducted all the experiments, and help perfect the manuscript. THE helped design the experiments and wrote the first draft of the manuscript. Both authors read and approved the final manuscript.

#### Competing interests

The authors declare that they have no competing interests.

Received: 8 July 2011 Accepted: 29 September 2011

Published: 29 September 2011

#### References

- Laski FA, Rio DC, Rubin GM: Tissue specificity of *Drosophila* P element transposition is regulated at the level of mRNA splicing. *Cell* 1986, **44**:7-19.
- Chaboissier MC, Busseau I, Prosser J, Finnegan DJ, Bucheton A: Identification of a potential RNA intermediate for transposition of the LINE-like element I factor in *Drosophila melanogaster*. *EMBO J* 1990, **9**:3557-3563.
- Emmons SW, Yesner L: High-frequency excision of transposable element Tc1 in the nematode *Caenorhabditis elegans* is limited to somatic cells. *Cell* 1984, **36**:599-605.
- van den Hurk JA, Meij IC, Seleme MC, Kano H, Nikopoulos K, Hoefsloot LH, Siermans EA, de Wijs IJ, Mukhopadhyay A, Plomp AS, de Jong PTVM, Kazazian HH Jr, Cremers FPM: L1 retrotransposition can occur early in human embryonic development. *Hum Mol Genet* 2007, **16**:1587-1592.
- Kano H, Godoy I, Courtney C, Vetter MR, Gerton GL, Ostertag EM, Kazazian HH Jr: L1 retrotransposition occurs mainly in embryogenesis and creates somatic mosaicism. *Genes Dev* 2009, **23**:1303-1312.
- Muotri AR, Chu VT, Marchetto MCN, Deng W, Moran JV, Gage FH: Somatic mosaicism in neuronal precursor cells mediated by L1 retrotransposition. *Nature* 2005, **435**:903-910.
- Eickbush TH: R2 and related site-specific non-LTR retrotransposons. In *Mobile DNA II*. Edited by: Craig N, Craigie R, Gellert M, Lambowitz A. Washington DC: American Society of Microbiology Press; 2002:813-835.
- Kojima KK, Fujiwara H: Long-term inheritance of the 28S rDNA-specific retrotransposon R2. *Mol Biol Evol* 2005, **22**:2157-2165.
- Conconi A, Widmer AR, Koller T, Sogo JM: Two different chromatin structures coexist in ribosomal RNA genes throughout the cell cycle. *Cell* 1989, **57**:753-761.
- Ye J, Eickbush TH: Chromatin structure and transcription of R1- and R2-inserted rRNA genes of *Drosophila melanogaster*. *Mol Cell Biol* 2006, **26**:8781-8790.
- Zhang X, Eickbush TH: Characterization of active R2 retrotransposition in the rDNA locus of *Drosophila simulans*. *Genetics* 2005, **170**:195-205.
- Eickbush DG, Ye J, Zhang X, Burke WD, Eickbush TH: Epigenetic regulation of retrotransposons within the nucleolus of *Drosophila*. *Mol Cell Biol* 2008, **28**:6452-6461.
- Zhou J, Eickbush TH: The pattern of R2 retrotransposon activity in natural populations of *Drosophila simulans* reflects the dynamic nature of the rDNA locus. *PLoS Genetics* 2009, **5**:e1000386.
- Eickbush DG, Eickbush TH: R2 retrotransposons encode a self-cleaving ribozyme for processing from an rRNA co-transcript. *Mol Cell Biol* 2010, **30**:3142-3150.
- Zhang X, Zhou J, Eickbush TH: Rapid R2 retrotransposition leads to the loss of previously inserted copies via large deletions of the rDNA locus. *Mol Biol Evol* 2008, **25**:229-237.
- Endow SA, Glover DM: Differential replication of ribosomal gene repeats in polytene nuclei of *Drosophila*. *Cell* 1979, **17**:597-605.
- Perez-Gonzalez CE, Eickbush TH: Dynamics of R1 and R2 elements in the rDNA locus of *Drosophila*. *Genetics* 2001, **158**:1557-1567.
- Lohe AR, Roberts PA: An unusual Y chromosome of *Drosophila simulans* carrying amplified rDNA spacer without rRNA genes. *Genetics* 1990, **125**:399-406.
- Stage DE, Eickbush TH: Origin of nascent lineages and the mechanisms used to prime second-strand DNA synthesis in the R1 and R2 retrotransposons of *Drosophila*. *Genome Biol* 2009, **10**:R49.
- Ostertag EM, Kazazian HH Jr: Biology of mammalian L1 retrotransposons. *Annu Rev Genet* 2001, **35**:501-538.
- Technau GM: A single cell approach to problems of cell lineage and commitment during embryogenesis of *Drosophila melanogaster*. *Development* 1987, **100**:1-12.
- Cohen SM: Imaginal disc development. In *The Development of Drosophila melanogaster*. Volume 2. Edited by: Bate M, Arias AM. Cold Spring Harbor, NY, USA: Cold Spring Harbor Laboratory Press; 1993.
- Klebes A, Biehs B, Cifuentes F, Kornberg TB: Expression profiling of *Drosophila* imaginal discs. *Genome Biol* 2002, **3**:RESEARCH0038.
- Nasiadka A, Dietrich BH, Krause HM: Anterior-posterior patterning in the *Drosophila* embryo. *Adv Develop Biol Biochem* 2002, **12**:155-186.
- McKnight SL, Miller OL Jr: Ultrastructural patterns of RNA synthesis during early embryogenesis of *Drosophila melanogaster*. *Cell* 1976, **8**:305-319.

doi:10.1186/1759-8753-2-11

**Cite this article as:** Eickbush and Eickbush: Retrotransposition of R2 elements in somatic nuclei during the early development of *Drosophila*. *Mobile DNA* 2011 **2**:11.