

# Evolution of the *S*-Locus Region in *Arabidopsis* Relatives<sup>1[C][W]</sup>

Ya-Long Guo<sup>2\*</sup>, Xuan Zhao<sup>2</sup>, Christa Lanz, and Detlef Weigel

Department of Molecular Biology, Max Planck Institute for Developmental Biology, 72076 Tuebingen, Germany

The *S* locus, a single polymorphic locus, is responsible for self-incompatibility (SI) in the Brassicaceae family and many related plant families. Despite its importance, our knowledge of *S*-locus evolution is largely restricted to the causal genes encoding the *S*-locus receptor kinase (SRK) receptor and *S*-locus cysteine-rich protein (SCR) ligand of the SI system. Here, we present high-quality sequences of the genomic region of six *S*-locus haplotypes: *Arabidopsis* (*Arabidopsis thaliana*; one haplotype), *Arabidopsis lyrata* (four haplotypes), and *Capsella rubella* (one haplotype). We compared these with reference *S*-locus haplotypes of the self-compatible *Arabidopsis* and its SI congener *A. lyrata*. We subsequently reconstructed the likely genomic organization of the *S* locus in the most recent common ancestor of *Arabidopsis* and *Capsella*. As previously reported, the two SI-determining genes, *SCR* and *SRK*, showed a pattern of coevolution. In addition, consistent with previous studies, we found that duplication, gene conversion, and positive selection have been important factors in the evolution of these two genes and appear to contribute to the generation of new recognition specificities. Intriguingly, the inactive pseudo-*S*-locus haplotype in the self-compatible species *C. rubella* is likely to be an old *S*-locus haplotype that only very recently became fixed when *C. rubella* split off from its SI ancestor, *Capsella grandiflora*.

Self-incompatibility (SI) is an important mechanism used by many angiosperm species for preventing self-fertilization (selfing), which can be considered as an evolutionary dead end because it limits adaptive potential and causes inbreeding depression by the expression of recessive deleterious mutations (Lynch et al., 1995; Charlesworth et al., 2005; Newbigin and Uyenoyama, 2005; Busch and Schoen, 2008). SI is often controlled by a single polymorphic locus, the *S* locus. In the Brassicaceae, the *S* locus contains two specificity-determinant genes: *S-LOCUS RECEPTOR KINASE* (*SRK*), which encodes the stigmatic receptor kinase; and *SP11/S-LOCUS CYSTEINE-RICH PROTEIN* (*SCR*) (hereafter *SCR*), which encodes a small Cys-rich protein localized in the pollen coat, which is the ligand for the *SRK* receptor (Nasrallah, 2002). Pollen inhibition occurs when the same *S*-locus specificity is expressed by both pollen and pistil (Nasrallah, 2002). The *S* locus is exceptional, with its many transspecies polymorphisms (Charlesworth, 2006), providing a paradigm for

frequency-dependent selection (Schierup and Vekemans, 2008; Leducq et al., 2011).

Transformation of *Arabidopsis* (*Arabidopsis thaliana*) with *S*-locus haplotypes from the self-incompatible *Arabidopsis lyrata* can restore SI in some accessions (Nasrallah et al., 2002). Therefore, it is likely that the initial inactivation of the *S* locus was a key step in the transition from SI to self-compatibility (SC) in this species. *Arabidopsis* is the model plant among SC species. The SI system is inactive in the selfing *Arabidopsis* reference strain Columbia (Col-0), and both *SRK* and *SCR* become pseudogenized (Kusaba et al., 2001). The inactivation of the *S* locus is correlated with the transition from outcrossing to selfing in both *Arabidopsis* and the *Arabidopsis* relative *Capsella rubella* (Tang et al., 2007; Guo et al., 2009; Tsuchimatsu et al., 2010). Transitions to SC are frequent in plants. This transition has occurred independently many times in several lineages (Barrett, 2002, 2010). In the Brassicaceae, most tribes include selfing species (Fobis-Loisy et al., 2004).

The molecular mechanism of action and functional polymorphisms at the *S* locus and in the SI system have been described extensively for *Arabidopsis* and related genera (Uyenoyama, 2000; Schierup et al., 2001, 2006; Kachroo et al., 2002; Sato et al., 2002; Takebayashi et al., 2003; Nasrallah et al., 2004; Prigoda et al., 2005; Bechsgaard et al., 2006; Paetsch et al., 2006; Liu et al., 2007; Mable and Adam, 2007; Sherman-Broyles et al., 2007; Tang et al., 2007; Busch et al., 2008; Shimizu et al., 2008; Guo et al., 2009; Castric et al., 2010; Tsuchimatsu et al., 2010).

Because of the difficulties with recovering the exceedingly diverse sequences at this locus by PCR,

<sup>1</sup> This work was supported by ERA-NET on Plant Genomics grant ARelatives from the Deutsch Forschungsgemeinschaft (to D.W.) and by the Max Planck Society (to D.W.).


<sup>2</sup> These authors contributed equally to the article.

\* Corresponding author; e-mail ya-long.guo@hotmail.com.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors ([www.plantphysiol.org](http://www.plantphysiol.org)) is: Detlef Weigel ([weigel@weigelworld.org](mailto:weigel@weigelworld.org)).

<sup>[C]</sup> Some figures in this article are displayed in color online but in black and white in the print edition.

<sup>[W]</sup> The online version of this article contains Web-only data. [www.plantphysiol.org/cgi/doi/10.1104/pp.111.174912](http://www.plantphysiol.org/cgi/doi/10.1104/pp.111.174912)



**Figure 1.** Comparison of *S*-locus haplotypes. **A**, Dot plot highlighting sequence similarity and rearrangement between each *S*-locus haplotype and *Ath-Col*. **B**, A VISTA plot indicating similarity between *Ath-Col* and the other *S*-locus haplotypes. **C**, A comparison of gene content among different *S*-locus haplotypes. Red arrows indicate the *SRK*-, *SCR*-, and *ARK3*-related truncated sequences. Red *N* symbols represent gaps, and black *N* symbols represent large fragments not drawn to scale (actual length in kb is given below). [See online article for color version of this figure.]

most analyses of the *S* locus have focused on *SCR* and *SRK*, and sometimes only on a partial *S* domain of the *SRK* gene. Other genes within the *S*-locus region have rarely been comprehensively examined (Kamau and

Charlesworth, 2005; Hagenblad et al., 2006; Kamau et al., 2007; Ruggiero et al., 2008). There is, however, strong suppression of recombination and linked loci, which may influence the evolution of SI systems

(Uyenoyama, 2005). Recently, it has been demonstrated that pseudo-SC can be caused by an S-locus-linked gene that can regulate *SRK* transcript levels (Liu et al., 2007). Detailed knowledge about S-locus sequences from additional relatives of Arabidopsis is also necessary to determine whether the S-locus polymorphisms in selfing species, such as Arabidopsis and *C. rubella*, reflect alleles that were present in a self-incompatible ancestor or whether they were produced by differential degradation of the S locus in different self-fertile populations (Sherman-Broyles et al., 2007). The age of S-locus allelic lineages in Arabidopsis relatives has thus far not been conclusively determined. Some allelic lineages at the S locus in *A. lyrata* are very old (Schierup et al., 2001; Paetsch et al., 2006; Edh et al., 2009): it is likely that they predate the divergence between the class I and class II allele lineages in *Brassica*, estimated to have occurred 40 million years ago (Uyenoyama, 1995). Finally, broader knowledge of S-locus sequence diversity is required for understanding how new specificities evolve in a two-gene system that relies on matching specificities between the two genes.

To address these questions, two factors must be considered. First, it is important to compare S-locus haplotypes between related SI and SC species. Second, both key genes and linked genes at the S locus should be analyzed. Therefore, we sequenced bacterial artificial chromosome clones (BACs) covering the S-locus region, including the core region from At4g21350 (*PLANT U-BOX8* [*PUB8*]) to At4g21380 (*ARK3*), and several genes on either side, in two accessions of the selfing species Arabidopsis (Ath-Cvi) and *C. rubella* (Cru) and from four chromosomes of the outcrossing species *A. lyrata* (Aly-S16, Aly-S38, Aly-S50, and Aly-Sb). One additional S-locus haplotype each had already been sequenced and therefore was available from reference genome sequencing projects in Arabidopsis (Ath-Col; Arabidopsis Genome Initiative, 2000) and *A. lyrata* (Aly-Sa, from reference strain MN47; Hu et al., 2011). Our comparison of the eight S-locus haplotypes provides new insights into the evolution of the S locus in the Brassicaceae.

## RESULTS

### Genomic Organization and Gene Content of the S Locus

For the scale of phylogenetic sampling, we covered geographic (*A. lyrata* from Europe and North America) and mating system (SC species Arabidopsis and *C. rubella*, SI species *A. lyrata*) diversity. There was only one BAC library of *C. rubella* available when we started this work a few years ago; thus, only a single *C. rubella* haplotype was sequenced. Six BACs covering the core S-locus region, from At4g21350 (*PUB8*) to At4g21380 (*ARK3*), were sequenced: Arabidopsis (one S-locus haplotype), *A. lyrata* (four S-locus haplotypes), and *C. rubella* (one S-locus haplotype). For *SRK* and *SCR*, the published *SRKa/SCRa* or *SRKb/SCRb* (Kusaba et al., 2001) is the same S-locus haplotype as Aly-Sa or Aly-Sb, respectively. Only for Aly-Sb, we could not completely fill all assembly gaps. Nevertheless, the four large contigs of Aly-Sb contained both *SCR* and *SRK* in addition to the other flanking genes of the S locus. All six sequenced S-locus haplotypes share the region from At4g21326 to At4g21430, except Aly-S50, where the sequence from exon 1 to part of exon 6 of At4g21326 is missing (Fig. 1C). A phylogenetic tree (Supplemental Fig. S1) indicated that the recovered S-locus haplotypes span several S-locus lineages across Arabidopsis and *Capsella* relatives.

Comparing the six sequenced S-locus haplotypes and the Arabidopsis and *A. lyrata* reference genome S-locus haplotypes, the length of the eight S-locus haplotypes (from At4g21326 to At4g21430) ranges from 56.1 kb (Ath-Cvi) to 116.5 kb (Aly-Sb; Supplemental Table S1). Sequences on either side of the core S-locus region are highly conserved and syntenic, while the core region is highly diverged (Fig. 1, A and B). Gene order and orientation are conserved (Fig. 1, B and C), except for Aly-Sa, which has an inversion including *SCR* and *SRK* (Fig. 1), as well as Aly-S50 and Aly-Sb, which have an inversion of the *SCR* gene (Fig. 1C). It is likely that in the majority of S-locus haplotypes, there is one *SCR* gene and one *SRK* gene in head-to-head orientation.

The 11 protein-coding genes found in the Arabidopsis reference S-locus haplotype Ath-Col are conserved in all S-locus haplotypes. Aly-S16, Aly-S50, Aly-Sa,




Figure 2. Alignment of *SCR*-related sequences.

and Aly-Sb, however, have one or two extra genes, which are not shared with any of the other *S*-locus haplotypes (Fig. 1C; Supplemental Table S1). In addition, there are some partially duplicated genes, including *SCR*-like sequences (*SCR2* and *SCR3*) and an *SRK*-like gene (*At4g21366*) in *Ath-Col*, an *SCR3*-like fragment in *Ath-Cvi*, and an *ARK3*-like fragment in *Aly-S38*.

The content of transposable elements in different *S*-locus haplotypes varies greatly, from 5.45% in *Ath-Cvi* to 31.27% in *Aly-Sb* (Supplemental Tables S2 and S3). *Arabidopsis* *S*-locus haplotypes have the fewest transposable elements, while *Cru* is intermediate. The number and insertion times of long terminal repeat (LTR) retrotransposons also differ (Supplemental Table S2).

### Transspecific Allele Sharing and Coevolution

In *Ath-Col*, *Aly-S38*, *Aly-S50*, and *Cru* (Fig. 2; Table I), *SCR* is represented by a truncated open reading frame (ORF), as inferred from deduced amino acid sequences. In *Ath-Col*, *Ath-Cvi*, and *Cru*, the *SRK* genes also have truncated ORFs (Kusaba et al., 2001; Tang et al., 2007; Shimizu et al., 2008; Guo et al., 2009). The amino acid sequences of the *SCR* gene are highly diverged, but the eight conserved Cys residues (Kusaba et al., 2001) were found in each *S*-locus haplotype with complete ORF (Fig. 2).


The topologies of the phylogenetic trees for *SCR* and *SRK* match. *Ath-Cvi* and *Aly-S16*, as well as *Aly-S38* and *Cru*, show close relationships, supporting a pattern of coevolution for *SCR* and *SRK* (Fig. 3; Supplemental Figs. S1 and S2). None of the other genes at the *S* locus showed the same transspecies pattern (Supplemental Fig. S3), not even *At4g21380* (*ARK3*), which is less than 3 kb away from the *SRK* gene in most *S*-locus haplotypes. Thus, linkage disequilibrium is apparently weak, and limited to a narrow region, from *SCR* to *SRK*.

The length of the single intron in *SCR* can be very different, even between closely related *S*-locus haplotypes such as *Ath-Cvi* (2,036 bp) and *Aly-S16* (1,044 bp; Table I). The length of the intergenic region between *SRK* and *SCR*, for *SCR* with only the first exon, which was calculated based on the distance from the end of

**Table I.** The *SCR* direction, intron length, and distance from *SRK*

–1 indicates that only the first exon or part of it was found. NA indicates that the distance from *SCR* to *SRK* cannot be estimated for the gaps within *Aly-Sb*.

BAC	Direction	Intron Length	Distance from <i>SRK</i>
			bp
<i>Ath-Col</i>	+	782	710
<i>Ath-Cvi</i>	+	2,036	227
<i>Aly-S16</i>	+	1,044	255
<i>Aly-S38</i>	+	–1	3,812
<i>Aly-S50</i>	–	–1	5,249
<i>Aly-Sa</i>	–	1,519	1,751
<i>Aly-Sb</i>	–	78	NA
<i>Cru</i>	+	–1	14,422



**Figure 3.** Matching phylogenetic trees of the *SCR* and *SRK* sequences.

the first exon of *SCR* to *SRK*, is also highly variable, from 227 bp in *Ath-Cvi* to 14,422 bp in *Cru* (Table I).

### Gene Conversion

There are frequent duplications in the core *S*-locus region, including duplications of *SCR*, *SRK*, and *ARK3*. *SRK* and *ARK3* are closely related in sequence, and five potential gene conversion events were detected, with the size of the converted fragment ranging from 31 to 391 bp (Table II). Gene conversion also appears to have occurred between *SRK* and *ARK3* in both *Aly-Sb* and *Ath-Col* as well as between *SRK*-like (*At4g21366*) and *SRK* and between *SRK*-like (*At4g21366*) and *ARK3* in *Ath-Col*. Affected were exon 7 of *SRK* and *ARK3*, except for the gene conversion event between *At4g21366* and *ARK3* in *Ath-Col*, which included exons 6 and 7.

### Evidence for Positive Selection

We tested whether the evolutionary divergence of the *S* locus has been affected by positive selection. Because pseudogenes confound such tests and show different evolutionary patterns than functional genes, we excluded them from our analyses. The ratio of nonsynonymous and synonymous mutation ( $d_N/d_S$ ) averaged across all sites and lineages ranged from 0.11 (*At4g21350*) to 1.21 (*SCR*; Table III). For three genes, *SCR*, *SRK*, and *ARK3*, the positive selection model fit the data significantly better than the neutral model (Table III), although the *SCR* gene had a much higher proportion of codons under positive selection than *SRK* and *ARK3* (Fig. 4).

## DISCUSSION

### Genomic Organization and Gene Content of the *S* Locus in the *Arabidopsis/Capsella* Lineage

Although the core *S*-locus region between the *PUB8* gene and *ARK3* is very fluid and entails many inde-

**Table II.** Gene conversions in *SRK*- and *ARK3*-related sequences

Species	Sequence 1	Sequence 2	<i>P</i> <sup>a</sup>	Begin <sup>b</sup>	End <sup>c</sup>	Length <sup>d</sup>
Arabidopsis	At4g21366	SRK_Ath-Col	0.0000	4,145	4,560	391
	ARK3_Ath-Col	At4g21366	0.0034	4,392	4,422	31
	ARK3_Ath-Col	At4g21366	0.0450	4,516	4,546	31
	ARK3_Ath-Col	SRK_Ath-Col	0.0407	4,392	4,422	31
<i>A. lyrata</i>	ARK3_Aly-Sb	SRK_Aly-Sb	0.0015	4,490	4,532	43

<sup>a</sup>Simulated *P* values of the global inner fragments obtained from 10,000 permutations. <sup>b</sup>First nucleotide of the converted region. <sup>c</sup>Last nucleotide of the converted region. <sup>d</sup>Length of the converted region without alignment gaps.

pendent insertions, duplications, and inversions, genes flanking the outermost *PUB8* and *ARK3* genes were found to be conserved with respect to gene orientation and order in the eight analyzed *S*-locus haplotypes. The more common head-to-head orientation of *SCR* and *SRK*, however, is changed in *Aly-S50*, *Aly-Sb*, and *Ath-C24* (Fig. 5). *Aly-Sa* has an inversion including the *SCR-SRK* region, but the relative orientation is similar to the other *S*-locus haplotypes. We conclude that the ancestral organization of the *S* locus in the *Arabidopsis/Capsella* lineage appears to be one *SCR* gene and one *SRK* gene in head-to-head orientation, flanked by the *PUB8* and *ARK3* genes transcribed from the same strand as *SRK* (Fig. 5). The extra genes, which are not shared with any of the other *S*-locus haplotypes, must have originated independently in specific lineages.

### Transspecies Polymorphism and Coevolution

Transspecies patterns of polymorphism are typical for plant SI systems (Sato et al., 2002; Bechsgaard et al., 2006; Paetsch et al., 2006), a fungal SI system (May

et al., 1999), and the vertebrate animal major histocompatibility complex system (Adams et al., 2000). In the *Brassica* genus, some *S*-locus haplotypes have been estimated to be about 20 to 40 million years old (Uyenoyama, 1995). In the Solanaceae, the *S*-RNase, which is the female specificity determinant of SI in this family, has been estimated to be nearly 70 million years old (Xue et al., 1996). Some allelic lineages of the major histocompatibility complex system of animals appear to have persisted for at least 20 million years (Klein et al., 1998). Because the *S*-locus genes appear to be under selection, one cannot easily determine the age of the *S*-locus haplotype with a molecular clock. The close relationship of the *Ath-Cvi* and *Aly-S16* haplotypes between *Arabidopsis* and *A. lyrata* as well as of the *Aly-S38* and *Cru* haplotypes between *A. lyrata* and *C. rubella* (Paetsch et al., 2006) suggest that these *S*-locus haplotypes are at least 13 and 22 million years old, respectively (Fig. 5).


In the Brassicaceae, recombination events disrupting the linkage of matched *SRK* and *SCR* alleles are expected to result in the breakdown of SI. The pattern of coevolution between *SRK* and *SCR* suggests that they

**Table III.** Evidence for adaptive evolution of genes located in the *S*-locus region

M0 is the  $d_N/d_S$  ratio averaged across all sites and lineages. M7 versus M8 indicates the outcome of the likelihood test.  $-1$  means that the value of  $2\Delta$  (twice the log likelihood difference between two models) is 0 or less than 0.0001 and there is no significant difference between the two models. Estimates of parameters indicate the proportion of amino acids predicted to be under adaptive evolution (and their corresponding  $d_N/d_S$  ratio) from model M8. NA indicates not applicable because there is no significant difference between the positive selection model and the neutral model. Positively selected sites are amino acids under positive selection with Bayesian posterior  $P > 0.95$  based on model M8 using the Bayes Empirical Bayes method. Ns is the number of sequences in the analysis. LA indicates the codon in the alignment. For the *SRK* gene, *Aly-S16* is the reference sequence, and for the *SCR* gene, *Ath-Cvi* is the reference sequence. For *ARK3* and other genes, *Ath-Col* is the reference. \*  $P < 0.05$ , \*\*  $P < 0.001$ , \*\*\*  $P < 0.0001$ .

Gene	M0	M7 Versus M8 ( $2\Delta$ )	Estimates of Parameters	Positively Selected Sites	Ns	LA
At4g21326	0.198	-1	NA	NA	8	700
At4g21330	0.217	3.894	NA	NA	8	210
At4g21340	0.312	0.069	NA	NA	8	309
At4g21350 ( <i>PUB8</i> )	0.106	-1	NA	NA	8	374
<i>SCR</i>	1.210	7.896*	44.2% ( $\omega = 2.955$ )	32L, 34K, 43T, 48R, 58H, 63K, 67V	4	95
At4g21370 ( <i>SRK</i> )	0.340	26.137***	4.3% ( $\omega = 3.300$ )	275R	5	868
At4g21380 ( <i>ARK3</i> )	0.198	11.882*	0.3% ( $\omega = 9.127$ )	None with $P > 0.95$	8	854
At4g21390	0.113	-1	NA	NA	8	851
At4g21400	0.325	-1	NA	NA	7	718
At4g21410	0.185	0.195	NA	NA	7	680
At4g21430	0.314	2.965	NA	NA	8	940

**Figure 4.** Positive selection and variation in  $d_N/d_S$  among codons. A, *SRK*. B, *SCR*. C, *ARK3*. Single asterisks represent codons with a Bayesian posterior probability of positive selection greater than 75%, and double asterisks represent codons with a Bayesian posterior probability of positive selection greater than 95%.




are in strong linkage disequilibrium, consistent with the previous population genetics study in *A. lyrata* (Kamau et al., 2007). In *Brassica*, the tree topology of *SRK* and *SCR* is basically similar (Sato et al., 2002; Edh et al., 2009). The structural and sequence heterogeneity between *SCR* and *SRK* sequences is likely the most important mechanism suppressing recombination and maintaining SI. This conclusion is strongly supported by results from Castric et al. (2010), who found that recombination could occur when individuals were homozygous for recessive alleles, indicating that there is no general suppression of recombination in this region. It is interesting that coevolution and linkage are limited to a really small region around *SCR* and *SRK*.

**Duplication, Gene Conversion, Positive Selection, and Evolution of New Specificities**

There are several hypotheses for the generation of new SI specificities (Uyenoyama et al., 2001; Chookajorn et al., 2004; Charlesworth et al., 2005). In one hypothesis (Uyenoyama et al., 2001), a self-compatible intermediate is generated in which partial breakdown of SI

occurs because of a mutation in one of the SI specificity determinants. Subsequent compensatory mutations in the other specificity determinant then cause the pistil to reject the new pollen type, thus restoring the SI system. In the outcrossing species *A. lyrata*, many self-compatible accessions have been found in North America. SC, therefore, is likely to have originated multiple times (Mable et al., 2005; Mable and Adam, 2007; Mable, 2008; Foxe et al., 2010), and these self-compatible haplotypes might also be precursors of new SI haplotypes. Another model suggests that new SI specificities arise through gradual modification of *SRK-SCR* affinities (Chookajorn et al., 2004). We found that *SCR* appears to be under much stronger diversifying selection than *SRK*, consistent with a structure-function analysis of *SCR* that showed this protein to be unusually tolerant to sequence changes (Chookajorn et al., 2004), which suggests that *SCR* is the major engine for the modification of *SRK-SCR* affinities.

An additional factor might be gene conversion, which has been found in *Brassica* between *SLG* and *SRK* (Sato et al., 2002; Fujimoto et al., 2006) and between *SRK* and its possible paralogs in *A. lyrata*



**Figure 5.** Proposed history of the *S* locus. Divergence times (million years ago [MYA]) on the right are calculated based on the synonymous substitutions among the *ADH* gene orthologs extracted from the reference genomes (Col-0 and MN47) and GenBank (AF110435; *C. rubella*), using the spontaneous mutation rate estimated in *Arabidopsis* (Ossowski et al., 2010). The divergence time between *A. lyrata* and *Arabidopsis* is consistent with a recent study using fossil evidence (Beilstein et al., 2010). Gray arrows indicate truncated ORFs. Letters A to F represent inserted genes (relative to *Ath-Col*): A, AT4G04400-like; B, AT5G32590-like; C, AT4G07320-like; D, AT3G42410-like; E, AT3G13270-like; F, AT2G07760-like.

(Charlesworth et al., 2003). We have identified gene conversion events among *SRK*, *ARK3*, and duplicated *SRK*-like (At4g21366) genes. Interestingly, we found that gene conversion was largely restricted to the portion of the region encoding the kinase domain. Although the kinase domain of the *SRK* protein does not seem to directly contribute to the specificity of pollen/pistil recognition (Rea et al., 2010), it might indirectly influence the specificity of *SRK* by affecting protein conformation. In addition, we found that only three out of 11 genes (*SCR*, *SRK*, and *ARK3*) in the *S*-locus region show evidence of positive selection. Previous studies have found many sites in the *S* domain of *SRK* to be under positive selection in these species (Sainudiin et al., 2005; Castric and Vekemans, 2007). We also found that *ARK3*, which is the *S*-domain receptor-like kinase that is most closely related to *SRK*, has a history of gene duplication, gene conversion, and positive selection. These features of *ARK3* could be a motor for the evolution of *SRK*, given the apparent gene conversion between *ARK3* and *SRK* that we found here. In summary, there is a strong indication that duplication, gene conversion, and positive selection all contribute to the evolution of new *S*-locus specificities, as has been suggested previously.

#### Relationship between the *S* Locus and Selfing

One hypothesis for the labile nature of SI is that the transition from outcrossing to selfing can provide a

selective advantage at a given time and place (Stebbins, 1957; Busch et al., 2010; Goldberg et al., 2010; Wright and Barrett, 2010). A single inactivated *S*-locus haplotype was fixed in *C. rubella*, representing the key step in the recent transition to selfing for this species (Guo et al., 2009). In contrast, *SC* has apparently originated several times independently in *Arabidopsis* (Bechsgaard et al., 2006; Sherman-Broyles et al., 2007; Tang et al., 2007; Shimizu et al., 2008; Boggs et al., 2009b; Tsuchimatsu et al., 2010). In accessions with a Col-0-like *SRK* allele, *SCR* became pseudogenized after the gene was disrupted by an inversion (Boggs et al., 2009b; Tsuchimatsu et al., 2010). In the accession Cvi-0, *SCR* is intact but *SRK* has a truncated ORF (Tang et al., 2007; Shimizu et al., 2008; this work). Both *SCR* and *SRK* have truncated ORFs in Col-0 (Kusaba et al., 2001). C24, which also has a truncated *SRK* ORF, lacks *SCR* altogether (Sherman-Broyles et al., 2007). In *C. rubella*, the ORFs of all *SCR* alleles and some *SRK* alleles are truncated (Guo et al., 2009). Interestingly, one *A. lyrata* haplotype, Aly-S38, is very similar to *C. rubella*, with a closely related *SCR* with a truncated ORF and an *SRK* with a complete ORF (Fig. 3; Supplemental Fig. S1). This indicates that the truncated *S*-locus haplotype in *C. rubella* could be very old, despite it having been fixed only very recently. Although the simultaneous maintenance of both nonfunctional and functional *S*-locus haplotypes is rare (Uyenoyama et al., 2001; Porcher and Lande, 2005), two of the *S*-locus haplotypes we re-



covered for *A. lyrata* appear to contain a *SCR* allele with a truncated ORF, which spans two different *S*-locus lineages across Arabidopsis and *Capsella* relatives (Supplemental Fig. S1). It will be interesting to investigate the activity of those *S*-locus haplotypes with truncated ORFs (such as Aly-S38 and Aly-S50), given that the *SCR* gene of *Cvi* may not be functional even though it has a complete ORF (Boggs et al., 2009a).

Unfortunately, the sequences of Aly-S38 and Aly-S50 come from a BAC library generated a decade ago, and the individual plants with these haplotypes are no longer available. Because of the great diversity of *S*-locus haplotypes, finding the same haplotypes in extant samples would be very difficult, if not impossible. On the other hand, the distinction of functional/nonfunctional *S*-locus haplotypes was not a main focus of our study. The strength of our work is that it is, to our knowledge, the first sequencing study of the *S*-locus region in such a large number of Arabidopsis relatives. With additional high-quality sequences of the *S*-locus region from *A. lyrata* and other related species, one can address the mechanisms of the evolution of the *S*-locus region, determine whether *S*-locus haplotypes with truncated ORFs are functional or not, and also address the question of the age of inactive *S*-locus haplotypes with more confidence.

## CONCLUSION

Our results are consistent with previous studies of the *S* locus in that we found that the two SI-determining genes, *SCR* and *SRK*, show patterns of coevolution. Duplication, gene conversion, and positive selection have been important factors in the evolutionary history of these two genes and appear to contribute to the generation of new recognition specificities. An unexpected result is that the inactive pseudo-*S*-locus haplotype in the SC species *C. rubella* might be an old *S*-locus haplotype that only very recently became fixed when *C. rubella* split off from its SI ancestor, *Capsella grandiflora*. Finally, our data provide not only an important resource for understanding *S*-locus evolution but can also guide future functional studies of the SI system.

## MATERIALS AND METHODS

### BAC Screen, Sequencing, and Assembly

We generated a 400-bp probe derived from *PUB8* (At4g21350) by PCR amplification of Arabidopsis (*Arabidopsis thaliana*) Col-0 using primers G10720 (5'-TTAATCTCCACTCTCATCTCTC-3') and G10721 (5'-CTGATTCCTTCTCTCCCTATC-3') and used it to screen BAC libraries of Arabidopsis *Cvi*-0, *Arabidopsis lyrata*, and *Capsella rubella* by filter hybridization. The *A. lyrata* individuals used in constructing the BAC library were from the Pech area in Bavaria (Germany), and the *C. rubella* individuals were from Gargano, Italy. End sequencing was used to map the borders of positive BAC clones, and clones that include the region from At4g21326 to At4g21430 were analyzed by shotgun sequencing.

For Arabidopsis *Cvi*-0, BAC35E6 (Ath-*Cvi*) was analyzed. For the out-crossing *A. lyrata*, we first sequenced part of the *SRK* locus (using G12957 primer; 5'-CTTCGAGCTTGTTTCTTCA-3') to differentiate different *S*-locus haplotypes. We then used these *SRK* sequences and other available *SRK* sequences from GenBank to construct a phylogenetic tree for the different *S*-locus haplotypes. Among 16 positive BAC clones, there were three distinct *S*-locus haplotypes (named after the similarity to well-known *SRK* sequences of *A. lyrata*) besides Aly-Sb, located on the *SRKb* BAC of *A. lyrata* reported by Kusaba and colleagues (2001), hereafter represented as Aly-Sb, and Aly-Sa from the reference strain MN47 (Supplemental Fig. S1). Both Aly-Sa and Aly-Sb are derived from accessions of *A. lyrata* collected in Michigan (Kusaba et al., 2001). In the end, four independent *A. lyrata* BAC clones, BAC6K22 (Aly-S16), BAC39O17 (Aly-S38), BAC27M5 (Aly-S50), and Aly-Sb (Kusaba et al., 2001), were selected for sequencing. From the *C. rubella* library, we selected BAC15P16 (Cru).

BAC DNA was prepared using the Large Construct DNA Preparation Kit (Qiagen), physically sheared, and shotgun libraries were constructed using the TOPO Shotgun Subcloning Kit (Invitrogen). Shotgun clones were sequenced on an ABI 3730XL DNA Analyzer using T7 and SP6 primers. Sequences were processed using Phred for base calling, CROSS\_MATCH for removing vector sequences, and Phrap and Consed for assembly (Ewing and Green, 1998; Ewing et al., 1998; Gordon et al., 1998). Primer walking and PCR were used to fill or polish gaps and ambiguous regions. The sequences and the assembled contigs were in accordance with the Bermuda standards of sequencing (<http://www.genome.gov/10001812>). The *S*-locus region of Col-0 was extracted from the Arabidopsis reference genome (<http://www.arabidopsis.org/>), and the *S*-locus region of the MN47 strain of *A. lyrata* was extracted from the reference genome as well (<http://genome.jgi-psf.org/Araly1/Araly1.home.html>).

### Sequence Annotation and Phylogenetic Analysis

Annotation of the BAC sequences was performed using FGENESH (<http://linux1.softberry.com/berry.phtml>). The annotation was confirmed by comparison with Arabidopsis (The Arabidopsis Information Resource 8). The *SCR* gene was annotated by comparing it with *SCR1*, *SCR2*, and *SCR3* of Arabidopsis (Col-0), *SCRa* (AB052753) and *SCRb* (AB052754) of *A. lyrata*, *SP11-6* (AF195625) of *Brassica oleracea*, and *SP11-9* of *Brassica rapa* (AB022078). Spidey (<http://www.ncbi.nlm.nih.gov/spidey/>) was used to confirm the annotation based on the alignment to Arabidopsis mRNA when needed. VISTA (<http://genome.lbl.gov/vista/index.shtml>) and Mummer version 3.19 (Kurtz et al., 2004) were used to compare and view BAC sequences.

Sequences were aligned using ClustalX version 1.81 (Thompson et al., 1997) and then refined manually. PAUP\* version 4.0b10 (Swofford, 2003) was used to reconstruct the phylogenetic tree. For the maximum parsimony tree, a heuristic search was performed with the MULPARS option, tree bisection-reconnection branch swapping. The neighbor-joining trees were constructed using the two-parameter model (Kimura, 1980; Saitou and Nei, 1987). Topological robustness was assessed by bootstrapping with 1,000 replicates (Felsenstein, 1985).

### Transposable Element Analysis

RepeatMasker version 3.2.5 (<http://www.repeatmasker.org/>) was used to detect the content of transposable elements using the Arabidopsis library of Repbase (RM database version 20080611). LTR\_STRUC (McCarthy and McDonald, 2003) was used to identify full-length LTR retrotransposons. Each LTR retrotransposon was classified (gypsy-like, copia-like, others) based on BLASTN and T-BLAST searches of the Brassicaceae repeat database (<http://tigrblast.tigr.org/euk-blast/index.cgi?project=plant.repeats>). The sequence divergence between the two LTRs of each intact LTR retrotransposon was used to estimate the insertion time ( $T$ ) using the formula  $T = K/(2)(r)$ , where  $K$  is the number of substitutions per site between two LTRs, estimated with the two-parameter model using the distmat program implemented in the EMBOSS package (<http://emboss.sourceforge.net/>), and  $r$  is the spontaneous mutation rate estimated in Arabidopsis,  $7 \times 10^{-9}$  mutations per site per generation (year; Ossowski et al., 2010).

### Detection of Gene Conversion

Gene conversion was analyzed using GENECONV version 1.81a (Sawyer, 1989) with default settings. Based on an alignment of DNA sequences,



GENECONV detects gene conversion by looking for sufficiently similarly aligned segments between a pair of sequences. A global value of  $P \leq 0.05$  was used as a criterion for further analysis of the conversion tracts detected by GENECONV. Given that GENECONV could not distinguish between gene conversion and unequal crossing over, significant GENECONV results are indicated as conversion events or tracts (Mondragon-Palomino and Gaut, 2005).

### Positive Selection Detection Using Likelihood Ratio Tests

Potentially positive selection was analyzed with the program codeml of PAML version 3.15 (Yang, 1997). The first step was to find sites with  $\omega$  ( $d_N/d_S$ )  $> 1$  by comparing a null model (M7) that does not allow for sites with  $\omega > 1$  and a general model that does (M8) by performing a likelihood ratio test using codeml. If model M7 was rejected in favor of model M8, we took this as evidence against the  $\omega$  ratio being confined to the interval (0, 1) for all sites. The second step was to use a likelihood ratio test to identify residues (with Bayesian posterior probability) under positive selection with the Bayes Empirical Bayes tool (Yang et al., 2005). We assumed that sites with  $\omega > 1$  and a high probability ( $P \geq 95\%$ ) were likely to be under positive selection.

Sequence data from this article can be found in the GenBank/EMBL data libraries under accession numbers HQ376928-HQ376932 and EF637083.

### Supplemental Data

The following materials are available in the online version of this article.

**Supplemental Figure S1.** Selection of BAC clones for shotgun sequencing based on the phylogeny of *SRK* using the most parsimonious method.

**Supplemental Figure S2.** A single most parsimonious tree based on *SCR* gene amino acid sequences.

**Supplemental Figure S3.** Phylogenetic trees using the neighbor-joining method with two-parameter model based on the coding regions of nine protein-coding genes flanking the S locus, from At4g21326 to At4g21430, and the *SRK* gene to detect transspecies polymorphism.

**Supplemental Table S1.** BAC information and gene density.

**Supplemental Table S2.** Repeat content and full-length LTR retrotransposon content and insertion times.

**Supplemental Table S3.** Transposable element content in S-locus haplotypes.

### ACKNOWLEDGMENTS

We thank June B. Nasrallah for giving the *A. lyrata* Aly-Sb BAC clone; Maarten Koornneef for providing access to the Cvi-0 BAC library; Norman Warthmann, Kirsten Bomblies, Stefan Henz, and Namita Tripathi for technical advice; Stanley A. Sawyer for helpful suggestions on the usage of GENECONV for gene conversion analysis; Beth Rowan for greatly improving the manuscript; and Carmen Melchers for proofreading.

Received February 22, 2011; accepted August 1, 2011; published August 2, 2011.

### LITERATURE CITED

- Adams EJ, Cooper S, Thomson G, Parham P (2000) Common chimpanzees have greater diversity than humans at two of the three highly polymorphic MHC class I genes. *Immunogenetics* **51**: 410–424
- Arabidopsis Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**: 796–815
- Barrett SC (2002) The evolution of plant sexual diversity. *Nat Rev Genet* **3**: 274–284
- Barrett SC (2010) Understanding plant reproductive diversity. *Philos Trans R Soc Lond B Biol Sci* **365**: 99–109

- Bechsgaard JS, Castric V, Charlesworth D, Vekemans X, Schierup MH (2006) The transition to self-compatibility in *Arabidopsis thaliana* and evolution within S-haplotypes over 10 Myr. *Mol Biol Evol* **23**: 1741–1750
- Beilstein MA, Nagalingum NS, Clements MD, Manchester SR, Mathews S (2010) Dated molecular phylogenies indicate a Miocene origin for *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **107**: 18724–18728
- Boggs NA, Dwyer KG, Shah P, McCulloch AA, Bechsgaard J, Schierup MH, Nasrallah ME, Nasrallah JB (2009a) Expression of distinct self-incompatibility specificities in *Arabidopsis thaliana*. *Genetics* **182**: 1313–1321
- Boggs NA, Nasrallah JB, Nasrallah ME (2009b) Independent S-locus mutations caused self-fertility in *Arabidopsis thaliana*. *PLoS Genet* **5**: e1000426
- Busch JW, Joly S, Schoen DJ (2010) Does mate limitation in self-incompatible species promote the evolution of selfing? The case of *Leavenworthia alabamica*. *Evolution* **64**: 1657–1670
- Busch JW, Schoen DJ (2008) The evolution of self-incompatibility when mates are limiting. *Trends Plant Sci* **13**: 128–136
- Busch JW, Sharma J, Schoen DJ (2008) Molecular characterization of Lal2, an SRK-like gene linked to the S-locus in the wild mustard *Leavenworthia alabamica*. *Genetics* **178**: 2055–2067
- Castric V, Bechsgaard JS, Grenier S, Noureddine R, Schierup MH, Vekemans X (2010) Molecular evolution within and between self-incompatibility specificities. *Mol Biol Evol* **27**: 11–20
- Castric V, Vekemans X (2007) Evolution under strong balancing selection: how many codons determine specificity at the female self-incompatibility gene *SRK* in Brassicaceae? *BMC Evol Biol* **7**: 132
- Charlesworth D (2006) Balancing selection and its effects on sequences in nearby genome regions. *PLoS Genet* **2**: e64
- Charlesworth D, Bartolomé C, Schierup MH, Mable BK (2003) Haplotype structure of the stigmatic self-incompatibility gene in natural populations of *Arabidopsis lyrata*. *Mol Biol Evol* **20**: 1741–1753
- Charlesworth D, Vekemans X, Castric V, Glémin S (2005) Plant self-incompatibility systems: a molecular evolutionary perspective. *New Phytol* **168**: 61–69
- Chookajorn T, Kachroo A, Ripoll DR, Clark AG, Nasrallah JB (2004) Specificity determinants and diversification of the *Brassica* self-incompatibility pollen ligand. *Proc Natl Acad Sci USA* **101**: 911–917
- Edh K, Widén B, Ceplitis A (2009) The evolution and diversification of S-locus haplotypes in the Brassicaceae family. *Genetics* **181**: 977–984
- Ewing B, Green P (1998) Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res* **8**: 186–194
- Ewing B, Hillier L, Wendl MC, Green P (1998) Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res* **8**: 175–185
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**: 783–791
- Fobis-Loisy I, Miede C, Gaude T (2004) Molecular evolution of the S locus controlling mating in the Brassicaceae. *Plant Biol (Stuttg)* **6**: 109–118
- Foxe JP, Stift M, Tedder A, Haudry A, Wright SI, Mable BK (2010) Reconstructing origins of loss of self-incompatibility and selfing in North American *Arabidopsis lyrata*: a population genetic context. *Evolution* **64**: 3495–3510
- Fujimoto R, Sugimura T, Nishio T (2006) Gene conversion from *SLG* to *SRK* resulting in self-compatibility in *Brassica rapa*. *FEBS Lett* **580**: 425–430
- Goldberg EE, Kohn JR, Lande R, Robertson KA, Smith SA, Igić B (2010) Species selection maintains self-incompatibility. *Science* **330**: 493–495
- Gordon D, Abajian C, Green P (1998) Consed: a graphical tool for sequence finishing. *Genome Res* **8**: 195–202
- Guo YL, Bechsgaard JS, Slotte T, Neuffer B, Lascoux M, Weigel D, Schierup MH (2009) Recent speciation of *Capsella rubella* from *Capsella grandiflora*, associated with loss of self-incompatibility and an extreme bottleneck. *Proc Natl Acad Sci USA* **106**: 5246–5251
- Hagenblad J, Bechsgaard J, Charlesworth D (2006) Linkage disequilibrium between incompatibility locus region genes in the plant *Arabidopsis lyrata*. *Genetics* **173**: 1057–1073
- Hu TT, Pattyn P, Bakker EG, Cao J, Cheng JF, Clark RM, Fahlgren N, Fawcett JA, Grimwood J, Gundlach H, et al (2011) The *Arabidopsis lyrata* genome sequence and the basis of rapid genome size change. *Nat Genet* **43**: 476–481
- Kachroo A, Nasrallah ME, Nasrallah JB (2002) Self-incompatibility in the

- Brassicaceae: receptor-ligand signaling and cell-to-cell communication. *Plant Cell (Suppl)* **14**: S227–S238
- Kamau E, Charlesworth B, Charlesworth D** (2007) Linkage disequilibrium and recombination rate estimates in the self-incompatibility region of *Arabidopsis lyrata*. *Genetics* **176**: 2357–2369
- Kamau E, Charlesworth D** (2005) Balancing selection and low recombination affect diversity near the self-incompatibility loci of the plant *Arabidopsis lyrata*. *Curr Biol* **15**: 1773–1778
- Kimura M** (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* **16**: 111–120
- Klein J, Sato A, Nagl S, O’Huigin C** (1998) Molecular trans-species polymorphism. *Annu Rev Ecol Syst* **29**: 1–21
- Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL** (2004) Versatile and open software for comparing large genomes. *Genome Biol* **5**: R12
- Kusaba M, Dwyer K, Hendershot J, Vrebalov J, Nasrallah JB, Nasrallah ME** (2001) Self-incompatibility in the genus *Arabidopsis*: characterization of the S locus in the outcrossing *A. lyrata* and its autogamous relative *A. thaliana*. *Plant Cell* **13**: 627–643
- Leducq JB, Llaurens V, Castric V, Saumitou-Laprade P, Hardy OJ, Vekemans X** (2011) Effect of balancing selection on spatial genetic structure within populations: theoretical investigations on the self-incompatibility locus and empirical studies in *Arabidopsis halleri*. *Heredity* **106**: 319–329
- Liu P, Sherman-Broyles S, Nasrallah ME, Nasrallah JB** (2007) A cryptic modifier causing transient self-incompatibility in *Arabidopsis thaliana*. *Curr Biol* **17**: 734–740
- Lynch M, Conery J, Bürger R** (1995) Mutation accumulation and the extinction of small populations. *Am Nat* **146**: 489–518
- Mable BK** (2008) Genetic causes and consequences of the breakdown of self-incompatibility: case studies in the Brassicaceae. *Genet Res (Camb)* **90**: 47–60
- Mable BK, Adam A** (2007) Patterns of genetic diversity in outcrossing and selfing populations of *Arabidopsis lyrata*. *Mol Ecol* **16**: 3565–3580
- Mable BK, Robertson AV, Dart S, Di Berardo C, Witham L** (2005) Breakdown of self-incompatibility in the perennial *Arabidopsis lyrata* (Brassicaceae) and its genetic consequences. *Evolution* **59**: 1437–1448
- May G, Shaw F, Badrane H, Vekemans X** (1999) The signature of balancing selection: fungal mating compatibility gene evolution. *Proc Natl Acad Sci USA* **96**: 9172–9177
- McCarthy EM, McDonald JF** (2003) LTR\_STRUC: a novel search and identification program for LTR retrotransposons. *Bioinformatics* **19**: 362–367
- Mondragon-Palomino M, Gaut BS** (2005) Gene conversion and the evolution of three leucine-rich repeat gene families in *Arabidopsis thaliana*. *Mol Biol Evol* **22**: 2444–2456
- Nasrallah JB** (2002) Recognition and rejection of self in plant reproduction. *Science* **296**: 305–308
- Nasrallah ME, Liu P, Nasrallah JB** (2002) Generation of self-incompatible *Arabidopsis thaliana* by transfer of two S locus genes from *A. lyrata*. *Science* **297**: 247–249
- Nasrallah ME, Liu P, Sherman-Broyles S, Boggs NA, Nasrallah JB** (2004) Natural variation in expression of self-incompatibility in *Arabidopsis thaliana*: implications for the evolution of selfing. *Proc Natl Acad Sci USA* **101**: 16070–16074
- Newbigin E, Uyenoyama MK** (2005) The evolutionary dynamics of self-incompatibility systems. *Trends Genet* **21**: 500–505
- Ossowski S, Schneeberger K, Lucas-Lledó JI, Warthmann N, Clark RM, Shaw RG, Weigel D, Lynch M** (2010) The rate and molecular spectrum of spontaneous mutations in *Arabidopsis thaliana*. *Science* **327**: 92–94
- Paetsch M, Mayland-Quellhorst S, Neuffer B** (2006) Evolution of the self-incompatibility system in the Brassicaceae: identification of S-locus receptor kinase (SRK) in self-incompatible *Capsella grandiflora*. *Heredity* **97**: 283–290
- Porcher E, Lande R** (2005) Loss of gametophytic self-incompatibility with evolution of inbreeding depression. *Evolution* **59**: 46–60
- Prigoda NL, Nassuth A, Mable BK** (2005) Phenotypic and genotypic expression of self-incompatibility haplotypes in *Arabidopsis lyrata* suggests unique origin of alleles in different dominance classes. *Mol Biol Evol* **22**: 1609–1620
- Rea AC, Liu P, Nasrallah JB** (2010) A transgenic self-incompatible *Arabidopsis thaliana* model for evolutionary and mechanistic studies of crucifer self-incompatibility. *J Exp Bot* **61**: 1897–1906
- Ruggiero MV, Jacquemin B, Castric V, Vekemans X** (2008) Hitch-hiking to a locus under balancing selection: high sequence diversity and low population subdivision at the S-locus genomic region in *Arabidopsis halleri*. *Genet Res (Camb)* **90**: 37–46
- Sainudiin R, Wong WS, Yogeewaran K, Nasrallah JB, Yang Z, Nielsen R** (2005) Detecting site-specific physicochemical selective pressures: applications to the class I HLA of the human major histocompatibility complex and the SRK of the plant sporophytic self-incompatibility system. *J Mol Evol* **60**: 315–326
- Saitou N, Nei M** (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**: 406–425
- Sato K, Nishio T, Kimura R, Kusaba M, Suzuki T, Hatakeyama K, Ockendon DJ, Satta Y** (2002) Coevolution of the S-locus genes SRK, SLG and SP11/SCR in *Brassica oleracea* and *B. rapa*. *Genetics* **162**: 931–940
- Sawyer S** (1989) Statistical tests for detecting gene conversion. *Mol Biol Evol* **6**: 526–538
- Schierup MH, Bechsgaard JS, Nielsen LH, Christiansen FB** (2006) Selection at work in self-incompatible *Arabidopsis lyrata*: mating patterns in a natural population. *Genetics* **172**: 477–484
- Schierup MH, Mable BK, Awadalla P, Charlesworth D** (2001) Identification and characterization of a polymorphic receptor kinase gene linked to the self-incompatibility locus of *Arabidopsis lyrata*. *Genetics* **158**: 387–399
- Schierup MH, Vekemans X** (2008) Genomic consequences of selection on self-incompatibility genes. *Curr Opin Plant Biol* **11**: 116–122
- Sherman-Broyles S, Boggs N, Farkas A, Liu P, Vrebalov J, Nasrallah ME, Nasrallah JB** (2007) S locus genes and the evolution of self-fertility in *Arabidopsis thaliana*. *Plant Cell* **19**: 94–106
- Shimizu KK, Shimizu-Inatsugi R, Tsuchimatsu T, Purugganan MD** (2008) Independent origins of self-compatibility in *Arabidopsis thaliana*. *Mol Ecol* **17**: 704–714
- Stebbins GL** (1957) Self-fertilization and population variability in the higher plants. *Am Nat* **41**: 337–354
- Swofford DL** (2003) PAUP\*: Phylogenetic Analysis Using Parsimony (\*and Other Methods), Version 4. Sinauer Associates, Sunderland, MA
- Takebayashi N, Brewer PB, Newbigin E, Uyenoyama MK** (2003) Patterns of variation within self-incompatibility loci. *Mol Biol Evol* **20**: 1778–1794
- Tang C, Toomajian C, Sherman-Broyles S, Plagnol V, Guo YL, Hu TT, Clark RM, Nasrallah JB, Weigel D, Nordborg M** (2007) The evolution of selfing in *Arabidopsis thaliana*. *Science* **317**: 1070–1072
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG** (1997) The ClustalX Windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **24**: 4876–4882
- Tsuchimatsu T, Suwabe K, Shimizu-Inatsugi R, Isokawa S, Pavlidis P, Städler T, Suzuki G, Takayama S, Watanabe M, Shimizu KK** (2010) Evolution of self-compatibility in *Arabidopsis* by a mutation in the male specificity gene. *Nature* **464**: 1342–1346
- Uyenoyama MK** (1995) A generalized least-squares estimate for the origin of sporophytic self-incompatibility. *Genetics* **139**: 975–992
- Uyenoyama MK** (2000) Evolutionary dynamics of self-incompatibility alleles in *Brassica*. *Genetics* **156**: 351–359
- Uyenoyama MK** (2005) Evolution under tight linkage to mating type. *New Phytol* **165**: 63–70
- Uyenoyama MK, Zhang Y, Newbigin E** (2001) On the origin of self-incompatibility haplotypes: transition through self-compatible intermediates. *Genetics* **157**: 1805–1817
- Wright SI, Barrett SC** (2010) Evolution: the long-term benefits of self-rejection. *Science* **330**: 459–460
- Xue Y, Carpenter R, Dickinson HG, Coen ES** (1996) Origin of allelic diversity in *Antirrhinum* S locus RNases. *Plant Cell* **8**: 805–814
- Yang Z** (1997) PAML: a program package for phylogenetic analysis by maximum likelihood. *Comput Appl Biosci* **13**: 555–556
- Yang Z, Wong WS, Nielsen R** (2005) Bayes empirical Bayes inference of amino acid sites under positive selection. *Mol Biol Evol* **22**: 1107–1118