



Published in final edited form as:

*Eur J Cancer*. 2012 March ; 48(5): 728–736. doi:10.1016/j.ejca.2011.06.022.

## Child Maltreatment and Breast Cancer Survivors: Social Support Makes a Difference for Quality of Life, Fatigue, and Cancer Stress

Christopher P. Fagundes<sup>a,\*</sup>, Monica E. Lindgren<sup>a,b</sup>, Charles L. Shapiro<sup>c</sup>, and Janice K. Kiecolt-Glaser<sup>a,e,f</sup>

<sup>a</sup>Institute for Behavioral Medicine Research, The Ohio State University College of Medicine, The Ohio State University, Columbus, OH

<sup>b</sup>Department of Psychology, The Ohio State University, Columbus, OH

<sup>c</sup>Department of Internal Medicine, The Ohio State University College of Medicine, The Ohio State University, Columbus, OH

<sup>d</sup>Department of Molecular Virology, Immunology and Medical Genetics, Department of Internal Medicine, The Ohio State University College of Medicine, The Ohio State University, Columbus, OH

<sup>e</sup>Comprehensive Cancer Center, The Ohio State University College of Medicine, The Ohio State University, Columbus, OH

<sup>f</sup>Department of Psychiatry, The Ohio State University College of Medicine, The Ohio State University, Columbus, OH

### Abstract

**Purpose**—To identify how child maltreatment is associated with quality of life (QOL) among breast cancer survivors.

**Patients and Methods**—One hundred and thirty two women who had completed treatment for stage 0-IIIa breast cancer within the past two years (except for tamoxifen/aromatase inhibitors) and were at least two months post surgery, radiation, or chemotherapy completed questionnaires including the Childhood Trauma Questionnaire, the Impact of Events Scale, the Multidimensional Fatigue Symptom Inventory-Short Form (MFSI-SF), and the Fact-B breast cancer quality of life questionnaire.

**Results**—Women who were abused or neglected as children reported more cancer-related psychological distress, more fatigue, and poorer physical, emotional, functional, and breast cancer specific well-being after treatment. These relations were partially explained by the fact that breast cancer survivors reported receiving less support as adults.

---

© 2011 Elsevier Ltd. All rights reserved.

\*Christopher P. Fagundes, Ph.D., American Cancer Society Postdoctoral Fellow, Institute for Behavioral Medicine Research, The Ohio State University, 460 Medical Center Drive, Room 144B, Columbus, OH 43210.

Conflict of Interest Statement: The authors have no financial interests or relationships that pose potential conflicts of interest with this article.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Conclusion**—The findings suggest that child maltreatment is an important predictor of QOL among breast cancer survivors. One reason why this association exists is because those who are maltreated as children report less support as adults. A better understanding of how child maltreatment contributes to breast cancer survivor QOL will help in tailoring and therefore enhancing the efficacy of interventions aimed at improving QOL.

Being diagnosed and treated for cancer is emotionally and physically challenging.<sup>1</sup> Breast cancer treatment contributes to mental and physical health problems.<sup>2</sup> Even when treatment-related problems subside, many breast cancer survivors report quality of life (QOL) difficulties including psychological distress, fatigue, occupational disruption, and loss of physical functioning.<sup>3</sup> Clinically, understanding why some breast cancer survivors are more vulnerable to poorer QOL after treatment than others is important.

Women who experienced past traumas are at increased risk for psychological distress when confronted with new traumatic experiences.<sup>4</sup> Breast cancer patients who experienced a serious accident, illness, or death of a close loved one during the year before their diagnosis were more likely to develop PTSD symptoms.<sup>5</sup> Breast cancer survivors who reported severe emotional, physical, or sexual trauma over the course of their lifetime were more susceptible to cancer related emotional distress than those who did not have these experiences.<sup>6</sup> Holocaust survivors experienced significantly more psychological distress than non-Holocaust survivors after a cancer experience.<sup>7</sup>

Child maltreatment is a common experience; approximately 50% of adults report experiencing some type of abuse or neglect as children.<sup>8</sup> Those who were abused or neglected as children are more susceptible to a host of mental and physical health problems in adulthood, especially following a life threatening experience.<sup>9</sup> For example, war veterans with a history of childhood abuse were more likely to have PTSD than their non-abused counterparts.<sup>10</sup> Child abuse has also been linked to somatic symptoms in otherwise healthy people.<sup>11</sup>

Women who have experienced abuse or neglect as children may be at increased risk for poorer QOL after a cancer experience. Newly diagnosed breast cancer patients who were emotionally abused as children had more psychological distress compared to those who were not abused.<sup>12</sup> Breast cancer patients who recalled one or more forms of abuse as children were more likely to experience emotional difficulties two days after cancer surgery.<sup>13</sup>

In sum, child maltreatment has been linked to cancer-related psychological distress. However, we do not know if child maltreatment also contributes to other QOL factors affecting breast cancer survivors such as fatigue, occupational disruption, loss of physical functioning, and problems specifically related to breast cancer. Furthermore, we do not know the mechanisms underlying why child maltreatment leads to these poorer QOL outcomes.<sup>13</sup>

Social support plays an important role in the QOL of cancer survivors.<sup>14</sup> Cancer survivors who report receiving less social support have poorer mental health outcomes than those who report receiving more social support.<sup>15</sup> Breast cancer survivors who received less support from family reported higher levels of depressive symptoms, less positive and hopeful outlooks for the future, less marital satisfaction, less self-esteem, lower levels of role functioning, more sexual problems, and higher levels of hostility than those who reported more support.<sup>16–19</sup>

People who were abused or neglected as children report receiving less social support as adults.<sup>20</sup> Children who have troubled relationships with parents and other adults are less

likely to develop social and emotional skills that are crucial for establishing supportive close relationships in adulthood.<sup>21</sup> Compared to people with positive early relationships, those with troubled early relationships are more likely to report receiving less social support later in life.<sup>22</sup> Accordingly, social support may play an important role linking child maltreatment to the QOL of breast cancer survivors.

The current study examined relationships between child maltreatment and QOL in breast cancer survivors. We hypothesized that those who experienced neglect or abuse as children would have more cancer-related distress, fatigue, and poorer QOL after breast cancer treatment. We also hypothesized that these associations would be partially explained by the fact that those who experienced neglect or abuse as children would report receiving less social support as adults.

## Participants

The study data were drawn from the baseline sample of 132 women who participated in a clinical trial addressing the potential benefits of yoga for breast cancer survivors. Participants were recruited through breast cancer clinics and media announcements. Eligible women had completed treatment for stage 0-IIIa breast cancer within the past two years (except for tamoxifen/aromatase inhibitors) and were at least two months post surgery, radiation, or chemotherapy (whichever occurred last). Screening exclusions included a prior history of breast or any other cancer except basal or squamous cell, more than five hours a week of vigorous physical exercise, a body mass index (BMI) of 40 or greater, diabetes, chronic obstructive pulmonary disease, uncontrolled hypertension, evidence of liver or kidney failure, and symptomatic ischemic heart disease. The Ohio State Biomedical Research Review Committee approved the project; all subjects gave written informed consent prior to participation.

## Measures

In order to assess cancer-related psychological distress, we used the 15-item **Impact of Events Scale (IES)**, which assessed women's avoidant and intrusive thoughts about the cancer experience.<sup>23</sup> The current investigation used the total score. Cronbach's alpha was .88.

The **Functional Assessment of Cancer Therapy-Breast (FACT-B)** is a self-report inventory that provides a multidimensional assessment of QOL.<sup>24</sup> The items assess 5 areas of well-being (physical, social/family, emotional, and functional), while 19 breast-cancer-specific items include breast cancer-related emotional concerns (e.g., worried about cancer risk in family members, worried about the effects of stress on illness), physical concerns (e.g., feeling short of breath, being bothered by swollen/tender arms), body image, and sexual functioning. Widely used in oncology trials and clinical practice, extensive data support its reliability and validity.<sup>24, 25</sup> For the purpose of this study, we adopted the physical, emotional, functional, and cancer-specific scales. We excluded the social/family scale given its considerable conceptual and measurement overlap with the ISEL.

The **Multidimensional Fatigue Symptom Inventory-Short Form (MFSI-SF)** is a 30-item scale that assesses five dimensions of cancer-related fatigue.<sup>26</sup> The total score represents the sum of four subscales (general fatigue, physical fatigue, emotional fatigue, and mental fatigue) minus the vigor scale. Alphas for individual subscales ranged from .86-.92. Alpha for the total score was .90.

The **Interpersonal Support Evaluation List (ISEL)** provided a comprehensive measure of perceived social support.<sup>27</sup> Items are rated on a four-point scale (i.e. definitely false,

probably false, probably true, and definitely true). The ISEL measures the perceived availability of the following kinds of support: emotional (someone to confide in), belonging (people with whom one can do things with), self-esteem (positive social comparison), and tangible (provision of material aid). For the current analyses, we used the total ISEL score. Alpha was .93.

**The Charlson index**<sup>28</sup>, the most widely used comorbidity index for predicting mortality, was used to assess comorbidities. The measure assigns weights to 19 comorbid conditions based on their potential influence on one-year mortality in breast cancer patients. Originally developed for predicting mortality in breast cancer patients, it has now been widely used with both cancer and noncancer populations.<sup>29</sup>

The **Childhood Trauma Questionnaire** provided data on early childhood abuse and neglect. Widely used, it has excellent normative data for its 5 scales: Physical, Sexual, and Emotional Abuse, and Physical and Emotional Neglect.<sup>30</sup> We adopted the Walker cuts<sup>8</sup> to make categorical cut-offs (with sensitivity and specificity >.85 for each scale). Then, we created a categorical indicator variable representing any maltreatment (exceeding  $\geq$  CTQ cut point threshold), and a continuous variable representing number of maltreatment categories.<sup>8</sup>

## Analytic Method

Using separate ordinary least squares general linear models, we first addressed the question of whether child maltreatment predicted cancer-related psychological distress, as well as the following Fact-B QOL subscales: physical well-being, emotional well-being, functional well-being, and the breast cancer subscale. We modeled child maltreatment as categorical (1= any abuse or neglect, 0= no abuse or neglect), and continuous (the number of abusive or neglectful categories) across separate analyses. For all significant associations between child maltreatment and adjustment outcomes, we examined whether social support mediated the association. We used Barron and Kenny's<sup>31</sup> four step regression approach to establish mediation. First, the initial variable (i.e. child maltreatment) should be associated with the outcome. Second, the initial variable (i.e. child maltreatment) should be associated with the mediator (i.e. perceived support). Third, the mediator variable (i.e. perceived support) should be associated with the outcome. Fourth, the association between the initial variable and the outcome variable should be reduced when the mediator is added to the model with the initial variable. Subsequent research on mediation have revealed that only steps 2 and 3 are essential for partial mediation to exist as long as there is a significant mediated effect.<sup>32</sup> In order to test whether there was a significant mediated effect (indirect effect), we employed bias-correct bootstrap estimates (2000) to obtain a confidence interval and a corresponding p-value. Bias-correct bootstrapping is superior to the traditional sobel test for testing indirect effects.<sup>33</sup> All independent variables were grand mean centered. We examined residuals to confirm that they distributed normally.

All models were adjusted for age, cancer stage, and time since last treatment. Time since last treatment was highly correlated with time since diagnosis ( $r = .90 p < .001$ ), accordingly we could not put both in the model simultaneously. In ancillary analyses, we controlled for cancer treatment rather than cancer stage; none of the results presented below changed.

## Results

Table 1 reports descriptive information for all participants. Almost half (48%) of our sample had at least one form of maltreatment, consistent with the broader literature on child abuse and neglect.<sup>8</sup> Maltreated and non-maltreated participants did not differ by treatment type, cancer stage, time since diagnosis, time since last treatment, or age. Less than 5% of our sample had any Charlson-rated comorbidities other than their breast cancer diagnosis, and

thus we did not control for them in our analyses. In ancillary analyses that included Charlson scores, the models did not substantially change.

We first present results for the analyses when maltreatment was modeled categorically. As seen in table 2, those who were abused or neglected as children had more cancer-related psychological distress (as indexed by the IES), more fatigue, poorer physical, emotional, functional, and breast cancer specific well-being. Second, the hypothesized mediator, social support, also predicted all of these outcomes. Third, those who were abused or neglected as children had lower social support. Finally, in every regression, when social support was included in the same regression model as child neglect/abuse, the association between child neglect/abuse and the QOL outcome was attenuated. When social support was added to the model along with child maltreatment, the associations between child maltreatment and emotional well-being, physical well-being, functional well being, and breast cancer specific well being were attenuated to non-significance. Importantly, in all six models, the bootstrap procedure showed the indirect effect of social support was significant. Accordingly, social support partially mediated the association between child maltreatment and each outcome.

We then present data from the analyses when maltreatment was modeled continuously. As can be seen in Table 3, those who experienced more types of abuse/neglect as children had more cancer-specific psychological distress, more fatigue, and poorer physical, emotional, functional, and breast cancer specific well-being. Second, the hypothesized mediator, social support, also predicted all of these outcomes. Third, those who experienced more types of abuse/neglect as children had lower social support. Finally, in every regression model, when social support was included in the same regression model as child neglect/abuse, the association between child neglect/abuse and the QOL outcome was attenuated. Importantly, in all six models, the bootstrap procedure showed the indirect effect of social support was significant. Accordingly, social support partially mediated the association between child maltreatment and each outcome.

## Discussion

With more women surviving breast cancer, health professionals have focused on why some breast cancer survivors are more vulnerable to poorer post-treatment QOL than others.<sup>34</sup> The current study examined relationships between child maltreatment and QOL among breast cancer survivors. Those who were abused or neglected as children experienced more cancer-specific psychological distress, more fatigue, and poorer physical, emotional, functional, and breast cancer specific well-being after treatment. Those who were maltreated as children also reported receiving less social support, and those who had reported receiving less social support also had poorer QOL across all of the aforementioned components. Furthermore, social support partially explained the link between child maltreatment and these quality of life outcomes.

The association between child abuse/neglect and each QOL outcome is notable. Previous studies have shown that childhood abuse predicts PTSD and emotional distress after a traumatic life event.<sup>12, 13</sup> In addition to replicating these associations, we also demonstrated relationships between maltreatment and fatigue, and poorer physical, emotional, functional, and breast cancer specific well being. This suggests that child abuse/neglect affects facets of breast cancer survivor QOL beyond emotional distress. Health care professionals should devote increased attention to a breast cancer patient's abuse history when addressing both emotional *and* somatic problems.

The finding that child maltreatment predicted fatigue is particularly notable. Fatigue is the most common problem among long-term cancer survivors<sup>35</sup>, as well as the symptom that

interferes most with their daily life.<sup>3</sup> Fatigue adversely affects overall QOL.<sup>36</sup> In general, neither disease type nor treatment variables have demonstrated reliable associations with fatigue in cancer survivors. Thus, understanding the psychological characteristics that predict cancer-specific fatigue is important.

Our findings also show that those who were maltreated as children report receiving less social support, which contributes to the association between child maltreatment and QOL outcomes. Improving women's social support networks is one of the best documented ways to improve breast cancer survivor QOL.<sup>37</sup> Future work examining whether interventions targeting those with a history of child maltreatment should differ from general support interventions is needed.

Child maltreatment and social support may have implications beyond QOL. Epidemiological research has linked lower levels of social support with greater breast cancer mortality.<sup>14</sup> For example, in a study of 2,835 breast cancer survivors, women who reported less social support before diagnosis were two times as likely to die of breast cancer over a 10 year period compared with women who had greater support.<sup>38</sup> Furthermore, in a recent study of over 13,000 adults, those who were physically abused as children had 49% higher odds of having a cancer diagnosis than those who were not abused.<sup>39</sup>

This study has limitations. First, it is possible that people were biased when reporting abuse or neglect as children. However, people generally under-report rather than over report childhood abuse and neglect.<sup>40</sup> We focused exclusively on women who were newly diagnosed with breast cancer; thus, we do not know if our findings generalize to men. Future work assessing cancers that predominately affect males are needed in order to generalize our results to men. Additionally, our sample was predominately white, another limitation of our study that could be addressed in future work with a more diverse sample.

Well after treatment-related problems subside, many breast cancer survivors report QOL difficulties. Our findings suggest that child maltreatment is related to poorer QOL among breast cancer survivors, and social support contributes to the link. A better understanding of how child maltreatment contributes to breast cancer survivor QOL will help in tailoring and therefore enhancing the efficacy of interventions aimed at improving these outcomes.

## Acknowledgments

Work on this paper was supported in part by NIH grants R01CA126857, R01 CA131029, NCRR Grant UL1RR025755, which funds the Clinical Research Center, the Ohio State Comprehensive Cancer Center Core Grant CA16058, and an American Cancer Society Postdoctoral Fellowship Grant PF-11-007-01-CPPB awarded to the first author.

## References

1. Palmer SC, Kagee A, Coyne JC, DeMichele A. Experience of trauma, distress, and posttraumatic stress disorder among breast cancer patients. *Psychosomatic Medicine*. 2004; 66(2):258. [PubMed: 15039512]
2. Beyer K. Depression in patients with cancer receiving adjuvant chemotherapy. 2009
3. Ganz PA, Desmond KA, Leedham B, Rowland JH, Meyerowitz BE, Belin TR. Quality of life in long-term, disease-free survivors of breast cancer: a follow-up study. *Journal of the National Cancer Institute*. 2002; 94(1):39–49. [PubMed: 11773281]
4. Brewin CR, Andrews B, Valentine JD. Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *Journal of Consulting and Clinical Psychology*. 2000; 68(5):748–66. [PubMed: 11068961]
5. Tjemsland L, Søreide JA, Malt UF. Traumatic distress symptoms in early breast cancer. II: Outcome six weeks post surgery. *Psycho-oncology*. 1996; 5(4):295–303.



6. Green BL, Krupnick JL, Rowland JH, et al. Trauma history as a predictor of psychologic symptoms in women with breast cancer. *Journal of Clinical Oncology : Official Journal of the American Society of Clinical Oncology*. 2000; 18(5):1084–93. [PubMed: 10694561]
7. Peretz T, Baider L, Ever-Hadani P, De-Nour AK. Psychological distress in female cancer patients with Holocaust experience *General Hospital Psychiatry*. 1994; 16(6):413–18.
8. Walker EA, Unutzer J, Rutter C, et al. Costs of health care use by women HMO members with a history of childhood abuse and neglect. *Archives of General Psychiatry*. 1999; 56(7):609. [PubMed: 10401506]
9. Zielinski DS, Bradshaw CP. Ecological influences on the sequelae of child maltreatment: A review of the literature. *Child Maltreatment*. 2006; 11(1):49. [PubMed: 16382091]
10. Zaidi LY, Foy DW. Childhood abuse experiences and combat-related PTSD. *Journal of Traumatic Stress*. 1994; 7(1):33–42. [PubMed: 8044441]
11. Walker EA, Keegan D, Gardner G, Sullivan M, Bernstein D, Katon WJ. Psychosocial factors in fibromyalgia compared with rheumatoid arthritis: II. Sexual, physical, and emotional abuse and neglect. *Psychosomatic Medicine*. 1997; 59(6):572. [PubMed: 9407574]
12. Goldsmith RE, Jandorf L, Valdimarsdottir H, et al. Traumatic stress symptoms and breast cancer: The role of childhood abuse. *Child Abuse & Neglect*. 2010
13. Salmon P, Hill J, Krespi R, Clark L, Fisher J, Holcombe C. The role of child abuse and age in vulnerability to emotional problems after surgery for breast cancer. *European Journal of Cancer*. 2006; 42(15):2517–23. [PubMed: 16934976]
14. Nausheen B, Gidron Y, Peveler R, Moss-Morris R. Social support and cancer progression: a systematic review. *Journal of Psychosomatic Research*. 2009; 67(5):403–15. [PubMed: 19837203]
15. Bolger N, Amarel D. Effects of social support visibility on adjustment to stress: Experimental Evidence. *Journal of Personality and Social Psychology*. 2007; 92(3):458–75. [PubMed: 17352603]
16. Bloom JR, Spiegel D. The relationship of two dimensions of social support to the psychological well-being and social functioning of women with advanced breast cancer. *Social Science & Medicine* (1982). 1984; 19(8):831–37.
17. Jamison KR, Wellisch DK, Pasnau RO. Psychosocial aspects of mastectomy: I. the women's perspective. *The American Journal of Psychiatry*. 1978; 135(4):432–36. [PubMed: 637137]
18. Zemore R, Shepel LF. Effects of breast cancer and mastectomy on emotional support and adjustment. *Social Science & Medicine* (1982). 1989; 28(1):19–27.
19. Wimberly SR, Carver CS, Laurenceau JP, Harris SD, Antoni MH. Perceived partner reactions to diagnosis and treatment of breast cancer: impact on psychosocial and psychosexual adjustment. *Journal of Consulting and Clinical Psychology*. 2005; 73(2):300–11. [PubMed: 15796638]
20. Crouch JL, Milner JS, Thomsen C. Childhood physical abuse, early social support, and risk for maltreatment: current social support as a mediator of risk for child physical abuse. *Child Abuse & Neglect*. 2001; 25(1):93–107. [PubMed: 11214816]
21. Hazan C, Shaver PR. Romantic love conceptualized as an attachment process. *Journal of Personality and Social Psychology*. 1987; 59:511–24. [PubMed: 3572722]
22. Uchino BN. Understanding the links between social support and physical health: A life-span perspective with emphasis on the separability of perceived and received support. *Perspectives on Psychological Science*. 2009; 4(3):236–55.
23. Horowitz M, Wilner N, Alvarez WA. Impact of Events Scale: A measure of subjective stress. *Psychosomatic Medicine*. 1979; 41:209–18. [PubMed: 472086]
24. Brady MJ, Cella DF, Mo F, et al. Reliability and validity of the functional assessment of cancer therapy - breast quality-of-life instrument. *Journal of Clinical Oncology*. 1997; 15:974–86. [PubMed: 9060536]
25. Cella, D. Quality of Life. In: Holland, JC.; Breitbart, W., editors. *Psycho-Oncology*. New York: Oxford University Press; 1998. p. 1135–45.
26. Stein KD, Martin SC, Hann DM, Jacobsen PB. A multidimensional measure of fatigue for use with cancer patients. *Cancer Practice*. 1998; 6(3):143–52. [PubMed: 9652245]
27. Cohen S, Hoberman HM. Positive events and social supports as buffers of life change stress. *Journal of Applied Social Psychology*. 1983; 13:99–125.

28. Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index. *Journal of Clinical Epidemiology*. 1994; 47(11):1245–51. [PubMed: 7722560]
29. Dobnig H, Pilz S, Scharnagl H, et al. Independent association of low serum 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D levels with all-cause and cardiovascular mortality. *Archives of Internal Medicine*. 2008; 168(12):1340–49. [PubMed: 18574092]
30. Bernstein, DP.; Fink, L. *Childhood Trauma Questionnaire: A retrospective self-report*. San Antonio, Texas: The Psychological Corporation; 1998.
31. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality in Social Psychology*. 1986; 51:1173–82.
32. MacKinnon DP, Fairchild AJ. Current directions in mediation analysis. *Current Directions in Psychological Science*. 2009; 18(1):16. [PubMed: 20157637]
33. MacKinnon DP, Lockwood CM, Williams J. Confidence limits for the indirect effect: distribution of the product and resampling methods. *Multivariate Behavioral Research*. 2004; 39(1):99–128. [PubMed: 20157642]
34. Ganz PA, Rowland JH, Desmond K, Meyerowitz BE, Wyatt GE. Life after breast cancer: understanding women's health-related quality of life and sexual functioning. *Journal of Clinical Oncology*. 1998; 16(2):501. [PubMed: 9469334]
35. Bower JE, Ganz PA, Desmond KA, et al. Fatigue in long-term breast carcinoma survivors: a longitudinal investigation. *Cancer*. 2006; 106(4):751–58. [PubMed: 16400678]
36. Bower JE, Ganz PA, Aziz N, Fahey JL. Fatigue and proinflammatory cytokine activity in breast cancer survivors. *Psychosomatic Medicine*. 2002; 64(4):604–11. [PubMed: 12140350]
37. Ganz PA, Desmond KA, Leedham B, Rowland JH, Meyerowitz BE, Belin TR. Quality of life in long-term, disease-free survivors of breast cancer: a follow-up study. *JNCI Journal of the National Cancer Institute*. 2002; 94(1):39.
38. Kroenke CH, Kubzansky LD, Scherhammer MD, Kawachi I. Social networks, social support, and survival after breast cancer diagnosis. *Journal of Clinical Oncology*. 2006; 24(7):1105–11. [PubMed: 16505430]
39. Fuller Thomson E, Brennenstuhl S. Making a link between childhood physical abuse and cancer. *Cancer*. 2009; 115(14):3341–50. [PubMed: 19472404]
40. Dill DL, Chu JA, Grob MC, Eisen SV. The reliability of abuse history reports: a comparison of two inquiry formats. *Comprehensive Psychiatry*. 1991; 32(2):166–69. [PubMed: 2022116]



Table 1

Sample Characteristics		
Characteristic	(n= 132)	
	No	%
Age, years		
Mean (SD)	51.70 (9.488)	
Abuse		
Emotional Neglect	16	12.1
Physical Neglect	36	27.3
Emotional Abuse	31	23.5
Physical Abuse	19	14.4
Sexual Abuse	19	14.4
Ethnicity		
Asian	3	2.3
Black	11	8.3
Latino	4	3.0
White	117	88
Marital Status		
Single	18	13.6
Married	97	73.5
Separated/ Divorced	15	11.4
Widowed	2	1.5
Education level		
High school or less	11	8.3
Some College	33	25.0
College or University Graduate	40	30.3
Postgraduate	48	36.4
Employment Status		
Employed full or part time	90	68.2
Unemployed	22	16.7
Retired	20	15.2
Income Level		
\$0-\$25,000	4	3.1
\$25,000-\$50,000	20	15.2
\$50,000-\$75,000	26	19.7
\$75,000-\$100,000	35	26.5
>\$100,000	35	26.5
No Report	12	9.1

Sample Characteristics		
Characteristic	(n= 132)	
	No	%
Type of Treatment		
Surgery Only	14	10.6
Surgery + Radiation	30	22.7
Surgery + Chemotherapy	34	25.8
Surgery + Radiation + Chemotherapy	54	40.9
Cancer Stage		
Stage 0	9	6.8
Stage I	57	43.2
Stage IIA	37	28.0
Stage IIB	15	11.4
Stage IIIA	14	10.6
Months since diagnosis		
Mean (SD)	17.682 (7.953)	
Months since last treatment		
Mean (SD)	11.26 (7.777)	
Impact of Events		
Mean (SD)	27.864 (14.734)	
Physical Well-Being		
Mean (SD)	22.212 (4.7083)	
Emotional Well-Being		
Mean (SD)	18.667 (4.190)	
Functional Well-Being		
Mean (SD)	19.750 (5.4217)	
Breast Cancer Specific Well Being		
Mean (SD)	23.705 (5.9473)	
ISEL		
Mean (SD)	93.697 (14.538)	

**Table 2**

Multiple regression analyses predicting PTSD symptoms, scores on the FACT-B quality of life subscales and the MFSL-SF fatigue scale from childhood maltreatment (categorical) and social support.

Predictor	Cancer-related Psychological Distress			FACT-B Emotional			FACT-B Physical			FACT-B Functional			FACT-B Breast Cancer			MFSL-SF Fatigue			Social Support			
	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	
Step 1 Control Variables	.022			.055*			.046			.069**			.024									
Step 2 Childhood Maltreatment	.062**	.256	.004	.049**	-.226	.010	.046**	-.220	.012	.068**	-.267	.003	.029**	-.173	.047	.131***	.372	.000			-.277	.002
Step 3 Childhood Maltreatment	.066**	.180	.042	.061**	-.153	.080	.077***	-.138	.113	.262***	-.116	.138	.065**	-.098	.262	.069***	.294	.001				
Social Support		-.274	.002		.262	.003		.295	.001		.546	.000		.272	.002		-.281	.001				
Test of Indirect Effect (Bias Corrected Bootstrap)			.001			.002		.002			.001			.004			.004					
Total $R^2$	.150			.164			.169			.336			.163			.225					.120	
<i>n</i>	132			132			132			132			132			132					132	

\*  $p < 0.10$ ;  
 \*\*  $p < 0.05$ ;  
 \*\*\*  $p < 0.001$

**Table 3**

Multiple regression analyses predicting PTSD symptoms, scores on the FACT-B quality of life subscales and the MFSL-SF fatigue scale from childhood maltreatment (continuous) and social support.

Predictor	Cancer-related Psychological Distress			FACT-B Emotional			FACT-B Physical			FACT-B Functional			FACT-B Breast Cancer			MFSL-SF Fatigue			Social Support			
	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	$\Delta R^2$	$\beta$	<i>p</i>	
Step 1 Control Variables	.022			.055*			.006			.069**			.024			.047						
Step 2 Childhood Maltreatment	.052**	.230	.009	.042**	-.208	.016	.088***	-.301	.001	.099***	-.319	.000	.161***	.406	.000	.114***	-.343	.000				
Step 3 Childhood Maltreatment	.064**	.136	.132	.058**	-.118	.185	.242***	-.117	.139	.038**	-.246	.005	.052**	.321	.000							
Social Support		-.28	.003	.263	.004			.537	.000		.213	.015		-.25	.004							
Test of Indirect Effect (Bias Corrected Bootstrap)			.001	.002					.001						.004							
Total <i>R</i> <sup>2</sup>	.138			.156			.336			.206			.237			.161						
<i>n</i>	132			132			132			132			132			132						

\* *p* < 0.10;  
 \*\* *p* < 0.05;  
 \*\*\* *p* < 0.001