

# Activation of Sirt1 by Resveratrol Inhibits TNF- $\alpha$ Induced Inflammation in Fibroblasts

Xiaoxia Zhu<sup>1,2</sup>, Qiong Liu<sup>2,3</sup>, Meimei Wang<sup>4</sup>, Minrui Liang<sup>1,2</sup>, Xue Yang<sup>1,2</sup>, Xue Xu<sup>1,2</sup>, Hejian Zou<sup>1,2\*</sup>, Jianhua Qiu<sup>5\*</sup>

**1** Division of Rheumatology, Shanghai Medical College, Fudan University, Huashan Hospital, Shanghai, China, **2** Institute of Rheumatology, Immunology and Allergy, Shanghai Medical College, Fudan University, Huashan Hospital, Shanghai, China, **3** Department of Human Anatomy, Histology and Embryology, Shanghai Medical College, Fudan University, Shanghai, China, **4** Division of Rheumatology, Dongnan University, Zhongda Hospital, Nanjing, China, **5** Department of Neurology and Radiology, Harvard Medical School, Massachusetts General Hospital, Charlestown, Massachusetts, United States of America

## Abstract

Inflammation is one of main mechanisms of autoimmune disorders and a common feature of most diseases. Appropriate suppression of inflammation is a key resolution to treat the diseases. Sirtuin1 (Sirt1) has been shown to play a role in regulation of inflammation. Resveratrol, a potent Sirt1 activator, has anti-inflammation property. However, the detailed mechanism is not fully understood. In this study, we investigated the anti-inflammation role of Sirt1 in NIH/3T3 fibroblast cell line. Upregulation of matrix metalloproteinases 9 (MMP-9), interleukin-1beta (IL-1 $\beta$ ), IL-6 and inducible nitric oxide synthase (iNOS) were induced by tumor necrosis factor alpha (TNF- $\alpha$ ) in 3T3 cells and resveratrol suppressed overexpression of these pro-inflammatory molecules in a dose-dependent manner. Knockdown of Sirt1 by RNA interference caused 3T3 cells susceptible to TNF- $\alpha$  stimulation and diminished anti-inflammatory effect of resveratrol. We also explored potential anti-inflammatory mechanisms of resveratrol. Resveratrol reduced NF- $\kappa$ B subunit RelA/p65 acetylation, which is notably Sirt1 dependent. Resveratrol also attenuated phosphorylation of mammalian target of rapamycin (mTOR) and S6 ribosomal protein (S6RP) while ameliorating inflammation. Our data demonstrate that resveratrol inhibits TNF- $\alpha$ -induced inflammation via Sirt1. It suggests that Sirt1 is an efficient target for regulation of inflammation. This study provides insight on treatment of inflammation-related diseases.

**Citation:** Zhu X, Liu Q, Wang M, Liang M, Yang X, et al. (2011) Activation of Sirt1 by Resveratrol Inhibits TNF- $\alpha$  Induced Inflammation in Fibroblasts. PLoS ONE 6(11): e27081. doi:10.1371/journal.pone.0027081

**Editor:** Stefan Bereswill, Charité-University Medicine Berlin, Germany

**Received:** October 3, 2011; **Accepted:** October 9, 2011; **Published:** November 1, 2011

**Copyright:** © 2011 Zhu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by Program of Shanghai Subject Chief Scientist (NO. 11XD1401100) and National Natural Science Foundation of China (No. 81072463). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: jqiu@partners.org (JQ); hjzou@fudan.edu.cn (HZ)

These authors contributed equally to this work.

## Introduction

Autoimmune diseases such as rheumatoid arthritis and systemic sclerosis are characterized by aseptic inflammation manifested with upregulation of pro-inflammatory cytokines [1,2]. Increase of cytokines further enhances and sustains inflammatory processes and causes tissue damage [3,4]. Inhibition or neutralization of cytokines suppresses inflammatory cascades and improves functional recovery in experimental models [5,6]. In clinic, blockade of TNF- $\alpha$  by anti-TNF- $\alpha$  antibodies notably reduces inflammation and ameliorates clinical outcomes [7,8,9,10], suggesting that cytokines play a central role in autoimmune diseases. Reduction of cytokines production or suppression of their signaling is an efficient therapeutic target.

Sirtuin 1 (Sirt1), a mammalian homolog of Sir2, is a NAD<sup>+</sup>-dependent class III histone deacetylase. It has been shown to be involved in a variety of pathophysiological processes, such as anti-inflammation, cell growth and metabolism modulation, anti-carcinogen [11,12,13]. Sirt1 regulates pro-inflammatory mediator [14,15,16]. Knockout or knockdown of Sirt1 gene leads to increase of cytokines release whereas Sirt1 activation by its activators inhibits productions of TNF- $\alpha$ , monocyte chemoattractant protein

1 (MCP-1) and IL-8 [14,15,16]. Furthermore, Sirt1 has inhibitory effects in experimental chronic inflammatory diseases such as chronic obstructive pulmonary disease and colitis [14,16,17]. Suppression of pro-inflammatory cytokines production by Sirt1 is highly related to its negative regulation of NF- $\kappa$ B activity by deacetylating of RelA/p65 subunit at lysine 310 [15].

Resveratrol (trans-3,4',5-trihydroxystilbene), a polyphenolic phytoalexins, is a potent activator of Sirt1 [18]. Increase of evidence indicates that resveratrol exerts an anti-inflammatory property [19,20,21]. Resveratrol has a chondroprotective capacity through suppressing the production of IL-1 $\beta$  and reactive oxygen species (ROS) [22]. In human primary airway epithelial cells, resveratrol inhibits cytokine-stimulated iNOS expression and nitrite production [23]. Resveratrol also protects cartilage against the development of experimentally induced inflammatory arthritis [24]. Recent multiple lines of evidence demonstrate that resveratrol inhibits inflammation via blockade of NF- $\kappa$ B transcriptional activity [25,26,27]. Resveratrol decreases the expression of NF- $\kappa$ B subunit RelA/p65 or attenuates translocation of p65 from the cytosol to the nucleus with stabilization of inhibitory I $\kappa$ B, and further downregulates levels of TNF- $\alpha$  and cyclooxygenase-2 (COX-2) [19,28].

Sirt1 may be a promising target for anti-inflammation therapy [29]. In the present study, we investigated the inhibitory role of Sirt1 in TNF- $\alpha$  induced cytokine production in fibroblast cells through activating Sirt1 with resveratrol or downregulating Sirt1 by RNA interference. We further demonstrated that resveratrol inhibited inflammation via a Sirt1-dependent manner.

## Methods

### Cell culture and treatment

Mouse embryonic 3T3/NIH fibroblasts (obtained from the American Type Culture Collection) were cultured in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with antibiotics (100 U/ml penicillin, 100  $\mu$ g/ml streptomycin) and 10% fetal bovine serum, at 37°C in a humidified incubator with 5% CO<sub>2</sub>. Recombinant mouse TNF- $\alpha$  (R&D System), resveratrol (Sigma) or rapamycin (EMD4Bioscience) were used in this study.

### Gelatin zymography

Gelatin zymography was done as previous described [30,31]. Briefly, culture media were collected after treatment and subjected to SDS-PAGE in 10% polyacrylamide gels copolymerized with 1 mg/ml gelatin. After electrophoresis, gels were washed in renature buffer to remove the SDS and further incubated with developing buffer (Invitrogen) at 37°C for 24 hours. The gels then were stained with Coomassie blue R-250 (Bio-Rad) for 15 minutes followed by destaining in deionized water with 10% acetic acid and 20% methanol. MMP-9 expression and proteolytic activity were evidenced as clear bands against the background of stained gelatin.

### Western blotting

Cells were lysed in RIPA buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, proteinase inhibitor (Roche) and phosphatase inhibitor (Calbiochem)). Protein concentration was detected using the DC™ protein assay (Bio-Rad). Protein (30  $\mu$ g) was loaded to 10% SDS-PAGE and semi-dry transferred onto a Polyvinylidene fluoride (PVDF) membrane. After blocking with 5% non-fat milk, membranes were incubated overnight at 4°C with primary antibodies against IL-1 $\beta$  (1:500, Santa Cruz), Sirt1 (1:500, Millipore), acetyl-NF- $\kappa$ Bp65(Lys310) (1:500, Cell Signaling), phospho(ser245/236)-S6RP (1:1000, Cell Signaling), phospho(ser2448)-mTOR (1:1000, Cell Signaling). Horseradish peroxidase-conjugated secondary antibodies were used for ECL-plus (GE Healthcare) detection. The results were normalized to  $\beta$ -actin (1:5000, Abcam).

### Real-time RT-PCR

Total RNA was extracted using RNeasy Mini Isolation Kit (GE Healthcare), and reverse-transcribed into cDNA using the SuperScripts III First-Strand Synthesis System (Invitrogen). The genes were examined by real-time PCR. Primers were purchased from Applied Biosystems (MMP-9, Mm00442991\_m1; IL-6, Mm00446190\_m1; iNOS, Mm00440502\_m1). Endogenous 18S ribosomal RNA was used as internal control for normalizing gene expression. Results then were assessed by t-test or ANOVA;  $P \leq 0.05$  was considered to be statistically significant.

### RNA interference

siRNA for mouse Sirt1 (ON-TARGETplus SMARTpool) was purchased from Thermo Fisher Scientific and diluted in RNase-free double distilled water (DDW). Cells were seeded on 6-well plates. After 24 hours, cells at 60~70% confluency were

transfected with 20 nM Sirt1 siRNA, 20 nM control siRNA, or the same volume of DDW for 6~8 hours using Lipofectamine 2000 according to the manufacturer's protocol. Then cells were switched into DMEM and incubated for 72 hours before treatment.

## Results

### TNF- $\alpha$ induced overexpression of MMP-9 and other inflammatory factors in fibroblasts


TNF- $\alpha$  induces inflammation in various cell types [20,32]. In current study, we examined expressions of MMP-9, IL-6, iNOS and IL-1 $\beta$  in TNF- $\alpha$  treated 3T3/NIH mouse embryonic fibroblasts. Within 0.5 ng/ml to 10 ng/ml of dose range, no cell death was observed up to 72 hours after TNF- $\alpha$  treatment (data not shown). As shown in figure 1, TNF- $\alpha$  induced notably MMP-9 upregulation in a dose- and time- dependent manner. The concentration of 10 ng/ml of TNF- $\alpha$  was chosen for investigating the response of fibroblasts to cytokine stimulation. Messenger RNA expressions of MMP-9, iNOS and IL-6 were significantly induced by TNF- $\alpha$ . IL-6 was increased strikingly as early as 30 minutes post treatment; while MMP-9 and iNOS were elevated 3 hours after treatment, and reached to the peak 6 hours after treatment. IL-1 $\beta$  expression was also examined by western blotting analysis. A remarkable upregulation of IL-1 $\beta$  was detected 3 hours after TNF- $\alpha$  treatment. These results are consistent with the reports from other investigators [33,34].

### Resveratrol blocked TNF- $\alpha$ induced inflammation

Resveratrol is a potent Sirt1 agonist and increases Sirt1 activity [18]. We employed resveratrol to investigate the anti-inflammatory function of Sirt1. 3T3 fibroblasts were pretreated with resveratrol followed by TNF- $\alpha$  challenge. Release of MMP-9 in the culture media was observed in TNF- $\alpha$  treated cells and was greatly inhibited by resveratrol in a dose dependent manner (figure 2A). To test the time course of inflammatory inhibition by resveratrol, resveratrol was administered 1, 0 hour before or 1, 3 hours post TNF- $\alpha$  treatment, the relative mRNA levels of inflammatory factors were detected by real-time RT-PCR analysis at 9 hours after TNF- $\alpha$  treatment. Resveratrol inhibited the pro-inflammatory factors in a clearly time-dependent fashion (figure 2B). Upregulation of MMP-9, IL-6 and iNOS were attenuated by resveratrol treated when resveratrol was employed before or at the same time of TNF- $\alpha$  treatment. Induced IL-6 upregulation was inhibited even when resveratrol was employed 3 hours after TNF- $\alpha$  stimulation. Interestingly, lack of significant inhibitory effect was observed when resveratrol was treated 1 hour (for iNOS) and 3 hours (for MMP-9) after TNF- $\alpha$  stimulation. Furthermore, the TNF- $\alpha$  induced IL-1 $\beta$  expression was also strikingly inhibited by resveratrol pretreatment for 1 hour (figure 2C). No cell death was observed within 72 hours under these conditions (data not shown).


### Resveratrol Inhibited inflammation via Sirt1

Since resveratrol is a pharmacological activator of Sirt1 and may have off-target effects, we further examined whether Sirt1 is required for anti-inflammatory activity of resveratrol. Sirt1 expression is induced by TNF- $\alpha$  or inflammation [35]. And Sirt1 activity is highly related to its expression [36,37]. Here we used RNA interference technique to knock down Sirt1 expression. Expression of Sirt1 was reduced 72 hours after treatment of small interference RNA (siRNA) targeting Sirt1 gene (figure 3A). Inhibitory effect of resveratrol on the upregulation of MMP-9, IL-6 and iNOS was attenuated in the cells in which Sirt1




**Figure 1. TNF- $\alpha$  induced overexpression of MMP-9, iNOS, IL-6 and IL-1 $\beta$  in 3T3/NIH fibroblasts.** **A.** 3T3/NIH cells were incubated for 6, 24 or 48 hours (h) with varying concentrations of TNF- $\alpha$ , gelatin zymography were performed to detect MMP-9 expression in the medium. **B.** Cells were treated by TNF- $\alpha$  (10 ng/ml) for 0, 1/2, 3, 6, 18 h, relative mRNA expressions of MMP-9, iNOS and IL-6 were examined by real-time RT-PCR analysis (n=3 per group, \* $p$ <0.05 vs. 0 h; \*\* $p$ <0.01 vs. 0 h). **C.** Cells were treated by TNF- $\alpha$  for 0, 1/2, 3, 9, 24 h, IL-1 $\beta$  was measured by western blotting analysis.

doi:10.1371/journal.pone.0027081.g001



**Figure 2. Resveratrol inhibited TNF- $\alpha$  induced inflammation.** **A.** 3T3/NIH cells were pretreated 1 h with varying concentrations of resveratrol (Res) before TNF- $\alpha$  (10 ng/ml) treatment. MMP-9 expression in the medium was examined by gelatin zymography after 24 h treatment. **B.** Resveratrol (50 nM) was used to treat cells 1, 0 hour before or 1, 3 hours post TNF- $\alpha$  (10 ng/ml), the relative mRNA levels of MMP-9, iNOS and IL-6 were detected by real-time RT-PCR analysis at 6 hours post TNF- $\alpha$  treatment. (n=3 per group, \* $p$ <0.05 vs. TNF- $\alpha$ +Res-; \*\* $p$ <0.01 vs. TNF- $\alpha$ +Res-). **C.** Cells were pretreated by resveratrol 1 h before TNF- $\alpha$ , western blotting analysis was used to detect IL-1 $\beta$  expression after 9 h treatment.

doi:10.1371/journal.pone.0027081.g002



**Figure 3. Resveratrol inhibited inflammation in a Sirt1 dependent manner.** **A.** Western blotting analysis shows Sirt1 siRNA (20 nM) transfection for 72 hours strikingly blocked Sirt1 expression, compared to control (Cont) siRNA (20 nM) or DDW transfection. **B.** After Sirt1 or control siRNA transfection for 72 h, cells were pretreated with resveratrol (50 nM) in the presence of TNF- $\alpha$  (10 ng/ml) treatment for 24 h, MMP-9 expression in the medium were examined by gelatin zymography. **C.** Cells were processed as (B) for 6 h, real-time RT-PCR analysis was used to examine mRNA expression of MMP-9, IL-6, iNOS (n=3 per group, \* $p$ <0.05 vs. Sirt1 siRNA-; \*\* $p$ <0.01 vs. Sirt1 siRNA-). doi:10.1371/journal.pone.0027081.g003

expression was knocked down (figure 3B-C). These results suggest that anti-inflammation of resveratrol is mainly dependent on Sirt1 and Sirt1 exerts a negative regulatory effect on the inflammation.

### Resveratrol decreased acetylated RelA/p65

NF- $\kappa$ B plays a central role in inflammation [38]. Acetylation of NF- $\kappa$ B subunit RelA/p65 at Lysine310 is involved in the activity of NF- $\kappa$ B and inflammatory factors transcription [39]. We found that TNF- $\alpha$  increased acetyl-NF- $\kappa$ Bp65 (Lysine310) and resveratrol suppressed NF- $\kappa$ Bp65 acetylation in 3T3 cells (figure 4A-B). To explore whether the deacetylation function of resveratrol is also Sirt1 dependent, Sirt1 was knocked down by siRNA. Interestingly, basal level of acetyl-NF- $\kappa$ Bp65 (Lys310) was increased after Sirt1 knockdown. Inhibitory effect of resveratrol on the acetyl-NF- $\kappa$ Bp65 was diminished by Sirt1 knockdown (figure 4C). The results indicate that Sirt1 is required for inhibiting NF- $\kappa$ Bp65 acetylation by resveratrol. It also suggests that Sirt1 plays a role in regulation of RelA/p65 acetylation.

### Resveratrol inhibited activation of mTOR induced by TNF- $\alpha$


Mammalian target of rapamycin is related to TNF- $\alpha$  induced inflammation [40]. In this study, upregulation of phosphorylated mTOR and S6RP were detected as early as 30 minutes and returned back to control level within 24 hours after TNF- $\alpha$  stimulation (figure 5A). We also found that resveratrol suppressed phosphorylated mTOR and S6RP (figure 5B) and further reduced the expression of IL-1 $\beta$  and MMP-9 (figure 5 B-C). Moreover, rapamycin inhibited phosphorylation of mTOR and S6RP, and suppressed MMP-9 and IL-1 $\beta$  upregulation (figure 5 B-C).

## Discussion

Previous studies have revealed that resveratrol activates Sirt1 and suppresses the inflammation of both *in vivo* and *in vitro* experiment models [20,21,36,41]. In the present study, we found that TNF- $\alpha$  induced upregulation of cytokines and MMP-9 in 3T3 fibroblasts. Sirt1 was constitutively expressed in the cells and upregulated after TNF- $\alpha$  treatment. Resveratrol inhibited TNF- $\alpha$  induced overexpression of cytokines, iNOS and MMP-9 via Sirt1 dependent manner. Knockdown of Sirt1 caused increase of IL-6, iNOS and MMP-9 expression. The anti-inflammation function of resveratrol was blocked after Sirt1 knockdown. This study suggests that resveratrol ameliorates inflammation via activating Sirt1. Reduction of acetylated NF- $\kappa$ B and suppression of mTOR/S6RP phosphorylation may be involved in the mechanisms.

TNF- $\alpha$  has been demonstrated to be one of the main inflammatory mediators that involved in autoimmune diseases, such as rheumatoid arthritis, systemic sclerosis [2,42]. TNF- $\alpha$  initiates or aggravates inflammation by activating NF- $\kappa$ B and producing cytokines, chemokines, MMPs, and other inflammatory molecules in different types of cells [20,32]. In the present study, we found that TNF- $\alpha$  induced MMP-9, IL-6, and iNOS in a dose- and time-dependent fashion without causing cell death in fibroblasts. These findings are consistent with other reports [20,21]. We applied this *in vitro* model to further investigate the role of Sirt1 in TNF- $\alpha$  induced inflammation.


Sirt1 plays a critical role in regulation of several transcription factors such as p53, NF- $\kappa$ B and FoxOs and has many important functions in metabolism, anti-cancer, anti-ageing and anti-inflammation [12,13,43,44]. Overexpression of Sirt1 or increase of Sirt1 activity significantly suppresses cytokines production and reduces inflammation in different animal models [20,21]. On the



**Figure 4. Resveratrol decreased acetylated RelA/p65.** **A.** 3T3/NIH cells were treated with TNF- $\alpha$  (10 ng/ml) for 0, 1/2, 3, 9, 24 h, and acetyl-NF- $\kappa$ Bp65(Lys310) (acetyl-p65) expression was detected by western blotting analysis in cell lysis. **B.** Cells were pretreated by resveratrol (50 nM) in the presence of TNF- $\alpha$  treatment for 3 h, acetyl-NF- $\kappa$ Bp65(Lys310) expression were examined by western blotting analysis. **C.** Cells were transfected with Sirt1 or control siRNA for 72 h, then pretreated with resveratrol (50 nM) in the presence of TNF- $\alpha$  (10 ng/ml) treatment for 3 h. Western blotting analysis was used to detect acetyl-NF- $\kappa$ Bp65(Lys310) expression.  
doi:10.1371/journal.pone.0027081.g004

other hand, reduction of Sirt1 activity results in increase of inflammatory response [45,46]. It has been shown that TNF- $\alpha$  induces inflammation through activation of cathepsin B followed by cleavage of Sirt1 [47]. More severe encephalomyelitis and autoimmunity induced by myelin oligodendroglia glycoprotein (MOG) peptide is detected in Sirt1-deficient mice compared to the controls [48]. MMP-9 expression in Sirt1 $^{-/-}$  mouse embryonic fibroblasts (MEF) is significant higher than that in wild type cells [49]. Our data showed that Sirt1 was constitutively expressed in the 3T3 fibroblasts (figure 3A). Knockdown of Sirt1 caused increase of MMP-9, iNOS and IL-6 expression. Similar phenomenon is also observed in vascular smooth muscle cells. The inhibitory effect on AP-1, c-Jun or c-Fos by Sirt1 might be the potential mechanism [48,50]. These results suggest that Sirt1 plays a role in suppressing expression of inflammatory molecules even under normal conditions.

Resveratrol is a potent activator of Sirt1 and has multiple effects on metabolism, anti-cancer, anti-ageing and anti-inflammation [11,19,51,52]. Treatment of resveratrol suppresses cyclooxygenase-2 activity [53] and modulates interferon-gamma (IFN- $\gamma$ ), TNF- $\alpha$ , IL-6 and MCP-1 expression in experimental ileitis [54]. In cultured neural cells, resveratrol prevents increase of cytokines, chemokines and iNOS/NO induced by lipopolysaccharide (LPS) [55]. In IL-1 $\beta$  treated human chondrocytes, resveratrol inhibits the cytokine production [22]. Furthermore, resveratrol ameliorates LPS induced inflammatory arthritis *in vivo* [24]. In this study, we observed that resveratrol significantly inhibited the TNF- $\alpha$  induced increase of MMP-9, IL-6, iNOS and IL-1 $\beta$  in the fibroblasts. The inhibitory effect was in a dose- and time-



**Figure 5. mTOR/S6RP is also involved in the inhibitory function of resveratrol on TNF- $\alpha$  induced inflammation.** **A.** 3T3/NIH cells were treated with TNF- $\alpha$  (10 ng/ml) for 0, 1/2, 3, 9, 24 h, phosphorylation of mTOR and S6RP was detected by western blotting analysis. **B.** Cells were pretreated by rapamycin (Rapa, 10 ng/ml) or resveratrol (50 nM) for 1 h in the presence of TNF- $\alpha$  treatment for 24 h, MMP-9 released in medium was measured by gelatin zymography. **C.** Cells were processed as (B) for 3 h, IL-1 $\beta$  and phosphorylation of mTOR and S6RP were examined by western blotting analysis in cell lysis.  
doi:10.1371/journal.pone.0027081.g005

dependent fashion. The evidence indicates that resveratrol is an efficient inhibitor of inflammation in fibroblasts.

Although resveratrol is a pharmacological agonist of Sirt1 and may have multiple targets, several studies indicate that the anti-inflammatory function of resveratrol is highly dependent on Sirt1 [51,56]. Resveratrol upregulates Sirt1 expression at both transcriptional and translational level [57,58]. Furthermore, by binding to Sirt1, resveratrol alters Sirt1 structure and upregulates its activity as much as 8 folds, and lowers the  $K_m$  value for acetylated substrate [11,18]. However, other studies suggest that resveratrol may not directly activate Sirt1 [59,60]. Resveratrol lacks inhibition of upregulation of hypoxia induced factor 1 alpha (HIF-1 $\alpha$ ) and AMP-activated protein kinase (AMPK) in the cells in which Sirt1 is deficit [61]. Reduction of MMP-9 expression by resveratrol is remarkably attenuated in Sirt1 $^{-/-}$  mouse embryonic fibroblasts (MEF) [49]. But the inhibitory effect of resveratrol on Smad2/3 phosphorylation was reported to be independent of Sirt1 [62,63]. In the present study, suppression of TNF- $\alpha$ -induced inflammatory response by resveratrol was attenuated in the cells in which Sirt1 was knocked down by RNA interference. Our findings support that Sirt1 is required for anti-inflammation of resveratrol and suggest that Sirt1 is an important target for treatment of autoimmune disorders.

Several lines of evidence indicate that NF- $\kappa$ B is one of Sirt1 targets. NF- $\kappa$ B acetylation is critical for its activity, especially in cytokines induced inflammation [64]. Acetylation of NF- $\kappa$ B at

specific lysine residues results in uncoiling of the DNA and increased accessibility to transcription factor binding which contributes to pro-inflammatory factors production [65,66]. Sirt1 physically interacts with the RelA/p65 subunit of NF- $\kappa$ B and inhibits transcription by deacetylation of RelA/p65 at lysine 310 [15,67]. Acetylation of RelA/p65 at lysine 310 has been reported to be required for full transcriptional activity of NF- $\kappa$ B in inflammation [68]. In this study, we found that TNF- $\alpha$  increased acetylation at Lysine 310 of RelA/p65 subunit. Resveratrol inhibited the TNF- $\alpha$  induced acetylation of RelA/p65 via a Sirt1-dependent manner (figure 4). Our findings further support the elucidation of the mechanism that the inhibitory effect of Sirt1 is due to suppressing NF- $\kappa$ B activity.

Mammalian target of rapamycin (mTOR) is a serine/threonine protein kinase which regulates protein synthesis [69,70]. Studies report that activation of mTOR plays a role in TNF- $\alpha$ -induced inflammatory cascades [71], and is also implicated in inflammation related diseases [72]. TNF- $\alpha$  increases phosphorylation of mTOR and its downstream targets p70S6K and S6RP [73,74,75]. Rapamycin ameliorates inflammation by inhibiting mTOR and further decreasing expression of cytokines and chemokines, improves outcomes in inflammation related disease models [76,77]. These results suggest that mTOR is an important target for treatment of inflammation. In the present study, mTOR and

S6RP were phosphorylated and activated in TNF- $\alpha$  stimulated fibroblasts. Resveratrol also notably downregulated the phosphorylated mTOR and S6RP while ameliorating the inflammation in fibroblasts, suggesting that inhibition of mTOR may be one of the mechanisms of its anti-inflammatory effect.

In summary, our results indicate that Sirt1 plays a key role in inflammation. Activation of Sirt1 by its agonists such as resveratrol efficiently suppresses inflammatory cascades. This study provides an insight for developing therapeutic approaches for inflammation related diseases or autoimmune disorders. Further investigation in animal models is warranted to confirm the findings from *in vitro* studies.

## Acknowledgments

We thank China Scholarship Council for awarding Xiaoxia Zhu the state scholarship to support her in studying in the United States. We also appreciate the excellent assistance from Laiming Yung and Yi Zheng.

## Author Contributions

Conceived and designed the experiments: HZ JQ. Performed the experiments: XZ QL. Analyzed the data: XZ MW. Contributed reagents/materials/analysis tools: XZ ML XY XX. Wrote the paper: XZ HZ JQ.

## References

- Kang SY, Kim MH, Lee WI (2010) Measurement of inflammatory cytokines in patients with rheumatoid arthritis. *Korean J Lab Med* 30: 301–306.
- Scala E, Pallotta S, Frezzolini A, Abeni D, Barbieri C, et al. (2004) Cytokine and chemokine levels in systemic sclerosis: relationship with cutaneous and internal organ involvement. *Clin Exp Immunol* 138: 540–546.
- Hu X, Ho HH, Lou O, Hidaka C, Ivashkiv LB (2005) Homeostatic role of interferons conferred by inhibition of IL-1-mediated inflammation and tissue destruction. *J Immunol* 175: 131–138.
- Hernandez-Pando R, Rook GA (1994) The role of TNF-alpha in T-cell-mediated inflammation depends on the Th1/Th2 cytokine balance. *Immunology* 82: 591–595.
- Shahrara S, Proudfoot AE, Park CC, Volin MV, Haines GK, et al. (2008) Inhibition of monocyte chemoattractant protein-1 ameliorates rat adjuvant-induced arthritis. *J Immunol* 180: 3447–3456.
- Christodoulou C, Choy EH (2006) Joint inflammation and cytokine inhibition in rheumatoid arthritis. *Clin Exp Med* 6: 13–19.
- Suryaprasad AG, Prindiville T (2003) The biology of TNF blockade. *Autoimmun Rev* 2: 346–357.
- Ancuta C, Ancuta E, Miu S, Iordache C, Belibou C, et al. (2009) Adalimumab therapy in patients with active rheumatoid arthritis. *Rev Med Chir Soc Med Nat Iasi* 113: 710–715.
- Yount S, Sorensen MV, Cella D, Sengupta N, Grober J, et al. (2007) Adalimumab plus methotrexate or standard therapy is more effective than methotrexate or standard therapies alone in the treatment of fatigue in patients with active, inadequately treated rheumatoid arthritis. *Clin Exp Rheumatol* 25: 838–846.
- Keystone EC, Kavanaugh AF, Sharp JT, Tannenbaum H, Hua Y, et al. (2004) Radiographic, clinical, and functional outcomes of treatment with adalimumab (a human anti-tumor necrosis factor monoclonal antibody) in patients with active rheumatoid arthritis receiving concomitant methotrexate therapy: a randomized, placebo-controlled, 52-week trial. *Arthritis Rheum* 50: 1400–1411.
- Howitz KT, Bitterman KJ, Cohen HY, Lamming DW, Lavu S, et al. (2003) Small molecule activators of sirtuins extend *Saccharomyces cerevisiae* lifespan. *Nature* 425: 191–196.
- Kojima K, Ohhashi R, Fujita Y, Hamada N, Akao Y, et al. (2008) A role for SIRT1 in cell growth and chemoresistance in prostate cancer PC3 and DU145 cells. *Biochem Biophys Res Commun* 373: 423–428.
- Ford J, Jiang M, Milner J (2005) Cancer-specific functions of SIRT1 enable human epithelial cancer cell growth and survival. *Cancer Res* 65: 10457–10463.
- Yang SR, Wright J, Bauter M, Seweryniak K, Kode A, et al. (2007) Sirtuin regulates cigarette smoke-induced proinflammatory mediator release via RelA/p65 NF-kappaB in macrophages *in vitro* and in rat lungs *in vivo*: implications for chronic inflammation and aging. *Am J Physiol Lung Cell Mol Physiol* 292: L567–576.
- Yeung F, Hoberg JE, Ramsey CS, Keller MD, Jones DR, et al. (2004) Modulation of NF-kappaB-dependent transcription and cell survival by the SIRT1 deacetylase. *EMBO J* 23: 2369–2380.
- Rajendrasozhan S, Yang SR, Kinnula VL, Rahman I (2008) SIRT1, an antiinflammatory and antiaging protein, is decreased in lungs of patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 177: 861–870.
- Singh UP, Singh NP, Singh B, Hofseth LJ, Price RL, et al. (2010) Resveratrol (trans-3,5,4'-trihydroxystilbene) induces silent mating type information regulation-1 and down-regulates nuclear transcription factor-kappaB activation to abrogate dextran sulfate sodium-induced colitis. *J Pharmacol Exp Ther* 332: 829–839.
- Borra MT, Smith BC, Denu JM (2005) Mechanism of human SIRT1 activation by resveratrol. *J Biol Chem* 280: 17187–17195.
- Bishayec A, Waghray A, Barnes KF, Mbimba T, Bhatia D, et al. (2010) Suppression of the inflammatory cascade is implicated in resveratrol chemoprevention of experimental hepatocarcinogenesis. *Pharm Res* 27: 1080–1091.
- Knobloch J, Sibbing B, Jungck D, Lin Y, Urban K, et al. (2010) Resveratrol impairs the release of steroid-resistant inflammatory cytokines from human airway smooth muscle cells in chronic obstructive pulmonary disease. *J Pharmacol Exp Ther* 335: 788–798.
- Chung EY, Kim BH, Hong JT, Lee CK, Ahn B, et al. (2011) Resveratrol down-regulates interferon-gamma-inducible inflammatory genes in macrophages: molecular mechanism via decreased STAT-1 activation. *J Nutr Biochem* 22: 902–909.
- Csaki C, Keshishzadeh N, Fischer K, Shakibaei M (2008) Regulation of inflammation signalling by resveratrol in human chondrocytes *in vitro*. *Biochem Pharmacol* 75: 677–687.
- Donnelly LE, Newton R, Kennedy GE, Fenwick PS, Leung RH, et al. (2004) Anti-inflammatory effects of resveratrol in lung epithelial cells: molecular mechanisms. *Am J Physiol Lung Cell Mol Physiol* 287: L774–783.
- Elmali N, Baysal O, Harma A, Esenkaya I, Mizrak B (2007) Effects of resveratrol in inflammatory arthritis. *Inflammation* 30: 1–6.
- Holmes-McNary M, Baldwin AS Jr. (2000) Chemopreventive properties of trans-resveratrol are associated with inhibition of activation of the I-kappaB kinase. *Cancer Res* 60: 3477–3483.
- Manna SK, Mukhopadhyay A, Aggarwal BB (2000) Resveratrol suppresses TNF-induced activation of nuclear transcription factors NF-kappa B, activator protein-1, and apoptosis: potential role of reactive oxygen intermediates and lipid peroxidation. *J Immunol* 164: 6509–6519.
- Csaki C, Mobasheri A, Shakibaei M (2009) Synergistic chondroprotective effects of curcumin and resveratrol in human articular chondrocytes: inhibition of IL-1beta-induced NF-kappaB-mediated inflammation and apoptosis. *Arthritis Res Ther* 11: R165.
- Kumar A, Sharma SS (2010) NF-kappaB inhibitory action of resveratrol: a probable mechanism of neuroprotection in experimental diabetic neuropathy. *Biochem Biophys Res Commun* 394: 360–365.
- Olholm J, Paulsen SK, Cullberg KB, Richelsen B, Pedersen SB (2010) Anti-inflammatory effect of resveratrol on adipokine expression and secretion in human adipose tissue explants. *Int J Obes (Lond)* 34: 1546–1553.
- Gursoy-Ozdemir Y, Qiu J, Matsuoka N, Bolay H, Bernpohl D, et al. (2004) Cortical spreading depression activates and upregulates MMP-9. *J Clin Invest* 113: 1447–1455.

31. Qiu J, Xu J, Zheng Y, Wei Y, Zhu X, et al. (2010) High-mobility group box 1 promotes metalloproteinase-9 upregulation through Toll-like receptor 4 after cerebral ischemia. *Stroke* 41: 2077–2082.
32. Ding GJ, Fischer PA, Boltz RC, Schmidt JA, Colaianni JJ, et al. (1998) Characterization and quantitation of NF-kappaB nuclear translocation induced by interleukin-1 and tumor necrosis factor-alpha. Development and use of a high capacity fluorescence cytometric system. *J Biol Chem* 273: 28897–28905.
33. Shao XT, Feng L, Yao HP, Sun WJ, Zhang LH (2004) Effect of Triptolide on TNFalpha-induced activation of NF-kappaB and expression of COX-2 and iNOS in human rheumatoid arthritis synovial fibroblasts. *Zhejiang Da Xue Xue Bao Yi Xue Ban* 33: 160–165.
34. Boussetta T, Raad H, Letteron P, Gougerot-Pocidal MA, Marie JC, et al. (2009) Punicic acid a conjugated linolenic acid inhibits TNFalpha-induced neutrophil hyperactivation and protects from experimental colon inflammation in rats. *PLoS One* 4: e6458.
35. Zhang HN, Li L, Gao P, Chen HZ, Zhang R, et al. (2010) Involvement of the p65/RelA subunit of NF-kappaB in TNF-alpha-induced SIRT1 expression in vascular smooth muscle cells. *Biochem Biophys Res Commun* 397: 569–575.
36. Lee SJ, Kim MM (2011) Resveratrol with antioxidant activity inhibits matrix metalloproteinase via modulation of SIRT1 in human fibrosarcoma cells. *Life Sci* 88: 465–472.
37. Niederer F, Ospelt C, Brentano F, Hottiger MO, Gay RE, et al. (2011) SIRT1 overexpression in the rheumatoid arthritis synovium contributes to proinflammatory cytokine production and apoptosis resistance. *Ann Rheum Dis* 70: 1866–1873.
38. Tak PP, Firestein GS (2001) NF-kappaB: a key role in inflammatory diseases. *J Clin Invest* 107: 7–11.
39. Ito K (2007) Impact of post-translational modifications of proteins on the inflammatory process. *Biochem Soc Trans* 35: 281–283.
40. Lee DF, Kuo HP, Chen CT, Hsu JM, Chou CK, et al. (2007) IKK beta suppression of TSC1 links inflammation and tumor angiogenesis via the mTOR pathway. *Cell* 130: 440–455.
41. Zhang C, Feng Y, Qu S, Wei X, Zhu H, et al. (2011) Resveratrol attenuates doxorubicin-induced cardiomyocyte apoptosis in mice through SIRT1-mediated deacetylation of p53. *Cardiovasc Res* 90: 538–545.
42. Lo SF, Huang CM, Lin HC, Chen WC, Tsai CH, et al. (2008) Cytokine (IL-6) and chemokine (IL-8) gene polymorphisms among rheumatoid arthritis patients in Taiwan. *Clin Exp Rheumatol* 26: 632–637.
43. Jung-Hynes B, Ahmad N (2009) Role of p53 in the anti-proliferative effects of Sirt1 inhibition in prostate cancer cells. *Cell Cycle* 8: 1478–1483.
44. Salminen A, Ojala J, Huuskonen J, Kauppinen A, Suuronen T, et al. (2008) Interaction of aging-associated signaling cascades: inhibition of NF-kappaB signaling by longevity factors FoxOs and SIRT1. *Cell Mol Life Sci* 65: 1049–1058.
45. Kim YS, Lee YM, Park JS, Lee SK, Kim EC (2010) SIRT1 modulates high-mobility group box 1-induced osteoclastogenic cytokines in human periodontal ligament cells. *J Cell Biochem* 111: 1310–1320.
46. Ohguchi K, Itoh T, Akao Y, Inoue H, Nozawa Y, et al. (2010) SIRT1 modulates expression of matrix metalloproteinases in human dermal fibroblasts. *Br J Dermatol* 163: 689–694.
47. Dvir-Ginzberg M, Gagarina V, Lee EJ, Booth R, Gabay O, et al. (2011) Tumor necrosis factor alpha-mediated cleavage and inactivation of Sirt1 in human osteoarthritic chondrocytes. *Arthritis Rheum* 63: 2363–2373.
48. Zhang J, Lee SM, Shannon S, Gao B, Chen W, et al. (2009) The type III histone deacetylase Sirt1 is essential for maintenance of T cell tolerance in mice. *J Clin Invest* 119: 3048–3058.
49. Gao Z, Ye J (2008) Inhibition of transcriptional activity of c-JUN by SIRT1. *Biochem Biophys Res Commun* 376: 793–796.
50. Ishinaga H, Jono H, Lim JH, Kweon SM, Xu H, et al. (2007) TGF-beta induces p65 acetylation to enhance bacteria-induced NF-kappaB activation. *EMBO J* 26: 1150–1162.
51. Lagouge M, Argmann C, Gerhart-Hines Z, Meziane H, Lerin C, et al. (2006) Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1alpha. *Cell* 127: 1109–1122.
52. Sun W, Wang W, Kim J, Keng P, Yang S, et al. (2008) Anti-cancer effect of resveratrol is associated with induction of apoptosis via a mitochondrial pathway alignment. *Adv Exp Med Biol* 614: 179–186.
53. Banerjee S, Bueso-Ramos C, Aggarwal BB (2002) Suppression of 7,12-dimethylbenz(a)anthracene-induced mammary carcinogenesis in rats by resveratrol: role of nuclear factor-kappaB, cyclooxygenase 2, and matrix metalloproteinase 9. *Cancer Res* 62: 4945–4954.
54. Bereswill S, Munoz M, Fischer A, Plickert R, Haag LM, et al. (2010) Anti-inflammatory effects of resveratrol, curcumin and simvastatin in acute small intestinal inflammation. *PLoS One* 5: e15099.
55. Lu X, Ma L, Ruan L, Kong Y, Mou H, et al. (2010) Resveratrol differentially modulates inflammatory responses of microglia and astrocytes. *J Neuroinflammation* 7: 46.
56. Chen CJ, Yu W, Fu YC, Wang X, Li JL, et al. (2009) Resveratrol protects cardiomyocytes from hypoxia-induced apoptosis through the SIRT1-FoxO1 pathway. *Biochem Biophys Res Commun* 378: 389–393.
57. Costa Cdos S, Rohden F, Hammes TO, Margis R, Bortolotto JW, et al. (2010) Resveratrol upregulated SIRT1, FOXO1, and adiponectin and downregulated PPARgamma1-3 mRNA expression in human visceral adipocytes. *Obes Surg* 21: 356–361.
58. Zhang HS, Zhou Y, Wu MR, Zhou HS, Xu F (2009) Resveratrol inhibited Tat-induced HIV-1 LTR transactivation via NAD(+)-dependent SIRT1 activity. *Life Sci* 85: 484–489.
59. Beher D, Wu J, Cumine S, Kim KW, Lu SC, et al. (2009) Resveratrol is not a direct activator of SIRT1 enzyme activity. *Chem Biol Drug Des* 74: 619–624.
60. Pacholec M, Bleasdale JE, Chrnyk B, Cunningham D, Flynn D, et al. (2010) SIRT1720, SIRT2183, SIRT1460, and resveratrol are not direct activators of SIRT1. *J Biol Chem* 285: 8340–8351.
61. Zhang Z, Lowry SF, Guarente L, Haimovich B (2010) Roles of SIRT1 in the acute and restorative phases following induction of inflammation. *J Biol Chem* 285: 41391–41401.
62. Kitada M, Kume S, Imaizumi N, Koya D (2011) Resveratrol improves oxidative stress and protects against diabetic nephropathy through normalization of Mn-SOD dysfunction in AMPK/SIRT1-independent pathway. *Diabetes* 60: 634–643.
63. Lan D, Lu M, Sharma S, Mellon PL, Olefsky JM, et al. (2011) Trans-resveratrol inhibits phosphorylation of Smad2/3 and represses FSHbeta gene expression by a Sirt1-independent pathway in LbetaT2 gonadotrope cells. *Reprod Toxicol* 32: 85–92.
64. Pan WW, Li JD, Huang S, Papadimos TJ, Pan ZK, et al. (2010) Synergistic activation of NF-kappaB by bacterial chemoattractant and TNF-alpha is mediated by p38 MAPK-dependent RelA acetylation. *J Biol Chem* 285: 34348–34354.
65. Quivy V, Van Lint C (2004) Regulation at multiple levels of NF-kappaB-mediated transactivation by protein acetylation. *Biochem Pharmacol* 68: 1221–1229.
66. Chen LF, Greene WC (2003) Regulation of distinct biological activities of the NF-kappaB transcription factor complex by acetylation. *J Mol Med* 81: 549–557.
67. Chen LF, Williams SA, Mu Y, Nakano H, Duerr JM, et al. (2005) NF-kappaB RelA phosphorylation regulates RelA acetylation. *Mol Cell Biol* 25: 7966–7975.
68. Chen LF, Mu Y, Greene WC (2002) Acetylation of RelA at discrete sites regulates distinct nuclear functions of NF-kappaB. *EMBO J* 21: 6539–6548.
69. Schaeffer V, Abrass CK (2010) Mechanisms and control of protein translation in the kidney. *Am J Nephrol* 31: 189–201.
70. Wullschlegel S, Loewth R, Hall MN (2006) TOR signaling in growth and metabolism. *Cell* 124: 471–484.
71. Lorne E, Zhao X, Zmijewski JW, Liu G, Park YJ, et al. (2009) Participation of mammalian target of rapamycin complex 1 in Toll-like receptor 2- and 4-induced neutrophil activation and acute lung injury. *Am J Respir Cell Mol Biol* 41: 237–245.
72. Lisi L, Navarra P, Feinstein DL, Dello Russo C (2011) The mTOR kinase inhibitor rapamycin decreases iNOS mRNA stability in astrocytes. *J Neuroinflammation* 8: 1.
73. Plaisance I, Morandi C, Murigande C, Brink M (2008) TNF-alpha increases protein content in C2C12 and primary myotubes by enhancing protein translation via the TNF-R1, PI3K, and MEK. *Am J Physiol Endocrinol Metab* 294: E241–250.
74. Glantschnig H, Fisher JE, Wesolowski G, Rodan GA, Reszka AA (2003) M-CSF, TNFalpha and RANK ligand promote osteoclast survival by signaling through mTOR/S6 kinase. *Cell Death Differ* 10: 1165–1177.
75. Lang CH, Pruznak AM, Frost RA (2005) TNFalpha mediates sepsis-induced impairment of basal and leucine-stimulated signaling via S6K1 and eIF4E in cardiac muscle. *J Cell Biochem* 94: 419–431.
76. Bonogio RG, Fuhro R, Wang Z, Valeri CR, Andry C, et al. (2005) Rapamycin ameliorates proteinuria-associated tubulointerstitial inflammation and fibrosis in experimental membranous nephropathy. *J Am Soc Nephrol* 16: 2063–2072.
77. Liu M, Agreda P, Crow M, Racusen L, Rabb H (2009) Effects of delayed rapamycin treatment on renal fibrosis and inflammation in experimental ischemia reperfusion injury. *Transplant Proc* 41: 4065–4071.