

Naturally Occurring Motility-Defective Mutants of *Salmonella enterica* Serovar Enteritidis Isolated Preferentially from Nonhuman Rather than Human Sources^{∇†}

Lucía Yim,^{1*} Laura Betancor,^{1,3} Arací Martínez,^{1,3} Clare Bryant,²
Duncan Maskell,² and José A. Chabalgoity¹

Department of Biotechnology, School of Medicine, Universidad de la República, Instituto de Higiene, Av. A. Navarro 3051, Montevideo CP 11600, Uruguay¹; Department of Veterinary Medicine, University of Cambridge, Madingley Road, Cambridge CB3 0ES, United Kingdom²; and Department of Bacteriology and Virology, School of Medicine, Universidad de la República, Instituto de Higiene, Av. A. Navarro 3051, Montevideo CP 11600, Uruguay³

Received 30 April 2011/Accepted 5 September 2011

Salmonellosis represents a worldwide health problem because it is one of the major causes of food-borne disease. Although motility is postulated as an important *Salmonella* virulence attribute, there is little information about variation in motility in natural isolates. Here we report the identification of a point mutation (T551 → G) in *motA*, a gene essential for flagellar rotation, in several *Salmonella enterica* serovar Enteritidis field isolates. This mutation results in bacteria that can biosynthesize structurally normal but paralyzed flagella and are impaired in their capacity to invade human intestinal epithelial cells. Introduction of a wild-type copy of *motA* into one of these isolates restored both motility and cell invasiveness. The *motA* mutant triggered higher proinflammatory transcriptional responses than an aflagellate isolate in differentiated Caco-2 cells, suggesting that the paralyzed flagella are able to signal through pattern recognition receptors. A specific PCR was designed to screen for the T551 → G mutation in a collection of 266 *S. Enteritidis* field isolates from a nationwide epidemic, comprising 194 from humans and 72 from other sources. We found that 72 of the 266 (27%) isolates were nonmotile, including 24.7% (48/194) of human and 33.3% (24/72) of food isolates. Among nonmotile isolates, 15 carried the T551 → G mutation and, significantly, 13 were recovered from food, including 7 from eggs, but only 2 were from human sources. These results suggest that the presence of paralyzed flagella may impair the ability of *S. Enteritidis* to cause disease in the human host but does not prevent its ability to colonize chickens and infect eggs.

Salmonella enterica serovar Enteritidis represents a major cause of food-borne human gastroenteritis worldwide, with a significant impact on public health. Infection usually occurs by ingestion of contaminated water or food, particularly eggs and other poultry-derived products (14). In humans, the disease is characterized by acute intestinal inflammation and diarrhea, which is normally self-limiting. In chickens, *S. Enteritidis* can asymptotically colonize the gastrointestinal tract and deeper organs of the animal and can be transmitted to the forming eggs through the transovarian route and to laid eggs through the eggshell (14, 18, 21).

Although rarely isolated in Uruguay before 1994, *S. Enteritidis* was the etiological agent of an epidemic of human gastroenteritis that occurred in the country between 1995 and 2004, peaking in 2001–2002 (5). Starting in 2009, *S. Enteritidis* reemerged, overtaking *Salmonella enterica* serovar Typhimurium as the primary cause of salmonellosis in Uruguay (data from the National *Salmonella* Center, NSC, Institute of Hygiene, Uruguay).

Similarly, FoodNet surveillance for 2009 in 10 U.S. states reported that salmonellae were the most common causes of food-borne infections, and among them, *S. Enteritidis* was the serovar most commonly isolated (<http://jama.ama-assn.org/cgi/content/full/303/21/2130>).

Motility and flagella are considered to be important virulence factors contributing to gastrointestinal disease caused by *Salmonella*. Previous studies in streptomycin-pretreated mice have shown that during infection, *S. Typhimurium* flagella and motility contribute to early cecal inflammation (37, 38). A flagellated but nonmotile mutant of *S. Enteritidis* showed reduced attachment to rat ileal explants compared to that of the corresponding wild-type strain in the presence of an intact mucus layer, suggesting that active flagella are important for *S. Enteritidis* to penetrate this layer (31). *In vitro*, it has been found that aflagellate *S. Typhimurium* mutants can attach to, but are defective in entering, cultured intestinal epithelial cells (32). Similarly, aflagellate mutants or mutants of *S. Enteritidis* with paralyzed flagella showed reduced kinetics of invasion of cultured intestinal epithelial cells compared to findings for the wild-type strain, even if the adhesion levels were similar (41). Moreover, flagella are the main stimuli for triggering of host proinflammatory responses through activation of Toll-like receptor 5 (TLR5) and Ipaf (26, 30). Although these data indicate important roles for flagella and motility in *Salmonella* virulence, most of the studies so far rely on work done with

* Corresponding author. Mailing address: Department of Biotechnology, Instituto de Higiene, School of Medicine, Av. A. Navarro 3051, Montevideo CP 11600, Uruguay. Phone: 5982 487 12 88, ext. 1120. Fax: 5982 487 30 73. E-mail: lyim@higiene.edu.uy.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

∇ Published ahead of print on 16 September 2011.

TABLE 1. Bacterial strains and plasmid used in this study

| Strain or plasmid | Description | Source and/or reference |
|-------------------------------|---|--------------------------------------|
| S. Enteritidis strains | | |
| 251/01 | Chicken egg isolate [<i>motA</i> (T551G)] | 4 |
| 8/02 | Human gastroenteritis isolate | 4 |
| LVR18 | 251/01, pLVR10 (Ap ^r) | This work |
| LVR19 | 251/01, pBAD22 (Ap ^r) | This work |
| LVR20 | 251/01 derivative, <i>motA::cat</i> (Cm ^r) | This work |
| LVR21 | 8/02 derivative, <i>motA::cat</i> (Cm ^r) | This work |
| LVR22 | <i>motA</i> ⁺ derivative of LVR20 (Cm ^s) | This work |
| LVR23 | <i>motA</i> ⁺ derivative of LVR21 (Cm ^s) | This work |
| PT4 P125109 | Wild-type, sequenced strain | Wellcome Trust Sanger Institute (39) |
| S. Typhimurium SL5338 | | |
| | <i>galE</i> r ⁻ m ⁺ | 9 |
| E. coli strains | | |
| DH5α | F ⁻ <i>endA1 hsdR17 supE44 thi1 recA1 gyrA relA1 Δ(lacZYA-argF) U169 (φ80lacZΔM15)</i> | 16 |
| SY327λpir | <i>Δ(lac pro) argE(Am) rif 20 nalA recA56 (λpir)</i> | 27 |
| Plasmids | | |
| pBAD22 | Expression vector with P _{BAD} and AraC control (Ap ^r) | 15 |
| pKD3 | Template plasmid, pANTS _γ derivative (Ap ^r Cm ^r) | 10 |
| pKD46 | Red recombinant expression plasmid, pINT-ts derivative (Ap ^r) | 10 |
| pLVR10 | <i>motA</i> inserted between sites EcoRI and HindIII of pBAD22 | This work |

artificially constructed mutant strains, but little is known about variations in flagellar function among field isolates and their effects on *Salmonella*-host interactions.

In *Salmonella*, about 50 genes are needed for flagellar assembly and function, but among these, only a few are directly involved in torque generation (24). MotA is one of the two proteins composing the stator responsible for torque generation in the proton-driven flagellar motor. Together with MotB, it forms a complex embedded in the cytoplasmic membrane, which is composed of four copies of MotA and two copies of MotB, which is essential for the proton translocation process that fuels the flagellar motion (3, 22, 28, 36).

Here we report the identification of a point mutation in *motA* in 15 *S. Enteritidis* isolates from a comprehensive collection of clinical and environmental samples. This mutation results in bacteria that harbor paralyzed flagella and are impaired for invasion of Caco-2 cells. All but two of the isolates identified as carrying this mutation were isolated from nonhuman sources.

MATERIALS AND METHODS

Bacterial strains, media, and growth conditions. A set of 266 *S. Enteritidis* strains isolated before, during, and after the epidemics of this serovar in Uruguay (covering 1988 to 2006) was obtained from the NSC collection. These included 194 human clinical isolates and 72 from animals or food as previously reported (4). Specific bacterial strains used in this study are listed in Table 1. LB broth and LB agar, supplemented with antibiotics when required (ampicillin, 100 μg/ml; chloramphenicol, 25 μg/ml), were used for routine cultures of *Escherichia coli* or *S. enterica* at 37°C in an orbital shaking incubator (200 rpm). *E. coli* DH5α was used as a host for cloning.

All *Salmonella* isolates were confirmed biochemically and serologically at the NSC and were stored in replicates at -80°C in Luria-Bertani (LB) broth containing 25% glycerol. They were resuscitated by gently scraping the surface of the frozen content with a sterile loop and streaked on LB agar plates. After overnight incubation, isolated colonies were inoculated in LB broth for further analysis.

S. Enteritidis PT4 P125109 (NCTC 13349, here referred to as PT4 [39]), was obtained from The Wellcome Trust Sanger Institute (Hinxton, United Kingdom) and used as a reference in phenotypic assays. Strain *S. Typhimurium* SL5338 (r⁻

m⁺) was used as an intermediate host for plasmid constructs to be transferred from *E. coli* to *Salmonella*.

S. Enteritidis strains LVR18 and LVR19 are isolate 251/01 transformed with plasmids pLVR10 and pBAD22, respectively (see below).

S. Enteritidis strains LVR20 and LVR21 are *motA::cat* derivatives of isolates 251/01 and 8/02, respectively, and were constructed using the Lambda Red method (10). Briefly, plasmid pKD3 was used as a template for PCR amplification of a DNA fragment containing the chloramphenicol resistance gene cassette, flanked by regions of homology to *motA*, using the primers motAP1b and motAP2b (see Table S1 in the supplemental material). Both primers have a 5' 47- to 50-mer region exhibiting perfect homology with the *motAB* locus of the *S. Enteritidis* PT4 chromosome and a 3' region homologous to priming site 1 or priming site 2 of pKD3, respectively. Three hundred nanograms of the resulting 1.1-kb PCR fragment was gel purified and electroporated into PT4 carrying pKD46 for insertion into its chromosome. Loss of the helper plasmid in the obtained chloramphenicol-resistant colonies was verified by growth on LB plates at 37°C, and sensitivity to ampicillin was checked by replica plating on LB plates containing ampicillin. Then, the *motA::cat* gene was introduced into the chromosome of isolate 251/01 or 8/02 by P22 transduction and selection of Cm^r and nonmotile transductants, resulting in strains LVR20 and LVR21, respectively. All chromosomal insertions were verified by PCR using primers C1 and C2 (see Table S1 in the supplemental material).

Strains LVR22 and LVR23 were obtained by P22 transduction of wild-type *motA* from PT4 into strains LVR20 and LVR21, respectively, and selection of motile and Cm^s transductants. This was achieved by spot plating the transduction mix onto soft agar LB plates (containing 0.3% agar) for 6 h at 37°C and picking bacteria with an inoculation loop about 2 cm away from the plating spot. The loopful of bacteria was streaked out onto LB plates, and after overnight incubation at 37°C, isolated colonies were replica plated on LB plates with or without chloramphenicol to verify sensitivity. Plating the transduction mix on LB plates without the recipient strain resulted in no colonies after overnight incubation at 37°C. The absence of *cat* insertion into *motA* was verified by PCR using primers motA1 and motB2.

All transductants were streaked out on Evans blue uridine agar to confirm the absence of contaminating phages.

Plasmid constructs and DNA manipulation. Genetic techniques were performed using standard laboratory methods. Plasmid and chromosomal DNA purifications were done according to protocols recommended by the supplier (Qiagen, Germany). DNA was digested with restriction endonucleases or ligated with T4 DNA ligase under standard conditions as per the manufacturer's instructions (Fermentas, Invitrogen). Preparation of electrocompetent *E. coli* and *S. enterica* cells and DNA transformation were performed as previously described (12).

Plasmids and primers used in this study are listed in Table 1 and in Table S1 in the supplemental material, respectively. pLVR10 contains the *motA* gene cloned into pBAD22 under the control of the arabinose-inducible P_{BAD} promoter and was constructed as follows. *motA* was PCR amplified from purified genomic PT4 DNA using the primers *motA5'* and *motA3'* and a 10:1 mixture of *Taq-Pfu* DNA polymerases (Fermentas). The resulting amplicon was digested with EcoRI and HindIII and ligated into pBAD22 previously cut with the same restriction enzymes. The plasmid construct was verified by sequencing of the cloned insert.

DNA sequencing. For sequencing of flagellar genes, genomic DNA was extracted from the bacterial strains using the DNeasy blood and tissue kit (Qiagen). Specific genes or regions were PCR amplified using Proof Start DNA polymerase (Qiagen) and the primers listed in Table S1 in the supplemental material. *fliC* (including regulatory regions) was amplified using the primers *fliC1/fliC4*, and the resulting 1.71-kb PCR product was sequenced with the primers *fliC1*, -2, -3, and -4, GFor, and GRev. For *fliG* sequencing, its coding region was amplified using the primers *fliG1/fliG3* and sequenced with *fliG1*, -2, and -3. *fliM* and *fliN* were amplified using primers *fliM1/fliN2*, and the resulting 1.55-kb PCR product was sequenced with *fliM1*, -2, -3 and *fliN1* and -2. For sequencing of *motA* and *motB*, the primers *motA1/motB2* were used to PCR amplify a 1.96-kb fragment, which was sequenced using the primers *motA1*, -2, and -3 and *motB1* and -2. Before sequencing, all PCR products were purified using a QIAquick PCR purification kit (Qiagen).

Motility tests. Motility tests were performed as described by Yim et al. (43). Briefly, 2 μ l of overnight cultures grown in LB broth were spotted onto the surface of an LB plate containing 0.3% agar (and arabinose when indicated) and incubated for 6 h at 37°C. Those isolates showing no halo of growth from the inoculation spot after 6 h of incubation (indistinguishable from a nonflagellated strain) were considered nonmotile. Values are expressed as a percentage of the diameter of growth (in mm) obtained for PT4. The assays were repeated three times, and the results were confirmed by phase-contrast microscope visualization of mid-log-phase bacterial cultures grown in LB broth.

Bacterial cell fractionation and protein analysis. For evaluation of bacterial surface protein profiles, heat extracts (HE), which are composed predominantly of flagella and other surface proteins, were obtained as described by Nicholas (29), with modifications. Briefly, mid-log-phase bacteria were harvested by centrifugation, resuspended in phosphate-buffered saline (PBS), and incubated for 1 h at 65°C. The samples were then centrifuged for 25 min at 1,000 \times g, and supernatants were quantified by Bradford assays (7). Equivalent amounts of protein (20 μ g) were loaded into 12% SDS-PAGE gels and visualized with Coomassie blue R250 staining. Protein extracts from all strains were prepared at the same time and by the same procedure.

Secreted proteins were obtained from the supernatant of mid-log-phase bacterial cultures, filtered through 0.22- μ m filters, and subsequently precipitated with trichloroacetic acid (TCA) 25% (final concentration). For preparation of total protein extracts, mid-log-phase bacterial cultures were centrifuged and resuspended in PBS, sonicated, and centrifuged again to remove unbroken cells. The supernatants (cleared lysates) were quantified by Bradford assays.

For Western blot analysis, 10 μ g of HE extracts and secreted proteins or 80 μ g of total extracts were loaded onto a 12% SDS-PAGE gel and analyzed by Western blotting using rabbit anti-FliC (Hm) antiserum (*Salmonella* H antiserum m; Difco).

Flagellar staining. For detection of flagella in live cells, we performed a previously described method using Alexa Fluor 594 carboxylic acid succinimidyl ester (Molecular Probes) (40). Briefly, overnight cultures of bacteria grown in LB broth at 37°C and 200 rpm were diluted 1/100 in fresh medium and grown in the same conditions to mid-log phase (optical density at 600 nm [OD_{600}] = 0.4 to 0.6). Then, the protocol was followed exactly as described previously (40). The samples were visualized on a Leica DM6000B fluorescence microscope using the Cy3.5 filter, and the images were acquired and manipulated using Leica AF6000 software. A minimum of 10 fields was recorded for each isolate.

Cell lines, medium, and growth conditions. The human colon carcinoma Caco-2 cell line was obtained from the American Type Culture Collection. The cells were maintained in minimal essential medium with Earle's Salts (high glucose, 4.5 g/liter), supplemented with 4 mM L-glutamine and 20% fetal calf serum at 37°C in 5% CO₂, at up to 80% confluence.

To polarize Caco-2 cells, 5 \times 10⁴ cells were seeded in Costar transwells (6.5-mm diameter, 4 μ m pore, polycarbonate) in 24-well plates and grown for 19 days, changing the culture medium every other day, until transepithelial electrical resistance (TEER) stabilization (TEER > 200 Ω · cm²). Impermeability of the monolayer, indicative of the differentiation level of the cells, was verified by a diffusion test using fluorescent beads (Fluospheres carboxylate-modified microspheres, 0.2 μ m, yellow/green fluorescent [505/515]; Molecular Probes).

Cell invasion assays. Caco-2 invasion assays were performed as previously described (43). Briefly, log-phase-grown bacteria were added to the cells at an MOI of ~30:1, the plates were centrifuged for 5 min at 200 \times g, and invasion was allowed to proceed for 1 h. Then, the medium was changed to gentamicin-containing medium (100 μ g/ml), and after 1.5 h the cells were washed and lysed with 0.1% Triton X-100 for bacterial release and counting. Data are expressed as the percentage of the initial inoculum. Each isolate was tested in duplicate in two independent experiments.

Quantitative real-time PCR. Polarized Caco-2 cells were infected apically at an MOI of ~30:1 and processed as for the invasion assays, but after 1.5 h of incubation with culture medium supplemented with 100 μ g/ml of gentamicin, the antibiotic concentration was changed to 10 μ g/ml and the cells were incubated for an additional 1.5 h. Then, at 3 h postinfection, the cells were gently washed 3 times with prewarmed PBS and resuspended in TRIzol (Invitrogen) for extraction of total RNA. After reverse transcription with random hexamers and real-time PCR using specific primers, threshold cycle (C_T) values were normalized with values of 18S RNA and referred to values of uninfected cells. Total RNA extraction, reverse transcription, and quantitative real-time PCR (qRT-PCR) were carried out as previously reported (43). Mean results of four independent experiments are shown.

PCR screening for *motA*(T551G) mutation. From the 266 field isolates subjected to motility tests, those totally devoid of motility were studied further using a specific PCR designed to detect the presence of the *motA*(T551G) mutation. Isolates with reduced motility compared to the reference were not included in the analysis. In brief, genomic DNA was extracted by resuspending one bacterial colony in 200 μ l of sterile milliQ water and boiling the suspension for 5 min. After that, the extract was centrifuged for 5 min at 10,000 \times g, and 0.5 μ l of the supernatant used as a template in a PCR using the primer *motA2* as a forward primer and *motAWT2* as a reverse primer (see Table S1 in the supplemental material). The cycling conditions were as follows: 3 min at 94°C and 30 cycles of 45 s at 95°C, 30 s at 60°C, and 30 s at 72°C. Those strains that gave no product carried a nucleotide in position 551 different from that of the wild type and were further analyzed by PCR with the primers *motA2* and *motAmut2*. In this case the cycling conditions were as described above but the annealing temperature was raised to 66°C to increase specificity. Obtaining a product of 113 bp in size indicated the presence of the *motA*(T551G) mutation. Those DNAs positive for the mutation were sequenced with primer *motA3* for verification of the T551 \rightarrow G change.

Statistical analysis. For analysis of differences in motility, invasiveness for Caco-2 cells, and the transcriptional response to the infection, we used the Mann-Whitney U test (GraphPad Prism 4.0 software program), considering a *P* value of <0.05 (two-tailed) to be statistically significant. For analysis of the frequency of strains positive for the *motA*(T551G) mutation, we used Fisher's exact test (GraphPad Prism software), considering alpha to be <0.05.

RESULTS

Motility analysis of *S. Enteritidis* isolates. Previously, we characterized a collection of 29 Uruguayan *S. Enteritidis* isolates from human, animal, and environmental sources (43). Ten isolates were impaired for motility, and this correlated with diminished invasion of Caco-2 cells in centrifuge-assisted assays. In this work, we performed a wider screening of motility in 266 natural isolates of *S. Enteritidis*, including 194 from humans and 72 from other sources, and found that 72 of them (48 from humans and 24 from nonhuman origins) were devoid of motility. Thus, 24.7% of human isolates were nonmotile, compared to 33.3% of the nonhuman isolates. However, this difference was not statistically significant (*P* = 0.16 in Fisher's exact test).

Characterization of the nonmotile phenotype of isolate 251/01. One of the nonmotile isolates (strain 251/01), obtained from a nationwide microbiological survey of chicken eggs (5), was selected for further comparative studies with one motile isolate (strain 8/02), obtained from a case of human gastroenteritis. Both strains were isolated during the peak of the *S. Enteritidis* epidemic in Uruguay and had previously revealed identical genetic profiles with all of the typing methods utilized




FIG. 1. (A) SDS-PAGE analysis of heat extracts from isolates 8/02 and 251/01. Location of FliC (53 kDa) in the gel is indicated to the right. Western blot analysis of heat extracts (B), secreted proteins (C), or total protein extracts (D) of isolates using anti-flagellin antibody. Fla-, a *Salmonella* isolate lacking flagella. (E) Fluorescent labeling of flagellar filaments in live cells of strain 8/02, 251/01, or PT4.

(4), yet they differed significantly in their ability to invade epithelial cells, survive in egg albumen, and colonize chicken organs (43). The *in vitro* growth rates of the two strains in rich medium were similar between them and to that of the PT4 reference strain (data not shown).

The lack of motility could be due to defects in flagellin synthesis, export, or assembly (in which case no flagellar filament would be observed at all) or in the mechanisms of flagellar motion (in which case the bacteria would harbor structurally normal but paralyzed flagella). To gain further information about this phenotype, we analyzed the protein content of heat extract (HE) fractions of the isolates, which are composed predominantly of flagellar filaments and other surface proteins. As seen in Fig. 1A, the two isolates contained similar amounts of flagellin (FliC) protein on their surfaces, suggesting no differences in FliC production or in assembly of the flagellar filament. FliC levels were further verified by Western blot analyses of HE, secreted proteins, and total protein extracts (Fig. 1B to D). Furthermore, the nucleotide sequences of the *fliC* gene and its promoter region were identical in both isolates and identical to that of the sequenced *S. Enteritidis* PT4 strain (data not shown).

We further investigated the presence, appearance, and movement of flagella by directly labeling the flagellar filaments with an amino-specific fluorescent dye in live cells (Fig. 1E). The flagellar filaments of isolate 251/01 were readily visualized, and the shape, length, and number were indistinguishable from those of 8/02 or PT4. However, while 8/02 and PT4 cells were actively swimming, no movement of the 251/01 cells was observed (data not shown). This result demonstrates that isolate

251/01 is flagellated but nonmotile, suggesting that the lack of motility in this strain is probably due to impaired function of motor proteins.

Sequence analysis of motor protein-coding genes. Since *motA* and *motB* encode essential components of the flagellar motor, we analyzed the sequence of both genes from isolates 8/02 and 251/01. While *motB* sequences were identical between the two isolates and to that of the PT4 strain, a T → G substitution was found in nucleotide 551 of *motA* from isolate 251/01 which changes an amino acid residue of the protein (Val184Gly). This residue is located in a highly conserved region of MotA, comprising one of the four membrane-spanning segments of the protein (Tm3), where several mutations that render the protein nonfunctional were previously described (Fig. 2) (6, 8). We also analyzed the nucleotide sequences of other motor genes (*fliG*, *fliM*, and *fliN*) from isolate 251/01, but no further differences were found compared to PT4. These results indicate that the motility impairment of isolate 251/01 is most likely due to a nonfunctional MotA.

Genetic complementation of the *motA*(T551G) mutant. In order to confirm that the lack of motility in isolate 251/01 is due to the *motA*(T551G) mutation, we performed genetic complementation. Isolate 251/01 was electroporated with plasmid pLVR10, carrying wild-type *motA* into an arabinose-inducible expression vector, and the resulting strain, LVR18, was subjected to motility assays with increasing concentrations of inducer (Fig. 3A). LVR18 displayed motility in an arabinose-dependent fashion, whereas the 251/01 isolate transformed with the empty vector did not. This demonstrates that the absence of motility in isolate 251/01 is due exclusively to the

CCL20 (MIP3A), and several proinflammatory chemokines (44), that recruits neutrophils and dendritic cells into the sub-epithelial compartment (35, 44). In polarized Caco-2 cells used as model epithelia, it has been demonstrated that upregulation of transcription of the *CCL20* and *IL8* genes in response to *Salmonella* infection depends on pattern recognition of flagellin by TLR5 (35). Thus, we tested if the isolate with paralyzed flagella would be able to trigger a response similar to that induced by the motile one. For this, we measured the *CCL20* and *IL8* mRNA levels from polarized Caco-2 cells infected with isolate 251/01 or 8/02 in comparison with those of uninfected cells (2, 44). Infection with the *motA* mutant induced levels of transcription of *CCL20* similar to those with the *motA*⁺ isolate, while a *Salmonella* isolate completely lacking flagella induced significantly reduced levels of *CCL20* transcript (Fig. 5A). Concerning *IL8* expression, isolate 251/01 induced lower levels of transcription than isolate 8/02 and higher levels than the aflagellate strain, but these differences were not statistically significant (Fig. 5B). These results suggest that the paralyzed flagella can still drive proinflammatory responses in Caco-2 cells, although with reduced efficiency compared to those with motile flagella.

PCR screening for *motA*(T551G) mutation. To investigate the prevalence of the *motA*(T551G) mutation in the strains circulating in Uruguay, we developed a specific PCR using a forward primer complementary to nucleotides 459 to 478 of *motA* and two reverse primers complementary to nucleotides 551 to 572, containing either the T551 → G change or the wild-type sequence at the 3' end, to specifically amplify the mutant or the wild-type sequence, respectively (Fig. 6). We analyzed the 72 nonmotile *S. Enteritidis* isolates identified in the motility screening (see above) and found in total 15 isolates carrying this mutation (including 251/01). Interestingly, 13 of these isolates were obtained from nonhuman sources, including 7 from eggs, and only 2 were from human sources (Table 2). This corresponds to 18.1% versus 1.0% of the total number of strains isolated from nonhuman or human sources, respectively ($P < 0.0001$ in Fisher's exact test). This difference is also significant if we consider only the nonmotile isolates from each origin ($P < 0.0001$). To rule out the possibility that the mutation occurred during prolonged storage of the isolates, we analyzed both replicates of each isolate stored, and the result was the same in both of them.

All isolates positive for the mutation in the PCR carried the T551 → G mutation, as confirmed by DNA sequencing. We tested all the *motA*(T551G) isolates in Caco-2 invasion assays, and all except one, the nonhuman isolate 31/04, were hypoinvasive compared to PT4 (Fig. 7).

DISCUSSION

Here we report the identification of a missense mutation in *motA* from *S. Enteritidis* field isolates, which is unevenly distributed among isolates according to their source. We show that the prevalence of this mutation is significantly higher in isolates from nonhuman sources than in those from clinical human sources. To our knowledge, this is the first time that a mutation in this gene has been identified in natural isolates of *Salmonella*. Since many of these bacteria have been stored for more than 10 years, it could be argued that the mutation may




FIG. 5. Analysis of the Caco-2 transcriptional response to infection by *S. Enteritidis* motile or paralyzed isolates. Polarized Caco-2 cells were infected with isolate 251/01 or 8/02, and at 3 h postinfection, the levels of mRNA transcripts for *CCL20* (A) or *IL8* (B) were measured by qRT-PCR. Values were normalized with values of 18S RNA and referenced to values of uninfected cells. Fla⁻, a *Salmonella* isolate lacking flagella; ns, not statistically significant ($P > 0.05$); *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

have appeared during the storage period. However, this would not explain why it is found with much more frequency among isolates derived from nonhuman sources than in those from human sources. Further, we confirmed the presence of the mutation in both replicate cryovials of each isolate, which suggests that the possibility that the mutation arose during storage is negligible. Since undirected genetic mutation arises in *Salmonella* in response to environmental stressors (25), this mutation could have occurred spontaneously, and its presence did not prevent the mutant bacteria from surviving in the environment or from infecting chickens. In fact, the avian-adapted *Salmonella enterica* Gallinarum serotype is nonmotile. The fact that the same mutation was found in strains isolated during various periods of time (Table 2) suggests that its ap-




FIG. 6. PCR screening for *motA*(T551G) mutation. Colony-extracted DNA was used as a template in a PCR specific for wild-type (A) or *motA*(T551G) (B) alleles. Amplicon size is 113 bp. (A) PCR using primers *motA*2/*motA*WT2, amplifying only wild-type *motA* alleles. (B) PCR using primers *motA*2/*motA*mut2, amplifying only *motA*(T551G) alleles. Only a fraction of isolates analyzed are shown. Sample 7, isolate 18/97; 8, 19/97; 9, 20/97; 10, 21/97; 14, PT4; 15, 8/02; 16, 251/01.

pearance is not an unlikely event or alternatively that the same clone has persisted for long periods of time.

An isolate carrying this mutation assembled normal flagella but had totally lost motility and showed reduced invasion of Caco-2 epithelial cells in culture compared to a motile isolate. However, the ability to attach to cells was previously shown to be comparable between both isolates (43). These results are in line with those reported for nonmotile insertional mutant strains of *S. Enteritidis* (41) and *S. Typhimurium* (32) and suggest a role for motile flagella in invasion but not in adhesion to epithelial cells.

It has been previously reported that in nonflagellated *S. Typhimurium*, the impairment in cell invasion can be largely reversed by applying a centrifugal force upon infection (19, 20), while no increase in invasiveness after centrifugation is observed for *Salmonella enterica* serovar Typhi that is lacking flagella or motility (23), suggesting the existence of differences in the entry mechanisms between these serovars. For *S. Enteritidis*, it was demonstrated that promoting contact between bacteria and cells by application of centrifugal force greatly enhances invasion but that this is not enough to bring the levels of invasion of a nonmotile strain up to those of a motile one (11, 41, 43). Moreover, the presence of paralyzed flagella

seems to be more detrimental for epithelial cell invasion than a total absence of flagella (41), which could be explained by steric hindrance caused by the paralyzed flagella (19). All these data suggest a role for functional flagella in addition to the facilitation of approximation of bacteria to the apical surface of the host cell.

We were able to assign the paralyzed phenotype of isolate 251/01 to a nonfunctional mutation in *motA* [*motA*(T551G)]. The introduction of a wild-type copy of *motA* in the chromosome of isolate 251/01 reverted the paralyzed phenotype and rendered the bacteria fully invasive in Caco-2 cells, indicating that the absence of motility is responsible for the impairment of invasion seen in this isolate.

One crucial step in the enteropathogenesis of *Salmonella* is its ability to trigger a proinflammatory response from the intestinal epithelium. We found that the *motA* mutant isolate elicited proinflammatory transcriptional responses in polarized Caco-2 cells, indicating that the paralyzed flagella, although being associated with a significant reduction in the invasiveness of the strain, are still able to signal through TLR5 in this model epithelium. These results are consistent with those reported by

TABLE 2. *S. Enteritidis* isolates positive for the *motA*(T551G) mutation in the NSC collection, isolated between 1988 and 2006

| Isolate ^a | Period of isolation | Source ^b |
|----------------------|---------------------|------------------------|
| 18/97 | Epidemic | Food |
| 19/97 | Epidemic | Food |
| 20/97 | Epidemic | Food |
| 21/97 | Epidemic | Food |
| 11/99 | Epidemic | Human gastroenteritis |
| 48/01 | Epidemic | Food |
| 117/01 | Epidemic | Egg |
| 118/01 | Epidemic | Egg |
| 251/01 | Epidemic | Egg |
| 252/01 | Epidemic | Egg |
| 253/01 | Epidemic | Egg |
| 254/01 | Epidemic | Egg |
| 255/01 | Epidemic | Egg |
| 31/04 | Postepidemic | Food |
| 93/04 | Postepidemic | Human invasive disease |

^a Strain designations adhere to the following rule: number of isolate/year of isolation.

^b "Food" refers to any product for human consumption (e.g., cake or sandwich), with the exception of eggs.




FIG. 7. Nonpolarized Caco-2 invasion assays of *motA*(T551G) isolates. Isolate 8/02, carrying wild-type *motA*, was included for comparison, as was PT4, used as a reference strain. Data are expressed as percentages of intracellular bacteria related to the initial inoculum and further normalized to the result for PT4 (considering this value equal to 100%). Means \pm SEM are shown. **, significant difference compared to result for PT4 ($P < 0.01$).

Winter et al. (42), who demonstrated that an *S. Typhimurium* *flgK* mutant, which is unable to assemble flagella and is consequently nonmotile but can still secrete flagellin, is as invasive as an *flgK fliC fljB* mutant (a non-flagellin-expressing mutant) and is significantly less invasive than the wild type in bovine ligated ileal loops. However, the *flgK* mutant elicited significantly more fluid secretion and MIP3A (CCL20) gene expression than the *flgK fliC fljB* mutant, suggesting that invasion-independent flagellin pattern recognition contributes to diarrhea during the early phase of *S. Typhimurium* infection in calves (42).

Data about epidemiology and motility in *Salmonella* are scarce. Grossman et al. (13) reported an association between decreased motility and decreased severity of illness in clinical isolates of *S. Typhi* from Indonesia. More recently, it was reported that impaired motility in four poultry-associated isolates of *S. Enteritidis* was associated with low invasiveness in differentiated Caco-2 cells and reduced virulence in the mouse typhoid model (33). In the present work, we screened a comprehensive collection of isolates derived from human clinical, animal, and environmental sources and found that 72 out of 266 (27%) showed total absence of motility. A specific survey for the *motA*(T551G) mutation revealed its presence in 15 isolates obtained during the epidemic and postepidemic periods in Uruguay, with it being recovered mainly from animal sources, including eggs, rather than from human sources. Thus, it can be hypothesized that the presence of proinflammatory but paralyzed flagella impairs the ability of *S. Enteritidis* to cause intestinal disease in the human host but does not prevent *Salmonella* infections of chickens nor egg contamination. Supporting this view, it has been reported that a *motAB::cat* mutant of *S. Enteritidis*, with paralyzed flagella, was recovered from livers and spleens of chickens in numbers similar to those of the wild type (1). Further, our previous results demonstrated that two *motA*(T551G) *S. Enteritidis* isolates (251/01 and 254/01) were capable of colonizing the spleens and reproductive tracts of 3 day-old-chicks (43). However, both isolates differed extensively in their chicken invasiveness (43), demonstrating that with natural isolates the picture is more complex than that obtained with artificially constructed mutants. To the best of our knowledge, there are no reports studying the capability of *Salmonella*-paralyzed mutants to infect eggs, but the fact that 7 out of the 15 *motA* mutants were isolated from eggs supports the hypothesis that this mutation does not prevent *S. Enteritidis* from contaminating them.

In mice, it has been reported that motility allows *S. Typhimurium* to benefit from the nutrients released in the context of an inflamed gut by increasing the growth rate and improving pathogen fitness to compete with the intestinal microbiota (37). We speculate that in the human host, the *motA* mutants would not be able to cause disease, although they may elicit a proinflammatory response, because the lack of motility prevents them from efficient access to the localized high-energy nutrients released upon inflammation, avoiding efficient gut colonization. In addition, the diminished ability of the *motA* isolate to enter intestinal epithelial cells may account for a reduced proinflammatory response in the intestine and reduced pathogenesis.

The hypothesis that paralyzed *S. Enteritidis* mutants, though able to colonize chickens and infect eggs, would be impaired in

causing disease in humans is challenging. It could be the result of the different ways the mammalian and avian hosts respond to *Salmonella* infection. In this regard, it has been reported that oral infection with *S. Typhimurium* induces upregulation of the TLR15 receptor in chicken ceca (17). Interestingly, TLR15 is an avian-specific TLR which is therefore absent in mammals, suggesting that different pathways are involved in the host response to *Salmonella* infection.

In conclusion, in this study we identified 72 natural isolates of *S. Enteritidis* lacking motility from a comprehensive collection of 266 epidemic-spanning isolates and determined the presence of paralyzed flagella as the cause of this phenotype in 15 of them, derived mainly from nonhuman sources. The cause of the absence of motility in the remaining 57 nonmotile isolates remains to be determined.

ACKNOWLEDGMENTS

This work was supported by a project grant from the Wellcome Trust (078168/Z/05/Z) and from Comisión Sectorial para la Investigación Científica (CSIC), Uruguay, I+D 2008.

REFERENCES

- Allen-Vercoe, E., A. R. Sayers, and M. J. Woodward. 1999. Virulence of *Salmonella enterica* serotype Enteritidis aflagellate and afimbriate mutants in a day-old chick model. *Epidemiol. Infect.* **122**:395–402.
- Anderle, P., et al. 2005. Novel markers of the human follicle-associated epithelium identified by genomic profiling and microdissection. *Gastroenterology* **129**:321–327.
- Berg, H. C. 2003. The rotary motor of bacterial flagella. *Annu. Rev. Biochem.* **72**:19–54.
- Betancor, L., et al. 2009. Genomic and phenotypic variation in epidemic-spanning *Salmonella enterica* serovar Enteritidis isolates. *BMC Microbiol.* **9**:237.
- Betancor, L., et al. 2010. Prevalence of *Salmonella enterica* in poultry and eggs in Uruguay during an epidemic due to *Salmonella enterica* serovar Enteritidis. *J. Clin. Microbiol.* **48**:2413–2423.
- Blair, D. F., and H. C. Berg. 1991. Mutations in the MotA protein of *Escherichia coli* reveal domains critical for proton conduction. *J. Mol. Biol.* **221**:1433–1442.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**:248–254.
- Braun, T. F., L. Q. Al-Mawsawi, S. Kojima, and D. F. Blair. 2004. Arrangement of core membrane segments in the MotA/MotB proton-channel complex of *Escherichia coli*. *Biochemistry* **43**:35–45.
- Brown, A., et al. 1987. An attenuated *araA Salmonella typhimurium* vaccine elicits humoral and cellular immunity to cloned beta-galactosidase in mice. *J. Infect. Dis.* **155**:86–92.
- Datsenko, K. A., and B. L. Wanner. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* **97**:6640–6645.
- Dibb-Fuller, M. P., E. Allen-Vercoe, C. J. Thorns, and M. J. Woodward. 1999. Fimbriae- and flagella-mediated association with and invasion of cultured epithelial cells by *Salmonella enteritidis*. *Microbiology* **145**(Pt. 5):1023–1031.
- Dower, W. J., J. F. Miller, and C. W. Ragsdale. 1988. High efficiency transformation of *E. coli* by high voltage electroporation. *Nucleic Acids Res.* **16**:6127–6145.
- Grossman, D. A., et al. 1995. Flagellar serotypes of *Salmonella typhi* in Indonesia: relationships among motility, invasiveness, and clinical illness. *J. Infect. Dis.* **171**:212–216.
- Guard-Petter, J. 2001. The chicken, the egg and *Salmonella enteritidis*. *Environ. Microbiol.* **3**:421–430.
- Guzman, L. M., D. Belin, M. J. Carson, and J. Beckwith. 1995. Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. *J. Bacteriol.* **177**:4121–4130.
- Hanahan, D. 1983. Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**:557–580.
- Higgs, R., et al. 2006. Induction of a novel chicken Toll-like receptor following *Salmonella enterica* serovar Typhimurium infection. *Infect. Immun.* **74**:1692–1698.
- Humphrey, T. J. 1994. Contamination of egg shell and contents with *Salmonella enteritidis*: a review. *Int. J. Food Microbiol.* **21**:31–40.
- Jones, B. D., C. A. Lee, and S. Falkow. 1992. Invasion by *Salmonella typhi*

- murium* is affected by the direction of flagellar rotation. *Infect. Immun.* **60**:2475–2480.
20. Jones, G. W., L. A. Richardson, and D. Uhlman. 1981. The invasion of HeLa cells by *Salmonella typhimurium*: reversible and irreversible bacterial attachment and the role of bacterial motility. *J. Gen. Microbiol.* **123**:351–360.
 21. Keller, L. H., C. E. Benson, K. Krotec, R. J. Eckroade. 1995. *Salmonella enteritidis* colonization of the reproductive tract and forming and freshly laid eggs of chickens. *Infect. Immun.* **63**:2443–2449.
 22. Kojima, S., and D. F. Blair. 2004. The bacterial flagellar motor: structure and function of a complex molecular machine. *Int. Rev. Cytol.* **233**:93–134.
 23. Liu, S. L., T. Ezaki, H. Miura, K. Matsui, and E. Yabuuchi. 1988. Intact motility as a *Salmonella typhi* invasion-related factor. *Infect. Immun.* **56**:1967–1973.
 24. Macnab, R. M. 1992. Genetics and biogenesis of bacterial flagella. *Annu. Rev. Genet.* **26**:131–158.
 25. Massey, R. C., P. B. Rainey, B. J. Sheehan, O. M. Keane, and C. J. Dorman. 1999. Environmentally constrained mutation and adaptive evolution in *Salmonella*. *Curr. Biol.* **9**:1477–1480.
 26. Miao, E. A., E. Andersen-Nissen, S. E. Warren, and A. Aderem. 2007. TLR5 and Ipaf: dual sensors of bacterial flagellin in the innate immune system. *Semin. Immunopathol.* **29**:275–288.
 27. Miller, V. L., and J. J. Mekalanos. 1988. A novel suicide vector and its use in construction of insertion mutations: osmoregulation of outer membrane proteins and virulence determinants in *Vibrio cholerae* requires *toxR*. *J. Bacteriol.* **170**:2575–2583.
 28. Minamino, T., K. Imada, and K. Namba. 2008. Molecular motors of the bacterial flagella. *Curr. Opin. Struct. Biol.* **18**:693–701.
 29. Nicholas, R. A., and G. A. Cullen. 1991. Development and application of an ELISA for detecting antibodies to *Salmonella enteritidis* in chicken flocks. *Vet. Rec.* **128**:74–76.
 30. Ramos, H. C., M. Rumbo, and J. C. Sirard. 2004. Bacterial flagellins: mediators of pathogenicity and host immune responses in mucosa. *Trends Microbiol.* **12**:509–517.
 31. Robertson, J. M., et al. 2000. Adhesion of *Salmonella enterica* var *Enteritidis* strains lacking fimbriae and flagella to rat ileal explants cultured at the air interface or submerged in tissue culture medium. *J. Med. Microbiol.* **49**:691–696.
 32. Schmitt, C. K., et al. 2001. Absence of all components of the flagellar export and synthesis machinery differentially alters virulence of *Salmonella enterica* serovar Typhimurium in models of typhoid fever, survival in macrophages, tissue culture invasiveness, and calf enterocolitis. *Infect. Immun.* **69**:5619–5625.
 33. Shah, D. H., et al. 2011. Cell invasion of poultry-associated *Salmonella* Enteritidis isolates is associated with pathogenicity, motility and secretion of type-three secretion system secreted proteins. *Microbiology* **157**:1428–1445.
 34. Sharp, L. L., J. Zhou, and D. F. Blair. 1995. Features of MotA proton channel structure revealed by tryptophan-scanning mutagenesis. *Proc. Natl. Acad. Sci. U. S. A.* **92**:7946–7950.
 35. Sierro, F., et al. 2001. Flagellin stimulation of intestinal epithelial cells triggers CCL20-mediated migration of dendritic cells. *Proc. Natl. Acad. Sci. U. S. A.* **98**:13722–13727.
 36. Sowa, Y., and R. M. Berry. 2008. Bacterial flagellar motor. *Q. Rev. Biophys.* **41**:103–132.
 37. Stecher, B., et al. 2008. Motility allows *S. Typhimurium* to benefit from the mucosal defence. *Cell Microbiol.* **10**:1166–1180.
 38. Stecher, B., et al. 2004. Flagella and chemotaxis are required for efficient induction of *Salmonella enterica* serovar Typhimurium colitis in streptomycin-pretreated mice. *Infect. Immun.* **72**:4138–4150.
 39. Thomson, N. R., et al. 2008. Comparative genome analysis of *Salmonella* Enteritidis PT4 and *Salmonella* Gallinarum 287/91 provides insights into evolutionary and host adaptation pathways. *Genome Res.* **18**:1624–1637.
 40. Turner, L., W. S. Ryu, and H. C. Berg. 2000. Real-time imaging of fluorescent flagellar filaments. *J. Bacteriol.* **182**:2793–2801.
 41. van Asten, F. J., H. G. Hendriks, J. F. Koninkx, and J. E. van Dijk. 2004. Flagella-mediated bacterial motility accelerates but is not required for *Salmonella* serotype Enteritidis invasion of differentiated Caco-2 cells. *Int. J. Med. Microbiol.* **294**:395–399.
 42. Winter, S. E., et al. 2009. Contribution of flagellin pattern recognition to intestinal inflammation during *Salmonella enterica* serotype Typhimurium infection. *Infect. Immun.* **77**:1904–1916.
 43. Yim, L., et al. 2010. Differential phenotypic diversity among epidemic-spanning *Salmonella enterica* serovar Enteritidis isolates from humans or animals. *Appl. Environ. Microbiol.* **76**:6812–6820.
 44. Zeng, H., et al. 2003. Flagellin is the major proinflammatory determinant of enteropathogenic *Salmonella*. *J. Immunol.* **171**:3668–3674.