

Isolation of *Serratia marcescens* SR₁ as a Source of Chitinase Having Potentiality of Using as a Biocontrol Agent

K. Parani · G. P. Shetty · B. K. Saha

Received: 16 January 2008 / Accepted: 29 May 2009 / Published online: 7 May 2011
© Association of Microbiologists of India 2011

Abstract *Serratia marcescens*, strain SR₁ was isolated from the local soil of a cultivated farm and it was screened as potent strain for chitinase production. Maximum chitinase production ($77.3 \text{ u Mh}^{-1} 100^{-1}$) was observed after 96 h of incubation period with pH 5.5 at 30°C under shake conditions (120 rpm). Compare to still flasks, shake culture with prawn fish colloidal chitin of 0.5% (w/v) concentration, showed a better enzyme yield. Crude enzyme showed antifungal activity against plant pathogens.

Keywords Chitinase enzyme · Colloidal chitin · Plant pathogens · *Serratia marcescens*

Introduction

Chemical pesticides are widely used to control the plant diseases. However, most pesticides are hazardous to the ecological systems. Biological control promises to be a useful alternative approach in the control of plant pathogens in the sustainable agriculture system [1]. Biological pesticides are mainly formulated with the microbial metabolites. Among them, chitinase is a secondary metabolite novel enzyme, which effective against a number of plant pathogens. Chitinase catalyses the hydrolysis of chitin between the C₁ and C₄ of two consecutive N-Acetylglucosamines. The enzyme is regarded as important for the extensive carbon and nitrogen recycle in nature. Chitinase occurs widely in soil microorganisms and in some plants, fulfilling a possible defence role in them [2].

Chitinase is found widely in the soil microorganisms, including *Trichoderma* sp., *Bacillus* sp., and *Serratia* sp. have a chitinase producing ability [3–5]. It showed pesticidal effects against a number of plant pathogens including *Rhizoctonia solani* and *Fusarium oxysporum* causing wilt disease [6]. *Serratia marcescens*, a gram negative bacterium, soil inhabitant, is very efficient in degradation of chitin because of its ability to produce different chitinolytic enzymes [7]. Chitinase production and its activity depends on a number of limiting factors viz., culture state, temperature and pH of media etc.

In the present study, an attempt was made to optimize the production of chitinase enzyme from the isolated strain *S. marcescens* (SR₁) and screening of its biological activity against a number of fungal pathogens.

Materials and Methods

Isolation of Chitinolytic Microbes from Soil

A number of local strains of *Bacillus* sp., *Serratia* sp. and *Trichoderma* sp. were isolated from the soil of Northern parts of Bangalore, India. The isolated strains were purified according to the standard microbiological techniques. Strains were identified as described [8]. The purified isolates were maintained in the nutrient agar and MEA slants at ±4°C in refrigerator.

Screening for Potent Strains

Purified strains were plated on the chitin plates for the determination of their chitinase activity. Chitin plates were made with use of chitin agar medium containing 2% colloidal chitin (prawn shell), 0.5% yeast extract, 0.05%

K. Parani · G. P. Shetty · B. K. Saha (✉)
Research and Development, Multiplex Biotech Private Limited,
Peenya 1st Stage, Bangalore 560058, India
e-mail: bksaha2@rediffmail.com

MgSO_4 , 0.2% sodium nitrate, 0.05% KCl, FeSO_4 pinch, 0.1% K_2HPO_4 and 2% agar (w/v), adjusted to pH 6.0 using 1 N NaOH/HCl. Colloidal chitin was processed using prawn shell [9]. The medium was autoclaved at 121°C for 15–20 min. The purified isolates were inoculated onto chitin plates and incubated for 3–5 days at 30 ± 2°C. The potent strain was selected on basis of the formation clear zones on the plates.

To standardize the enzyme production, a log phase culture of the potent strain (10^9 cells/ml) was inoculated into 100 ml (working volume) of sterile synthetic broth with colloidal chitin as substrate. The flasks were incubated at different time intervals, pH, temperature in both still and shake conditions. All experiments were carried out in triplicates.

Optimization studies for enzyme production and activity was continued through the fermentation of SR₁ using a 5 l glass fermentor with a provision of air at 0.8 vvm. The fermentor along with 3 l media with pH 5.0 was sterilized at 121°C for 15–20 min. The fermentor was inoculated with 300 ml log phase seed culture of SR₁ and fermentation was carried for a period of 8 days of incubation.

Assay of Cultures

At different incubation time intervals, cultures were clarified by filtration through Whatmann No.1 filter paper and centrifuged (1,800×g for 15 min at 1°C). Enzyme activities were determined after dialysis at 4°C for 24 h against about 200 vol. of distilled water, pH 6.0. The assay was based on the estimation of reducing sugars released during the hydrolysis of colloidal chitin. The reaction mixture, containing 1.0 ml of 0.5% colloidal chitin (suspended in 50 mM sodium acetate buffer, pH 5.2) and 1.0 ml of enzyme solution was incubated for 1 h at 40°C with intermittent shaking and then centrifuged at 4,000 rpm for 5 min. The amount of reducing sugar released in the supernatant was determined by using *N*-acetylglucosamine as standard [10].

Biological Activity Test

The cell free culture filtrate of the potent strain was tested against *Fusarium oxysporum*, *Rhizoctonia solani*, *Helminthosporium*, *Xanthomonas putida* by cup-plate method.

Results and Discussion

Identification of the Potent Strain

Comparative studies for chitinase activity showed that the strain SR₁, *Serratia marcescens* was the potent among the

other isolates. Chitinase production in *S. marcescens* are induced by the inducer, are soluble oligomers derived from chitin, but not the monomer (*N*-acetylglucosamine) [11].

Effect on Age of the Culture

The chitinase activity of SR₁ was maximum at 96 h of incubation ($77.3 \mu\text{M h}^{-1} 100^{-1}$) when it was grown in liquid synthetic media, supplemented with prawn shell

Table 1 Effect of incubation time (h) on chitinase activity ($\mu\text{M/h}/100 \text{ ml}$) by *Serratia marsecens* strain (SR₁) grown in still and shake cultures

S. no.	Incubation time (h)	Chitinase activity ($\mu\text{M/h}/100 \text{ ml}$)	
		Still culture	Shake culture
1.	0	0	0
2.	24	17.3 ± 0.8	25.8 ± 1.3
3.	48	43.7 ± 2.1	50.1 ± 2.1
4.	72	59.4 ± 2.5	61.2 ± 2.5
5.	96	65.3 ± 3.2	77.3 ± 4.2
6.	120	48.6 ± 2.2	54.3 ± 2.5
7.	144	11.3 ± 0.9	21.2 ± 1.3

Table 2 Effect of temperature (°C) on chitinase activity ($\mu\text{M/h}/100 \text{ ml}$) by *Serratia marsecens* strain (SR₁) grown in still and shake cultures

S. no.	Temperature (°C)	Chitinase activity ($\mu\text{M/h}/100 \text{ ml}$)	
		Still culture	Shake culture
1.	0	0	0
2.	20	8.2 ± 0.42	18.7 ± 0.8
3.	25	23.4 ± 1.3	51.6 ± 2.3
4.	30	56.7 ± 2.5	76.7 ± 4.3
5.	35	52.1 ± 2.3	65.5 ± 3.7
6.	40	42.5 ± 2.1	55.3 ± 3.1
7.	45	35.3 ± 3.3	44.7 ± 2.1
8.	50	23.8 ± 1.3	35.2 ± 1.8
9.	55	14.5 ± 1.3	23.6 ± 1.3

Table 3 Effect of pH on chitinase activity ($\mu\text{M/h}/100 \text{ ml}$) by *Serratia marsecens* strain (SR₁) grown in still and shake cultures

S. no.	pH	Chitinase activity ($\mu\text{M/h}/100 \text{ ml}$)	
		Still culture	Shake culture
1.	0	0	0
2.	3.0	17.4 ± 0.8	26.4 ± 1.3
3.	5.0	57.3 ± 2.5	76.6 ± 4.1
4.	7.0	40.5 ± 2.1	60.3 ± 2.5
5.	9.0	29.3 ± 1.3	48.4 ± 2.1

Table 4 Effect of cell free culture filtrate of *Serratia marsecens* strain (SR₁) on growth of fungal pathogens

Culture filtrate (ml) of <i>Serratia marsecens</i> (SR ₁)	<i>Fusarium oxysporum</i>		<i>Sclerotium rolfsii</i>		<i>Rhizoctonia solani</i>		<i>Alternaria alternata</i>	
	Radial growth (mm)	% Inhibition zone	Radial growth (mm)	% Inhibition zone	Radial growth (mm)	% Inhibition zone	Radial growth (mm)	% Inhibition zone
Control	45.0 ± 2.1	0	45.0 ± 2.1	0	45.0 ± 2.1	0	45.0 ± 2.1	0
1.0	15 ± 0.7	66.6 ± 3.1	12.0 ± 0.81	73.3 ± 3.4	17.0 ± 0.8	62.2 ± 2.6	18.0 ± 0.73	60.0 ± 2.5
3.0	10 ± 0.52	77.7 ± 3.64	8.0 ± 0.34	82.2 ± 4.1	13.0 ± 1.1	71.1 ± 5.5	14.0 ± 0.63	68.8 ± 3.2
5.0	5.0 ± 0.23	88.8 ± 4.1	3.0 ± 0.12	93.3 ± 4.62	6.0 ± 0.24	86.6 ± 3.2	5.0 ± 0.23	88.8 ± 4.2

colloidal chitin, as the sole carbon source, at 30°C with pH 5.0 in shake culture (120 rpm) (Table 1). Compare to the shake culture, the still culture showing less activity (65.3 μM h⁻¹ 100⁻¹). Similar results were observed in *Trichoderma harzianum* (39.1 μM h⁻¹ 100⁻¹) [12].

Effect of pH and Temperature

A wide range of effect of pH on chitinase activity was shown by SR₁ (Table 3). The rate of hydrolysis of chitin was increased almost linearly between pH 3.5–5.0 in both still and shake cultures. A fairly broad optimum pH occurred with maximal activity at pH 5.0. Beyond which the activity slowly declined. Initial pH value of the culture medium of pH 5.0 showed the highest chitinolytic activity in 7-day-old shaken culture using *Aspergillus carneus* strain [13]. Studies over a different periods of incubation of SR₁, determined the optimum temperature of chitinase activity was at 30°C (Table 2). The optimum temperature reflects a balance between the effect on the activity and the maximum yield of the enzyme protein. This was positively correlated on the chitinase production by *Trichoderma harzianum* in submerged fermentation [14, 15].

Fermentations

The chitinase activity was increased at the beginning of the cultivation and reached to a maximum level at the log phase. Subsequently, the chitinase activity kept as plateau/decreased. The highest chitinase activity appeared on the fourth day from the day of inoculation. Started with fifth day, the production of chitinase was going to decreased. The decrease in the chitinase yield may be due to shear inactivation of the enzyme [16]. Similarly, the highest chitinase activity (18.2 μM/ml) was obtained under the optimal cultivation conditions of aeration rate at 0.6 vvm, pH 4.0, agitation rate at 150 rpm and 24°C with a 5 l stirred tank bioreactor (STR) [17]. The analysis on fermentation process indicated that chitinase belonged to the growth associated enzyme and also an inducible enzyme. Accordingly, the synthesis of this kind of enzyme can be induced [18]. However, when the inducer is removed/the

cell is in stationary phase, the enzyme synthesis will be blocked, indicating that, the mRNA of this kind of enzyme is unstable. Within a certain pH range would enhance the stability of the mRNA and thus prolonged the duration of enzyme production [19].

Effect of *S. marsecens* strain (SR₁) cell free filtrate on fungal pathogens: in order to find out, whether the bacterial component was responsible for the antagonistic/antifungal activity, the effect of culture filtrate on growth of fungal pathogens viz., *Fusarium oxysporum*, *Sclerotium rolfsii*, *Rhizoctonia solani*, *Alternaria alternata* were studied. The addition of 1 ml cell free culture filtrate of *Serratia marsecens* was sufficient to inhibit the fungal growth by 60.0–73.3% for fungal pathogens with 3–4 days of incubation (Table 4). Increasing the amount of culture filtrate to 3 and 5 ml reduced the fungal biomass by 85–93% for fungal pathogens respectively.

Major objective of this study was to isolate a potentially useful bacterial antagonist for biocontrol of fungal pathogens. Chitinase plays a major role in degrading fungal cell walls [20]. Micro organisms capable of producing and excreting chitinase have been shown to be efficient biocontrol agents [21–23]. This study has demonstrated the presence of cell wall degrading enzymes in *Serratia marsecens* which is mainly responsible for antifungal activity.

References

- Weller DM (1988) Biological control of soil borne plant pathogens in the rhizosphere with bacteria. Annu Rev Phytopathol 26:379–407
- Liu BL, Kao PM, Tzeng YM, Feng KC (2003) Production of chitinase from *Verticillium lecanii* FO91 using submerged fermentation. Enzyme Microb Technol 13:410–415
- Felse PA, Panda T (1999) Self-directing optimization of parameters for extracellular chitinase production by *Trichoderma harzianum* in batch mode. Process Biochem 34:563–566
- Frandsberg E, Schnurer J (1994) Chitinolytic properties of *Bacillus pabuli* K1. J Appl Bacteriol 76:361–367
- Young ME, Bell RL, Carroll PA (1985) Kinetics of chitinase production II. Relation between bacterial growth, chitin hydrolysis and enzyme synthesis. Biotechnol Bioeng 27:776–780

6. Someya N, Kataoka N (2000) Biological control of cyclamen soilborne diseases by *Serratia marcescens* strain B2. Plant Dis 84:334–340
7. Brurberg MB, Eijisink VGH, Venema G, Nes IF (1995) Chitinase B from *Serratia marcescens* BJL200 is exported to the periplasm without processing. Microbiology 141:123–131
8. Aneja KR (2003) Experiments in microbiology, plant pathology and biotechnology, 4th edn. New Age International (P) Ltd, New Delhi, pp 276–281
9. Leger ST, Charnley J, Cooper RM (1986) Cuticle degrading enzymes of entomopathogenic fungi; mechanisms of interactions between pathogen enzymes and insect cuticle. J Invertebr Pathol 47:295–302
10. Miller GL (1959) Use of ditrosalicylic acid reagent for determination of reducing sugars. Anal Chem 31:426–428
11. Monreal J, Reese ET (1969) The chitinase of *Serratia marcescens*. Can J Microbiol 15:689–696
12. Ulhoa CJ, Peberedy JF (1991) Regulation of chitinase synthesis in *Trichoderma harzianum*. J Gen Microbiol 137:2163–2169
13. Sherief A, El-Sawah MMA, Abd El-Naby MA (1991) Some properties of chitinase produced by a potent *Aspergillus carneus* strain. Appl Microbiol Biotechnol 35:228–230
14. El-Katatny MH, Somtsch W, Robra KH, Giigitz GM (2000) Production of chitinase and 1, 3-glucanase by *Trichoderma harzianum* for control of the phytopathogenic fungus *Sclerotium rolfsii*. Food Technol Biotechnol 38:173–180
15. Sandya C, Adapa LK, Nampoothiri MK, Szakacs G, Pandey A (2004) Extracellular chitinase production by *Trichoderma harzianum* in submerged fermentation. J Basic Microbiol 44:49–58
16. Felse PA, Panda T (2000) Submerged culture production of chitinase by *Trichoderma harzianum* in stirred-tank bioreactors: the influence of agitator speed. Biochem Eng J 4:115–120
17. Jacob F, Monod J (1961) Genetic regulatory mechanisms in the synthesis of proteins. Mol Biol (USSR) 3:318–356
18. Feng Y, Heb Z, Inga SL, Hua J, Zhang Z, Nga WJ (2003) Optimization of agitation, aeration and temperature conditions for maximum β -mannase production. Enzyme Microb Technol 32: 282–289
19. Mitchell R, Alezander M (1963) Lysis of soil fungi by bacteria. Can J Microbiol 9:169–177
20. Lim H, Kim Y, Kim S (1991) *Pseudomonas stutzeri* YPL-I genetic transformation and antifungal mechanism against *Fusarium solani*, an agent of plant root-rot. Appl Environ Microbiol 57:510–516
21. Schirmbock M, Lorito M, Wang Y, Hayes CK, Arisan-Atac I, Scala F, Harman GE, Kubieck CP (1994) Parallel formation and synergism of hydrolytic enzymes and peptaibol antibiotics, molecular mechanisms involved in the antagonistic action of *Trichoderma harzianum* against phytopathogenic fungi. Appl. Environ Microbiol 60:4364–4370
22. Kobayashi DY, Reedy RM, Bick JA, Qudemans PV (2002) Characterization of a chitinase gene from *Stenotrophomonas maltophilia* strain 34SI and its involvement in biological control. Appl. Environ Microbiol 68:1047–1054
23. Zhang Z, Yuen GY (2000) The role of chitinase production by *Stenotrophomonas maltophilia* strain C₃ in biological control of *Bipolaris sorokiniana*. Phytopathology 90:384–389