



IL28B SNP genotyping among Iranian HCV-infected patients: A preliminary report

Nastaran Mahboobi ¹, Bita Behnava ^{2*}, Seyed Moayed Alavian ²

¹ Tehran Hepatitis Center, Tehran, IR Iran

² Baqiyatallah Research Center for Gastroenterology and Liver Diseases, Baqiyatallah University of Medical Sciences, Tehran, IR Iran

ARTICLE INFO

Article Type:
Letter to Editor

Article history:
Received: 07 Mar 2011
Revised: 17 Apr 2011
Accepted: 2 May 2011

Keywords:
IL28B
Genotype
Hepatitis C virus
Iran

► *Please cite this paper as:*

Mahboobi N, Behnava B, Alavian SM. IL28B SNP genotyping among Iranian HCV-infected patients: A preliminary report. *Hepat Mon.* 2011;11(5):386-387.

© 2011 Kowsar M.P.Co. All rights reserved.

Dear Editor,

At the beginning of the third millennium, Hepatitis C virus (HCV) infection continues to be a major health hazard, with 170 million chronically infected people worldwide (1). Recent research has found that the prevalence of HCV infection in Iran is 1% and 0.1% in males and females, respectively (2). Although a combination of pegylated interferon (PEG-IFN) and ribavirin (RBV) has been established as a standard anti-HCV therapy, treatment responses vary. Numerous factors, including gender, age, HCV RNA level prior to treatment, liver cirrhosis, and HCV genotypes have been determined to contribute to this discrepancy (1). In addition, a number of recently published, genome-wide association studies (GWAS) have shown that polymorphisms within the IL28B locus result in differences between patients' responses to anti-HCV treatment (3-6). In this regard, rs8099917 (8 kb upstream of the IL28B gene) and rs12979860 (3 kb upstream of the IL28B gene) have been identified as

relevant single nucleotide polymorphisms (SNPs) that significantly impact treatment response to HCV infection (7). Although both SNPs are known to be independent predictors of response to anti-HCV therapy, a recent, extensive meta-analysis has reported that rs12979860 is more closely related to treatment responses (8). The prevalence of these SNPs and the rate of sustained viral response (SVR) have been found to vary across Caucasian, African American, and Japanese ethnicities (1). Based on these important findings, this preliminary report presents data on IL28B SNP (rs12979860) in a group of Iranian HCV-infected patients. To our knowledge, this is the first such report published from this region, and we hope that the results will stimulate extensive research.

In this study, 68 HCV-infected patients were referred to and enrolled in 2009 at the Tehran Hepatitis Center, a primary referral hepatitis center in Iran. Of these patients, 46 (67.6%) were male and 22 (32.4%) were female, with a Mean \pm SD age of 44.57 ± 13.67 years (ranging from 10 to 75 years). Whole blood (2-ml samples) in EDTA tubes was collected from these patients for genotyping analysis. DNA was extracted from 200 μ l of whole blood using a pure gene Blood Core C kit (Qiagen, Hilden, Germany)

* Corresponding author at: Bita Behnava, Baqiyatallah Research Center for Gastroenterology and Liver Disease, Baqiyatallah University of Medical Sciences, Tehran, IR Iran. Tel: +98-2188945186, Fax: +98-2181262072.

E-mail: bita_behnava@yahoo.com

Table1. SNP genotyping results of the study group

	Gender		
	Male [No.(%)]	Female [No.(%)]	
IL28			
CC	24 (66.7%)	12 (33.3%)	36
CT	19 (70.4%)	8 (29.6%)	27
TT	3 (60%)	2 (40%)	5
Total	46	22	68

Table2. HCV genotypes of the study group according to SNP genotyping

	HCV Genotype						Total
	1	3	non-typable	3a and 1a	2	4	
IL28B							
CC	21	9	2	1	1	0	34
CT	21	3	0	1	0	1	26
TT	2	1	0	0	0	0	3
Total	44	13	2	2	1	1	63

according to the manufacturer's manual. The quality and the purity of extracted DNA were measured with a NanoDrop 1000 spectrophotometer. Real-time PCR was performed on IL28B SNP (rs12979860) using Roche LightCycler 480 II according to Ge, *et al.*'s publication (3). The genotyping results were reported as C/C, C/T, and T/T. The SNP genotyping results for the study group, as shown in Table 1, showed C/C in 52.9% (No.= 36), C/T in 39.7% (No. = 27) and T/T in 7.3% of patients (No. = 5). Table 2 shows HCV genotypes of the study group according to IL28B SNP genotyping. The table shows that C allele, which is known as the protective allele, was present in more than 90% of the HCV-infected patients in this Iranian sample. This is in accordance with what has been reported in a previous study among healthy individuals, in which the C allele was reported to be more frequent, especially in East Asia. In that study, conducted by Thomas, *et al.* the authors have suggested that IL28B is under active selection, especially in the abovementioned geographic region (1). Accordingly, our results support this hypothesis in Iran, a country located in southwestern Asia. Moreover, we believe that our findings can explain

previous observations demonstrating the higher rates of SVR among Iranian HCV-infected patients in comparison with similar patients from other countries (9). As previously described, Iranian HCV-infected patients with genotype 1 responded better than expected to anti-HCV therapy, and some hypotheses were made to justify this observation. The high prevalence of C/C and C/T genotypes of rs12979860 compared with the low prevalence of T/T among this sample of HCV-infected patients is one possible reason for the higher SVR rate. To draw a more robust conclusion in this regard, a cohort study involving a large number of HCV-infected patients is under way in our center. The study is investigating the prevalence of both rs12979860 and rs8099917 and their probable influences on the SVR rate among Iranian HCV-infected patients, and the results will be published soon.

References

- Ahlenstiel G, Booth DR, George J. IL28B in hepatitis C virus infection: translating pharmacogenomics into clinical practice. *J Gastroenterol.* 2010;**45**(9):903-10.
- Merat S, Rezvan H, Nouraei M, Jafari E, Abolghasemi H, Radmard AR, et al. Seroprevalence of hepatitis C virus: the first population-based study from Iran. *Int J Infect Dis.* 2010;**14**(Suppl 3):e113-6.
- Ge D, Fellay J, Thompson AJ, Simon JS, Shianna KV, Urban TJ, et al. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature.* 2009;**461**(7262):399-401.
- Suppiah V, Moldovan M, Ahlenstiel G, Berg T, Weltman M, Abate ML, et al. IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet.* 2009;**41**(10):1100-4.
- Afdhal NH, McHutchison JG, Zeuzem S, Mangia A, Pawlotsky JM, Murray JS, et al. Hepatitis C pharmacogenetics: state of the art in 2010. *Hepatology.* 2011;**53**(1):336-45.
- Thomas DL, Thio CL, Martin MP, Qi Y, Ge D, O'Huigin C, et al. Genetic variation in IL28B and spontaneous clearance of hepatitis C virus. *Nature.* 2009;**461**(7265):798-801.
- Balogopal A, Thomas DL, Thio CL. IL28B and the control of hepatitis C virus infection. *Gastroenterology.* 2010;**139**(6):1865-76.
- Li S, Hu P, Zhang Q, Liu YH, Hu HD, Zhang DZ, et al. Single nucleotide polymorphisms of the IL28B and sustained virologic response of patients with chronic hepatitis C to PEG-interferon/ribavirin therapy: A meta-analysis. *Hepat Mon.* 2011;**11**(3):163-72.
- Alavian SM. Are the real HCV infection features in Iranian patients the same as what is expected? *Hepat Mon.* 2005;**5**(1):3-5.