

Errata

Assessment of GFR by four methods in adults in Ashanti, Ghana: the need for an eGFR equation for lean African populations

John B. Eastwood¹, Sally M. Kerry², Jacob Plange-Rhule^{1,3}, Frank B. Micah³, Sampson Antwi⁴, Frances G. Boa⁵, Debasish Banerjee¹, Lynsey Emmett², Michelle A. Miller⁶ and Francesco P. Cappuccio⁶

¹Departments of Renal Medicine and Transplantation, St Georges, University of London, Cranmer Terrace, London SW17 0RE, UK, ²Community Health Sciences, St George's, University of London, Cranmer Terrace, London SW17 0RE, UK, ³Department of Medicine, Komfo Anokye Teaching Hospital, P.O. Box 1934, Kumasi, Ghana, ⁴Department of Paediatrics, Komfo Anokye Teaching Hospital, P.O. Box 1934, Kumasi, Ghana, ⁵Department of Clinical Blood Sciences, St George's Hospital, Blackshaw Road, London SW17 0QT, UK and ⁶Clinical Sciences Research Institute, Warwick Medical School, UHCW Campus, Clifford Bridge Road, 14 Coventry CV2 2DX, UK

Nephrol Dial Transpl 2010; 25: 2178–2187. doi: 10.1093/ndt/gfp765

The following two authors were missing from the original manuscript:

Lynsey Emmett²

Community Health Sciences, St George's, University of London, Cranmer Terrace, London SW17 0RE, UK

Michelle A. Miller⁶

Clinical Sciences Research Institute, Warwick Medical School, UHCW Campus, Clifford Bridge Road, Coventry CV2 2DX, UK

The author apologises for this omission.