

Congenital Cataract Causing Mutants of α A-Crystallin/sHSP Form Aggregates and Aggresomes Degraded through Ubiquitin-Proteasome Pathway

Ilangovan Raju, Edathara C. Abraham*

Department of Biochemistry and Molecular Biology, University of Arkansas for Medical Sciences, Little Rock, Arkansas, United States of America

Abstract

Background: Mutations of human α A-crystallin cause congenital cataract by protein aggregation. How mutations of α A-crystallin cause disease pathogenesis through protein aggregation is not well understood. To better understand the cellular events leading to protein aggregation, we transfected cataract causing mutants, R12C, R21L, R21W, R49C, R54C, R116C and R116H, of human α A-crystallin in HeLa cells and examined the formation of intracellular protein aggregates and aggresomes by confocal microscopy.

Methodology/Principal Findings: YFP-tagged human α A-wild-type (α A-wt) was sub-cloned and the mutants were generated by site-directed mutagenesis. The α A-wt and the mutants were individually transfected or co-transfected with CFP-tagged α A-wt or α B-wild-type (α B-wt) in HeLa cells. Overexpression of these mutants forms multiple small dispersed cytoplasmic aggregates as well as aggresomes. Co-expression of α B-wt with these mutants significantly inhibited protein aggregates where as co-expression with α A-wt enhanced protein aggregates which seems to be due to co-aggregation of the mutants with α A-wt. Aggresomes were validated by double immunofluorescence by co-localization of γ -tubulin, a centrosome marker protein with α A-crystallin. Furthermore, increased ubiquitination was detected in R21W, R116C and R116H as assessed by western blot analyses. Immunostaining with an ubiquitin antibody revealed that ubiquitin inclusions in the perinuclear regions were evident only in R116C transfected cells. Pulse chase assay, after cycloheximide treatment, suggested that R116C degraded faster than the wild-type control.

Conclusions/Significance: Mutants of α A-crystallin form aggregates and aggresomes. Co-expression of α A-wt with the mutants increased aggregates and co-expression of α B-wt with the mutants significantly decreased the aggregates. The mutant, R116C protein degraded faster than wild-type control and increased ubiquitination was evident in R116C expressing cells.

Citation: Raju I, Abraham EC (2011) Congenital Cataract Causing Mutants of α A-Crystallin/sHSP Form Aggregates and Aggresomes Degraded through Ubiquitin-Proteasome Pathway. PLoS ONE 6(11): e28085. doi:10.1371/journal.pone.0028085

Editor: Mark R. Cookson, National Institutes of Health, United States of America

Received: August 19, 2011; **Accepted:** October 31, 2011; **Published:** November 29, 2011

Copyright: © 2011 Raju, Abraham. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by National Institute of Health grant 5R01EY001352-15 from National Eye Institute to ECA. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ecabraham@uams.edu

Introduction

Cataract of the eye lens is the leading cause of blindness worldwide [1]. Pediatric cataract of the congenital type is the most common form of childhood blindness and it is clinically and genetically heterogeneous. About 30-50% of all bilateral pediatric cataracts have a genetic basis [2]. All three forms of Mendelian inheritance have been observed, the most frequently observed type seen in non-consanguineous population being the autosomal dominant transmission. At least 34 loci in the human genome have been reported to be associated with various forms of pediatric cataracts. Autosomal dominant and recessive forms of cataracts have been caused by mutations in 22 different genes [2]. More than half of the mutations occur in crystallins (α -, β -, and γ -crystallins) and the remaining in connexins, intrinsic membrane proteins and intermediate filament proteins. Most interestingly, a total of 12 mutants belong to α -crystallin, 8 for α A-crystallin and 4

for α B-crystallin., it points to a major role α -crystallin mutants play in the development of congenital cataracts.

The α -crystallin gene family consists of two similar genes coding for α A-crystallin, CRYAA located on chromosome 21q22.3, and for α B-crystallin, CRYAB located on chromosome 11q22.1 [3]. The first exon of each gene encodes 60 amino acids consisting of a repeat of 30 amino acid motif and the second and the third exons code for regions homologous for the sHsps [4]. Three α A-crystallin missense mutations have been reported recently which are: base 104 C>T (R12C), base 130 C>T (R21W) and base 230 C>T (R54C) [5]. The affected members of the three families had autosomal dominant bilateral congenital nuclear cataract in association with microcornea, all detected at the time of birth. Affected members of one of this family (R21W) were also diagnosed with microphthalmia. R12C and R21W cases showed zonular opacification with varying involvement of the anterior and posterior pole. It is noteworthy that these mutations occurred

outside the α -crystallin/sHsp core domain. Moreover, the arginines at positions 12, 21, and 54 are highly conserved in α A-crystallin. The other α A mutants reported earlier with autosomal dominant congenital cataracts are: R21L [6], R49C [7], G98R [8], R116C [9] and R116H [10].

A recent report [11] on the biophysical as well as the hydrodynamic properties of the mutants of α A-crystallin have prompted us to further investigate the actual mechanism by which these mutations can lead to early onset of cataract. In all the seven mutants, arginine residues were mutated to mostly cysteine, leucine, tryptophan or histidine. In this study [11], the quaternary structural parameters (hydrodynamic properties) were determined by dynamic light scattering measurements. As compared to α A-wt, average molar mass, polydispersity, and hydrodynamic radius increased several fold in R116C and R116H, moderately increased in R12C, R21W, and R54C, and not increased in R21L and R49C. With regard to secondary and tertiary structural changes, all the mutants showed varying degree of secondary and tertiary structural changes, R21W, R116C, and R116H consistently showing the largest changes. Such changes can lead to protein unfolding/misfolding and subsequently forming protein aggregates.

Since mutants of α A-crystallin contribute to the development of congenital cataract through the formation of aggregated proteins precipitated in the cells of eye lens, we evaluated the expression of mutants of α A-crystallin in mammalian cells (HeLa cells) in terms of identifying the cells having aggregates and aggresomes as the general cellular response to having over expressed mutant proteins [12]. Aggresomes are thought to immobilize protein aggregates and render them susceptible to proteolysis by a component known as proteasomes and/or autophagy [13]–[15]. In view of the propensity of α A-crystallin mutants to aggregate in cells, we also explored whether there is any involvement of ubiquitin-proteasomal pathway (UPP) contributing to the degradation of unfolded proteins of mutants of α A-crystallin. The ubiquitin-proteasome system (UPS) plays an essential role in degrading damaged or unfolded proteins [16]. Unfolded proteins and protein fragments generated by proteolysis are polyubiquitinated by ubiquitin ligases, a process that targets the substrate proteins to the proteasome for degradation [17]. The 26S proteasome which consists of a catalytic 20S core particle and a 19S regulatory particle selectively degrades ubiquitinated proteins [18]; our results suggest that the mutants, R21W, R116C and R116H have elevated polyubiquitinated species.

Materials and Methods

Site directed mutagenesis

To generate mutants, QuickChange site directed mutagenesis kit (Agilent Technologies Inc, CA) was used. Appropriate mutagenic primers of human α A-crystallin for the mutants, R12C, R21L, R21W, R49C, R54C, R116C and R116H were designed and used for PCR. The PCR products were amplified by using YFP-tagged α A-wt as a template DNA with the following PCR conditions, the mix was initially denatured at 95°C for 1 min followed by 95°C for 50 sec, 60°C for 50 sec and 68°C for 5 minutes for 16 cycles and followed by overall extension at 68°C for 7 minutes. The PCR product was digested with *Dpn* I for 1 hour at 37°C and 1 μ l of PCR product was transformed with XL-10 Gold competent cells. The transformants were selected on LB agar medium plates containing 50 μ g/ml Kanamycin. The mutant constructs were sequenced and confirmed by DNA sequence analysis. Untagged pCDNA3.1- α A-wt and the mutant R116C were PCR amplified from appropriate cDNA templates and sub-cloned into *Xho* I and *Eco* RI sites of pCDNA3.1 (-) vector

(Invitrogen, CA). The constructs were validated by restriction digestion and DNA sequence analyses.

Cell culture and transfection

HeLa cells purchased from ATCC, Manassas, VA were grown in 35 mm dishes and 80–90% confluent cells were transfected with Lipofectamine 2000 (Invitrogen, CA) and a total of 2 μ g of plasmid DNA encoded for α A-wt and α B-wt and or mutated constructs fused with either CFP or YFP were used. In one set of experiments, individual constructs for α A-wt and the mutants and for co-expression studies, equal amount of both α A-wt and α B-wt with mutated α A constructs were transfected. Transfected cells showing aggregates were typically counted at x40 magnification. Fields were randomly chosen and about 300 cells were counted per experiment and repeated at least three times and counts were blindly performed.

Laser scanning confocal microscopic studies

An LSM 510 Laser Scanning Microscope (Carl Zeiss Inc., Thornwood, NY) with 63x oil-immersion objective (plan Apochromat, NA 1.4) (University of Arkansas for Medical Sciences core facility) was utilized. To visualize CFP and YFP fluorescence, cells expressing fluorescent proteins were excited at appropriate laser beam and filtered with both dichromatic band pass filters, captured at 12 bit 512 x 512, multitrack channel images with CCD cameras with the following configurations: for CFP channel, the cells were excited with 458 nm filter by argon-ion laser and the emission intensity was collected using band pass (BP) 475–525 nm filters and for YFP channel, the cells were excited with 514 nm filter by argon-ion laser and the emission intensity was collected using BP 530–600 nm filters. Both the CFP and YFP was excited using argon-ion laser at 25 mW, 2.0 and 0.5% exposure respectively.

Cycloheximide chase assay

Degradation of α A wild-type and the mutant α A-R116C proteins was assessed using cycloheximide-chase assay. HeLa cells were grown in 35 mm dishes and transfected with untagged α A-wt and the mutant R116C. After 24 h transfection, cells were treated with 20 μ g/mL of cycloheximide (Sigma) for the indicated time period and lysed with lysis buffer containing 50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 0.02% sodium azide, 0.1% SDS, 1% NP-40, 0.5% sodium deoxycholate and 0.1 mM EDTA supplemented with cock-tail protease inhibitors (Roche Diagnostics) and 3M Urea. For immunoblot analysis, 5 μ g of total protein was loaded into 12% SDS-PAGE and the western blot was probed with a rabbit polyclonal anti- α A-crystallin antibody (Enzo Life Sciences Inc, SPA-221) at a dilution of 1 in 6000.

SDS-PAGE and Western blot analysis

After 48 hours transfection, cells were lysed with lysis buffer containing 50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 0.02% sodium azide, 0.1% SDS, 1% NP-40, 0.5% sodium deoxycholate and 0.1 mM EDTA supplemented with cock-tail protease inhibitors and 3M Urea. Further, the cell lysate was sonicated and the protein concentration was measured by BCA assay method. For each sample, 5 μ g of total protein was loaded into 12% SDS-PAGE and electroblotted to PVDF membrane. The blots were blocked with 5% non-fat dry milk prepared in TBST (Tris-buffered saline supplemented with 0.1% Tween 20) and subsequently incubated with primary antibody for α A-crystallin (monoclonal, Abcam, ab78439, 1:2000), α B-crystallin (rabbit polyclonal, Abcam, ab13497, 1:2000) for one hour at room temperature. Blots were washed with TBST for three times and incubated with appropriate HRP-conjugated secondary antibodies

(1 in 10000, Santa Cruz Biotechnology Inc, CA) for one hour at room temperature. Enhanced Chemiluminescence substrate was used and the signal was detected by exposing the blots on films. For loading control, blots were stripped with Restore Western Blot stripping buffer (Thermo Scientific Inc, IL) and re-probed with a rabbit polyclonal antibody against β -actin (Abcam, ab8227, 1:10000) for 1 hour at room temperature.

Immunofluorescence microscopy for aggresome detection

Cells were grown on 35-mm cover glass bottom dishes. After 48 hours transfection, cells were washed with PBS, fixed with 4% paraformaldehyde for 20 minutes at room temperature (RT) and permeabilized with 0.5% Triton X-100 for 10 minutes at RT. Cells were blocked with 5% normal goat serum (NGS) for one hour at

RT. The cells were simultaneously incubated with α A-crystallin mouse monoclonal antibody (1 in 200 in 5% NGS, Abcam, ab78439) and a rabbit polyclonal antibody for γ -tubulin (Abcam, ab16504; 1 in 200 dilution in 5% NGS) for overnight at 4°C. The cells were stained with Alexa Fluor 594 (mouse) and Alexa Fluor 488 (rabbit) (Molecular Probes) for one hour at room temperature. Nuclei were counter stained with Hoechst 33342. The images were acquired with an LSM 510 Meta Carl Zeiss Confocal microscope at x63 objective and analyzed using AIM Imaging Software.

Data Analysis and Statistics

In all the experiments, values were expressed as mean \pm SD. Two-tailed Student's t-test was used for statistical analysis. The p value < 0.05 was considered as significant.




Figure 1. YFP- α A-wt and the mutant constructs, R12C, R21L, R21W, R49C, R54C, R116C and R116H were individually expressed in HeLa cells. A: LSM Images were captured after 48 hours transfection. HeLa cells were individually transfected with 2 μ g of YFP-tagged α A-wt and mutants of α A-crystallin. A homogenous expression of α A-crystallin was evident in α A-wt transfected cells. Cytoplasmic aggregates were evident in α A-crystallin mutants transfected cells. The YFP signal was excited at 514 nm and the images were collected by BP 530-600 nm filter. The images represent one of the four similar images obtained in three independent experiments. B: Graph represents per cent of cells with aggregates. The results obtained after 48 hours transfection for the individually expressed α A-wt or its mutants in HeLa cells. Cells containing aggregates were counted in 10 random fields each field with 30 cells. The mutant, R116C showed a high per cent (~47) of cells having aggregates and the mutant R21L showed least per cent (~10) of cells containing aggregates. The results were presented as means \pm SD obtained in three independent experiments. All the mutants were statistically significant, p < 0.01. doi:10.1371/journal.pone.0028085.g001

Results

Individual expression of YFP-tagged α A-crystallin wild-type and mutants in HeLa cells

To investigate whether the mutants of α A-crystallin forms aggregates in cells, YFP-tagged wild-type and the mutants of α A-crystallins were transfected individually in HeLa cells. Cells transfected with CFP or YFP alone showed a homogenous expression of the fluorescent protein in both nucleus and cytoplasm (data not shown). Cells transfected with α A-wt showed a homogenous distribution of its expression in the cytoplasm alone and there was a little or no aggregation was observed in these cells (Fig. 1A). Cells transfected with the mutants, R12C, R21L, R21W, R49C, R54C, R116C, R116H showed significant number of cells having protein aggregates, 37.2 ± 3.8 , 10.5 ± 2.6 , 30.2 ± 4.7 ,

24.5 ± 1.1 , 18.5 ± 3.1 , 47 ± 5.8 , 28.1 ± 3.4 (Fig. 1B) respectively. Here, the intracellular aggregation is referred to as the clumped particles predominantly localized in the cytoplasm. Cells containing more than three such particles were considered as positive for cells having aggregates and scored in this assay. Moreover, the morphology of cells was altered in cells expressing R21W and R116C mutants (Fig. 1A).

Co-expression of CFP- α A-wt and YFP-tagged mutants of α A-crystallins

To investigate whether co-expression of α A-wt can inhibit aggregates caused by mutants of α A-crystallin in cells, we transfected YFP-tagged α A-mutants with CFP-tagged α A-wt. Co-expression of CFP- α A-wt with YFP- α A-wt constructs in HeLa cells did not show any aggregates and a homogenous expression of




Figure 2. YFP- α A-wt and its mutants were co-expressed with CFP-tagged α A-wt. *A:* Laser scanning confocal microscope images. HeLa cells were transfected with 1 μ g each of CFP-tagged α A-wt and YFP-tagged mutant constructs. After 48 h transfection, cells were analyzed with confocal microscope. Cells showed more aggregates when co-expressed with α A-wt. The CFP signal was excited at 458 nm and the images were collected by BP 475–525 nm filter, YFP was excited at 514 nm and the images were collected by BP 530–600 nm filter. The images represent one of the four similar images obtained in three independent experiments. *B&C:* Graph represents per cent of cells with aggregates. Cells were transfected with YFP-tagged α A-wt or mutants and CFP-tagged α A-wt. Cells having aggregated were counted in 10 random fields each field with 30 cells after 48 h transfection. The results were presented as means \pm SD obtained in three independent experiments. For α A-wt + α A-R12C: $p < 0.03$; for α A-wt + α A-R21L: < 0.05 ; for α A-wt + α A-R21W: < 0.04 , for α A-wt + α A-R49C, is not significant; for α A-wt + α A-R54C: < 0.0001 ; for α A-wt + α A-R116C: < 0.01 and for α A-wt + α A-R116H, is not significant. doi:10.1371/journal.pone.0028085.g002

protein was evident in the cytoplasm (Fig. 2A). However, in α A-wt co-expression with the mutants, R12C and R54C showed a few nuclear foci. Cells co-expressed with all the other mutants except R21L and R116H significantly increased the number of cells having aggregates. The proportion of the cells with aggregates were 53, 15, 51, 35, 40, 68 and 30% in mutants, R12C, R21L, R21W, R49C, R54C, R116C and R116H respectively (Fig. 2B&C). Cells co-expressed with R12C, R49C and R116C distorted the cellular morphology (Fig. 2A). It is possible that the increase in the percent of cells having aggregates in the mutants expressing cells was due to co-aggregation with native α A-wt.

Co-expression of CFP- tagged α B-wt and YFP-tagged mutants of α A-crystallin

To investigate, whether α B-wt co-expression can inhibit protein aggregates caused by mutants of α A-crystallin, cells were transfected with CFP- α B-wt and YFP-tagged α A-crystallin mutants. The results indicate that a significant decrease (52–72%, as compared to the data in Fig. 2) of the number of cells having aggregates was observed in cells transfected with all the mutants but it is not statistically significant with the mutants, R21L and R54C. Specifically, the mutants, R12C, R21W, R116C and R116H showed the largest effect (61–72%) (Fig. 3A, B, C). Thus,




Figure 3. YFP-tagged α A-wt and each of the mutated α A-crystallin co-expressed with CFP- α B-wt. A: LSM images of HeLa cells after 48 h transfection. HeLa cells were transfected with 1 μ g each of CFP-tagged α B-wt and YFP-tagged α A-wt and the mutants. After 48 h transfection, images were captured with an LSM confocal microscope. The CFP signal was excited at 458 nm and the images were collected by BP 475–525 nm filter, YFP was excited at 514 nm and the images were collected by BP 530–600 nm filter. Co-expression of α B-wt significantly inhibited aggregates in cells transfected with the α A- mutants. Images represent one of the four similar images obtained in three independent experiments. B&C: Graph represents per cent of cells containing aggregates. The results obtained after 48 h transfection. Cells containing aggregates were counted in 10 random fields each field with 30 cells. In all mutants, co-expression of α B-wt except with R21L and R54C significantly decreased the number of cells having aggregates. The results were presented as means \pm SD obtained in three independent experiments. The p value for α B-wt + α A-R12C is < 0.02 ; for α B-wt + α A-R21L is not significant; for α B-wt + α A-R21W is < 0.05 ; for α B-wt + α A-R49C is < 0.02 ; for α B-wt + α A-R54C is not significant, for α B-wt + α A-R116C is < 0.01 and for α B-wt + α A-R116H is < 0.001 . doi:10.1371/journal.pone.0028085.g003

co-expression of α B-wt with the mutants decreases protein aggregates. These results strongly suggest that α B-wt is a potential chaperone to protect the cells from aggregates caused by the α A-mutants.

Validation of the expression of CFP or YFP-tagged α A-wt and/or α B-wt with mutants of α A-crystallin in HeLa cells

To validate that there was no discrepancy in the transfection efficiency of the either individually transfected or co-transfected with α A-wt and or α B-wt, total cell lysate were subjected to immunoblot and probed with α A and or α B-crystallin antibodies. A similar level of expression was detected in individually expressed α A-wt or mutant constructs (Fig. 4A). The level of expression was nearly equal in mutants co-expressed with α A-wt (Fig. 4B) and α B-wt constructs (Fig. 4C).

Mutant α A-crystallins in HeLa cells form aggresomes

To demonstrate whether the large perinuclear structures are aggresomes as seen in the cells transfected with the mutants, R21W, R116C and R116H (three major aggregate forming mutants) and to validate these inclusions as aggresomes, transfected cells were subjected to double immunostaining with a mouse monoclonal α A-crystallin (red) and rabbit polyclonal γ -tubulin, (a centrosome marker protein) (green) antibodies. For this

study, untagged pCDNA3.1 constructs of both wild-type and the mutants R116C (severely affected cells) and R21L (mildly affected cells) were used in order to eliminate the false positive signal by overlapping of YFP signal with the Alexa Fluor 488, both of these signals being acquired at argon-ion laser line in confocal microscopy. The results showed that only in R116C, the colocalization of α A-crystallin with γ -tubulin occurred as yellow punctate signals in the perinuclear region validated these structures are aggresomes (Fig. 5) whereas, there were no aggresomes in R21L expressing cells (Fig. 5).

Degradation of aggregate-prone α A-crystallin mutants

Since our finding that mutants of α A-crystallin form intracellular aggregates in cells and this aggregate formation may affect protein turn over which may contribute to the pathogenesis of the cataract, we next asked whether mutation in α A-crystallin may affect protein's turn over by faster degradation, cells were treated with cycloheximide (for inhibition of protein synthesis) and at different time points. After 24 h transfection, the cells lysed and a total protein of 5 μ g from each of the sample was subjected to immunoblot probed with α A-crystallin antibody (rabbit polyclonal, Enzo Life Sci, SPA-221). For this study, we used untagged constructs of α A, i.e. pCDNA3.1/ α A-wt and pCDNA3.1/ α A-R116C. As shown in Fig. 6A and Fig. 6B, pulse chase experiments




Figure 4. Western blot analysis of α A-crystallin-wt and its mutants, R12C, R21L, R21W, R49C, R54C, R116C and R116H expressed in HeLa cells. A. Western blot analysis of HeLa cells individually expressed with α A-Crystallin: Cells were individually transfected with YFP-tagged α A-wt and mutants. After 48 h transfection, cells were lysed and 5 μ g of total protein was subjected to western blot. The blot was probed with human anti- α A antibody. The same blot was stripped and re-probed for β -actin to serve as a loading control. Nearly a similar level of expression of α A was evident in each of the transfected cells. The * indicates the non-specific band. B: Western blot analysis of HeLa cells co-expressed with CFP-tagged α A-wt and YFP-tagged α A-wt or its mutants. Cells were co-transfected with YFP-tagged α A-wt and mutants with CFP-tagged α A-wt. After 48 h transfection, cells were lysed and 5 μ g of total protein was subjected to western blot. The blot was probed with an antibody against human α A and the same blot was stripped and re-probed for actin. A similar level of expression of α A was detected in each group. C: Western blot analysis of HeLa cells co-expressed with CFP-tagged α B-wt and YFP-tagged α A-wt or its mutants. After 48 h transfection, 5 μ g of total protein was subjected to western blot. The blot was probed with an antibody against human α A and the same blot was stripped and re-probed for anti- α B. The same blot was again stripped with anti- β -actin for loading control. The level of both α A and α B were nearly equal in each of the transfected cells. doi:10.1371/journal.pone.0028085.g004




Figure 5. Detection of Aggresomes: Untagged pCDNA3.1 constructs of α A-wt and the mutants, R21L and R116C were used in this study and compared. After 48 h transfection, cells were fixed, permeabilized with 0.5% Triton X-100 and double immunostained with a mouse monoclonal α A-crystallin (red) antibody and a rabbit polyclonal antibody for γ -tubulin (green). A strong degree of overlapping signal (yellow) was evident only in R116C transfected cells (arrow). Co-localization of γ -tubulin, a centrosome maker protein with α A-crystallin in the perinuclear region validated these inclusions were aggresomes. There was no co-localization in cells transfected with either α A-wt or the mutant, R21L. Goat anti-mouse Alexa Fluor 594 (red) antibody was used to stain and visualize the localization of α A-wt and the mutants, R21L and R116C. A rabbit secondary antibody, Alexa Fluor 488 (green) was used to stain and visualize the γ -tubulin. The nuclear stain Hoechst was used to counter stain the nuclei. The images were representative of one of four such images obtained in three independent experiments.
doi:10.1371/journal.pone.0028085.g005

followed by cycloheximide treatment showed that wild-type protein is stable at least for 24 h. But the level of the mutant protein R116C decreased to $75.6 \pm 3.2\%$, $53 \pm 5.5\%$, and $51.3 \pm 2.5\%$ compared to control after 6, 12 and 24 h of treatment respectively (Fig. 6C and 6D) As expected the mutant protein has lower half-life and degraded faster than the wild-type protein.

Accumulation of polyubiquitinated conjugates in α A-crystallin mutants

A number of reports implicated UPS dysfunction in a range of aggregation prone mutant proteins in neurodegenerative diseases.

To explore any effects of aggregation-prone α A-crystallin mutants might have on UPS proteolytic function, we measured the level of polyubiquitinated proteins in whole cell lysate subjected to western blot probed with anti-ubiquitin antibody (FK2, Mouse Monoclonal, Enzo Life Sciences, PW8810). Distinct polyubiquitin conjugated proteins accumulated in cells transfected with mutants, R21L, R21W, R116C and R116H (Fig. 7, lanes 3, 4, 7 and 8). In a separate experiment, to further show ubiquitin inclusions in transfected cells, cells were subjected to immunostaining with a mouse monoclonal ubiquitin antibody (FK2; Enzo Life Sci.Inc., PW 8810). The ubiquitin inclusions in the perinuclear regions




Figure 6. Degradation of aggregate-prone α A-crystallin mutant, R116C. *A and C:* Western blot analysis of cycloheximide treated cells: HeLa cells were transfected with 2 μ g of untagged pCDNA3.1 constructs of α A-wt and the mutant, R116C. After 24 h transfection, cells were treated with 20 μ g/ml of cycloheximide and lysed with lysis buffer at indicated time points. For each of the sample, 5 μ g of total protein was loaded and western blot probed with an anti- α A-crystallin (rabbit polyclonal, Enzo Life Sciences Inc., Catalog # SPA-221). The mutant R116C protein level has decreased after 6 hours treatment with cycloheximide compared to wild-type protein which demonstrated R116C protein instability. The β -Actin blot serves as a loading control. *B and D:* Quantification of wild-type and the mutant, R116C protein at different time points. The density of the band was quantified using NIH Image J software and plotted. Values represent as means \pm SD as obtained in three independent experiments. The mutant, R116C Vs wild-type control is significant at $p < 0.001$. doi:10.1371/journal.pone.0028085.g006

were evident only in the mutant R116C transfected cells (Fig. 8). Altogether, these results suggest that mutants of α A-crystallin proteins were conjugated with ubiquitin for degradation.

Discussion

In the present study, we have demonstrated that overexpression of the cataract causing mutants of α A-crystallin in HeLa cells led to the formation of multiple intracellular protein aggregates. There was no evidence for the endogenous expression of both α A- and α B-crystallins in these cells as shown in our earlier study [19] and thus serves as a perfect model to study the role of cataract causing mutants of α A-crystallin in mammalian cells including the eye lens cells. Compared to α A-wt, expression of all the mutants in these cells showed significant increase in protein aggregation after 48 hours of transfection, R21L showing the least increase although higher than the control α A-wt. It is likely that the protein aggregation in the cytoplasm was due to protein conformational changes the mutants undergo [11]. When these mutants were expressed individually, they formed aggregates probably due to stress in the absence of any protective mechanism such as the presence of α B-crystallin, a powerful sHsp. sHsps, in general, can inhibit protein aggregation and can reverse or refold aggregated proteins in conjunction with Hsp 70, a major molecular

chaperone. As shown in an earlier study of the C-terminal truncated α A-crystallins [19], association with native α A-crystallin significantly increased the number of cells containing aggregates in all mutants. These results suggest that α A-crystallin is not a potential chaperone to protect the cells from protein aggregation. It is interesting to note that most of the mutants of α A-crystallin involve arginine residues and lead in a dominant fashion. Co-expression with α B-crystallin, on the other hand, significantly diminished the aggregation, R12C, R21L, R116C and R116H showing the most effect and R49C and R54C showing the least effect. α B-crystallin is known to be a better molecular chaperone than α A-crystallin and as it has been shown earlier with C-terminal truncated α A-crystallins [20], it readily recognizes partially unfolded structures and prevent them from aggregation.

Also, this study provides the first evidence for cataract causing mutants of α A-crystallin forming aggresomes in cells. Accumulation of misfolded proteins results from saturation of protein degradation system observed in conformational diseases like Huntington disease [21] and cystic fibrosis [22] leading to the formation of inclusion bodies also known as aggresomes. The inclusion bodies concentrated in the perinuclear region of α A-crystallin mutants expressed in HeLa cells suggests that they have the characteristic features of aggresomes. They are cytoplasmic globular structures formed due to protein misfolding in the cytosol




Figure 7. Overexpression of α A-Crystallin mutants showed increased ubiquitination. *A: Western blot analysis:* After 48 h transfection, cells were lysed and 5 μ g of total protein was subjected to immunoblot probed with an anti-ubiquitin (FK2) monoclonal antibody. The pattern of polyubiquitinated species were dramatically increased in the mutants, R21W, R116C and R116H compared to other mutants and wild-type. * indicates a non-specific band. The same blot was stripped and re-probed for α A-crystallin and β -actin for loading controls. The blot shown here is a representative blot of three independent experiments. *B: Quantitative data for the western blot:* Using NIH Image J software, densitometric measurements were normalized against β -actin. The mutants, R21L, R21W, R116C and R116H showed increased ubiquitination as obtained in three independent experiments and plotted. The results were expressed as mean \pm SD. The p value for α A-wt Vs α A-R12C is not significant, for α A-wt Vs α A-R21L is < 0.05 ; for α A-wt Vs α A-R21W is < 0.01 ; for α A-wt Vs α A-R49C is not significant; for α A-wt Vs α A-R54C is not significant; for α A-wt Vs α A-R116C is < 0.003 and for α A-wt Vs α A-R116H is < 0.001 . doi:10.1371/journal.pone.0028085.g007

and these structures are delivered to the microtubule organizing center (MTOC) by retrograde transport along microtubules [12]. These aggresomes are not merely a storage site for misfolded proteins; they can facilitate the degradation of protein aggregates [12] and are the pathological hallmark of conformational diseases that results from protein misfolding. In the present study, double immunofluorescence results (Fig. 5) validate the co-localization of α A-crystallin with a centrosome marker protein, γ -tubulin, in aggresomes. The γ -tubulin has been previously shown to co-localize with aggresomes in MTOC [23–25]. They are normally formed when ubiquitin-proteasome degradation pathway is impaired and disease-associated proteins inefficiently fold [26], [27]. Our findings on aggresomes detection in cells transfected with α A-crystallin mutants are very similar to the previous studies on a myopathy causing mutant of human α B-crystallin, R120G, which forms aggresomes in CCL39 cells [28]. Aggresomes are special protective structures that fundamentally differ from other multiple aggregates, that some of which cause cellular toxicity. They are formed around MTOC, a sub-cellular region which is robustly enriched with chaperones and components of UPS [12]. It has been reported that

multiple aggregates or pre-aggresome particles may be an intermediate step in aggresome formation which can proceed further upon proteasome inhibition [15]. The present study documented both multiple aggregates and typical perinuclear localized aggresomes. Typical aggresomes were detected only in R116C mutant but not in R21L, which suggest that the multiple aggregates in cells expressing this mutant did not develop as aggresomes.

It has been suggested that unfolded proteins develop into insoluble form that cannot be degraded, their sequestration in one large mass may facilitate their removal by autophagy [23]. Degradation of C-terminal truncated α A-162 through ubiquitin-proteasome pathway has been reported previously using recombinant proteins [29]. Proteins with short half-lives are mostly degraded by the proteasome [30]. Our results on the mutant-R116C protein degradation upon cycloheximide treatment are consistent with a previous report [31] on a truncated α B-crystallin, 450delA protein which has lower half-life [31]. The Ubiquitin-Proteasome System (UPS) degrades short-lived nuclear, misfolded cytosolic proteins extruded from the endoplasmic reticulum [16]. Substrates of UPS need to unfold to pass through the narrow barrel-




Figure 8. Overexpression of α A-crystallin mutants shows accumulation of ubiquitin inclusions in HeLa cells. YFP-tagged α A-wt and mutant, R116C were transfected in HeLa cells. After 48 h transfection, cells were fixed and immunostained with a mouse monoclonal ubiquitin (FK2) antibody and further stained with a fluorescent conjugated secondary antibody, Alexa Fluor 594 (red). The arrows indicate ubiquitinated cytoplasmic inclusions only in the mutant, R116C expressing cells. Hoechst 33342 was used to counter stain the nuclei. The images were representative of three similar images obtained in three independent experiments. doi:10.1371/journal.pone.0028085.g008

References

- Reddy MA, Francis JP, Bery V, Bhattacharya S, Moore AT (2004) Molecular genetic basis of inherited cataract and associated phenotypes. *Surv Ophthalmol* 49: 300–315.
- De Jong WW, Terwindt EC, Bloemendal H (1975) The amino acid sequence of the A chain of human alpha-crystallin. *FEBS Lett* 58: 310–313.
- Horwitz J (1992) Alpha-crystallin can function as a molecular chaperone. *Proc Natl Acad Sci USA* 89: 10449–10453.
- Hansen L, Yao W, Eiberg H, Kjaer KW, Baggesen K, et al. (2007) Genetic heterogeneity in microcornea-cataract: five novel mutations in CRYAA, CRYGD and GJA8. *Invest Ophthalmol Vis Sci* 48: 3937–3944.
- Xia C, Liu H, Chang B, Cheung D, Wang M, et al. (2006) Arginine 54 and Tyrosine 118 Residues of α A-Crystallin are Crucial for Lens Formation and Transparency. *Invest Ophthalmol Vis Sci* 47: 3004–3010.
- Graw J, Klopp N, Illig T, Preising MN, Lorenz B (2006) Congenital cataract and macular hypoplasia in humans associated with a de novo mutation in CRYAA and compound heterozygous mutations in P Graefes Arch Clin Exp Ophthalmol 244: 912–919.
- Mackay DS, Andley UP, Shiels A (2003) Cell death triggered by a novel mutation in the alphaA-crystallin gene underlies autosomal dominant cataract linked to chromosome 21q. *Eur J Hum Genet* 11: 784–793.
- Santhiya ST, Soker T, Klopp N, Illig T, Prakash MV, et al. (2006) Identification of a novel, putative cataract-causing allele in CRYAA (G98R) in an Indian family. *Mol Vis* 12: 768–773.
- Litt M, Kramer P, LaMorticella DM, Murphey W, Lovrien EW, et al. (1998) Autosomal dominant congenital cataract associated with a missense mutation in the human alpha crystallin gene CRYAA. *Hum Mol Genet* 7: 471–474.
- Vanita V, Singh JR, Hejtmancik JF, Nuernberg P, Hennies HC, et al. (2006) A novel fan-shaped cataract-microcornea syndrome caused by a mutation of CRYAA in an Indian family. *Mol Vis* 12: 518–522.
- Kore RA, Santhoshkumar P, Sharma KK, Abraham EC (2010) Structural, Functional And Hydrodynamic Properties of alphaA-crystallin Mutants Causing Congenital Cataracts. *ARVO Abstract* 4617/ D687.
- Kopito RR (2000) Aggresomes, inclusion bodies and protein aggregation. *Trends Cell Biol* 10: 524–530.
- Taylor JP, Tanaka F, Robitschek J, Sandoval JM, Taye A, et al. (2003) Aggresomes protects cells by enhancing the degradation of toxic-polyglutamine containing protein. *Hum Mol Genet* 12: 749–757.
- Kawaguchi Y, Kovacs JJ, McLaurin A, Vance JM, Ito A, et al. (2003) The deacetylase HDAC6 regulates aggresome formation and cell viability in response to misfolded protein stress. *Cell* 115: 727–738.
- Junn E, Lee SS, Suhr UT, Mouradian MM (2002) Parkin Accumulation in Aggresomes Due to Proteasome Impairment. *J Biol Chem* 277: 47870–47877.
- Hershko A, Ciechanover A (1998) The ubiquitin system. *Annu Rev Biochem* 67: 425–479.
- Pickart CM, Cohen RE (2004) Proteasomes and their kin: proteases in the machine age. *Nat Rev Mol Cell Biol* 5: 177–187.
- DeMartino GN, Gillette TG (2007) Proteasomes: machine for all reasons. *Cell* 129: 659–662.
- Raju I, Kumarasamy A, Abraham EC (2011) Multiple Aggregates and Aggresomes of C-terminal truncated human α A-Crystallins in mammalian cells and Protection by α B-Crystallin. *PLoS ONE* 6: e19876.
- Kumarasamy A, Abraham EC (2008) Interaction of C-terminal truncated human alphaA-crystallins with target proteins. *PLoS ONE* 3: e3175.
- Ren PH, Laueckner JE, Kachirskaja I, Heuser JE, Melki R, et al. (2009) Cytoplasmic penetration and persistent infection of mammalian cells by polyglutamine aggregates. *Nat Cell Biol* 11: 219–225.
- Luciani A, Vilella VR, Esposito S, Brunetti-Pierri N, Medina D, et al. (2010) Defective CFTR induces aggresome formation and lung inflammation in cystic fibrosis through ROS-mediated autophagy inhibition. *Nat Cell Biol* 12: 863–875.
- Iwata A, Riley BE, Johnston JA, Kopito RR (2005) HDAC6 and microtubules are required for autophagic degradation of aggregated huntingtin. *J Biol Chem* 280: 40282–40292.
- Wileman T (2007) Aggresomes and pericentriolar sites of virus assembly: cellular defense or viral design? *Ann Rev Microbiol* 61: 149–167.
- Tanaka M, Kim YM, Lee G, Junn E, Iwatsubo T, et al. (2004) Aggresomes formed by alpha-synuclein and synphilin-1 are cytoprotective. *J Biol Chem* 279: 4625–4631.
- Zaarur N, Merrin AB, Gabai VL, Sherman MY (2008) Triggering aggresome formation. Dissecting aggresome-targeting and aggregation signals in synphilin-1. *J Biol Chem* 283: 25575–25584.
- Arrasate M, Mitra S, Schweitzer ES, Segal MR, Finkbeiner S (2004) Inclusion body formation reduces levels of mutant huntingtin and the risk of neuronal death. *Nature* 431: 805–810.
- Chavez Zobel AT, Loranger A, Marceau N, Theriault JR, Lambert H, et al. (2003) Distinct chaperone mechanisms can delay formation of aggresomes by the myopathy-causing R120G alphaB-crystallin mutant. *Hum Mol Genet* 12: 1609–1620.
- Zhang X, Dudek EJ, Liu B, Ding L, Fernandes AF, et al. (2007) Degradation of C-terminal truncated alpha A-Crystallins by the ubiquitin-proteasome pathway. *Invest Ophthalmol Vis Sci* 48: 4200–4208.
- Rechsteiner M, Hoffman L, Dubiel W (1993) The multicatalytic and 26S proteases. *J Biol Chem* 268: 6065–6068.
- Zhang H, Rajasekaran NS, Orosz A, Xiao X, Rechsteiner M, et al. (2010) Selective degradation of aggregate-prone CryAB mutants by HSPB1 is mediated by ubiquitin-proteasome pathways. *J Mol Cell Cardio* 49: 918–930.
- Glickman MH, Ciechanover A (2002) The ubiquitin-proteasome proteolytic pathway: destruction for the sake of construction. *Physiol Rev* 82: 373–428.
- Rubinsztein DC (2006) The roles of intracellular protein-degradation pathways in neurodegeneration. *Nature* 443: 780–786.