



## Liver Injury Induced by High-Dose Methylprednisolone Therapy: A Case Report and Brief Review of the Literature

Krzysztof Gutkowski<sup>1\*</sup>, Alina Chwist<sup>1</sup>, Marek Hartleb<sup>1</sup>

<sup>1</sup>Department of Gastroenterology and Hepatology, Medical University of Silesia, Katowice, Poland

### ARTICLE INFO

*Article Type:*  
Case Report

*Article history:*  
Received: 30 May 2011  
Revised: 13 Jun 2011  
Accepted: 20 Jun 2011

*Keywords:*  
Methylprednisolone  
Hepatotoxicity  
Adverse drug reactions  
Multiple sclerosis

### ABSTRACT

Corticosteroids are used widely to treat many types of disease. In general, these drugs are considered safe for the liver; however, recent reports have demonstrated that high-dose methylprednisolone (MT) may cause severe liver injury. Here, we report a case of a 24-year-old female who was given pulsed MT therapy for multiple sclerosis. MT induced icteric hepatitis and impaired liver synthetic function. Hepatotoxicity developed several weeks after drug exposure, and the causal association with MT was confirmed by unintentional rechallenge test. A brief review of the literature on corticosteroid-induced hepatotoxicity is presented.

© 2011 Kowsar M.P.Co. All rights reserved.

#### ► *Implication for health policy/practice/research/medical education:*

Corticosteroid-induced liver injury may develop several weeks after short term drug exposure, therefore patients who receive corticosteroid pulses should be screened for potential liver injury.

#### ► *Please cite this paper as:*

Gutkowski K, Chwist A, Hartleb M. Liver Injury Induced by High-dose Methylprednisolone Therapy: A Case Report and Brief Review of the Literature. *Hepat Mon.* 2011;11(8):656-661. [DOI:10.5812/kowsar.1735143X.713]

## 1. Introduction

Adverse drug reactions are frequent and remain underestimated causes of acute liver injury that sometimes lead to liver failure, requiring liver transplantation.

High-dose intravenous glucocorticosteroid treatment is one of the most efficient therapeutic options for severe exacerbations of many autoimmune diseases. A review of the literature shows that corticosteroids are not entirely safe for the liver and have been occasionally linked to severe hepatotoxicity (1-3). We present a case

of a 24-year-old woman who was treated with pulsed methylprednisolone (MT) for multiple sclerosis. MT induced serious liver injury, developing as icteric acute hepatitis with impaired prothrombin synthesis. Hepatotoxicity with prolonged latency appeared after the second MT pulse, and the causal relationship between MT and liver injury was confirmed by unintentional rechallenge test.

## 2. Case Report

A 24-year-old woman with a 3-year history of multiple sclerosis and a primary diagnosis of retrobulbar optic neuritis was referred to our department due to severe acute hepatitis. Three months earlier, she was admitted to the neurology department due to exacerbation of multiple sclerosis. Neurological treatment began with

\* Corresponding author at: Krzysztof Gutkowski, Department of Gastroenterology and Hepatology, Medical University of Silesia, ul. Medykow 14, 40-752, Katowice, Poland. Tel: +48-327894401, Fax: +48-327894402.

E-mail: kgutski@intertele.pl

DOI: 10.5812/kowsar.1735143X.713

Copyright © 2011, BRCCGL, Published by Kowsar M.P.Co. All rights reserved.

a 6-day course (0.5 g/d) of high-dose intravenous MT, resulting in full recovery of her left arm function. Routine laboratory examinations showed normal liver tests and no autoantibodies in the peripheral blood.

The patient was subsequently switched to beta-1b interferon (Betaferon) treatment and received 2 doses of this drug without any immediate or delayed adverse reactions. After 6 weeks, she developed a subsequent flare of MS, presenting as left-sided limb paralysis. A high-dose intravenous MT pulse (total dose 3.0 g) and 2 injections of Betaferon were given, effecting nearly a full recovery. Four weeks after the MT pulse and 1 week after her last Betaferon dose, the patient developed jaundice with elevated serum levels of aspartate aminotransferase (AST) 900 IU/L (n: 10-31 IU/L), serum alanine aminotransferase (ALT) 1740 IU/L (n: 9-34 IU/L), serum alkaline phosphatase 186 IU/L (n: 38-126 IU/L), and serum GGTP 50 IU/L (n: <38 IU/L). The serum bilirubin level was 17.9 mg/dL (n: 0.3-1.2 mg/dL), the direct bilirubin was 16.1 mg/dL, and the prothrombin index was 35.1% (INR 2.52). The serum AFP level was 17.18 ng/mL (n: <5 ng/mL). The patient had no history of hepatic disease and denied any use of alcohol. In the previous 6 months, the patient took acetaminophen (used only in a single dose before each Betaferon injection) and oral contraception, which was stopped immediately before the second MT pulse. Serological tests for hepatitis B (including antibodies to HBc), hepatitis C, hepatitis A, and infection with cytomegalovirus (CMV) were negative. Smooth muscle antibodies (SMA) were found in high titers (>1:320), and autoantibodies against mitochondria and nuclei were undetected by direct immunofluorescence tests. Her copper urinary excretion was not elevated. An ultrasound abdominal examination showed an intact liver and a normal-sized spleen. No liver biopsy was done due to a low prothrombin level. With a MELD (model for end-stage liver disease) score of 26, the patient was a potential liver transplantation candidate. Fortunately, her liver function improved spontaneously, and the patient was discharged after 3 weeks with normal aminotransferase levels. Considering the potential hepatotoxicity of interferons, we cautiously suspected the Betaferon of inducing liver injury. Three months later, the patient developed a consecutive flare of multiple sclerosis, for which she received a 6-day intravenous MT pulse (total dose 3.0 g) without Betaferon. Her neurological symptoms resolved, but 4 weeks later, she was readmitted to our department with symptoms of acute hepatitis. The serum level of AST was 1488 IU/L, ALT was 1129 IU/L, alkaline phosphatase was 164 IU/L, GGTP was 168 IU/L, prothrombin index was 47% (INR 1.71), and serum bilirubin was 7.3 mg/dL. No drugs, as before, were used to treat the liver disease, and after several weeks, aminotransferases and bilirubin reached reference levels.

### 3. Discussion

We report two serious liver injuries that were related

to pulsed MT treatment. The diagnosis was confirmed by the dechallenge-rechallenge relationship between drug administration and hepatitis. Based on a literature review, we identified 12 case reports of corticosteroid-induced hepatotoxicity (Table). The clinical course of liver injury varied from asymptomatic hypertransaminasemia to fulminant hepatic failure (3 deaths). In vivo liver histopathology was performed in only 8 cases, showing a large spectrum of lesions. Fernandez *et al.* reported recurrent acute hepatitis that was characterized by necrosis on histopathological examination, related to intravenous MT that was given on 3 occasions for the management of relapsing multiple sclerosis (4). Loraschi *et al.* reported 2 cases of liver damage that was related to high-dose MT therapy for demyelinating disease (5). The first patient, a 33-year-old man, developed a histologically recognized acute steatohepatitis 5 weeks after last exposure to MT. The second patient, a 27-year-old woman, presented with moderate and asymptomatic augmentation of liver enzymes 6 days after withdrawal of MT. Hypersensitivity reactions were not observed in any patient. Despite their anti-inflammatory and antiallergic properties, corticosteroids also trigger immunoallergic liver injuries. Das *et al.* reported recurrent liver injuries that occurred 6 weeks and 3 weeks following the second and third course of intravenous MT, respectively, for multiple sclerosis (6). Liver biopsy showed lobular, primarily perivenular, infiltration with activated lymphocytes, eosinophils, and plasma cells. Moreover, Japanese authors reported the occurrence of autoimmune hepatitis, confirmed by liver histology, in a patient with multiple sclerosis who was treated with MP pulses (7). In their opinion, autoimmune hepatitis was a consequence of an immune rebound phenomenon after pulsed MP.

The mechanisms of corticosteroid-induced liver injury are unclear and only occasionally are related to reactivation of HBV infection or to the excipient of the MT preparation (6, 8, 9). Though low doses of corticosteroids are considered safe for the liver, chronic administration of these drugs may be associated with steatosis or steatohepatitis (2, 10). Intrinsic hepatotoxicity of high doses of corticosteroids is rather unlikely, as serious hepatic injuries that are related to MT occur rarely and are unpredictable. The majority of hepatotoxic drugs causes idiosyncratic reactions. There are two types of idiosyncrasy: immunoallergic and nonallergic (metabolic). The essence of an immunoallergic reaction is a complex interaction between a parent drug or its metabolites with immunologically competent cells, leading to necrosis and apoptosis of hepatocytes. Released cytokines additionally damage liver cells or have immune-modulating effects (11). In metabolic idiosyncrasy, the liver injury is caused by aberrant hepatic metabolism, leading to overproduction of reactive metabolites from a parent compound. In general, idiosyncratic drug-induced hepatotoxicity is considered to be unpredictable and dose-independent (11); however,

Table. Glucocorticosteroid-Induced Hepatotoxicity: Review of 13 Cases

Reference	Age/ Sex	Principal Disease	Type of Steroid	Dose and Duration of Treatment	Max. ALT <sup>a</sup> /AST <sup>a</sup> , IU/L	Max. GGT <sup>a</sup> /ALP <sup>a</sup> , IU/L	Histology	Concomitant Treatment	Follow-up
Gerolami et al. (8)	27/F <sup>a</sup>	Crohn disease	MP <sup>a</sup>	50 mg daily IV, 2 days	7.5xN <sup>a</sup> /3.2xN	5.1xN/1.8xN	Biopsy not done	None	Normalization of liver tests after MP discontinuation
Nanki et al. (1)	53/F	Systemic lupus erythematosus	P	60 mg daily PO, 6 days 20 mg daily; PO, (105 days)	658/871	ND <sup>a</sup>	Macrovesicular steatosis and mild periportal PMN a infiltration (autopsy)	None	Death
Dourakis et al. (2)	67/F	Dermatomyositis	P	25 mg/t.i.d. IV, (26 days)	545/1229	2092/467	Macrovesicular steatosis and mild portal lymphocyte and PMN infiltration (autopsy)	None	Death
Weissel et al. (3)	71/F	Graves ophthalmopathy	MP + C <sup>a</sup>	1,0 g daily IV, 3 days- tapering to 0 within 10-14 days; 5 courses	ND	ND	Necrosis of liver parenchyme (autopsy)	Methimazole started 6 months before MP, continued until the last course	Death
Salvi et al. (17)	43/F	Thyroid associated ophthalmopathy	MP	7.5 mg/kg IV, every 2 weeks (4 courses)	1200/850	ND	Compatible with autoimmune hepatitis	Levothyroxine since 3 years	Normalization of liver tests
Hofstee et al. (18)	46/F	Multiple sclerosis	MP	0,5 g IV, 5 days 1 g IV, 3 days; 2 years later 1 g IV, 3 days; 4 years later 1 g IV, 3 days ; 9 years later	N	ND	Biopsy not done	None	Normalization of liver tests after MP discontinuation

Das et al. (6)	48/F	Multiple sclerosis	MP	ND; 3 courses	1650/1430	ND/590	Preserved architecture with lobular infiltration by lymphocytes, eosinophils and plasma cells	None	Normalization of liver tests after MP discontinuation
Topal et al. (19)	47/F	Vasculitis of the central nervous system	MP	ND; PO, 7 days course	2478/1600	242/138	Biopsy not done	Topiramate; since 1 year and during MP course	Normalization of liver tests after MP discontinuation
Rivero Fernandez et al. (4)	57/F	Multiple sclerosis	MP	1,0 g IV, 3 days	1223/543	71/113	Acute necrotic hepatitis with ceroid-laden macrophage hyperplasia	None	Normalization of liver tests 3 months after MP discontinuation
Takahashi et al. (7)	43/F	Multiple sclerosis	MP+P	1,0 g IV, 3 days followed by 50 mg/dL PO, for 1 month	Normal	Normal	Bridging perivenular necrosis with infiltration by inflammatory cells including eosinophils (first biopsy)	None	Normalization of liver tests several months after MP discontinuation. P tapered within 8 months
			MP+P	1 g IV, 3 days; 3 years later followed by 50 mg/d PO, plus	1067/1102	26/377	Bridging perivenular necrosis and interface hepatitis (second biopsy)	6 doses of Interferone beta-1b	Normalization of liver tests 3 months after MP discontinuation. P tapered within 3 months
			MP	1 g IV, 3 days; 5 days later	566/875	1785/214			
			MP	1 g IV, 3 days; 13 months later					

Loraschi et al. (5)	33/M <sup>a</sup>	Demyelinating encephalopathy	MP	Total dose 2.5 g IV, 4 days course	1042/349	ND/ND	Focal liver cell necrosis in acinar zones 2 and 3, monocyte/macrophage infiltration, Kupffer cell hyperplasia, acidophilic bodies and focal microvesicular steatosis	None	Normalization of liver tests 20 days after MP discontinuation
Loraschi et al. (5)	27/F	Retrolubar optic neuritis	MP	Total dose 4.5 g IV, 6 days course	122/39	ND/ND	Biopsy not done	None	Normalization of liver tests 4 days after MP discontinuation
Gutkowski et al. (present case)	24/F	Multiple sclerosis	MP	Total dose 3.0g IV, 6 days course	Normal	Normal	Biopsy not done	None	Normalization of liver tests 3 weeks after MP discontinuation
			MP	Total dose 3.0g IV, 6 days course; 6 weeks later	1740/900	186/186		4 doses of Interferone beta-1b 0.5 g acetaminophen; 4 times	Normalization of liver tests 6 weeks after MP discontinuation
			MP	Total dose 3.0g IV, 6 days course; 3 months later	1129/1488	168/164		None	Normalization of liver tests 6 weeks after MP discontinuation

<sup>a</sup> Abbreviations: ALP, alkaline phosphatase; ALT, alanine aminotransferase; AST, aspartate aminotransferase; C, cortisone; F, female; GGT,  $\gamma$ -glutamyltransferase; M, male; MP, methylprednisolone; N, normal; ND, not determined; P, prednisolone; PMN, polymorphonuclear leukocyte.

it has been suggested that there is dose-dependency for drugs that are extensively metabolized in the liver (12). MT is metabolized by cytochrome P450 3A4 (CYP3A4), and its metabolites undergo renal elimination.

In our patient, we did not observe any systemic hypersensitivity symptoms before or during liver injury. After two exposures to MT, the hepatitis episodes appeared 4 weeks after drug withdrawal. In metabolic idiosyncrasy, the latency periods vary considerably from days to months, and adverse reactions can occur even several weeks after drug discontinuation. Such examples are not common but have been reported for some antibiotics, such as amoxicillin clavulanate, midecamycin, trovafloxacin, and flucloxacillin (13-16). The duration of anti-inflammatory effects after a single intramuscular injection of 40-80 mg MT ranges from 4 days to 8 days. Thus, it seems unlikely that chemically active metabolites damage hepatocytes 4 weeks after drug discontinuation. Rather, this scenario suggests a delayed immune response to the metabolite that is bound to the host protein and successive presentation as a neoantigen to the immune cells following the death of hepatocytes. The appearance of SMA following exposure to MT might be a feature of the immunological response.

#### 4. Conclusion

MT pulses are increasingly used by neurologists, rheumatologists, and endocrinologists to treat various autoimmune diseases. The general awareness of the potential hepatotoxicity of high-dose corticosteroids is very low. Corticosteroid-induced liver injury may occur as acute hepatitis that develops several weeks after short-term drug exposure. We therefore feel that MT should be placed on the list of hepatotoxic drugs and that patients who receive corticosteroid pulses should be screened for potential liver injury.

#### Acknowledgements

None declared.

#### Financial Disclosures

None declared.

#### Funding/Support

None declared.

#### References

1. Nanki T, Koike R, Miyasaka N. Subacute severe steatohepatitis during prednisolone therapy for systemic lupus erythematosus. *Am J Gastroenterol.* 1999;**94**(11):3379.
2. Dourakis SP, Sevastianos VA, Kaliopi P. Acute severe steatohepatitis related to prednisolone therapy. *Am J Gastroenterol.* 2002;**97**(4):1074-5.
3. Weissel M, Hauff W. Fatal liver failure after high-dose glucocorticoid pulse therapy in a patient with severe thyroid eye disease. *Thyroid.* 2000;**10**(6):521.
4. Rivero Fernandez M, Riesco JM, Moreira VF, Moreno A, Lopez San Roman A, Arranz G, et al. Recurrent acute liver toxicity from intravenous methylprednisolone. *Rev Esp Enferm Dig.* 2008;**100**(11):720-3.
5. Loraschi A, Banfi P, Mauri M, Sessa F, Bono G, Cosentino M. Hepatotoxicity after high-dose methylprednisolone for demyelinating disease. *Clin Neuropharmacol.* 2010;**33**(1):52-4.
6. Das D, Graham I, Rose J. Recurrent acute hepatitis in patient receiving pulsed methylprednisolone for multiple sclerosis. *Indian J Gastroenterol.* 2006;**25**(6):314-6.
7. Takahashi A, Kanno Y, Takahashi Y, Sakamoto N, Monoe K, Saito H, et al. Development of autoimmune hepatitis type 1 after pulsed methylprednisolone therapy for multiple sclerosis: a case report. *World J Gastroenterol.* 2008;**14**(35):5474-7.
8. Gerolami R, Mambrini P, Barthet M, Jean-Pastor MJ, Salducci J, Grimaud JC. Acute hepatitis caused by Solupred in a patient with Crohn disease. *Gastroenterol Clin Biol.* 1997;**21**(3):236-7.
9. Hammond A, Ramersdorfer C, Palitzsch KD, Scholmerich J, Lock G. Fatal liver failure after corticosteroid treatment of a hepatitis B virus carrier. *Dtsch Med Wochenschr.* 1999;**124**(22):687-90.
10. Stravitz RT, Sanyal AJ. Drug-induced steatohepatitis. *Clin Liver Dis.* 2003;**7**(2):435-51.
11. Gunawan BK, Kaplowitz N. Mechanisms of drug-induced liver disease. *Clin Liver Dis.* 2007;**11**(3):459-75, v.
12. Kaplowitz N. Idiosyncratic drug hepatotoxicity. *Nat Rev Drug Discov.* 2005;**4**(6):489-99.
13. Andrade RJ, Lucena MI, Fernandez MC, Vega JL, Camargo R. Hepatotoxicity in patients with cirrhosis, an often unrecognized problem: lessons from a fatal case related to amoxicillin/clavulanic acid. *Dig Dis Sci.* 2001;**46**(7):1416-9.
14. Perez Moreno JM, Saldana Gonzalez FJ, Puertas Montenegro M, Baez Perea J. Cholestatic hepatitis caused by midecamycin. *Gastroenterol Hepatol.* 1996;**19**(9):459-61.
15. Lucena MI, Andrade RJ, Rodrigo L, Salmeron J, Alvarez A, Lopez-Garrido MJ, et al. Trovafloxacin-induced acute hepatitis. *Clin Infect Dis.* 2000;**30**(2):400-1.
16. Olsson R, Wiholm BE, Sand C, Zettergren L, Hultcrantz R, Myrhed M. Liver damage from flucloxacillin, cloxacillin and dicloxacillin. *J Hepatol.* 1992;**15**(1-2):154-61.
17. Salvi M, Vannucchi G, Sbrozzi F, Del Castello AB, Carnevali A, Fargion S, et al. Onset of autoimmune hepatitis during intravenous steroid therapy for thyroid-associated ophthalmopathy in a patient with Hashimoto's thyroiditis: case report. *Thyroid.* 2004;**14**(8):631-4.
18. Hofstee HM, Nanayakkara PW, Stehouwer CD. Acute hepatitis related to prednisolone. *Eur J Intern Med.* 2005;**16**(3):209-10.
19. Topal F, Ozaşlan E, Akbulut S, Kucukazman M, Yuksel O, Altiparmak E. Methylprednisolone-induced toxic hepatitis. *Ann Pharmacother.* 2006;**40**(10):1868-71.