



Published in final edited form as:

J Thorac Oncol. 2011 June ; 6(6): 1128–1131. doi:10.1097/JTO.0b013e3182161508.

Phase II Trial of Dasatinib for Patients with Acquired Resistance to Treatment with the Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors Erlotinib or Gefitinib

Melissa L. Johnson, MD¹, Greg J. Riely, MD, PhD¹, Naiyer A. Rizvi, MD¹, Christopher G. Azzoli, MD¹, Mark G. Kris, MD¹, Camelia S. Sima, MD, MS², Michelle S. Ginsberg, MD³, William Pao, MD, PhD⁴, and Vincent A. Miller, MD¹

¹Department of Medicine, Memorial Sloan-Kettering Cancer Center, New York, NY

²Department of Epidemiology and Biostatistics, Memorial Sloan-Kettering Cancer Center, New York, NY

³Department of Radiology, Memorial Sloan-Kettering Cancer Center, New York, NY

⁴Vanderbilt-Ingram Cancer Center, Vanderbilt University, Nashville, TN

Abstract

Introduction—Dual inhibition of *SRC* and *EGFR* -dependent pathways may overcome acquired resistance to EGFR-TKIs for patients with lung adenocarcinoma with *EGFR* mutations. The *SRC*-inhibitor dasatinib demonstrates anti-tumor activity in gefitinib-resistant cells lines and xenografts. Dasatinib is tolerable for patients with advanced non-small cell lung cancer, and in combination with erlotinib.

Methods—We conducted this phase II study of dasatinib 70 mg twice daily in patients with *EGFR*-mutant lung adenocarcinoma and acquired resistance to EGFR-TKIs. After a protocol amendment based upon evolving data about both drugs, patients received dasatinib at a dose of 100 mg daily with continued erlotinib after developing acquired resistance. Enrolled patients either harbored an activating mutation in *EGFR* or experienced clinical benefit with single-agent erlotinib or gefitinib, followed by RECIST documented progression while being treated with an EGFR-TKI.

Results—Twenty-one patients were enrolled, nine under the original trial design and 12 after the protocol amendments. We observed no complete or partial responses (0% observed rate, 95% CI 0–18%). The median time to progression was 0.5 months (range 0.2–1.8 months) in patients treated with dasatinib, and 0.9 months (range 0.4–5 months) for patients treated with dasatinib and erlotinib in combination. Pleural effusions and dyspnea were frequent toxicities.

Conclusions—Dasatinib has no activity in patients with *EGFR*-mutant lung adenocarcinoma with acquired resistance to erlotinib and gefitinib.

INTRODUCTION

Seventy percent of patients with lung adenocarcinoma harboring mutations in the epidermal growth factor receptor gene (*EGFR*) experience a partial response when treated with EGFR

Corresponding Author: Melissa L. Johnson, MD Robert H. Lurie Cancer Center of Northwestern University, 675 North St. Clair Street Suite 21-100 Chicago, IL 60611, melissa-johnson@northwestern.edu.

Work previously presented as a Poster at the 13th World Conference on Lung Cancer: *J Thor Oncol* 2009 4(9): S687 (suppl 1; abstr 6408)

tyrosine kinase inhibitors (TKIs) erlotinib or gefitinib.¹ However, the majority of patients progress within seventeen months of starting treatment.² At least 50% of lung cancer patients with acquired resistance to erlotinib or gefitinib develop a secondary T790M mutation within *EGFR*, and another 10–15% of patients demonstrate *MET* amplification.^{3–5} Therapies directed against these mechanisms of acquired resistance are desperately needed.

SRC is a non-receptor tyrosine kinase which demonstrates increased protein levels in *EGFR*-dependent tumors. *SRC* and *EGFR* are proteins capable of mutual phosphorylation that share downstream effectors such as phosphatidylinositol 3-kinase/PTEN/Akt and *STAT* proteins.⁶ Because of these functional associations, *SRC* kinase has been proposed as a target to overcome acquired resistance in *EGFR*-mutant tumors.

Preclinical models demonstrate *EGFR* mutant cell lines containing either L858R (H3255) or exon 19 deletions (PC9 or HCC827) undergo apoptosis when treated with the *SRC* inhibitor dasatinib.⁷ Gefitinib-resistant adenocarcinoma cells with T790M (PC9/ZD) or *MET* amplification (HCC827 GR5) undergo cell death when treated with dasatinib.⁸ Dasatinib also inhibits tumor growth in HCC827 GR5 nude mouse xenografts.⁸ Dasatinib has been studied in patients with advanced solid tumors, with pleural effusions dose-limiting.⁹ Dasatinib can be combined with erlotinib in unselected patients with advanced non-small cell lung cancer.¹⁰

Given its pre-clinical rationale and early clinical trial results, we conducted a phase II study of dasatinib in patients with *EGFR*-mutant lung adenocarcinoma and acquired resistance to the EGFR-TKIs erlotinib and gefitinib.

METHODS

Patients with lung adenocarcinoma meeting consensus criteria for acquired resistance were eligible.¹¹ All patients agreed to undergo a repeat tumor biopsy. This protocol was reviewed and approved by the Institutional Review Board at Memorial Sloan-Kettering Cancer Center.

Initial Study Design

Seven days after discontinuing erlotinib or gefitinib therapy, patients began dasatinib at a dose of 70 mg twice daily. Patients were evaluated by CT scan at four weeks, eight weeks, and at eight-week intervals. Modified Response Evaluation Criteria in Solid Tumors (version 1.1) were used to assess response. Toxicities were graded using the National Cancer Institute Common Terminology Criteria of Adverse Events (version 3.0).

A Simon two-stage design was employed to calculate an initial sample size of 12. Cohort expansion to 37 patients was planned if one or more partial responses were observed. If four or more of the first 12 patients developed grade 3 or 4 pleural effusions, the trial would be stopped.

Amended Design

Coincident with the start of this trial, we observed that patients with acquired resistance who discontinued EGFR-TKIs experienced symptomatic deterioration and accelerated tumor growth with increased FDG avidity on PET scans.¹² After restarting erlotinib or gefitinib in these patients, tumors decreased in size and SUV_{max} on repeat studies, and tumor-related symptoms improved. Given these observations, we now recommend continued erlotinib in patients with acquired resistance while also adding second-line treatment agents, and amended this protocol to allow patients to continue erlotinib in addition to beginning dasatinib. In addition, new data indicated that dasatinib 100 mg daily provided similar

efficacy with less pleural effusions when used to treat patients with chronic myelogenous leukemia (CML),¹³ and we further amended this trial to allow a dose of dasatinib 100 mg daily.

Mutational Analysis

Prior to initiating dasatinib, all patients underwent tumor biopsies, preferably at a site of growing or new disease. Genomic DNA was extracted from tumor specimens, and all *EGFR* mutations (exon 19 deletions, L858R and T790M substitutions) were identified by mutation-specific PCR-based methods.¹⁴ Tumor specimens were analyzed for *MET* amplification using dual-color fluorescent in situ hybridization (FISH) with a *MET* specific gene probe.⁵ *MET* amplification was defined as having a *MET*: CEP7 ratio of >2:1.

RESULTS

Dasatinib 70 mg twice daily

Nine patients were enrolled under the original trial design. The median age was 68 years, and 66% of patients were women (Table 1). Similar numbers of patients in this cohort harbored exon 19 deletions and L858R mutations in their tumors. One patient had insufficient tissue for analysis (Table 2).

Patients were treated for a median 16 months with primary EGFR-TKIs before developing acquired resistance. When re-biopsied at the time of study enrollment, 44% (4/9) of patients had developed T790M acquired resistance mutations; none of the patients with adequate tissue for FISH testing exhibited *MET* amplification (0/4 tested) (Table 2).

There were no complete or partial responses observed (0%, 95% confidence interval: 0–34%). All patients progressed within 2 months of starting dasatinib. The median time until progression was 0.5 months (range, 0.2–1.8 months). The median overall survival was 13 months.

The combination of rapid disease progression among these initial nine patients, the majority (6/9) of whom developed pleural effusions, prompted revisions to our protocol design, although still three patients away from its required sample size of 12. We decided the protocol revisions were necessary (see **Methods, Amended Design** above) in order to effectively evaluate our original study hypothesis.

Dasatinib 100 mg daily + Erlotinib

Twelve more patients were enrolled. The median age was 65 years, and 58% were women (Table 1). The majority (80%) of this cohort had tumors with exon 19 deletions (Table 2). Patients received a median of 21 months of primary EGFR-TKI therapy prior to the development of acquired resistance. At re-biopsy, 75% had T790M (Table 2). *MET* amplification was not identified in the 7 specimens tested.

Twelve patients were treated with dasatinib and erlotinib, and no complete or partial responses were observed (0% objective response rate, 95% confidence interval: 0–28%). Patients were treated with dasatinib 100 mg daily and erlotinib for a median of 0.9 months (range, 0.4 to 5.4 months).

Toxicities

The primary toxicity was the development and/or enlargement of pre-existing pleural effusions and dyspnea. Among patients treated with dasatinib alone, 3 patients required

hospitalization for thoracostomy tube placement. One patient receiving dasatinib with erlotinib required a similar intervention. Peripheral and facial edema were also reported.

Fatigue was another significant side effect, whether patients were treated with dasatinib alone or with erlotinib. Patients reported grade 3 fatigue in 2/9 (22%) and 2/12 (17%) of patients, respectively. Nausea and vomiting (one episode of grade 3 toxicity each), as well as grade 2 diarrhea were reported.

DISCUSSION

In this phase II trial of *SRC* inhibitor dasatinib, with and without erlotinib, in patients with lung adenocarcinoma and acquired resistance to erlotinib, no objective responses were observed. Pleural effusions and dyspnea were the most frequent and significant toxicities. Regardless of whether erlotinib was continued, patients were treated with dasatinib for a median of less than one month due to a combination of disease progression and toxicity.

Haura et al. studied dasatinib and erlotinib in unselected patients with advanced lung cancer previously untreated.¹⁰ Two partial responses were reported, one in a patient whose tumor harbored an *EGFR* exon 19 deletion. This response likely reflects solely the effect of erlotinib in a tumor with a sensitizing-*EGFR* mutation. Very recently, Johnson et al. also reported the results of a phase II trial of single-agent dasatinib as first-line treatment for unselected patients with advanced NSCLC. Neither *SRC* activity nor mutations in *EGFR* (nor *KRAS*) were associated with the modest response rates reported—overall disease control rate of 43% and progression-free survival of 1.36 months.¹⁵

Pleural effusions were observed in 66% of patients enrolled in this study. The pleural effusions observed were clinically more significant for the patients who received dasatinib 70 mg twice daily—50% of whom required thoracentesis or tube thoracostomy. Only one patient treated with dasatinib 100 mg daily and erlotinib required a tube thoracostomy. Prior studies in patients with CML and advanced solid tumors treated with dasatinib also observed pleural effusions to be dose-related.^{9,13} Johnson et al. reported that the presence of pleural effusions in patients with advanced non-small cell lung cancer prior to treatment with dasatinib predicted the development of clinically significant effusions during treatment.¹⁵

In many cases it was difficult to discern whether these new pleural effusions were dasatinib-related drug toxicity or evidence of progressive cancer. Unlike patients with CML or solid tumors other than lung cancer treated with dasatinib, in whom pleural effusions can easily be recognized as drug toxicity and managed with dose attenuation, diuretics, and corticosteroids without treatment delays, this complication proved very challenging to manage in our patients with lung adenocarcinoma.

The amended trial design also limited interpretation of the study's results. We simultaneously changed the dasatinib dosing schedule and added erlotinib to the treatment regimen to incorporate the latest information about both drugs. By making changes to two aspects of the trial design at once, we made identification of the true reason (or reasons) for the disappointing results more difficult.

We found the combination of dasatinib and erlotinib to be inactive, with significant toxicities. Our study suggests these drugs in combination merit no further study in patients with *EGFR*-driven lung adenocarcinoma. Therefore, these patients remain ideal candidates for clinical trials investigating new strategies to overcome acquired resistance.

Acknowledgments

NCI grant CA (130343-01)

ASCO Young Investigator Award—Greg Riely MD, PhD

Acknowledgement of Support: R21 CA 130343-01

References

1. Mok TS, Wu YL, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med.* 2009; 361:947–57. [PubMed: 19692680]
2. Janne PAWX, Socinski MA, Crawford J, Capelletti M, Edelman MJ, Villalona-Calero MA, Kratzke RA, Vokes EE, Miller VA. Randomized phase II trial of erlotinib (E) alone or in combination with carboplatin/paclitaxel (CP) in never or light former smokers with advanced lung adenocarcinoma: CALGB 30406. *J Clin Oncol.* 2010;28.
3. Pao W, Miller VA, Politi KA, et al. Acquired resistance of lung adenocarcinomas to gefitinib or erlotinib is associated with a second mutation in the EGFR kinase domain. *PLoS Med.* 2005; 2:e73. [PubMed: 15737014]
4. Engelman JA, Zejnullahu K, Mitsudomi T, et al. MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling. *Science.* 2007; 316:1039–43. [PubMed: 17463250]
5. Bean J, Brennan C, Shih JY, et al. MET amplification occurs with or without T790M mutations in EGFR mutant lung tumors with acquired resistance to gefitinib or erlotinib. *Proc Natl Acad Sci U S A.* 2007; 104:20932–7. [PubMed: 18093943]
6. Tice DA, Biscardi JS, Nickles AL, Parsons SJ. Mechanism of biological synergy between cellular Src and epidermal growth factor receptor. *Proc Natl Acad Sci U S A.* 1999; 96:1415–20. [PubMed: 9990038]
7. Song L, Morris M, Bagui T, Lee FY, Jove R, Haura EB. Dasatinib (BMS-354825) selectively induces apoptosis in lung cancer cells dependent on epidermal growth factor receptor signaling for survival. *Cancer Res.* 2006; 66:5542–8. [PubMed: 16740687]
8. Yoshida T, Okamoto I, Okamoto W, et al. Effects of Src inhibitors on cell growth and epidermal growth factor receptor and MET signaling in gefitinib-resistant non-small cell lung cancer cells with acquired MET amplification. *Cancer Sci.* 2010; 101:167–72. [PubMed: 19804422]
9. Johnson FM, Agrawal S, Burris H, et al. Phase I pharmacokinetic and drug-interaction study of dasatinib in patients with advanced solid tumors. *Cancer.* 2010; 116:1582–91. [PubMed: 20108303]
10. Haura EB, Tanvetyanon T, Chiappori A, et al. Phase I/II study of the Src inhibitor dasatinib in combination with erlotinib in advanced non-small-cell lung cancer. *J Clin Oncol.* 2010; 28:1387–94. [PubMed: 20142592]
11. Jackman D, Pao W, Riely GJ, et al. Clinical definition of acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors in non-small-cell lung cancer. *J Clin Oncol.* 2010; 28:357–60. [PubMed: 19949011]
12. Riely GJ, Kris MG, Zhao B, et al. Prospective assessment of discontinuation and reinitiation of erlotinib or gefitinib in patients with acquired resistance to erlotinib or gefitinib followed by the addition of everolimus. *Clin Cancer Res.* 2007; 13:5150–5. [PubMed: 17785570]
13. Shah NP, Kantarjian HM, Kim DW, et al. Intermittent target inhibition with dasatinib 100 mg once daily preserves efficacy and improves tolerability in imatinib-resistant and -intolerant chronic-phase chronic myeloid leukemia. *J Clin Oncol.* 2008; 26:3204–12. [PubMed: 18541900]
14. Pan Q, Pao W, Ladanyi M. Rapid polymerase chain reaction-based detection of epidermal growth factor receptor gene mutations in lung adenocarcinomas. *J Mol Diagn.* 2005; 7:396–403. [PubMed: 16049312]
15. Johnson FM, Bekele BN, Feng L, et al. Phase II study of dasatinib in patients with advanced non-small-cell lung cancer. *J Clin Oncol.* 2010; 28:4609–15. [PubMed: 20855820]

Table 1

Patient Characteristics

Characteristic	Dasatinib 70 mg twice daily (n=9)	Dasatinib 100 mg daily + Erlotinib (n=12)
Age (mean, range)	68 (35–80)	60 (51–73)
Female/Male	6/3	7/5
Smoking History		
Never	7	8
Former	1	3
Current	1	1
KPS (median)	80	80
Months (median) treated with EGFR-TKI prior to developing acquired resistance	16 (8–102)	21 (12–49)

Table 2

Molecular Studies of Tumor Specimens

	Dasatinib 70 mg twice daily (n=9)	Dasatinib 100 mg daily + Erlotinib (n=12)
Primary <i>EGFR</i> Mutations		
Exon 19 deletion	4	10
L858R	4	2
Unable to be tested*	1	0
Molecular abnormalities found at the time of developing acquired resistance		
T790M + Exon 19 del	2	8
T790M + L858R	2	0
T790M not found + Exon 19 del	2	2
T790M not found + L858R	1	2
T790M unable to be tested*	2	0
<i>MET</i> amplification present	0	0
<i>MET</i> amplification absent	4	7
<i>MET</i> amplification unable to be tested*	5	5

* due to insufficient tumor tissue