

**Sagar Chittori,^a H. S. Savithri^b
 and M. R. N. Murthy^{a*}**

^aMolecular Biophysics Unit, Indian Institute of Science, Bangalore, Karnataka 560 012, India, and ^bDepartment of Biochemistry, Indian Institute of Science, Bangalore, Karnataka 560 012, India

Correspondence e-mail: mnrn@mbu.iisc.ernet.in

Received 11 September 2011

Accepted 21 October 2011


Preliminary X-ray crystallographic studies on acetate kinase (AckA) from *Salmonella typhimurium* in two crystal forms

Acetate kinase (AckA) catalyzes the reversible transfer of a phosphate group from acetyl phosphate to ADP, generating acetate and ATP, and plays a central role in carbon metabolism. In the present work, the gene corresponding to AckA from *Salmonella typhimurium* (*StAckA*) was cloned in the IPTG-inducible pRSET C vector, resulting in the attachment of a hexahistidine tag to the N-terminus of the expressed enzyme. The recombinant protein was overexpressed, purified and crystallized in two different crystal forms using the microbatch-under-oil method. Form I crystals diffracted to 2.70 Å resolution when examined using X-rays from a rotating-anode X-ray generator and belonged to the monoclinic space group *C2*, with unit-cell parameters $a = 283.16$, $b = 62.17$, $c = 91.69$ Å, $\beta = 93.57^\circ$. Form II crystals, which diffracted to a higher resolution of 2.35 Å on the rotating-anode X-ray generator and to 1.90 Å on beamline BM14 of the ESRF, Grenoble, also belonged to space group *C2* but with smaller unit-cell parameters ($a = 151.01$, $b = 78.50$, $c = 97.48$ Å, $\beta = 116.37^\circ$). Calculation of Matthews coefficients for the two crystal forms suggested the presence of four and two protomers of *StAckA* in the asymmetric units of forms I and II, respectively. Initial phases for the form I diffraction data were obtained by molecular replacement using the coordinates of *Thermotoga maritima* AckA (*TmAckA*) as the search model. The form II structure was phased using a monomer of form I as the phasing model. Inspection of the initial electron-density maps suggests dramatic conformational differences between residues 230 and 300 of the two crystal forms and warrants further investigation.

1. Introduction

The human gastrointestinal tract provides a natural environment for a large and diverse community of microbes which carry out degradation of complex metabolites that cannot be normally digested by the host (Cummings & Macfarlane, 1991; Miller & Wolin, 1979). The majority of the microbes in the human gut are anaerobic bacteria, which generate short-chain fatty acids (SCFAs; chain length of up to six C atoms) as metabolic byproducts of fermentation (Cummings & Macfarlane, 1991). SCFAs have been shown to inhibit bacterial growth and thus can increase host resistance towards these organisms (Barnes *et al.*, 1979; Bohnhoff & Miller, 1962; Cherrington *et al.*, 1991). However, studies on many aerobic bacteria, fungi and anaerobic bacteria such as *Escherichia coli* and *Salmonella typhimurium* have shown that these organisms possess pathways for the utilization of SCFAs as a source of carbon and energy (Cummings *et al.*, 1987; Klein *et al.*, 1971). The pathways that enable microorganisms to catabolize SCFAs are essential for preventing toxic effects of these compounds on microbes and for increasing their virulence against the host (Rishi *et al.*, 2005).

The degradation of SCFAs is initiated by their conversion to the corresponding activated SCF acyl-CoA derivatives (Nunn, 1986). Acetate can enter into multiple metabolic pathways after its conversion to acetyl-CoA. Its entry into the tricarboxylic/citric acid (TCA) cycle leads to the generation of ATP and NADH. Acetyl-CoA is also a precursor for the synthesis of various biomolecules (Nunn, 1986). In *S. typhimurium* and *E. coli* acetate can be converted to acetyl-CoA by reversible tandem reactions catalyzed by acetate kinase (AckA; EC 2.7.2.1) and phosphotransacetylase (Pta; EC 2.3.1.8) (Wolfe, 2005). AckA catalyzes the reversible transfer of a



© 2011 International Union of Crystallography
 All rights reserved

phosphate group from acetyl phosphate to ADP, generating acetate and ATP, in the presence of a metal ion, while Pta catalyzes the reversible transfer of an acetyl group from acetyl-CoA to a phosphate, resulting in the formation of acetyl phosphate (Fig. 1).

Biochemical reactions involving the transfer of a phosphate group to an acceptor ligand, similar to the AckA reaction (Fig. 1), are responsible for several indispensable cellular functions such as signal transduction and regulation of cell division (Das *et al.*, 1997). Although extensive biochemical studies have been carried out on AckA (Fox & Roseman, 1986; Wanner & Wilmes-Riesenberg, 1992), the structure of the enzyme (from the thermophilic organism *Methanosarcina thermophila*) was only reported in 2001 (Buss *et al.*, 2001). In the present work, we report cloning, overexpression, purification and preliminary X-ray crystallographic studies of AckA from the mesophilic organism *S. typhimurium* (*StAckA*; NCBI reference code NP_461279.1). Unexpected structural differences between two crystal forms of *StAckA* were observed, warranting further investigations into the significance of the conformational changes.

2. Materials and methods

2.1. Plasmid construction and protein purification

The 1200 bp open reading frame encoding AckA was PCR-amplified from *S. enterica* serovar Typhimurium strain IFO12529 genomic DNA template and cloned into pRSET C vector (Invitrogen). The final plasmid construct encodes recombinant *StAckA* with 15 additional amino acids (MRGSHHHHHHGMAH-*StAckA*) including a hexahistidine tag. The recombinant plasmid was transformed into *E. coli* BL21 (DE3) pLysS competent cells. The transformed cells were inoculated into 1 l Terrific Broth (HiMedia) containing 4 ml glycerol and allowed to grow at 310 K until the OD at 600 nm reached 0.6. Expression of the recombinant *StAckA* was then induced by addition of 0.3 mM isopropyl β -D-1-thiogalactopyranoside (IPTG). The culture was further incubated overnight at 293 K. The overexpressed protein was purified using Ni-NTA affinity chromatography followed by gel-filtration chromatography using a Superdex S-200 column (Fig. 2). The purified enzyme was dialyzed for 24 h against 25 mM Tris buffer pH 8.0 containing 100 mM NaCl and concentrated using a 10 kDa molecular-weight cutoff Centricon (Amicon).

2.2. Crystallization and X-ray diffraction

Two distinct crystal forms of *StAckA* suitable for structural studies were obtained at 298 K using the microbatch method. The crystallization drops consisted of 3 μ l 10 mg ml⁻¹ protein and 1 μ l crystallization cocktail. In both experiments crystals were observed after two months. Crystals of form I were obtained using polyethylene glycol 4000 as the precipitant, while form II crystals were obtained in the presence of trisodium citrate. Prior to X-ray diffraction data collection, a crystal was briefly exposed to a drop containing crys-

tallization buffer with 20% (w/v) ethylene glycol as a cryoprotectant and flash-cooled at 100 K in a nitrogen-gas stream. A complete three-dimensional X-ray diffraction data set was collected from a form I crystal at a wavelength of 1.5418 Å on a Bruker Microstar Ultra rotating-anode X-ray generator at the X-ray Facility for Structural Biology at the Molecular Biophysics Unit, Indian Institute of Science (IISc). The exposure time and crystal oscillation angles were set to 10 min and 1.0°, respectively. The crystal-to-detector distance was maintained at 200 mm. The crystal diffracted X-rays to 2.70 Å resolution. A total of 247 images were recorded using a MAR 345 image-plate detector. Form II crystals diffracted X-rays to a higher resolution of 2.35 Å at the home source. X-ray diffraction data from a form II crystal were subsequently collected on the BM14 beamline at the European Synchrotron Radiation Facility (ESRF), Grenoble, France at a wavelength of 0.95 Å; the crystal diffracted to 1.90 Å resolution. The crystal-to-detector distance was maintained at 187.25 mm. The exposure time and crystal oscillation angles were set to 15 s and 1.0°, respectively. A total of 210 images were recorded using a MAR 345/Offline FUJI image-plate detector. Both data sets were processed and scaled using the programs *DENZO* and *SCALEPACK*, respectively, from the *HKL-2000* suite (Otwinowski & Minor, 1997) and were further analyzed with programs from *CCP4* (Winn *et al.*, 2011).




Figure 2
Purification of the recombinant *StAckA*. (a) SDS-PAGE analysis of *StAckA*. Lane 1, crude cell lysate before IPTG induction; lane 2, crude cell lysate after induction with 0.3 mM IPTG; lane 3, soluble fraction of the cell lysate; lane 4, 20 mM imidazole wash; lane 5, purified *StAckA*; lane M, molecular-mass markers (labelled in kDa). (b) Gel-filtration chromatography of the purified *StAckA*. The elution volume suggests that the enzyme is a dimer in solution.





Figure 1
Reactions catalyzed by AckA and Pta.

3. Results and discussion


S. typhimurium AckA was cloned in the pRSET C vector (Invitrogen) with an N-terminal hexahistidine tag, overexpressed in *E. coli* and purified using Ni-NTA affinity-column chromatography. SDS-PAGE analysis revealed a single band with the expected molecular mass of 45 kDa (Fig. 2a). Gel-filtration chromatography performed on the purified *StAckA* indicated that it is in a dimeric form in solution (Fig. 2b), which is consistent with observations on other members of the acetokinase family (Buss *et al.*, 2001; Simanshu *et al.*, 2005). *StAckA* crystals were obtained in two forms using the microbatch method. Form I crystals (approximate dimensions of $0.4 \times 0.4 \times 0.2$ mm) of *StAckA* diffracted X-rays to 2.70 Å resolution and belonged to the monoclinic space group *C2*, with unit-cell parameters $a = 283.16$, $b = 62.17$, $c = 91.69$ Å, $\beta = 93.57^\circ$ (Fig. 3). Form II crystals

(approximate dimensions of $0.6 \times 0.3 \times 0.2$ mm) diffracted to 2.35 Å resolution at the home source (Cu $K\alpha$ radiation, $\lambda = 1.5418$ Å; data not shown) and to 1.90 Å resolution when examined on the BM14 beamline at the ESRF, Grenoble at a wavelength of 0.95 Å (Fig. 4). The form II crystals also belonged to the monoclinic space group *C2*, but had smaller unit-cell parameters $a = 151.01$, $b = 78.49$, $c = 97.48$ Å, $\beta = 116.37^\circ$. Calculation of Matthews coefficients (Matthews, 1968) suggested the presence of four and two *StAckA* protomers in the asymmetric units of the form I and form II crystals, respectively. Details of the data-collection and processing statistics for both crystal forms are summarized in Table 1.

Initial phases for form I *StAckA* diffraction data were determined by molecular replacement using a polyalanine model of the *TmAckA*




(a)




(b)

Figure 3 Crystals and X-ray diffraction pattern of form I *StAckA*. (a) Form I *StAckA* crystals used for three-dimensional X-ray diffraction data collection using X-rays from a rotating-anode generator. (b) A typical X-ray diffraction image obtained from a form I *StAckA* crystal.



(a)



(b)

Figure 4 Crystals and X-ray diffraction pattern of form II *StAckA*. (a) Form II *StAckA* crystals used for three-dimensional X-ray diffraction data collection using synchrotron radiation on BM14 at the ESRF. (b) A typical X-ray diffraction image obtained from a form II *StAckA* crystal.

Table 1

Data-collection and processing statistics.

Values in parentheses are for the highest resolution shell.

	Form I	Form II
Crystallization conditions	0.1 M HEPES pH 7.5, 30% (w/v) PEG 4000, 0.2 M calcium chloride	0.1 M HEPES pH 7.5, 1.4 M trisodium citrate
Wavelength (Å)	1.54	0.95
Temperature (K)	100	100
Resolution range	50.0–2.70 (2.80–2.70)	50.0–1.90 (1.97–1.90)
Space group	C2	C2
Unit-cell parameters (Å, °)	$a = 283.16, b = 62.17,$ $c = 91.69, \beta = 93.57$	$a = 151.01, b = 78.50,$ $c = 97.48, \beta = 116.37$
Observed reflections	76858	153171
Unique reflections	42614	79119
Data completeness (%)	96.9 (84.5)	98.6 (90.4)
Multiplicity	2.8 (1.9)	4.3 (3.4)
Mean $\langle I/\sigma(I) \rangle^\dagger$	8.12 (2.06)	16.0 (2.20)
$R_{\text{merge}}^\ddagger$ (%)	13.8 (55.8)	6.2 (78.6)
Matthews coefficient (Å ³ Da ⁻¹)	2.40	2.88
Solvent content (%)	48.7	52.8
Protomers in the asymmetric unit	4	2

$^\dagger I$ is the integrated intensity and $\sigma(I)$ is the estimated standard deviation of that intensity. $^\ddagger R_{\text{merge}} = \sum_{hkl} \sum_i |I_i(hkl) - \langle I(hkl) \rangle| / \sum_{hkl} \sum_i I_i(hkl)$, where $I_i(hkl)$ is the i th measurement of the intensity of reflection hkl and $\langle I(hkl) \rangle$ is the mean intensity.

(PDB entry 2iir; S. Mukhopadhyay, M. S. Hasson & D. A. Sanders, unpublished work; 46% sequence identity to *StAckA*) monomer as the search model. The initial solution obtained using *Phaser* (McCoy *et al.*, 2007) showed significant values of log-likelihood gain (LLG = 1100) and Z scores for the rotation function (RFZ = 6.9) and translation function (TFZ = 14.8). This initial model was subjected to 25 cycles of rigid-body refinement and 25 cycles of restrained refinement using *REFMAC5* (Murshudov *et al.*, 2011), which resulted in R_{work} and R_{free} values of 38% and 42%, respectively. The refined polyalanine model of form I *StAckA* was used to determine the structure of the form II crystal. Molecular replacement performed using *Phaser* (McCoy *et al.*, 2007) provided an initial solution with an LLG of 1366 and Z scores RFZ = 20.1 and TFZ = 19.4. Rigid-body and restrained refinement of this initial model using *REFMAC5* (Murshudov *et al.*, 2011) resulted in R_{work} and R_{free} values of 32% and 36%, respectively. Inspection of the electron-density map (data not shown) showed unexpectedly large disorder of residues 230–300 (hereafter referred as the variable segment) in both monomers of form II *StAckA*. Therefore, these residues were deleted from form I and the resulting model was used for re-estimation of the phases by molecular replacement. This led to significant improvements in

LLG (5668), Z scores (RFZ = 53.2, TFZ = 32.8) and R factors ($R_{\text{work}} = 31\%$, $R_{\text{free}} = 34\%$), suggesting that the structure of the variable segment in form II is indeed very different from that in form I. Further refinement and model building are currently in progress and the large structural changes that have been observed in the initial maps and their possible biological implications need to be further investigated.

We thank James and Babu of IISc and Dr Hassan Belrhali and Dr Babu Manjasetty of the ESRF for their assistance during X-ray diffraction data collection. MRNM and HSS thank the Department of Science and Technology (DST) and Department of Biotechnology (DBT), Government of India for financial support. SC acknowledges the Council for Scientific and Industrial Research, Government of India for the award of a Research Fellowship.

References

- Barnes, E. M., Impey, C. S. & Stevens, B. J. (1979). *J. Hyg. (Lond.)*, **82**, 263–283.
- Bohnhoff, M. & Miller, C. P. (1962). *J. Infect. Dis.* **111**, 117–127.
- Buss, K. A., Cooper, D. R., Ingram-Smith, C., Ferry, J. G., Sanders, D. A. & Hasson, M. S. (2001). *J. Bacteriol.* **183**, 680–686.
- Cherrington, C. A., Hinton, M., Mead, G. C. & Chopra, I. (1991). *Adv. Microb. Physiol.* **32**, 87–108.
- Cummings, J. H. & Macfarlane, G. T. (1991). *J. Appl. Bacteriol.* **70**, 443–459.
- Cummings, J. H., Pomare, E. W., Branch, W. J., Naylor, C. P. & Macfarlane, G. T. (1987). *Gut*, **28**, 1221–1227.
- Das, S., Yu, L., Gaitatzes, C., Rogers, R., Freeman, J., Bienkowska, J., Adams, R. M., Smith, T. F. & Lindelien, J. (1997). *Nature (London)*, **385**, 29–30.
- Fox, D. K. & Roseman, S. (1986). *J. Biol. Chem.* **261**, 13487–13497.
- Klein, K., Steinberg, R., Fiethen, B. & Overath, P. (1971). *Eur. J. Biochem.* **19**, 442–450.
- Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.
- McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C. & Read, R. J. (2007). *J. Appl. Cryst.* **40**, 658–674.
- Miller, T. L. & Wolin, M. J. (1979). *Am. J. Clin. Nutr.* **32**, 164–172.
- Murshudov, G. N., Skubák, P., Lebedev, A. A., Pannu, N. S., Steiner, R. A., Nicholls, R. A., Winn, M. D., Long, F. & Vagin, A. A. (2011). *Acta Cryst. D* **67**, 355–367.
- Nunn, W. D. (1986). *Microbiol. Rev.* **50**, 179–192.
- Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307–326.
- Rishi, P., Pathak, S. & Ricke, S. C. (2005). *J. Environ. Sci. Health B*, **40**, 645–657.
- Simanshu, D. K., Savithri, H. S. & Murthy, M. R. N. (2005). *J. Mol. Biol.* **352**, 876–892.
- Wanner, B. L. & Wilmes-Riesenberg, M. R. (1992). *J. Bacteriol.* **174**, 2124–2130.
- Winn, M. D. *et al.* (2011). *Acta Cryst. D* **67**, 235–242.
- Wolfe, A. J. (2005). *Microbiol. Mol. Biol. Rev.* **69**, 12–50.