

First published on March 27, 2002: e0011. doi: 10.1199/tab.0011

# Molecular and Developmental Biology of Inorganic Nitrogen Nutrition

Nigel M. Crawford<sup>a1</sup> and Brian G. Forde<sup>b</sup>

<sup>a</sup> Section of Cell and Developmental Biology, Division of Biological Sciences, University of California at San Diego, 9500 Gilman Dr., La Jolla, CA 92093-0116

<sup>b</sup> Department of Biological Sciences, Lancaster University, Lancaster LA1 4YQ, UK

<sup>1</sup>Corresponding Author, phone/fx: 858/534-1637; email: ncrawford@ucsd.edu

## INTRODUCTION

Unique among the major mineral nutrients, inorganic N is available to plants in both anionic and cationic forms ( $\text{NO}_3^-$  and  $\text{NH}_4^+$ , respectively). The relative abundance of these two ions in natural soils is highly variable and to a large degree depends on the relative rates of two microbial processes: mineralisation (the release of  $\text{NH}_4^+$  from organic N) and nitrification (the conversion of  $\text{NH}_4^+$  to  $\text{NO}_3^-$ ) (Marschner, 1995). In well-aerated soils nitrification is rapid, so that  $\text{NH}_4^+$  concentrations are low and  $\text{NO}_3^-$  is the main N source, but in waterlogged or acidic soils nitrification is inhibited and  $\text{NH}_4^+$  accumulates. Most plants (including *Arabidopsis*) seem to be able to use either form of N, although exceptions to this rule are known (e.g. Kronzucker et al., 1997).

Nitrogen's importance in plant biology extends far beyond its role as a nutrient. It is now clear that several different N compounds, including  $\text{NO}_3^-$ ,  $\text{NH}_4^+$  and some of the products of their assimilation, exert strong regulatory effects on both metabolic and developmental pathways (Redinbaugh and Campbell, 1991; Crawford, 1995; Forde and Clarkson, 1999; Stitt, 1999; Zhang and Forde, 2000; Coruzzi and Bush, 2001; Coruzzi and Zhou, 2001). Both the biochemical and the regulatory aspects of inorganic N nutrition with emphasis on *Arabidopsis* will be considered in this chapter.

## NITRATE ASSIMILATION

The assimilation of  $\text{NO}_3^-$  involves just a few steps: uptake, reduction to nitrite, reduction to  $\text{NH}_4^+$  then incorporation into amino acids (Figure 1). Though simple, this pathway often provides the bulk of net nitrogen input, and, as a result, is tightly regulated to insure that  $\text{NO}_3^-$  assimilation is integrated into overall metabolism and growth of the plant. Many excellent reviews are available on this topic. These include recent reviews on  $\text{NO}_3^-$  uptake (Crawford and Glass, 1998; Daniel-Vedele et al., 1998; Forde and Clarkson, 1999; Crawford et al., 2000; Forde, 2000; Galvan and Fernandez, 2001; Glass et al., 2001; Williams and Miller, 2001) and on  $\text{NO}_3^-$  reduction and its regulation (Campbell, 1999; Kaiser et al., 1999; Mendel and Schwarz, 1999; Stitt, 1999; Zimmer and Mendel, 1999; Crawford et al., 2000; Tischner, 2000; Campbell, 2001; Coruzzi and Bush, 2001; MacKintosh and Meek, 2001). In this section, we focus on *Arabidopsis* after reviewing the major principles of  $\text{NO}_3^-$  assimilation that have been uncovered by work with plants and eukaryotic algae in general.




**Figure 1. Nitrate assimilation pathway.** NRT: nitrate transporters; NR: nitrate reductase; NiR; nitrite reductase.

## Nitrate uptake

The first step in the assimilation of  $\text{NO}_3^-$  is usually uptake from the soil solution (reviewed by Glass and Siddiqi, 1995; Crawford and Glass, 1998; Forde and Clarkson, 1999; Galvan and Fernandez, 2001; Glass et al., 2001; Williams and Miller, 2001). Nitrate concentrations vary widely with typical levels in agricultural soils ranging from 1-10 mM. Uptake occurs in the outer cell layers of the root by active transport processes mediated by  $\text{H}^+$  cotransport. In other words, the energy stored in the plasma membrane  $\text{H}^+$  gradient drives  $\text{NO}_3^-$  transport into epidermal and cortical cells. Such activity can be followed by measuring the depletion of  $\text{NO}_3^-$  from the media surrounding roots or by monitoring the changes in electrical potential across the plasma membrane with microelectrodes as  $\text{NO}_3^-$  uptake coincides with a depolarization of the plasma membrane, i.e. becomes more positive inside (Miller et al., 2001). An excellent and extensive electrophysiological study using Arabidopsis root hairs has been reported (Meharg and Blatt, 1995). Such experiments uncovered two general systems that account for  $\text{NO}_3^-$  uptake: a high affinity system, whose  $K_m$  for  $\text{NO}_3^-$  ranges from 10-100  $\mu\text{M}$ , and a low affinity system, which typically shows linear or nonsaturable kinetics and is most evident above 500  $\mu\text{M}$   $\text{NO}_3^-$ . Both systems show depolarization in response to  $\text{NO}_3^-$  indicating that both are active and involve cotransport. The high and low affinity systems correspond to the two classical mechanisms, Mechanism I and Mechanism

II, respectively, described by Epstein, who first showed ion uptake in plant roots obeyed Michaelis-Menten kinetics (Epstein, 1953; Epstein, 1972).

An important aspect of  $\text{NO}_3^-$  uptake is that it is regulated (reviewed by Crawford and Glass, 1998; Daniel-Vedele et al., 1998; Forde and Clarkson, 1999; Forde, 2000; Galvan and Fernandez, 2001; Glass et al., 2001; Williams and Miller, 2001). Various environmental and internal signals influence the rate of uptake into roots. Nitrate itself acts as an inducer. When plants are deprived of  $\text{NO}_3^-$  (either by starving them of nitrogen or by providing them with another form of nitrogen such as  $\text{NH}_4^+$  or an amino acid), they retain a basal or constitutive level of  $\text{NO}_3^-$  uptake that has both high and low affinity components (abbreviated as cHATS and cLATS, respectively). Upon exposure to  $\text{NO}_3^-$ , uptake rates increase. The characteristics of such induced activity (abbreviated iHATS) indicate that it is a high affinity system distinct from cHATS. In addition to  $\text{NO}_3^-$  induction, uptake shows feedback inhibition and responds to signals from the shoot. Nitrate uptake decreases when plants are fed  $\text{NH}_4^+$  or high levels of  $\text{NO}_3^-$ . Uptake also responds to circadian rhythms, reduced carbon, and shoot nitrogen demand. A classic example of shoot demand comes from split root experiments where one half of the roots from a single plant are placed in one chamber and the other half are placed in a second chamber (Figure 2). The first half is monitored for  $\text{NO}_3^-$  uptake in a solution with ample  $\text{NO}_3^-$ . If the remaining half of the roots is placed in a solution with low  $\text{NO}_3^-$ , uptake rates increase in the first half. It is thought that the shoots sense a reduction in overall nitrogen availability when the second group of roots is placed in low  $\text{NO}_3^-$  solutions. The shoots then respond by sending a signal to the roots to increase  $\text{NO}_3^-$  uptake.



**Figure 2. Schematic diagram showing a split-root experiment.** Figure was adapted from (Cerezo et al., 2001).

## Nitrate reduction

Once  $\text{NO}_3^-$  has been taken up into the cytosol, it can be (1) mobilized into the xylem for transport to the shoot, (2) transported into the vacuole for storage or (3) reduced to nitrite, a reaction that is catalyzed by  $\text{NO}_3^-$  reductase (NR), a cytosolic enzyme that uses NAD(P)H for reductant (Campbell, 1999; Campbell, 2001). Nitrate reduction is the first committed step in the  $\text{NO}_3^-$  assimilation pathway and is regulated at both transcriptional and post-translational levels. Both the  $\text{NO}_3^-$  reductase genes (NIA) and the NR protein respond to many signals (Crawford, 1995; Kaiser et al., 1999; Stitt, 1999; Tischner, 2000; MacKintosh and Meek, 2001). As is the case for  $\text{NO}_3^-$  uptake,  $\text{NO}_3^-$  serves a key inducer that rapidly activates transcription of the NIA

genes. Sucrose, light and cytokinin enhance NIA induction while  $\text{NH}_4^+$  and amino acids repress expression. NIA genes also respond to circadian rhythms.


Another well understood mechanism of NR control occurs at the protein level (Kaiser et al., 1999; MacKintosh and Meek, 2001). NR produces nitrite, which is toxic to plants if it accumulates. Nitrite is transported into the chloroplasts in green tissues (plastids in non-green tissues), where it is reduced to  $\text{NH}_4^+$  by nitrite reductase (NiR) using reduced ferredoxin. If nitrite reduction is slowed (e.g. caused by a reduction in light, reduced carbon or oxygen), nitrite levels could increase to toxic levels if NR activity is not inhibited. Plants have a rapid and reversible mechanism for doing just this. NR is phosphorylated by one of several kinases on a key serine in the first hinge region, which separates the Mo-cofactor and heme-binding regions of NR. NR has three redox centers (FAD, heme and the Mo-cofactor) that are bound by three distinct regions of the protein (Campbell, 1999; Campbell, 2001). The Mo-cofactor transfers electrons to  $\text{NO}_3^-$  while the heme shuttles electrons from a FAD prosthetic group to the Mo-factor. If NR is phosphorylated in the hinge 1 region, it can bind a 14-3-3 dimer, which in turn inactivates the enzyme. Removal of the 14-3-3 dimer or dephosphorylation of NR by a phosphatase results in reactivation of the enzyme. Exposure of plant to light, high  $\text{CO}_2$  and oxygen levels leads to most of the NR being active while darkness, low  $\text{CO}_2$  levels and hypoxia result in most of the NR being inactive.

### Nitrate assimilatory mutants of Arabidopsis

In the  $\text{NO}_3^-$  assimilation field, Arabidopsis was the first plant used for mutant isolation. In the early seventies, a graduate student Fietje Braaksma isolated chlorate resistant mutants (chl) of Arabidopsis in the lab of W.J. Feenstra (Braaksma and Feenstra, 1973). Chlorate is the chlorine analog of  $\text{NO}_3^-$ , which, when taken up and reduced to chlorite, is toxic to plants. Chlorate resistant mutants had been identified in fungi and were shown to be defective in  $\text{NO}_3^-$  uptake, reduction and regulation (Cove, 1979; Marzluf, 1981; Crawford and Arst, 1993), but no such mutants had been described in plants. In her initial report, Braaksma described three mutants, two of which were allelic. B1 and B3 (now called *chl1* or *nrt1.1*) had normal levels of NR activity but reduced levels of chlorine accumulation (a measure of  $\text{ClO}_3^-$  uptake). B2-1 (now called *chl2*) had much reduced NR activity. These data uncovered two genes, one involved in  $\text{NO}_3^-$  reduction and another in chlorate uptake. After Braaksma and Feenstra's initial

publication, additional chl mutants were described in Arabidopsis (Braaksma and Feenstra, 1982; Braaksma and Feenstra, 1982), in cell lines of Nicotiana (Muller and Grafe, 1978) and in barley plants (Kleinhofs et al., 1978). As these and other mutants were characterised in angiosperms, they were all found to fall into two classes: those having mutations in NIA (R) genes and those having mutations in CNX genes needed for the synthesis of the Mo-cofactor (Crawford, 1992; Warner and Kleinhofs, 1992; Crawford and Arst, 1993; Hoff et al., 1994; Mendel, 1997). Both classes have reduced NR activity. No mutations were found in  $\text{NO}_3^-$  uptake or regulatory genes except for *chl1*.

Further work on the molecular genetics of  $\text{NO}_3^-$  reduction in Arabidopsis revealed that it had two NIA genes and several CNX genes (Crawford, 1992). The NIA2 gene, which encodes for most of the NR activity in the plant (85-90%), corresponds to the *chl3* locus at position 56 in the middle of chromosome 1 (Crawford et al., 1988; Wilkinson and Crawford, 1991). The NIA1 gene encodes for 10-15% of the NR activity and is located near the end of chromosome 1 at position 116 near the GL2 gene (Cheng et al., 1988; Wilkinson and Crawford, 1993). Both NIA genes are regulated by  $\text{NO}_3^-$  and circadian rhythms and are expressed in leaves and roots (Cheng et al., 1988; Crawford et al., 1988; Cheng et al., 1991; Pilgrim et al., 1993). Such duplication of genes is found over much of the Arabidopsis genome (Blanc et al., 2000), and the NIA genes provided an early example of this. It is interesting that even though there is genetic redundancy, one can still isolate chlorate resistant mutants in which only the NIA2 gene is inactivated. This is because the two genes have diverged enough so that NIA2 accounts for most of the NR activity. NIA2 (*chl3*) mutants still show some chlorate sensitivity due to NIA1, but they are much less sensitive than wildtype plants. The residual NR activity (10-15% wildtype) from the NIA1 gene is sufficient for normal growth on  $\text{NO}_3^-$  as the sole nitrogen source (Crawford, 1995). For such a key enzyme, this result was surprising but was found to be true in other plants as well (reviewed in (Crawford, 1995)). When it was discovered that NR is post-translationally regulated, an explanation for this puzzling result arose. In wildtype tobacco plants, the NR is not 100% active when measured in the presence of magnesium but oscillates between 20-90% of full potential activity, which is measured in the presence of EDTA (Scheible et al., 1997a). In NR mutants, the extent of inactivation is reduced so that the reduced NR levels in the mutant are compensated by less inactivation of NR (Scheible et al., 1997a). It is not until NR activity levels drop below 10% of wildtype levels that one observes nitrogen deficiencies in the mutants. In Arabidopsis, a *nia1nia2* double mutant has been described with a deletion of the NIA2 gene and a point mutation in the NIA1 gene that has only 1% of wildtype level of activi-



**Figure 3. Schematic diagram showing the Moco synthesis pathway.** Picture taken from (Mendel, 1997).

ty (Wilkinson and Crawford, 1993). This mutant has grows poorly on  $\text{NO}_3^-$  alone but does show some growth.

The other class of NR-deficient mutants is *cnx*. Four *cnx* mutants have been identified in Arabidopsis: *chl2*, *chl4* (B25), *chl6* (B73) and *chl7* (Crawford, 1992). An interesting feature of these mutants is that three of them (*chl2*, *chl6* and *chl7*) are unable to grow on 0.1 mM tungstate, a useful tool for testing the loss of Mo-cofactor (LaBrie et al., 1992). Further work has lead to the identification of six genes required for synthesis of the Mo-cofactor in Arabidopsis (Mendel, 1997) (Figure 3). The early steps convert a guanosine nucleotide into an intermediate called Precursor Z. These steps are mediated by the products of the *CNX2* and *CNX3* genes (Hoff et al., 1995). Precursor Z is converted into molybdopterin by the *CNX5*, *CNX6* and *CNX7* products (Mendel, 1997). Molybdate is then added to molybdopterin by the two domain protein *CNX1*, which is homologous to the mammalian neuroprotein Gephyrin and binds the cell's cytoskeleton as does Gephyrin (Stallmeyer et al., 1995; Kuper et al., 2000; Schwarz et al., 2000). The *CNX1* gene is the same as *CHL6* (originally named B73) (Schwarz et al., 2000).

### The NRT1 family of nitrate transporters

Unlike  $\text{NO}_3^-$  reduction, the molecular genetics of  $\text{NO}_3^-$  uptake had been hampered by the fact that no uptake mutants were known in plants for sometime except for *chl1* in Arabidopsis (in the new nomenclature the gene is called *AtNRT1.1*). Ironically, this was the first chlorate-resistant mutant ever identified in plants, and mutations at this locus are found more frequently than in any other gene in chlorate selections. The original work on *chl1* indicated that it was strongly inhibited in chlorate uptake and partially defective in  $\text{NO}_3^-$  uptake (Doddema et al., 1978). The latter defect appeared to be a reduction in low affinity but not high affinity  $\text{NO}_3^-$  uptake activity (Doddema and Telkamp, 1979). Subsequent cloning of the gene from a T-DNA-tagged line revealed that it encoded a hydrophobic protein with twelve membrane spanning regions indicative of a cotransporter (Tsay et al., 1993a). Expression of the *CHL1* protein in *Xenopus* oocytes showed it was a  $\text{NO}_3^-$  transporter (Tsay et al., 1993a) with, surprisingly, biphasic kinetics (Huang et al., 1996; Liu et al., 1999). These experiments along with studies of  $\text{NO}_3^-$  uptake in whole plants demonstrated that *CHL1* is a dual affinity transporter contributing to both HATS and LATS (Huang et al., 1996; Touraine and Glass, 1997; Wang et al., 1998; Liu et al., 1999). *CHL1*'s contribution, however, depends on the environmental conditions used for plant growth. One key effector is  $\text{NH}_4^+$ . Overall  $\text{NO}_3^-$  uptake is reduced when plants are grown with  $\text{NH}_4\text{NO}_3$  as compared to  $\text{KNO}_3$ ; however, *CHL1*'s contribution is the greatest when plants are grown with  $\text{NH}_4^+$ ; that is, *chl1* mutants show the largest reduction in uptake when grown with  $\text{NH}_4^+$  than without (Huang et al., 1996; Touraine and Glass, 1997). Another critical parameter is pH. *CHL1* is more highly expressed at acidic pHs (Tsay et al., 1993a) and makes a greater contribution to cHATS at pH 5.5 than pH 7.0 (Wang et al., 1998). *CHL1* expression also shows diurnal variation, and the reduction in the dark can be reversed by supplying sucrose indicating that carbon signals affect *CHL1* (Lejay et al., 1999). Nitrogen metabolites, however, do not appear to repress *CHL1* as depriving Arabidopsis of nitrogen does not increase *CHL1* expression (Lejay et al., 1999).

Recent experiments have shown that *CHL1* expression is targeted to actively growing regions of roots and shoots (Guo et al., 2001). High levels of expression are found in root tips, lateral root primordia, young leaves and flower buds. Little expression over background is found in mature leaves, flowers and the apical meristem. In the mature parts of the root, intermediate expression is found in the stele and little in the outer layers of the root. *chl1* mutants also are defective in growth of nascent organs. Thus, the function of *CHL1* is not simply to take up nitrate from the

**Table 1. NRT1 family of nitrate transporters**

<b>NRT1 Gene</b>	<b>Organism</b>	<b>Substrate</b>	<b>Regulation<sup>a</sup></b>	<b>Identity<sup>b</sup></b>	<b>Accession#</b>
AtNRT1.1 (CHL1)	Arabidopsis	NO <sub>3</sub> <sup>-</sup>	Inducible	100%	L10357
AtNRT1.2 (NTL1)	Arabidopsis	NO <sub>3</sub> <sup>-</sup>	Constitutive	36%	AF073361
AtNRT1.3 (NTP3)	Arabidopsis	Unknown	Constitutive	49%	AJ131464
AtNRT1.4 (NTP2)	Arabidopsis	Unknown	Constitutive	54%	AJ011604
AtPTR2B	Arabidopsis	Peptides	N/A	39%	L39082
BnNRT1.2	Brassica napus	NO <sub>3</sub> <sup>-</sup> & basic AA	Inducible	91%	U17987

<sup>a</sup> Response to nitrate. <sup>b</sup> Percent identity to AtNRT1.1 (CHL1) at the amino acid level.  
N/A=not applicable

soil by the main body of the root but also to support the growth of nascent and actively growing organs in both roots and shoots.

After cloning CHL1, four additional genes with similar sequences were found in Arabidopsis (Table 1). *AtNRT1.2* (originally named *NTL1*) was cloned by homology to *CHL1* and found to be 36% identical at the amino acid level (Huang et al., 1999). Further work showed that it is a low affinity NO<sub>3</sub><sup>-</sup> transporter ( $K_m$  of 6 mM in oocytes) that is constitutively expressed in root epidermal and root hair cells. This finding indicates that AtNRT1.2 is a component of cLATS. A second gene AtPTR2B encodes a peptide transporter (Rentsch et al., 1995; Song et al., 1996). In fact, almost all the *CHL1*-related genes which have been described from bacteria, yeast and mammals have been identified as transporters of small oligopeptides including di- and tri-peptides (Steiner et al., 1995; Graul and Sadee, 1997; Crawford and Glass, 1998). Why transporters with specificity to NO<sub>3</sub><sup>-</sup> should be related to peptide transporters is not clear. A most interesting protein, NRT1.2 from *Brassica napus*, transports histidine and NO<sub>3</sub><sup>-</sup> (Zhou et al., 1998), indicating that this protein retains substrate specificities found in all members of the family. This protein is also the most similar to CHL1 being 91% identical. The entire gene family has been named PRT for peptide transporter and POT for H<sup>+</sup>-dependent oligopeptide transporter. This family was tentatively placed in the Major Facilitator Superfamily (MFS) of transporters (Pao et al., 1998) but now is its own family (Saier, 2000). The PTR and POT names are misleading as they do not include the NO<sub>3</sub><sup>-</sup> transporters in the family. Two other genes in this family have been cloned from Arabidopsis and named NTP2 (now called AtNRT1.4) and NTP3 (now called AtNRT1.3) (Hatzfeld and Saito, 1999), but their functions are not known, and they do not appear to be nitrate-regulated (Glass et al., 2001).

### The NRT2 family of transporters

The second family of NO<sub>3</sub><sup>-</sup> transporter genes (*NRT2*) was first discovered in *Aspergillus*. A chlorate-resistant mutant called *crnA* was identified with a defect in NO<sub>3</sub><sup>-</sup> uptake (Brownlee and Arst, 1983). It was subsequently cloned and shown to be NO<sub>3</sub><sup>-</sup>-regulated and NH<sub>4</sub><sup>+</sup>-repressed and to encode a transport protein (Unkles et al., 1991). Subsequently, two related genes (NAR-3 and NAR-4) were isolated from *Chlamydomonas* (Quesada et al., 1994). NAR-3 (CrNrt2.1) was shown to encode a high affinity, bi-specific NO<sub>3</sub><sup>-</sup>/NO<sub>2</sub><sup>-</sup> transporter while NAR-4 (CrNrt2.2) encodes a high affinity NO<sub>3</sub><sup>-</sup>-specific transporter (Galvan and Fernandez, 2001). The NAR-3 protein is about 30% identical to CRNA. Independently, a barley cDNA clone (BCH1) was identified by PCR amplification using degenerate primers corresponding to two conserved motifs found in a small group of MFS transporters that included CRNA (Trueman et al., 1996; Trueman et al., 1996). BCH1 (now called HvNRT2.1) is 32% identical to CRNA and 50% identical to CrNRT2.1. Subsequently, NRT2 genes have been identified in six plant species as well as *Chlamydomonas*, *Aspergillus*, *Hansenula* and *E. coli* (reviewed by Crawford and Glass, 1998; Daniel-Vedele et al., 1998; Forde, 2000; Galvan and Fernandez, 2001; Glass et al., 2001; Williams and Miller, 2001). These genes are found in the NO<sub>3</sub><sup>-</sup>/NO<sub>2</sub><sup>-</sup> porter (NNP) family within the major facilitator superfamily (Saier, 2000).

In Arabidopsis a total of 7 *NRT2* genes have been found (Table 2) (Forde, 2000; Glass et al., 2001; Okamoto et al., 2001). The first two identified were *AtNRT2.1* and *AtNRT2.2* (Filleur and Daniel-Vedele, 1999; Zhuo et al., 1999) and Genbank Accession # AF019748 and Z97058). These genes are about 1 kb apart in a tail to tail orientation on chromosome 1 at position 10-12 cM, not far from *CHL1*. *NRT2.1* shows strong regulated expression in roots (Filleur and Daniel-Vedele, 1999; Lejay et al., 1999; Zhuo et al., 1999; Cerezo et al., 2001) while little mRNA is detect-

**Table 2. Arabidopsis NRT2 family of nitrate transporters**

<b>NRT2 Gene</b>	<b>Chromosome</b>	<b>Substrate</b>	<b>Regulation<sup>a</sup></b>	<b>Identity<sup>b</sup></b>	<b>Accession#</b>
AtNRT2.1	1	NO <sub>3</sub> <sup>-</sup>	Inducible	100%	AF019748
AtNRT2.2	1	Unknown	Inducible	87%	AF019749
AtNRT2.3	5	Unknown	Transient Rep.	68%	AB015472
AtNRT2.4	5	Unknown	Constitutive	82%	AB015472
AtNRT2.5	5	Unknown	Constitutive	44%	AL163792
AtNRT2.6	3	Unknown	Constitutive	68%	AL353992
AtNRT2.7	1	Unknown	Repressed	56%	AC012187

<sup>a</sup> Response to nitrate. <sup>b</sup> Percent identity to AtNRT2.1 at amino acid level.

Data for this table was a kind gift of Mamoru Okamoto and Anthony Glass (Okamoto et al., 2001).

ed for *NRT2.2* (Zhuo et al., 1999). *NRT2.1* is induced by NO<sub>3</sub><sup>-</sup> or by nitrogen starvation and is repressed by NH<sub>4</sub><sup>+</sup>, amino acids (especially arg) and by growth on high levels of NO<sub>3</sub><sup>-</sup>. These properties are all characteristic of the inducible HATS system, which is NO<sub>3</sub><sup>-</sup>-inducible and feedback repressed by nitrogen metabolites or high NO<sub>3</sub><sup>-</sup> levels. NRT2 has been proposed to be a component of the iHATS in plants (reviewed by (Forde, 2000)).

Direct demonstration that *NRT2.1* is component of iHATS has come from mutant and heterologous expression studies. Arabidopsis transgenic lines containing antisense constructs of *AtNRT2.1* (Forde, 2000) and an Arabidopsis deletion mutant removing all of *NRT2.1* and part of *NRT2.2* (Filleur et al., 2001) have been constructed and shown to be defective in HATS but not LATS. Furthermore, the NRT2.1/2.2 mutant is defective in iHATS activities (i.e. in NO<sub>3</sub><sup>-</sup>-inducible, starvation-derepressible and NH<sub>4</sub><sup>+</sup>-repressible high affinity uptake) (Cerezo et al., 2001). The CRNA protein has been functionally expressed in *Xenopus* oocytes and found to have two transport mechanisms: a H<sup>+</sup>-coupled mechanism with a  $K_m$  for NO<sub>3</sub><sup>-</sup> of 24 μM and a passive mechanism with a  $K_m$  for NO<sub>3</sub><sup>-</sup> of 126 μM (Zhou et al., 2000b). It was suggested that under conditions of high external NO<sub>3</sub><sup>-</sup> concentrations the passive mechanism would provide an energetic advantage to the fungus by enabling it to absorb NO<sub>3</sub><sup>-</sup> without the energetically expensive coupling to H<sup>+</sup> fluxes. Whether plant NRT2 transporters function in similar ways is still unknown. The *Chlamydomonas* NRT2.1 protein expresses high affinity NO<sub>3</sub><sup>-</sup> transport activity in *Xenopus* oocytes but only if co-expressed the a second protein (NAR2), indicating that this system requires two components (Zhou et al., 2000a).

### Overview of nitrate transporter regulation and function

Additional aspects of overall transporter regulation have emerged from Arabidopsis. Both *CHL1* and *NRT2.1* display diurnal rhythms having low levels of mRNA at night (Lejay et al., 1999). The low levels at night can be enhanced by the addition of sucrose. These two genes are also NO<sub>3</sub><sup>-</sup>-responsive but *CHL1* does not display the feedback regulation that *NRT2.1* does as it is not repressed by NH<sub>4</sub><sup>+</sup> treatment nor induced by nitrogen starvation (Filleur and Daniel-Vedele, 1999; Lejay et al., 1999). *CHL1* does respond to pH as its mRNA levels are enhanced at more acidic pH (Tsai et al., 1993a).

With all this information, it has become possible to associate a given gene with a NO<sub>3</sub><sup>-</sup> uptake system. The two genes for which we have the most physiological data are *CHL1* and *NRT2.1*. Present data indicate that *NRT2.1* is a major component of iHATS that would be especially significant when plants are provided low levels or are deprived of nitrogen. The role of *CHL1* is more complicated. The expression patterns of this gene and the properties of the *CHL1* protein suggest that it could contribute to all three uptake systems in that it is expressed in the absence of NO<sub>3</sub><sup>-</sup>, its expression is enhanced by NO<sub>3</sub><sup>-</sup> and it has dual affinity uptake kinetics. Measurements of uptake in whole plants have indicated that *CHL1* does contribute to cHATS and LATS, but its role in iHATS is controversial (Wang et al., 1998; Liu et al., 1999; Cerezo et al., 2001; Glass et al., 2001). These experiments have also indicated that *CHL1*'s contribution depends heavily on the presence of NH<sub>4</sub><sup>+</sup>, which would also repress *NRT2.1* expression. In the presence of NH<sub>4</sub><sup>+</sup>, *chl1* mutants have significantly reduced

$\text{NO}_3^-$  uptake activity compared with wildtype plants for both *CHATS* and *LATS*. This is true for plants grown with or without  $\text{NO}_3^-$ . Thus, unlike *NRT2.1*, it is difficult to assign *CHL1* to one particular system. It is also appears that *CHATS*, *iHATS* and *LATS* are not static but dynamic systems whose composition changes depending on the environmental conditions in which it is grown and on the developmental stage of the plant. Other *Arabidopsis* transporters that are known to contribute to the  $\text{NO}_3^-$  uptake systems are *NRT2.1*, a component of *iHATS* (Filleur and Daniel-Vedele, 1999; Cerezo et al., 2001; Filleur et al., 2001), *NRT1.2*, whose properties indicate it is a constitutive component of *LATS* (Huang et al., 1999), and *CHL8*, whose gene has not been cloned but which contributes to *CHATS* (Wang and Crawford, 1996).

### Nitrate regulation of gene expression

As can be seen from the above discussion,  $\text{NO}_3^-$  is not only a substrate for the  $\text{NO}_3^-$  assimilatory pathway, but it is also a key signal. One of the most studied aspects of  $\text{NO}_3^-$  signalling is induction of gene expression (reviewed by (Redinbaugh and Campbell, 1991; Crawford, 1995; Koch, 1997; Stitt, 1999; Tischner, 2000; Wang et al., 2000; Coruzzi and Bush, 2001)). Many genes involved in different aspects of nitrogen and carbon metabolism are induced by  $\text{NO}_3^-$ . This effect can be observed by treating plants that are either starved for nitrogen or grown with a reduced source of nitrogen such as  $\text{NH}_4^+$  or amino acids. Genes that respond to  $\text{NO}_3^-$  have been grouped based on their metabolic niche as follows:  $\text{NO}_3^-$  assimilation,  $\text{NH}_4^+$  assimilation, ferredoxin reduction, oxidative pentose phosphate pathway and organic acid synthesis. Nitrate also mobilises carbon into organic acids by suppressing starch synthesis.

The primary  $\text{NO}_3^-$  assimilatory genes (*NIA*, *NII* and *NRT*) all respond to  $\text{NO}_3^-$  (Tang and Wu, 1957; Cheng et al., 1986; Crawford et al., 1986; Melzer et al., 1989; Cheng et al., 1991; Gowri et al., 1992; Aslam et al., 1993; Tischner et al., 1993; Koch, 1997; Stitt, 1999; Tischner, 2000). The *NIA* genes have been the most studied and respond within minutes to  $\text{NO}_3^-$  at concentrations as low as 10  $\mu\text{M}$ . In addition to *NII*, there are a group of responsive genes that contribute to nitrite reduction by providing reductant (encoding ferredoxin (Fd), ferredoxin NADP+ oxidoreductase (FNR), 6-phosphogluconate dehydrogenase (6PGDH) and 6-phosphogluconate dehydrogenase (6PGDH) and S-adenosyl-L-methionine-dependent uroporphyrinogen III methyltransferase (UPM1), which makes a prosthetic group for the NiR enzyme) (reviewed by (Stitt, 1999; Wang et al., 2000)). Genes involved in  $\text{NH}_4^+$  assimilation, encod-

ing specific isoforms of glutamine synthetase and glutamate synthase, are also induced (reviewed by (Lam et al., 1996; Koch, 1997; Stitt, 1999)). Responsive genes involved in starch and organic acid metabolism encode phosphoenolpyruvate carboxylase (PEPC, involved in organic acid metabolism), which is induced, and ADP-glucose pyrophosphorylase (*agpS2*, involved in starch synthesis), which is repressed by  $\text{NO}_3^-$  (Scheible et al., 1997b). Other genes that encode organic acid metabolic enzymes—cytosolic pyruvate kinase (PKc), citrate synthase (CS), and NADP-isocitrate dehydrogenase (*ICDH1*)—respond to  $\text{NO}_3^-$  (Scheible et al., 1997bb).

A microarray analysis of  $\text{NO}_3^-$ -responsive genes has been reported for *Arabidopsis* (Wang et al., 2000). Using glass slides with approximately 5500 *Arabidopsis* genes/clones, mRNA from plants treated with low  $\text{NO}_3^-$  (250  $\mu\text{M}$ ) for short times (20 min) and with high  $\text{NO}_3^-$  (5–10 mM) for longer times (2 hrs) was analysed to detect both rapid response genes and responsive genes in general. Besides detecting many of the genes described above, new responsive genes were identified. Two genes of the nonoxidative branch of the pentose phosphate pathway—transketolase and transaldolase—were induced. Other metabolic genes that were induced included malate dehydrogenase, asparagine synthetase, histidine decarboxylase and two methyltransferases. Several other very interesting genes were induced including a myb transcription factor gene, a tonoplast high capacity calcium antiporter gene *CAX1*, the senescence-associated gene *SAG21* and the nonsymbiotic hemoglobin gene *AHB1*. Several genes were found to be repressed by  $\text{NO}_3^-$  including the  $\text{NH}_4^+$  transporter gene *AMT1;1*. The general conclusions that came from this and other studies are the following. First, the most responsive genes are those that are involved directly or indirectly with nitrite reduction (encoding NiR, Fd, FNR, UMP1, G6PDH, 6PGDH, transketolase and transaldolase). Nitrite is toxic to plants so that it makes sense that the genes critical for nitrite reduction would be rapidly induced when  $\text{NO}_3^-$  is present. Second, different genes show different response patterns indicating that multiple signalling and regulatory events are occurring as plants take up and assimilate  $\text{NO}_3^-$ . For example, some genes show reduced expression after 2 hrs in low  $\text{NO}_3^-$  while others do not. Key signals that may be dictating expression levels are  $\text{NO}_3^-$ , reduced nitrogen and carbon metabolites. As  $\text{NO}_3^-$  gets depleted by assimilation or is accumulated in the vacuole, some genes will sense the drop in  $\text{NO}_3^-$  and begin to diminish their expression. In addition, the products of  $\text{NO}_3^-$  assimilation including  $\text{NH}_4^+$  and amino acids, especially glutamine and glutamate, are also thought to feedback repress genes in the pathway (Hoff et al., 1994; Dzuibany et al., 1998; Tischner, 2000). Carbon metabolites sucrose (Cheng et al., 1992) and

malate (Muller et al., 2001) also have been shown to serve as signals.

With such elaborate signalling, it is clear that regulatory networks exist to control  $\text{NO}_3^-$  assimilation and integrate it with nitrogen and carbon metabolism. Our understanding of these networks is in its infancy. The discussion below is restricted to  $\text{NO}_3^-$  signalling. Several promoter elements have been identified that are needed for  $\text{NO}_3^-$  induction of NiR (Rastogi et al., 1993; Rastogi et al., 1997) and NR (Lin et al., 1994; Hwang et al., 1997) genes in tobacco (reviewed by Rothstein and Sivasankar, 1999). With the use of inhibitors, it has been shown that  $\text{NO}_3^-$  induction is a primary response (i.e. not blocked by protein synthesis inhibitors (Gowri et al., 1992; Redinbaugh and Campbell, 1993; Ritchie et al., 1994; Redinbaugh and Campbell, 1998)) and involves calcium (Sakakibara et al., 1996; Sakakibara et al., 1997; Sueyoshi et al., 1999) as well as kinases and phosphatases (Champigny and Foyer, 1992; Sakakibara et al., 1997; Sueyoshi et al., 1999). There have been two reported attempts to find  $\text{NO}_3^-$  regulatory mutants in Arabidopsis. The first selected for mutants that were resistant to chlorate after a transient treatment with low  $\text{NO}_3^-$  (Lin and Cheng, 1997). A mutant (cr88) was found that was defective in light reception and was designated as a hy mutant (Lin and Cheng, 1997; Cao et al., 2000). The other was a screen using transgenic plants containing the tobacco NiR promoter fused to GUS (Leydecker et al., 2000). This screen produced *cnx* mutants that overproduced NiR mRNA in the absence of  $\text{NO}_3^-$ . Thus, no regulatory mutants defective in  $\text{NO}_3^-$  induction have been reported. However, a  $\text{NO}_3^-$ -regulated gene (*ANR1*) has been identified which encodes a MADS box transcription factor that is involved in nitrate regulation of root development. This gene is discussed in more detail below (Zhang and Forde, 1998).

## AMMONIUM UPTAKE

Arabidopsis, like many plants, does not grow well with  $\text{NH}_4^+$  as its sole source of N (e.g. (Wilkinson and Crawford, 1993)), probably reflecting its adaptation for growth in aerobic soils, where  $\text{NO}_3^-$  is the main form of inorganic N and  $\text{NH}_4^+$  concentrations rarely exceed 50  $\mu\text{M}$  (Marschner, 1995). In culture, however, Arabidopsis grows best on a mixed N source ( $\text{NH}_4\text{NO}_3$ ) and under these conditions it may even absorb the  $\text{NH}_4^+$  preferentially (Gazzarrini et al., 1999).

The relevance of the  $\text{NH}_4^+$  uptake system extends beyond acquiring  $\text{NH}_4^+$  from the soil; significant leakage of  $\text{NH}_4^+$  occurs from root cells, even when  $\text{NO}_3^-$  is the N


source, so that an uptake system with a high affinity for  $\text{NH}_4^+$  is important for conserving valuable N supplies. As discussed in detail elsewhere (von Wiren et al., 2001),  $\text{NH}_4^+$  transport is also important for a variety of processes in the shoot, which include unloading of  $\text{NH}_4^+$  from the xylem, absorption of atmospheric  $\text{NH}_3$  and retrieval of photorespiratory  $\text{NH}_3$  released into the leaf apoplast. A number of recent reviews covering all aspects of  $\text{NH}_4^+$  uptake in plants are available (Glass et al., 1997; Forde and Clarkson, 1999; Howitt and Udvardi, 2000; von Wiren et al., 2000a; von Wiren et al., 2001).

## Kinetics and Energetics of $\text{NH}_4^+$ Uptake

In common with other plants, the kinetics of  $\text{NH}_4^+$  uptake in Arabidopsis are biphasic, being separable into a high-affinity system (HATS) and a low-affinity system (LATS) (Rawat et al., 1999). The LATS only becomes evident at external  $\text{NH}_4^+$  concentrations above 1 mM (Rawat et al., 1999). The high-affinity system displays typical Michaelis-Menten kinetics (Rawat et al., 1999; Shelden et al., 2001), while the LATS has linear (non-saturable) kinetics (Rawat et al., 1999), consistent with it being mediated by a passive transport system such as a channel (see below). The affinity of the HATS for its substrate, as estimated from  $^{13}\text{NH}_4^+$  labelling studies, falls within the range reported in other plants (Rawat et al., 1999). For seedlings maintained on  $\text{NH}_4\text{NO}_3$ , the estimated  $K_m$  values for  $\text{NH}_4^+$  were from 85 to 168  $\mu\text{M}$ , compared with values in the range 20-360  $\mu\text{M}$  for other plant species (von Wiren et al., 2001). Using an electrophysiological method, a similar  $K_m$  value of  $46 \pm 5$   $\mu\text{M}$  was obtained for  $\text{NH}_4\text{NO}_3$ -grown Arabidopsis seedlings, but seedlings that had been N-starved or grown on  $\text{NO}_3^-$  alone for 3 d gave  $K_m$  values of 7-8  $\mu\text{M}$  (Shelden et al., 2001).

It is now generally accepted that  $\text{NH}_4^+$ , rather than the unprotonated species  $\text{NH}_3$ , is the transported form (Forde and Clarkson, 1999; von Wiren et al., 2001). However the precise mechanism of  $\text{NH}_4^+$  influx across the plasma membrane is still not fully resolved. Based on typical estimates for cytosolic  $\text{NH}_4^+$  concentrations and the electrical potential across the plasma membrane, thermodynamic considerations imply that in the range of external  $\text{NH}_4^+$  concentrations that roots are exposed to in well-aerated soils (i.e.  $<50$   $\mu\text{M}$ ) the uptake of  $\text{NH}_4^+$  will require the input of energy (Forde and Clarkson, 1999). This could be achieved, for example, by a high-affinity  $\text{H}^+/\text{NH}_4^+$  co-transporter that coupled  $\text{NH}_4^+$  influx to the  $\text{H}^+$  gradient generated by the plasma membrane  $\text{H}^+$  pump. The importance of the  $\text{H}^+$  gradient in driving  $\text{NH}_4^+$  influx by the





**Figure 4. Possible mechanisms for high- and low-affinity  $\text{NH}_4^+$  uptake by Arabidopsis roots.** High-affinity  $\text{NH}_4^+$  influx is thought to be catalysed by members of the AMT transporter family (e.g. AtAMT1.1) and is dependent on the H<sup>+</sup> gradient generated by the plasma membrane H<sup>+</sup>-ATPase. The membrane proteins responsible for low-affinity  $\text{NH}_4^+$  uptake are still unidentified but are likely to include K<sup>+</sup> channels and transporters that do not discriminate against  $\text{NH}_4^+$ . Diffusion of the uncharged species  $\text{NH}_3$  across the plasma membrane is unlikely to contribute significantly to influx, but may be a major route for  $\text{NH}_3/\text{NH}_4^+$  efflux. See text for further discussion.

HATS in rice roots has been demonstrated using protonophores and ATPase inhibitors (Wang et al., 1993), but direct evidence for H<sup>+</sup> / $\text{NH}_4^+$  symport in plant roots is still lacking. In the bacterium *Corynebacterium glutamicum*, the amt  $\text{NH}_4^+$  transporter (which is homologous to higher plant  $\text{NH}_4^+$  transporters) has been proposed to operate via an  $\text{NH}_4^+$  uniport mechanism that is driven by the membrane potential (Siewe et al., 1996).

The net uptake of any ion is the result of the opposing processes of influx and efflux, and under some conditions it is known that  $\text{NH}_4^+$  efflux can have a significant influence on net  $\text{NH}_4^+$  uptake (Forde and Clarkson, 1999). Rates of  $\text{NH}_4^+$  efflux in Arabidopsis (Rawat et al., 1999) and in other plants (Wang et al., 1993; Kronzucker et al., 1995) are positively correlated with the cytosolic  $\text{NH}_4^+$  concentration. When Arabidopsis seedlings were grown on  $\text{NH}_4^+$ -rich medium (1 mM  $\text{NH}_4^+$ ) the rate of  $\text{NH}_4^+$  efflux was estimated to be as much as 35% of the influx rate (Rawat et al.,

1999). As discussed elsewhere (Forde and Clarkson, 1999), because the efflux of the protonated form of  $\text{NH}_4^+$  is likely to be thermodynamically uphill, it is thought that it is actually  $\text{NH}_3$  that diffuses passively from the cells and that this is then protonated in the external solution giving the false impression that the ionic form had crossed the plasma membrane (Figure 4).

#### Regulation of $\text{NH}_4^+$ Uptake: the Physiological Evidence

It has been established for a wide range of plant species that the HATS for  $\text{NH}_4^+$  (but not the LATS) is highly regulated according to the N nutrition of the plant (Forde and

Clarkson, 1999; von Wiren et al., 2001). A number of studies have now confirmed that the same is true in Arabidopsis (Gazzarrini et al., 1999; Rawat et al., 1999; Shelden et al., 2001). For example, transferring Arabidopsis plants from high  $\text{NH}_4\text{NO}_3$  concentrations to low led to a 12-fold increase in HATS activity (Rawat et al., 1999), and when N-depleted plants were resupplied with  $\text{NH}_4\text{NO}_3$ , the  $\text{NH}_4^+$  influx rate declined rapidly (Rawat et al., 1999). In the same experiments, the LATS activity was independent of the external  $\text{NH}_4^+$  supply and the plant's N status.

As already noted above, the kinetic properties of the HATS can also be affected by changes in N nutrition. In rice, a positive correlation was found between the external

$\text{NH}_4^+$  supply and the affinity of the HATS for  $\text{NH}_4^+$  (Wang et al., 1993). While Rawat et al., working with Arabidopsis, found no simple relationship between the ambient N supply and the affinity of the HATS for  $\text{NH}_4^+$  (Rawat et al., 1999), a more recent study reported that when Arabidopsis seedlings were grown for 3 d in the absence of  $\text{NH}_4^+$  (whether N-depleted or with  $\text{NO}_3^-$  as sole N source) the  $K_m$  for  $\text{NH}_4^+$  declined from 46  $\mu\text{M}$  to <10  $\mu\text{M}$  (Shelden et al., 2001). A possibly critical difference between the two studies is that in the former case the 'low N' plants were grown on 0.1 mM  $\text{NH}_4\text{NO}_3$  and so were not fully  $\text{NH}_4^+$ -depleted. Perhaps the very high-affinity HATS is synthesised or activated only when the external  $\text{NH}_4^+$  concentration is very low or zero.

**Table 3. AMT1 family of  $\text{NH}_4^+$  transporters**

AMT1 Gene	Tissue specificity <sup>a</sup>	N regulation <sup>b</sup>	$K_m$ for methyl $\text{NH}_4^{+c}$	$K_i$ for $\text{NH}_4^{+c}$	AGI code <sup>d</sup>
AtAMT1.1	R≥S [1,2,4] <sup>c</sup>	+++ [1,3,4]	65 $\mu\text{M}$ [2] 8 $\mu\text{M}$ [1] 32 $\mu\text{M}$ [4]	5-10 $\mu\text{M}$ [2] 0.5 $\mu\text{M}$ [1] 1-10 $\mu\text{M}$ [4]	At4g13510
AtAMT1.2	R>>S [1,4]	constitutive [1,4]	24 $\mu\text{M}$ [1] 36 & 3000 $\mu\text{M}$ [4]	~35 $\mu\text{M}$ [1] 1-10 $\mu\text{M}$ [4]	At1g64780
AtAMT1.3	R [1]	+ [1]	11 $\mu\text{M}$ [1]	~35 $\mu\text{M}$ [1]	At3g24300
AtAMT1.4	?	?	?	?	At4g28700
AtAMT1.5	?	?	?	?	At3g24290
AtAMT2	R<S [5]	++ (in root) [5]	not transported [5]	?	At2g38290

<sup>a</sup>Based on distribution of RNA transcripts. R = roots, S = shoots.

<sup>b</sup>N regulation of mRNA abundance.

<sup>c</sup>Substrate affinities determined from yeast expression studies.

<sup>d</sup>AGI locus name (see the MIPS *Arabidopsis thaliana* database at <http://mips.gsf.de/proj/thal/>)

<sup>e</sup>References: [1] Gazzarrini et al. (1999); [2] Ninnemann et al. (1994); [3] Rawat et al. (1999); [4] Shelden et al. (2001); [5] Sohlenkamp et al. (2000).


$\text{NH}_4^+$  influx in *Arabidopsis* roots is diurnally regulated, reaching a maximum at the end of the light period and declining markedly with the onset of darkness (Gazzarrini et al., 1999). This suggests that the C status of the plant may have a regulatory effect on HATS activity that is somehow transmitted from shoot to root.

### Structure and Biochemical Properties of the AMT Family of $\text{NH}_4^+$ Transporters

The *Arabidopsis* AtAMT1.1 gene was the first  $\text{NH}_4^+$  transporter gene to be cloned from higher plants (Ninnemann et al., 1994). Subsequently, two closely related cDNAs (AtAMT1.2 and AtAMT1.3) were isolated by homology and shown to be able, like AtAMT1.1, to complement an  $\text{NH}_4^+$  transport mutant of yeast (Gazzarrini et al., 1999). The completed *Arabidopsis* genome sequence reveals that there are two further members of the AMT1 family (AtAMT1.4 and AtAMT1.5) (Table 3), but no further data are available on these. A sixth gene (AtAMT2), which is more distantly related to the other AMT genes, has been identified and partially characterised (Sohlenkamp et al., 2000).

The AMT genes encode hydrophobic proteins of 475–514 amino acids and belong to a probably ubiquitous gene family (the AMT/MEP family), which has known members in bacteria, archaea, fungi, plants and animals (Saier et al., 1999). All functionally characterised members of the family catalyse  $\text{NH}_4^+$  uptake and most also transport the  $\text{NH}_4^+$  analogue, methylammonium. Figure 5 illustrates the phylogenetic relationships between the *Arabidopsis* AMT transporters and selected members of the wider AMT/MEP family. The AtAMT1 genes seem to fall into three subfamilies, one that includes AtAMT1.1, AtAMT1.3 and AtAMT1.5 (the AtAMT1.1 subfamily), and two others that have single members, AtAMT1.2 and AtAMT1.4. The tomato LeAMT1.1 and the *Lotus japonicus* LjAMT1.1 genes belong to the AtAMT1.1 subfamily, while the tomato LeAMT1.2 gene is most closely related to AtAMT1.2. The close relationship between AtAMT1.3 and AtAMT1.5, together with their occurrence adjacent to each other within a 6 kb region of chromosome 3, suggests that they represent a relatively recent gene duplication. The AtAMT2 gene is the most divergent of the *Arabidopsis* AMT sequences, being more closely related to the yeast and bacterial and yeast AMT/MEP genes than to the plant ones (Figure 5).


The tomato LeAMT1.3 transporter has an N-terminus that is 20 amino acids shorter than the other AMT1 polypeptides and two untranslated open reading frames are found in its 5'-untranslated region, something often



**Figure 5. Phylogenetic tree for the *Arabidopsis* AMT transporters and selected members of the AMT/MEP family.** The sequences were aligned using the CLUSTAL W program (Thompson et al., 1994) and the rooted tree constructed using DRAWGRAM (Felsenstein, 1993). The accession numbers for the AtAMT sequences are given in Table 3. The other sequences, with their accession numbers, are from *Saccharomyces cerevisiae* (ScMEP1, PIR:S46225; ScMEP2 PIR:S51089; ScMEP3 PIR:S69027), *Schizosaccharomyces pombe* (SpAMT1, PIR:T50244; SpAMT2, GB:CAB83006), *Caenorhabditis elegans* (CeAMT1, GB:P54145; CeAMT2, GB:Q20605; CeAMT3, GB:Q21565; CeAMT4, GB:AAA83579), rice (OsAMT1, GB:AAB58937), *Corynebacterium glutamicum* (CgAMT, GB:CAA63770), *Lotus japonicus* (LjAMT1.1, GB:AAG24944), tomato (LeAMT1.1, GB:X92854; LeAMT1.2, GB:CAA64475; LeAMT1.3, GB:AAG11397) and *Brassica napus* (BnAMT1.2, GB:AAG28780).

associated with regulatory genes (von Wiren et al., 2000b). Surprisingly, there appears to be no direct ortholog of LeAMT1.3 in the *Arabidopsis* genome.

Computer-based predictions that the *E. coli* AMTB protein has 12 transmembrane domains, with the N- and C-termini located on the interior face of the membrane, were recently confirmed using fusions with marker proteins



**Figure 6. Schematic model for the membrane topology of AtAMT1.1.** The model (above) is redrawn from (Thomas et al., 2000) and the predicted locations of the eleven transmembrane domains are indicated in red on the AtAMT1.1 amino acid sequence (below). The precise start and end points of these transmembrane domains in AtAMT1.1 were taken from the predictions in the Arabidopsis Membrane Protein Library (<http://www.cbs.umn.edu/arabidopsis>). A conserved aspartate residue suggested as a possible  $\text{NH}_4^+$  binding site (Thomas et al., 2000) is indicated on the topological model..

(Thomas et al., 2000). However, sequence alignments indicate that the most N-terminal of the 12 transmembrane domains in AMTB is not present in its plant or fungal homologues, so that an 11 transmembrane topology is predicted for the eukaryotic members of the AMT/MEP family (Thomas et al., 2000). The implication that the N-terminus of the eukaryotic AMT/MEP proteins must be on the outside of the membrane has been confirmed experimentally for the yeast MEP2 protein (Marini and Andre, 2000). Figure 6 shows a 2-D representation of the likely membrane topology of AtAMT1.1 transporter. According to this model, the majority of positively charged amino acids on the AtAMT1.1 polypeptide are on the cytoplasmic side of the membrane (Thomas et al., 2000), consistent with the 'positive-inside' rule (Von Heijne, 1992).

The kinetic properties of some of the Arabidopsis AMT transporters have been studied in the yeast heterologous expression system (Ninnemann et al., 1994; Gazzarrini et al., 1999; Sohlenkamp et al., 2000; Shelden et al., 2001). The ability of most AMT proteins to transport methylam-

monium enables ( $^{14}\text{C}$ )methylammonium to be conveniently used to assay transport activity. The pH optimum for methylammonium uptake by yeast cells expressing AtAMT1.1 was 7, with only low rates of uptake at pH 9, consistent with  $\text{NH}_4^+$  being the substrate rather than  $\text{NH}_3$  (Ninnemann et al., 1994). Transport activity was dependent on an energy source (glucose) and was sensitive to protonophores (which collapse the  $\text{H}^+$  gradient across the plasma membrane) and to inhibitors of the  $\text{H}^+$  pump (Ninnemann et al., 1994). AtAMT1.1 displayed Michaelis-Menten kinetics for methylammonium uptake, with a  $K_m$  of 65  $\mu\text{M}$ . As expected,  $\text{NH}_4^+$  acted as a strong competitive inhibitor of methylammonium uptake ( $K_i$  of <10  $\mu\text{M}$ ), but AtAMT1.1 showed no affinity for  $\text{K}^+$  (Ninnemann et al., 1994). Thus AtAMT1.1 was demonstrated to be an energy-dependent high-affinity  $\text{NH}_4^+$  uptake system.

Subsequent studies have used the same yeast expression system to examine the kinetic properties of AtAMT1.1, AtAMT1.2, AtAMT1.3 and AtAMT2 (see Table 3 for a summary). Gazzarrini and colleagues reported that of

the three AtAMT1 proteins, AtAMT1.1 had the highest affinity for  $\text{NH}_4^+$  ( $K_m$  0.5  $\mu\text{M}$ ) (Gazzarrini et al., 1999). In contrast, a more recent study (Shelden et al., 2001) found no significant difference between AtAMT1.1 and AtAMT1.2 in their affinity for  $\text{NH}_4^+$  and reported that AtAMT1.2 showed biphasic kinetics for methylammonium uptake, with  $K_m$  values of 36  $\mu\text{M}$  and 3  $\mu\text{M}$ .

AtAMT2 expressed in yeast was unable to transport methylammonium, making analysis of its biochemical properties difficult. However, it was established that yeast cells expressing AMT2 were able to use  $\text{NH}_4^+$  as a N source and to deplete  $\text{NH}_4^+$  from medium containing concentrations <100  $\mu\text{M}$ , although at a slower rate than cells expressing AtAMT1.1 (Sohlenkamp et al., 2000).

### Regulation of the AMT Genes

In addition to their differing biochemical properties, the AMT genes also show contrasting patterns of mRNA expression (Table 1). All four AMT genes so far analysed are expressed in roots (Ninnemann et al., 1994; Gazzarrini et al., 1999; Rawat et al., 1999; Sohlenkamp et al., 2000; Shelden et al., 2001), but only AMT1.1 and AMT2 are also expressed strongly in shoots (Gazzarrini et al., 1999; Sohlenkamp et al., 2000; Shelden et al., 2001). In roots, AtAMT1.1 and AtAMT2 are highly responsive to the  $\text{NH}_4^+$  supply, both being strongly repressed in the presence of high external  $\text{NH}_4^+$  concentrations (Rawat et al., 1997; Gazzarrini et al., 1999) (Sohlenkamp et al., 2000; Shelden et al., 2001). AtAMT1.3 is similarly regulated, but to a lesser degree, while AtAMT1.2 is constitutively expressed (Gazzarrini et al., 1999; Shelden et al., 2001).

Of the three N-responsive genes, it is AtAMT1.1 that responds to the N supply in a manner that most closely parallels the changes in HATS activity (Gazzarrini et al., 1999; Rawat et al., 1999; Shelden et al., 2001). For example, when plants were transferred from high  $\text{NH}_4^+$  concentrations to low, AtAMT1 mRNA levels began to increase within 1 h and continued to increase over a 24 h period concurrently with an increase in  $^{13}\text{NH}_4^+$  influx rates (Rawat et al., 1999). Similarly AtAMT1.1 mRNA levels in roots increased markedly during a 2 d period after the N supply was withdrawn, as did the  $^{15}\text{NH}_4^+$  influx rate, but there was no significant increase in AtAMT1.3 mRNA abundance during the same period (Gazzarrini et al., 1999). Nevertheless, these correlations need to be treated with caution since there are indications that post-transcription-

al as well as transcriptional controls can operate in the regulation of  $\text{NH}_4^+$  transporter activity (see below).

There is evidence that glutamine is a key regulatory metabolite involved in the feedback regulation of AtAMT1.1 expression (Rawat et al., 1999). Strong negative correlations were found between glutamine concentrations in root extracts and both  $^{13}\text{NH}_4^+$  influx rates and AtAMT1.1 transcript levels. Furthermore, the repressive effect of  $\text{NH}_4^+$  on AtAMT1.1 expression could be relieved by blocking the conversion of  $\text{NH}_4^+$  to glutamine with methionine sulfoximine (MSX, an inhibitor of glutamine synthetase activity).

### Are Post-Transcriptional Controls Involved in the Regulation of AMT Transporters?

There are several lines of evidence pointing to the possibility that regulation of the  $\text{NH}_4^+$  transporters is complex and operates at multiple levels. Gazzarrini and colleagues found that when Arabidopsis plants were transferred from  $\text{NH}_4\text{NO}_3$  to  $\text{KNO}_3$ , the  $\text{NH}_4^+$  influx rate increased over the following three days without any change in AtAMT1.1, AtAMT1.2 or AtAMT1.3 mRNA levels (Gazzarrini et al., 1999). Under similar conditions it was reported that the  $K_m$  for  $\text{NH}_4^+$  of the HATS changed from 46  $\mu\text{M}$  to 7  $\mu\text{M}$  (Shelden et al., 2001). Only two explanations for these observations seem possible: either one of the other AMT genes is responsible for these changes in the activity and kinetic properties of HATS (and is regulated by the N supply in a very different way from AtAMT1.1, AtAMT1.2 or AtAMT1.3), or regulation of AMT expression at some post-transcriptional level is occurring.

In experiments where MSX was used to block  $\text{NH}_4^+$  assimilation and  $\text{NH}_4^+$  consequently failed to cause a significant decline in AtAMT1.1 transcript levels, there was nevertheless a 30% decrease in  $^{13}\text{NH}_4^+$  influx rates (Rawat et al., 1999). Further evidence for post-transcriptional controls comes from experiments where the resupply of  $\text{NH}_4\text{NO}_3$  to N-starved plants consistently led to a more rapid decline in  $\text{NH}_4^+$  influx rates than in AtAMT1.1 mRNA abundance (Rawat et al., 1997), even though AtAMT1.1 is the most highly responsive of the AMT gene family (Gazzarrini et al., 1999); (Rawat et al., 1999; Sohlenkamp et al., 2000; Shelden et al., 2001).

All of these observations are consistent with a role for the  $\text{NH}_4^+$  ion in the post-transcriptional regulation of the HATS (Rawat et al., 1999), perhaps by directly inhibiting AMT transporter activity or by inhibiting the synthesis or post-translational processing of the AMT proteins. A

model for the feedback regulation of AtAMT1.1 by glutamine and  $\text{NH}_4^+$  is shown in Figure 7.

### Concluding Remarks on AMT Transporter Function

The current evidence strongly suggests that one or more of the AtAMT genes encode the HATS for  $\text{NH}_4^+$  in Arabidopsis roots. Their differing affinities for  $\text{NH}_4^+$  and responsiveness to N, and the varied patterns of expression within the plant, suggest that they perform different, but perhaps overlapping, physiological functions. The high-affinity of AtAMT1.1 for  $\text{NH}_4^+$  and the close correlation between AtAMT1.1 mRNA levels and the HATS activity has made this member of the AMT family the strongest candidate for being a major component of the high-affinity  $\text{NH}_4^+$  uptake system, particularly in  $\text{NO}_3^-$ -fed and N-deprived plants. The very high affinity of the AtAMT1.1 transporter expressed in yeast (Table 3), and its strong derepression in N-deprived roots, suggests that it accounts for the highest affinity component of the HATS ( $K_m$  of 8  $\mu\text{M}$ ) that has only been detected in fully  $\text{NH}_4^+$ -deprived roots (Shelden et al., 2001).


Nevertheless, since AtAMT1.2, AtAMT1.3 and AtAMT2 are also expressed in roots, and since each has been shown to have the capacity to transport  $\text{NH}_4^+$  in the con-

centration range typical of the HATS (Table 3), it is likely that they too contribute to the HATS. The extent to which each contributes seems certain to vary with such factors as the rate and nature of the external N supply, the time of day and the N and C status of the plant. Examination of the TIGR Gene Index for Arabidopsis ESTs (Quackenbush et al., 2001) reveals that the Columbia root EST collection contains two cDNAs for AtAMT1.1, four for AtAMT1.2 and one for AtAMT1.4, which may give some indication of the relative levels of expression of the AMT genes in  $\text{NH}_4\text{NO}_3^-$  grown plants of this ecotype.

The physiological roles of the different AtAMT transporters will become clearer once we have more information about their subcellular localizations and their cellular distributions within the root and shoot. Most important will be the analysis of mutants and transgenic lines in which the expression of single and multiple AtAMT genes is modified.

### Low-affinity $\text{NH}_4^+$ uptake

The non-saturable kinetics of  $\text{NH}_4^+$  influx at external  $\text{NH}_4^+$  concentrations above 1 mM suggest that it is likely to be occurring by diffusion, but it is only at high external pH that the diffusion of  $\text{NH}_3$  is likely to contribute much to the low-



**Figure 7. Model for the feedback regulation of the AtAMT1.1 gene by N metabolites.** While glutamine may repress AtAMT1.1 expression at the transcriptional level,  $\text{NH}_4^+$  may act post-transcriptionally (e.g. on mRNA translation or targeting of the protein) or may have direct effects on the activity or kinetic properties of the AtAMT1.1 transporter. Model based on that proposed by (Rawat et al., 1999).

affinity phase. In yeast, one member of the AMT/MEP family (Mep3p) has a very low affinity for  $\text{NH}_4^+$  (1.4–2.1 mM), so that within this one family a full range of substrate affinities is possible. The only indication so far that a member of the AtAMT family may contribute to the LATS is the report that AtAMT1.2 has a dual affinity for methylammonium when expressed in yeast, the low affinity phase having a  $K_m$  of 3 mM (Shelden et al., 2001).

Low-affinity  $\text{NH}_4^+$  uptake might also occur as a by-product of the activity of  $\text{K}^+$  channels or  $\text{K}^+$  transporters (Forde and Clarkson, 1999; von Wieren et al., 2001).  $\text{K}^+$  uptake systems in maize roots included both  $\text{NH}_4^+$ -sensitive and  $\text{NH}_4^+$ -insensitive components (Vale et al., 1987). Analysis of a mutant carrying an insertion in the gene for an inward-rectifying  $\text{K}^+$  channel (AKT1) has confirmed that a similar situation exists in Arabidopsis (Spalding et al., 1999). Furthermore, it was shown that the  $\text{NH}_4^+$ -sensitive component corresponded to the non-AKT1 fraction of  $\text{K}^+$  transport activity (Spalding et al., 1999). One class of  $\text{K}^+$  channels that could contribute to low-affinity  $\text{NH}_4^+$  influx are the voltage-independent non-selective channels (VICs; (Amtmann and Sanders, 1999)). Evidence that these are permeable to  $\text{NH}_4^+$  has come from lipid bilayer studies on a voltage-independent  $\text{K}^+$  channel from rye roots (White, 1996) and from in situ analysis of VICs in the plasma membrane of maize root cortical cells (A. Amtmann, unpublished results). Which genes encode VICs are still unknown. Figure 4 summarises the possible pathways for high and low-affinity  $\text{NH}_4^+$  transport across the plasma membrane.

### Ammonium sensing


In yeast, the  $\text{NH}_4^+$  transporter Mep2p acts as a sensor for environmental  $\text{NH}_4^+$  in regulating the initiation of the process of pseudohyphal differentiation (Lorenz and Heitman, 1998). Thus Mep2p belongs to a class of membrane transporters in yeast that also serve as sensors for metabolites; other examples include the Snf3p and Rgt2p glucose transporters (Ozcan et al., 1996) and the Ssy1p and Ptr3p amino acid transporters (Klasson et al., 1999), which each sense the extracellular levels of their respective substrates. Often transporters that function as nutrient sensors have extended N- or C-terminal domains that may serve as signalling domains. None of the AMT sequences possess any additional domains that might suggest a role other than in solute transport, but since this also true of Mep2p it is still a possibility worthy of investigation that one or more of the AMT permeases function as  $\text{NH}_4^+$  sensors.

## NITROGEN REGULATION OF PLANT DEVELOPMENT


It is well known that the N supply can have major effects on many aspects of plant development (Redinbaugh and Campbell, 1991; Crawford, 1995; Bloom, 1997; Stitt, 1999). Root developmental processes often reported to be N-regulated include root branching, root diameter, root hair density and/or length, and (in legumes) numbers of N-fixing root nodules (Forde and Lorenzo, 2001). Above ground processes that in many species are subject to N regulation include leaf growth, shoot branching and flowering time (Stitt, 1999). The term ‘trophomorphogenesis’ has been proposed to describe the changes in plant morphology arising from variations in the availability or distribution of nutrients in the environment (Forde and Lorenzo, 2001). Trophomorphogenesis, like photomorphogenesis, is an example of developmental plasticity, which is a characteristic feature of plant development. Arabidopsis, like other fast-growing plant species, displays a high degree of developmental plasticity in response to environmental signals and is therefore an excellent subject in which to study the molecular basis of these morphogenetic responses, a subject that has until recently been neglected. In the past few years a number of studies have examined the effects of the N supply on Arabidopsis development, with an emphasis on roots, and some progress is now beginning to be made in identifying components of the signalling pathways involved and their relationship to signalling pathways for plant hormones. Some of this work has been reviewed in detail elsewhere (Zhang and Forde, 2000).

### Nitrate Stimulation of Lateral Root Growth

A series of classical studies in the 1970's examined the effects of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  on root development in barley (*Hordeum vulgare*) (Drew et al., 1973; Drew, 1975; Drew and Saker, 1975). Using sand cultures or flowing nutrient solutions, Drew and colleagues found that locally concentrated supplies of  $\text{NO}_3^-$ ,  $\text{NH}_4^+$  or phosphate stimulated root branching specifically within the nutrient-rich zones. The response consisted of a stimulation of both lateral root initiation and lateral root growth in the zone of treatment. In contrast, growth of the seminal roots was little affected by the external nutrient supply (Drew et al., 1973), showing



**Figure 8. Segmented agar plate (SAP) method for applying localised nutrient treatment to Arabidopsis roots.** Figure adapted from (Zhang and Forde, 1998). To prevent diffusion of nutrients between different treatment zones, agar plates are divided into three segments using a sterile scalpel. For localised  $\text{NO}_3^-$  treatments, the agar contains a basal supply of N ( $10 \mu\text{M NH}_4\text{NO}_3$ ) and the middle segment is supplemented with either  $1 \text{ mM KNO}_3$  or (as control)  $1 \text{ mM KCl}$ . At the start of the experiment (1), seedlings with roots *ca* 2 cm long are placed on the top segment, with just a few mm of the primary root apex resting on the middle segment. The plates are incubated vertically while the primary roots grow over the surface of the middle (treated) segment and onto the bottom segment (2). Around 10–12 d after seedling transfer (3), lateral root lengths in each segment are measured.



**Figure 9. Localised stimulation of lateral root growth by  $\text{NO}_3^-$ .** Seedling 1 (control): wild-type seedling grown on segmented agar plate containing low  $\text{NO}_3^-$  ( $10 \mu\text{M NH}_4\text{NO}_3$ ) in all three segments; Seedling 2: wild-type seedling exposed to a localised supply of  $1 \text{ mM KNO}_3$ ; Seedling 3: transgenic line S10 (in which expression of the *ANR1* MADS-box gene has been down-regulated by co-suppression) exposed to a localised supply of  $1 \text{ mM KNO}_3$  (Zhang and Forde, 1998).

that different types of root meristem within one root system can have very different sensitivities to nutrients.

The development of a technique for applying nutrients in a localised fashion to Arabidopsis seedlings growing on vertical agar plates (Figure 8) allowed this phenomenon to be investigated in a species more amenable to molecular genetics (Zhang and Forde, 1998). Initial studies showed that localised  $\text{NO}_3^-$  treatments did indeed stimulate lateral root elongation (2- to 3-fold), specifically within the  $\text{NO}_3^-$ -rich zone (Figure 9). The stimulation occurred even if the  $\text{NO}_3^-$  treatments were delayed until after the lateral roots had emerged (Zhang and Forde, 2000), showing that the  $\text{NO}_3^-$  does not need to be present during initiation or maturation of lateral root primordia for it to be effective, as has also been shown in wheat (Hackett, 1972). As in barley and other species, primary root growth in Arabidopsis is remarkably insensitive to changes in the  $\text{NO}_3^-$  supply (Zhang and Forde, 1998).

Neither the number of lateral roots initiated (Zhang and Forde, 1998), nor the rate of progression through the early stages of lateral root development (Zhang and Forde, 2000), are modified by localised  $\text{NO}_3^-$  treatments. The absence of any significant effect on lateral root initiation differs from the situation in barley and wheat, where the localised response to  $\text{NO}_3^-$  consisted of both an increase in elongation rates and an increase in numbers of laterals (Hackett, 1972; Drew et al., 1973). In Arabidopsis it was shown that even relatively low concentrations of  $\text{NO}_3^-$  ( $50$ – $100 \mu\text{M}$ ) in the  $\text{NO}_3^-$ -enriched zone were sufficient to elicit the response (Zhang et al., 1999). The absence of any significant increase in the size of mature cells in the  $\text{NO}_3^-$ -stimulated roots established that the increased growth rates were due to increased meristematic activity in the lateral root tip (Zhang et al., 1999).


When localised supplies of  $\text{NH}_4^+$  or glutamine were substituted for  $\text{NO}_3^-$ , no stimulation of lateral root growth was observed (Zhang et al., 1999), indicating that the roots were responding to the  $\text{NO}_3^-$  ion itself rather than to  $\text{NO}_3^-$  as a source of N. Further evidence for this was obtained using a *nia1nia2* mutant which is deficient in NR and hence unable to use  $\text{NO}_3^-$  as an effective N source (Wilkinson and Crawford, 1993); this mutant showed a wild-type response to a localised  $\text{NO}_3^-$  treatment (Zhang and Forde, 1998). Thus, just as  $\text{NO}_3^-$  is a signal for the induction of the nitrate assimilatory pathway (see above), it is also a signal for increased root proliferation. It seems that individual lateral root tips have a mechanism for monitoring the external  $\text{NO}_3^-$  concentration and for modifying their meristematic activity accordingly (Zhang and Forde, 1998; Zhang et al., 1999).



### Genes Involved in Nitrate Stimulation of Lateral Root Growth

A possible component of the signal transduction pathway linking external  $\text{NO}_3^-$  to changes in meristematic activity has been identified during a screen for  $\text{NO}_3^-$  regulated genes in Arabidopsis roots (Zhang and Forde, 1998). Amongst the novel  $\text{NO}_3^-$ -inducible sequences identified was ANR1, a member of the MADS box family of transcription factors (Zhang and Forde, 1998). There are at least 82 MADS box genes in the Arabidopsis genome (Riechmann et al., 2000) and most of those that have so far been functionally characterised are involved in controlling floral organogenesis or other processes related to the reproductive phase of growth (Ng and Yanofsky, 2001). ANR1 belongs to a group of nine MADS box genes so far known to be expressed in Arabidopsis roots (Alvarez-Buylla et al., 2000), but most of these are of unknown function. When transgenic lines were generated in which ANR1 expression was down-regulated (by antisense or co-suppression), their lateral roots were found to have lost the ability to respond to localised supplies of  $\text{NO}_3^-$  (Figure 9). It was proposed that ANR1 is a component of the sensory pathway linking  $\text{NO}_3^-$  to the stimulation of lateral root growth (Zhang and Forde, 1998).

Auxin is well known as an important regulator of root growth (Leyser, 1997), and experiments using auxin-response mutants of Arabidopsis have provided evidence



**Figure 10. Model for  $\text{NO}_3^-$  signalling pathway leading to localised stimulation of lateral root growth.** The external  $\text{NO}_3^-$  supply is monitored by individual lateral root tips and the signal is transduced via the  $\text{NO}_3^-$ -inducible ANR1 gene and the auxin-sensitivity gene AXR4 to produce increased meristematic activity. The identity and sub-cellular location of the  $\text{NO}_3^-$  receptor/sensor are unknown, as is the mechanism for induction of ANR1. ANR1 has tentatively been placed upstream of ANR1 in the signal transduction pathway. The model is based on that proposed by (Zhang et al., 1999). See text for further details.

for an overlap between  $\text{NO}_3^-$  signalling and auxin signalling in the control of lateral root elongation (Zhang et al., 1999). Three auxin-response mutants (aux1, aux2 and aux4) were tested for their sensitivity to localised supplies of  $\text{NO}_3^-$ : While aux1 and aux2 showed responses that were not significantly different from the wild-type, the aux4 mutant showed a phenotype similar to the ANR1 down-regulated lines (Zhang et al., 1999). Although the AXR4 gene has not been identified at the molecular level, its location on chromosome 1 (Hobbie and Estelle, 1995) establishes it as distinct from ANR1 (which is on chromosome 2). Unusual among auxin-response mutants, the phenotype of aux4 mutants is restricted to roots and does not involve cross-resistance to other plant hormones such as cytokinin or ethylene (Hobbie and Estelle, 1995). Apart from reduced gravitropism, the root phenotype of the aux4-2 mutant is remarkably like the wild-type, making its altered sensitivity to  $\text{NO}_3^-$  stimulation of lateral root elongation of particular interest. A model for the  $\text{NO}_3^-$  signalling pathway leading to increased meristematic activity in the lateral root tip is shown in Figure 10.

### Nitrate as an Inhibitor of Early Lateral Root Development


Nitrate has also been found to have an inhibitory effect on lateral root development (Scheible et al., 1997c; Zhang and Forde, 1998; Stitt, 1999; Zhang et al., 1999). In Arabidopsis this consists of a delayed outgrowth of the lateral root primordium when roots are exposed to  $\text{NO}_3^-$  concentrations above 1 mM (Zhang et al., 1999). At 50 mM, the highest  $\text{NO}_3^-$  concentration tested, two-week-old seedlings developed primary roots of the same length as at 1 mM, and had a similar number of lateral root primordia, but most or all of the laterals were less than 0.5 mm in length. This inhibitory effect of  $\text{NO}_3^-$  appears to be very different in nature from its stimulatory effect (Zhang et al., 1999; Zhang and Forde, 2000). Firstly, while the stimulatory effect acts on elongation of mature laterals, the inhibitory effect acts specifically on immature LRs during a discrete phase around the time of their emergence from the primary root. Secondly, the high external  $\text{NO}_3^-$  concentration does not act directly on the developing lateral; if only a portion of the primary root was exposed to 50 mM  $\text{NO}_3^-$ , the inhibitory effect was much diminished and what inhibitory effect remained was distributed along the root (Zhang et al., 1999). Thus the effect seems to depend on the  $\text{NO}_3^-$  absorbed by the plant rather than the external  $\text{NO}_3^-$  concentration per se.

Using the NR-deficient *nia1nia2* mutant it was shown that the assimilation of  $\text{NO}_3^-$  was not required for the inhibitory effect, and furthermore that lateral root development in the mutant was rather more sensitive to  $\text{NO}_3^-$  inhibition than the wild-type (Zhang et al., 1999). Nitrate reductase-deficient mutants growing on  $\text{NO}_3^-$  are known to accumulate high levels of tissue  $\text{NO}_3^-$ , particularly in the leaves e.g. Warner and Huffaker, 1989), so this observation is consistent with the notion that  $\text{NO}_3^-$  levels inside the plant are responsible for generating the inhibitory effect on lateral root outgrowth. A similar phenomenon has been reported in tobacco, where NR deficient lines growing on  $\text{NO}_3^-$  display a marked inhibition of root growth (Scheible et al., 1997b; Scheible et al., 1997c) due to a reduction in lateral root frequency (Stitt, 1999). Here the evidence from metabolite analysis and split root experiments has led to the conclusion that the inhibition of root growth is due to a signal from the shoot triggered by  $\text{NO}_3^-$  accumulation in the shoot (Scheible et al., 1997c). In Arabidopsis (but not in tobacco), an increase in the sucrose concentration in the medium led to a partial alleviation of the inhibitory effect of  $\text{NO}_3^-$  on root branching (Scheible et al., 1997c; Zhang et al., 1999). This would suggest that either the balance between the carbon and nitrogen status of the plant is being monitored, or that sucrose itself has a signalling role that acts antagonistically with the  $\text{NO}_3^-$  signal (Smeekens, 2000).

Why outgrowth of the lateral root primordium should be particularly sensitive to inhibition by  $\text{NO}_3^-$  is unclear. From other work it is known that the stage just after its emergence is a critical step in the development of a lateral root

(Cheng et al., 1995; Malamy and Benfey, 1997a; Malamy and Benfey, 1997b). Although the lateral root meristem appears to be fully formed prior to emergence, emergence itself takes place through cell expansion rather than cell division (Malamy and Benfey, 1997b). It is only after emergence that activation of the lateral root meristem occurs and growth of the mature lateral root begins (Cheng et al., 1995; Malamy and Benfey, 1997b). Thus there may be a developmental process between differentiation of the lateral root meristem and its activation that is particularly sensitive to high  $\text{NO}_3^-$  concentrations. An alternative explanation could lie in the timing of the establishment of symplastic connections between the phloem of the primary root and the lateral root primordium. Since a shoot-derived signal may be involved, these connections could be necessary for transmission of the signal to the developing primordium. A study using fluorescent dyes and confocal microscopy has elegantly demonstrated for Arabidopsis that early in its development the lateral root is symplastically isolated and that the earliest functional symplastic connections with the lateral root primordium occur around the time of emergence from the cortex (Oparka et al., 1995).

The finding that  $\text{NO}_3^-$  can either inhibit or stimulate lateral root production, depending on its external distribution and its accumulation within plant tissues, has led to a 'dual pathway' model for  $\text{NO}_3^-$  regulation of root branching in Arabidopsis (Zhang et al., 1999). This model, illustrated in Figure 11, depicts one signalling pathway (the localised stimulatory pathway) which responds to the external  $\text{NO}_3^-$  concentration at the lateral root tip and regulates increase



**Figure 11. 'Dual pathway' model for  $\text{NO}_3^-$  regulation of root branching by nitrate.** (1). Localised stimulatory pathway responding to changes in the external  $\text{NO}_3^-$  supply (see Figure 10). (2) Systemic inhibitory pathway, requiring the uptake and transport of  $\text{NO}_3^-$  to the shoot. (3) Maturation of the lateral root primordium is under the control of shoot-derived inhibitory signal that becomes prevalent at high shoot  $\text{NO}_3^-$  concentrations. (4) An increased supply of sucrose is able to alleviate the inhibitory effect of high  $\text{NO}_3^-$  supplies. The model is based on that proposed by (Zhang et al., 1999). See text for further details.

meristematic activity specifically in that root tip, and a second signalling pathway (the systemic inhibitory pathway) which responds to the internal  $\text{NO}_3^-$  status of the plant and regulates progress through the latter stages of lateral root maturation. These two antagonistic pathways for regulation of root branching can be seen as providing a mechanism by which the plant can modify its root architecture in a manner that integrates information about the spatial distribution of  $\text{NO}_3^-$  in the soil and the plant's current N status (Zhang et al., 1999). Thus the plant could modulate the intensity of its proliferative response to a localised  $\text{NO}_3^-$  supply according to its demand for N. This would allow the allocation of resources within the plant to be optimised by ensuring that the strongest response to a localised  $\text{NO}_3^-$  supply occurred when the plant was N-depleted (Zhang et al., 1999). Increased lateral root proliferation is only one component of a plant's response to localised  $\text{NO}_3^-$  supplies; a much faster response is seen at the metabolic level, where  $\text{NO}_3^-$  influx by roots located within a  $\text{NO}_3^-$ -rich patch is rapidly induced (Laine et al., 1995; Laine et al., 1998). The up-regulation of the  $\text{NO}_3^-$  influx system can be seen as a short-term adaptive response to localised  $\text{NO}_3^-$ , with stimulation of lateral root development being the longer-term component of the response. It is known that the  $\text{NO}_3^-$  influx system is regulated by signals originating in the shoot (Laine et al., 1995), and it remains to be seen whether the same signals are involved in regulating lateral root development.

### Ammonium Inhibition of Root Growth

Ammonium, when supplied to plants as the sole N source, often inhibits plant growth, particularly in unbuffered media when acidification of the rhizosphere becomes a problem (von Wiren et al., 2001). However, a novel aspect of  $\text{NH}_4^+$  inhibition of root growth has been reported in Arabidopsis (Cao et al., 1993). In these studies it was found that if  $\text{K}^+$  was omitted from the medium, and  $\text{NH}_4^+$  concentrations were high (>3 mM) seedling root growth was dramatically inhibited, even when  $\text{NO}_3^-$  was present. This inhibition could be reversed if low levels of  $\text{K}^+$ ,  $\text{Rb}^+$  or  $\text{Cs}^+$  ions were added, and the protective effect of  $\text{K}^+$  was shown not to be due to an inhibition of  $\text{NH}_4^+$  uptake. The further finding that three auxin-response mutants (*aux1*, *axr1* and *axr2*) were resistant to  $\text{NH}_4^+$  inhibition of root growth led to the suggestion that this unusual response to  $\text{NH}_4^+$  was mediated by effects on hormone metabolism or signalling. Since it is unlikely that an Arabidopsis plant will often be exposed in nature to the required combination of high  $\text{NH}_4^+$  and very low  $\text{K}^+$  concentrations, it is unclear what


the physiological significance of this intriguing phenomenon may be; however, it is an example in Arabidopsis showing the role of auxin in nutrient effects on root growth.

### Nitrogen Regulation of Shoot Growth and Development

It has been established for Arabidopsis that a higher rate of N supply leads to an increased allocation of resources to shoot growth (Schulze et al., 1994), leading to increased shoot:root ratios, which is the morphological response commonly seen in many other plant species (Ericsson, 1995). Analysis of metabolite pools showed that there was a striking correlation between the concentrations of  $\text{NO}_3^-$  and organic N in rosette leaves and shoot growth rates (Schulze et al., 1994). In tobacco, transgenic lines with low NR activity have helped to demonstrate the role of the shoot  $\text{NO}_3^-$  pool as a signal in regulating shoot-root allocation (Scheible et al., 1997c).

Other studies have highlighted the importance of the short-term N supply and root-to-shoot signalling in the regulation of leaf growth (McDonald and Davies, 1996). Recent studies with tobacco found that the rate of cell division and cell elongation in leaves fell markedly within 24 h of transferring plants from  $\text{NO}_3^-$  to  $\text{NH}_4^+$  (Walch-Liu et al., 2000). This effect was not due to limitation of carbohydrates or N, nor to a lack of osmotica or water. A possible role for  $\text{NO}_3^-$  as a signal was inferred from evidence that it was absence of  $\text{NO}_3^-$  rather than the presence of  $\text{NH}_4^+$  that was responsible for the reduction in leaf growth (Walch-Liu et al., 2000). A decline in the zeatin + zeatin riboside fraction in xylem exudates which paralleled that in leaf expansion rates led to the suggestion that  $\text{NO}_3^-$  may be involved in a signal transduction chain that regulates leaf morphogenesis by modulating the flux of cytokinin from root to shoot (Walch-Liu et al., 2000).

Although leaves of  $\text{NO}_3^-$ -grown Arabidopsis plants are larger than those of  $\text{NH}_4^+$ -grown plants (e.g. (Oostinder-Braaksma and Feenstra, 1973)), the possible short-term modulation of leaf expansion by the  $\text{NO}_3^-$  supply has not been studied in Arabidopsis. However it is interesting to note that some members of the Arabidopsis family of response regulators (ARRs) have been shown to be induced in leaves by either the direct application of cytokinins or the re-supply of  $\text{NO}_3^-$  to roots of N-starved plants (Taniguchi et al., 1998). It has been proposed that the ARR genes are part of a cytokinin-mediated signalling pathway which enables gene expression in leaves to be modulated by the inorganic N supply to the roots



**Figure 12. Model for the control of leaf expansion by the NO<sub>3</sub><sup>-</sup> supply to the root.** (1) The external supply of NO<sub>3</sub><sup>-</sup> stimulates cytokinin (CK) production in the root. (2) CKs are transported to the shoot via the xylem. (3) CKs stimulate expression of members of the ARR (Arabidopsis response regulator) family of transcription factors. (4) The ARR transcription factor(s) activate a set of target genes to stimulate cell division and cell expansion in the leaf. See text for further details.

(Sakakibara et al., 2000). The finding that one of the cytokinin-regulated genes (ARR5) is expressed in shoot apical meristems (D'Agostino et al., 2000), would be consistent with a possible role in regulating leaf expansion in response to the NO<sub>3</sub><sup>-</sup> supply. A tentative model for such a NO<sub>3</sub><sup>-</sup>–cytokinin signalling pathway is depicted in Figure 12.

In many species, increased N fertilisation is reported to delay flowering (Bernier et al., 1993). In one study, flowering time in *Arabidopsis* was found to be only slightly delayed by a high rate of N supply (6 mM compared to 0.1 mM NH<sub>4</sub>NO<sub>3</sub>) (Schulze et al., 1994). In this study, seed set was much more sensitive to the N supply than flowering time, with the number of seeds per plant being over 10-fold greater under the high N regime. Importantly, by switching plants from high N to low N and vice versa at the end of the vegetative phase of growth, it was shown that it was the N supply during the reproductive phase that was critical in determining seed numbers (Schulze et al., 1994). It is not yet known which are the key N pools responsible for regulating leaf growth and seed set in *Arabidopsis*.

## REFERENCES

- Alvarez-Buylla, E. R., Liljegren, S. J., Pelaz, S., Gold, S. E., Burgeff, C., Ditta, G. S., Vergara-Silva, F., and Yanofsky, M. F. (2000). MADS-box gene evolution beyond flowers: expression in pollen, endosperm, guard cells, roots and trichomes. *Plant J.* **24**, 457-466.
- Amtmann, A., and Sanders, D. (1999). Mechanisms of Na<sup>+</sup> uptake by plant cells. *Adv. Bot. Res.* **29**, 75-112.
- Aslam, M., Travis, R., and Huffaker, R. C. (1993). Comparative induction of nitrate and nitrite uptake and reduction systems by ambient nitrate and nitrite in intact roots of barley seedlings. *Plant Physiol.* **102**, 811-819.
- Bernier, G., Havelange, A., Houssa, C., Petitjean, A., and Lejeune, P. (1993). Physical signals that induce flowering. *Plant Cell* **5**, 1147-1155.
- Blanc, G., Barakat, A., Guyot, R., Cooke, R., and Delseny, M. (2000). Extensive duplication and reshuffling in the *Arabidopsis* genome. *Plant Cell* **12**, 1093-1101.
- Bloom, A. J. (1997). Interactions between inorganic nitrogen nutrition and root development. *Zeitschrift Pflanzenernahrung Bodenkunde* **160**, 253-259.
- Braaksma, F. J., and Feenstra, W. J. (1973). Isolation and characterization of chlorate resistant mutants of *Arabidopsis thaliana*. *Mutat. Res.* **19**, 175-185.
- Braaksma, F. J., and Feenstra, W. J. (1982). Isolation and characterization of nitrate reductase-deficient mutants of *Arabidopsis thaliana*. *Theor. Appl. Genet.* **64**, 83-90.
- Braaksma, F. J., and Feenstra, W. J. (1982). Nitrate reduction in the wildtype and a nitrate reductase deficient mutant of *Arabidopsis thaliana*. *Physiol. Plant.* **54**, 351-360.
- Brownlee, A. G., and Arst, H. N., Jr. (1983). Nitrate uptake in *Aspergillus nidulans* and involvement of the third gene of the nitrate assimilation gene cluster. *J. Bacteriol.* **155**, 1138-1146.
- Campbell, W. H. (1999). Nitrate reductase structure, function and regulation: Bridging the gap between biochemistry and physiology. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **50**, 277-303.
- Campbell, W. H. (2001). Structure and function of eukaryotic NAD(P)H:nitrate reductase. *Cell. Mol. Life Sci.* **58**, 194-204.
- Cao, D., Lin, Y., and Cheng, C.-L. (2000). Genetic interactions between the chlorate-resistant mutant *cr88* and the photomorphogenic mutants *cop1* and *hy5*. *Plant Cell* **12**, 199-210.
- Cao, Y., Glass, A. D. M., and Crawford, N. M. (1993). Ammonium inhibition of *Arabidopsis* root growth can be reversed by potassium and auxin resistance mutations *aux1*, *axr1* and *axr2*. *Plant. Physiol.* **102**, 983-989.
- Cerezo, M., Tillard, P., Filleur, S., Munos, S., Daniel-Vedele, R., and Gojon, A. (2001). Alterations of the regulation of root NO<sub>3</sub><sup>-</sup> uptake are associated with the mutation of *Nrt2.1* and *Nrt2.2* genes in *Arabidopsis*. *Plant Physiol.* in press.
- Champigny, M.-L., and Foyer, C. (1992). Nitrate activation of cytosolic protein kinases diverts photosynthetic carbon from sucrose to amino acid biosynthesis. *Plant Physiol.* **100**, 7-12.
- Cheng, C.-L., Acedo, G. N., Cristinsin, M., and Conkling, M. A. (1992). Sucrose mimics the light induction of *Arabidopsis* nitrate reductase gene transcription. *Proc. Natl. Acad. Sci. USA* **89**, 1861-1864.

- Cheng, C.-L., Acedo, G. N., Dewdney, J., Goodman, H. M., and Cankling, M. A.** (1991). Differential expression of the two *Arabidopsis* nitrate reductase genes. *Plant Physiol.* **96**, 275-279.
- Cheng, C.-L., Dewdney, J., Kleinhofs, A., and Goodman, H. M.** (1986). Cloning and nitrate induction of nitrate reductase mRNA. *Proc. Natl. Acad. Sci. USA* **83**, 6825-6828.
- Cheng, C.-L., Dewdney, J., Nam, H.-G., Den Boer, B. G. W., and Goodman, H. M.** (1988). A new locus (*NIA1*) in *Arabidopsis thaliana* encoding nitrate reductase. *EMBO J.* **7**, 3309-3314.
- Cheng, J., Seeley, K. A., and Sung, Z. R.** (1995). *RML1* and *RML2*, *Arabidopsis* genes required for cell proliferation at the root tip. *Plant Physiol.* **107**, 365-376.
- Coruzzi, G., and Bush, D. R.** (2001). Nitrogen and carbon nutrient and metabolite signaling in plants. *Plant Phys.* **125**, 61-64.
- Coruzzi, G. M., and Zhou, L.** (2001). Carbon and nitrogen sensing and signaling in plants: emerging 'matrix effects'. *Curr. Opin. Plant Biol.* **4**, 247-253.
- Cove, D. J.** (1979). Genetic studies of nitrate assimilation in *Aspergillus nidulans*. *Biol. Rev.* **54**, 291-327.
- Crawford, N. M.** (1992). Study of chlorate resistant mutants of *Arabidopsis*: insights into nitrate assimilation and ion metabolism of plants. In *Genetic Engineering, Principles and Methods*, J. K. Setlow, ed (New York: Plenum Press), pp. 89-98.
- Crawford, N. M.** (1995). Nitrate: Nutrient and signal for plant growth. *Plant Cell* **7**, 859-868.
- Crawford, N. M., and Arst, H. N. J.** (1993). The molecular genetics of nitrate assimilation in fungi and plants. *Annu. Rev. Genet.* **27**, 115-146.
- Crawford, N. M., Campbell, W. H., and Davis, R. W.** (1986). Nitrate reductase from squash: cDNA cloning and nitrate regulation. *Proc. Natl. Acad. Sci. USA* **83**, 8073-8076.
- Crawford, N. M., and Glass, A. D. M.** (1998). Molecular and physiological aspects of nitrate uptake in plants. *Trends Plant Sci.* **3**, 389-395.
- Crawford, N. M., Kahn, M. L., Leustek, T., and Long, S. R.** (2000). Nitrogen and Sulfur. In *Biochemistry & Molecular Biology of Plants*, B. Buchanan, W. Gruissem and R. L. Jones, ed (Rochville, MD: American Society of Plant Physiologists), pp. 786-849.
- Crawford, N. M., Smith, M., Bellissimo, D., and Davis, R. W.** (1988). Sequence and nitrate regulation of the *Arabidopsis thaliana* mRNA encoding nitrate reductase, a metalloflavoprotein with three functional domains. *Proc. Natl. Acad. Sci. USA* **85**, 5006-5010.
- D'Agostino, I. B., Deruere, J., and Kieber, J. J.** (2000). Characterization of the response of the *Arabidopsis* response regulator gene family to cytokinin. *Plant Physiol.* **124**, 1706-1717.
- Daniel-Vedele, F., Filleur, S., and Caboche, M.** (1998). Nitrate transport: a key step in nitrate assimilation. *Curr. Opin. Plant Biol.* **1**, 235-239.
- Doddema, H., Hofstra, J., and Feenstra, W.** (1978). Uptake of nitrate by mutants of *Arabidopsis thaliana*, disturbed in uptake or reduction of nitrate I. Effect of nitrogen source during growth on uptake of nitrate and chlorate. *Physiol. Plant.* **43**, 343-350.
- Doddema, H., and Telkamp, G. P.** (1979). Uptake of nitrate by mutants of *Arabidopsis thaliana*, disturbed in uptake or reduction of nitrate II. Kinetics. *Physiol. Plant.* **45**, 332-338.
- Drew, M. C.** (1975). Comparison of the effects of a localized supply of phosphate, nitrate, ammonium and potassium on the growth of the seminal root system, and the shoot, in barley. *New Phyt.* **75**, 479-490.
- Drew, M. C., and Saker, L. R.** (1975). Nutrient supply and the growth of the seminal root system of barley. II. Localized, compensatory increases in lateral root growth and rates of nitrate uptake when nitrate supply is restricted to only part of the root system. *J. Exp. Bot.* **26**, 79-90.
- Drew, M. C., Saker, L. R., and Ashley, T. W.** (1973). Nutrient supply and the growth of the seminal root system in barley. I. The effect of nitrate concentration on the growth of axes and laterals. *J. Exper. Bot.* **24**, 1189-1202.
- Dzuibany, C., Haupt, S., Fock, H., Biehler, K., Migge, A., and Becker, T. W.** (1998). Regulation of nitrate reductase transcript levels by glutamine accumulating in the leaves of a ferredoxin-dependent glutamate synthase-deficient *gluS* mutant of *Arabidopsis thaliana*, and by glutamine provided via the roots. *Planta* **206**, 515-522.
- Epstein, E.** (1953). Mechanism of ion absorption by roots. *Nature* **171**, 83-84.
- Epstein, E.** (1972). *Mineral Nutrition of Plants: Principles and Perspectives*. Wiley, New York.
- Ericsson, T.** (1995). Growth and shoot:root ratio of seedlings in relation to nutrient availability. *Plant Soil* **169**, 205-214.
- Felsenstein, J.** (1993). PHYLIP (Phylogeny Inference Package). Seattle, Department of Genetics, University of Washington.
- Filleur, S., and Daniel-Vedele, F.** (1999). Expression analysis of a high-affinity nitrate transporter isolated from *Arabidopsis thaliana* by differential display. *Planta* **207**, 461-469.
- Filleur, S., Dorbe, M.-F., Cerizo, M., Orsel, M., Granier, F., Gojon, A., and Daniel-Vedele, F.** (2001). An *Arabidopsis* T-DNA mutant affected in *Nrt2* genes is impaired in nitrate uptake. *FEBS Letters* **489**, 220-224.
- Forde, B. G.** (2000). Nitrate transporters in plants: structure, function and regulation. *Biochim. Biophys. Acta* **1465**, 219-235.
- Forde, B. G., and Clarkson, D. T.** (1999). Nitrate and ammonium nutrition of plants: Physiological and molecular perspectives. *Adv. Bot. Res.* **30**, 1-90.
- Forde, B. G., and Lorenzo, H.** (2001). The nutritional control of root development. *Plant and Soil in press*.
- Galvan, A., and Fernandez, E.** (2001). Eukaryotic nitrate and nitrite transporters. *Cell. Mol. Life Sci.* **58**, 225-233.
- Gazzarrini, S., Lejay, T., Gojon, A., Ninnemann, O., Frommer, W. B., and von Wiren, N.** (1999). Three functional transporters for constitutive, diurnally regulated, and starvation-induced uptake of ammonium into *Arabidopsis* roots. *Plant Cell* **11**, 937-947.
- Glass, A. D. M., Brito, D. T., Kaiser, B. N., Kronzucker, H. J., Kumar, A., Okamoto, M., Rawat, S. R., Siddiqi, M. Y., Silim, S. M., Vidmar, J. J., and Zhuo, D.** (2001). Nitrogen transport in plants, with an emphasis on the regulation of fluxes to match plant demand. *J. Plant Nutr. Soil Sci.* **164**, 199-207.

- Glass, A. D. M., Erner, Y., Kronzucker, H. J., Schjoerring, J. K., Siddiqi, M. Y., and Wang, M. Y.** (1997). Ammonium fluxes into plant roots: energetics, kinetics and regulation. *Zeitschrift Pflanzenernahrung Bodenkunde* **160**, 261-268.
- Glass, A. D. M., and Siddiqi, M. Y.** (1995). Nitrogen absorption by plant roots. In *Nitrogen Nutrition in Higher Plants*, H. S. Srivastava and R. P. Singh, ed (New Delhi, India: Associated Publishing Co.), pp. 21-56.
- Gowri, G., Kenis, J. D., Ingemarsson, B., Redinbaugh, M. G., and Campbell, W. H.** (1992). Nitrate reductase transcript is expressed in the primary response of maize to environmental nitrate. *Plant Mol. Biol.* **18**, 55-64.
- Graul, R. C., and Sadee, W.** (1997). Sequence alignments of the H<sup>+</sup>-dependent oligopeptide transporter family PTR: Inferences on structure and function of the intestinal PET1 transporter. *Pharm. Res.* **14**, 388-400.
- Guo, F.-Q., Wang, R., Chen, M., and Crawford, N. M.** (2001). The Arabidopsis dual-affinity nitrate transporter gene *AtNRT1.1 (CHL1)* is activated and functions in nascent organ development during vegetative and reproductive growth. *Plant Cell* in press.
- Hackett, C.** (1972). A method of applying nutrients locally to roots under controlled conditions, and some morphological effects of locally applied nitrate on the branching of wheat roots. *Aust. J. Biol. Sci.* **25**, 1169-1180.
- Hatzfeld, Y., and Saito, K.** (1999). Identification of two putative nitrate transporters highly homologous to CHL1 from Arabidopsis. *Plant Phys.* **119**, 805.
- Hobbie, L., and Estelle, M.** (1995). The *axr4* auxin-resistant mutants of *Arabidopsis thaliana* define a gene important for root gravitropism and lateral root initiation. *Plant J.* **7**, 211-220.
- Hoff, T., Schnorr, K. M., Meyer, C., and Caboche, M.** (1995). Isolation of two *Arabidopsis* cDNAs involved in early steps in molybdenum cofactor biosynthesis by functional complementation of *Escherichia coli* mutants. *J. Biol. Chem.* **270**, 6100-6107.
- Hoff, T., Truon, H.-M., and Caboche, M.** (1994). The use of mutants and transgenic plants to study nitrate assimilation. *Plant, Cell & Environment* **17**, 489-506.
- Howitt, S. M., and Udvardi, M. K.** (2000). Structure, function and regulation of ammonium transporters in plants. *Biochim. Biophys. Acta-Biomembr.* **1465**, 152-170.
- Huang, N.-C., Chiang, C.-S., Crawford, N. M., and Tsay, Y.-F.** (1996). CHL1 encodes a component of the low-affinity nitrate uptake system in *Arabidopsis* and shows cell type-specific expression in roots. *Plant Cell* **8**, 2183-2191.
- Huang, N.-C., Liu, K.-H., Lo, H.-J., and Tsay, Y.-F.** (1999). Cloning and functional characterization of an Arabidopsis nitrate transporter gene that encodes a constitutive component of low-affinity uptake. *Plant Cell* **11**, 1381-1392.
- Hwang, C.-F., Lin, Y., D'Souza, T., and Cheng, C.-L.** (1997). Sequences necessary for nitrate-dependent transcription of Arabidopsis nitrate reductase genes. *Plant Physiol.* **113**, 853-862.
- Kaiser, W. M., Weiner, H., and Huber, S. C.** (1999). Nitrate reductase in higher plants: A case study for transduction of environmental stimuli into control of catalytic activity. *Physiol. Plant.* **105**, 385-390.
- Klasson, H., Fink, G. R., and Ljungdahl, P. O.** (1999). Ssy1p and Ptr3p are plasma membrane components of a yeast system that senses extracellular amino acids. *Mol. Cell. Biol.* **19**, 5405-5416.
- Kleinhofs, A., Warner, R. L., Muehlbauer, F. J., and R.A., N.** (1978). Inducible and selctin of specific gene mutations of hordeum and pisum. *Mut. Res.* **51**, 29-35.
- Koch, K. E.** (1997). Molecular crosstalk and the regulation of C- and N-responsive genes. In *A Molecular Approach to Primary Metabolism in Higher Plants*, C. H. Foyer and W. P. Quick, ed (London: Taylor & Francis), pp. 105-124.
- Kronzucker, H. J., Siddiqi, M. Y., and Glass, A. D. M.** (1995). Compartmentation and flux characteristics of ammonium in spruce. *Planta* **196**, 691-698.
- Kronzucker, H. J., Siddiqi, M. Y., and Glass, A. D. M.** (1997). Conifer root discrimination against soil nitrate and the ecology of forest succession. *Nature* **385**, 59-61.
- Kuper, J., Palmer, T., Mendel, R. R., and Schwarz, G.** (2000). Mutations in the molybdenum cofactor biosynthetic protein Cnx1G from *Arabidopsis thaliana* define functions for molybdopterin binding, molybdenum insertion, and molybdenum cofactor stabilization. *Proc. Natl. Acad. Sci. USA* **97**, 6475-6480.
- LaBrie, S. T., Wilkinson, J. Q., Tsay, Y.-F., Feldmann, K. A., and Crawford, N. M.** (1992). Identification of two tungstate-sensitive molybdenum cofactor mutants, *chl2* and *chl7*, of *Arabidopsis thaliana*. *Mol. Gen. Genet.* **233**, 169-176.
- Laine, P., Ourry, A., and Boucaud, J.** (1995). Shoot control of nitrate uptake rates by roots of *Brassica napus* L.: Effects of localized nitrate supply. *Planta* **196**, 77-83.
- Laine, P., Ourry, A., Boucaud, J., and Salette, J.** (1998). Effects of a localized supply of nitrate on NO<sub>3</sub><sup>-</sup> uptake rate and growth of roots in *Lolium multiflorum* Lam. *Plant Soil* **202**, 61-67.
- Lam, H. M., Coschigano, K. T., Oliveira, I. C., Melooliveira, R., and Coruzzi, G. M.** (1996). The molecular-genetics of nitrogen assimilation into amino acids in higher plants. *Ann. Rev. Plant Phys. Plant Mol. Biol.* **47**, 569-593.
- Lejay, L., Tillard, P., Leptit, M., Olive, F. D., Filleur, S., Daniel-Vedele, F., and Gojon, A.** (1999). Molecular and functional regulation of two NO<sub>3</sub><sup>-</sup> uptake systems by N- and C-status of Arabidopsis plants. *Plant Journal* **18**, 509-519.
- Leydecker, M. T., Camus, I., Daniel-Vedele, F., and Truong, H. N.** (2000). Screening for *Arabidopsis* mutants affected in the *Nii* gene expression using the *Gus* reporter gene. *Physiol. Plant.* **108**, 161-170.
- Leyser, O.** (1997). Auxin: lessons from a mutant weed. *Physiologia Plantarum* **100**, 407-414.
- Lin, Y., and Cheng, C. L.** (1997). A chlorate-resistant mutant defective in the regulation of nitrate reductase gene expression in Arabidopsis defines a new *HY* locus. *Plant Cell* **9**, 21-35.
- Lin, Y., Hwang, C.-F., Brown, J. B., and Cheng, C.-L.** (1994). 5' proximal regions of *Arabidopsis* nitrate reductase genes direct nitrate-induced transcription in transgenic tobacco. *Plant Physiol.* **106**, 477-484.
- Liu, K.-H., Huang, C.-Y., and Tsay, Y.-F.** (1999). CHL1 is a dual-affinity nitrate transporter of *Arabidopsis* involving multiple phases of nitrate uptake. *Plant Cell* **11**, 865-874.

- Lorenz, M. C., and Heitman, J.** (1998). The MEP2 ammonium permease regulates pseudohyphal differentiation in *Saccharomyces cerevisiae*. *EMBO J.* **17**, 1236-1247.
- MacKintosh, C., and Meek, S. E. M.** (2001). Regulation of plant NR activity by reversible phosphorylation, 14-3-3 proteins and proteolysis. *Cell. Mol. Life Sci.* **58**, 205-214.
- Malamy, J. E., and Benfey, P. N.** (1997a). Down and out in *Arabidopsis*: The formation of lateral roots. *Trends Plant Sci.* **2**, 390-396.
- Malamy, J. E., and Benfey, P. N.** (1997b). Organization and cell differentiation in lateral roots of *Arabidopsis thaliana*. *Development* **124**, 33-44.
- Marini, A. M., and Andre, B.** (2000). In vivo N-glycosylation of the Mep2 high-affinity ammonium transporter of *Saccharomyces cerevisiae* reveals an extracytosolic N-terminus. *Mol. Microbiol.* **38**, 552-564.
- Marschner, H.** (1995). *Mineral Nutrition of Higher Plants*. Academic Press, San Diego, CA.
- Marzluf, G. A.** (1981). Regulation of nitrogen metabolism and gene expression in fungi. *Microbiol. Rev.* **45**, 437-461.
- McDonald, A. J. S., and Davies, W. J.** (1996). Keeping in touch: Responses of the whole plant to deficits in water and nitrogen supply. *Adv. Bot. Res.* **22**, 229-300.
- Meharg, A. A., and Blatt, M. R.** (1995). NO<sub>3</sub><sup>-</sup> transport across the plasma membrane of *Arabidopsis thaliana* root hairs: kinetic control by pH and membrane voltage. *J. Membrane Biol.* **145**, 49-66.
- Melzer, J. M., Kleinhofs, A., and Warner, R. L.** (1989). Nitrate reductase regulation: Effects of nitrate and light on nitrate reductase mRNA accumulation. *Mol. Gen. Genet.* **217**, 341-346.
- Mendel, R. R.** (1997). Molybdenum cofactor of higher plants: biosynthesis and molecular biology. *Planta* **203**, 399-405.
- Mendel, R. R., and Schwarz, G.** (1999). Molybdoenzymes and molybdenum cofactor in plants. *Critical Rev. Plant Sci.* **18**, 33-69.
- Miller, A. J., Cookson, S. J., Smith, S. J., and Wells, D. M.** (2001). The use of microelectrodes to investigate compartmentation and the transport of metabolized inorganic ions in plants. *J. Exp. Bot.* **52**, 541-549.
- Muller, A. J., and Grafe, R.** (1978). Isolation and characterization of cell lines of *Nicotiana tabacum* lacking nitrate reductase. *Mol. Gen. Genet.* **161**, 67-76.
- Muller, C., Scheible, W.-R., Stitt, M., and Krapp, A.** (2001). Influence of malate and 2-oxoglutarate on the *NIA* transcript level and nitrate reductase activity in tobacco leaves. *Plant Cell Environ.* **24**, 191-203.
- Ng, M., and Yanofsky, M. F.** (2001). Function and evolution of the plant MADS-box gene family. *Nat. Rev. Genet.* **2**, 186-195.
- Ninnemann, O., Jauniaux, J. C., and Frommer, W. B.** (1994). Identification of a high affinity NH<sub>4</sub><sup>+</sup> transporter from plants. *EMBO J.* **13**, 3464-3471.
- Okamoto, M., Vidmar, J. J., and Glass, A. D. M.** (2001). Expression of 11 nitrate transporter (*AtNRT*) gene in *Arabidopsis thaliana*. *Plant Cell Physiol.* **42**, as217.
- Oostinder-Braaksma, F. J., and Feenstra, W. J.** (1973). Isolation and characterization of chlorate-resistant mutants of *Arabidopsis thaliana*. *Mutat. Res.* **19**, 175-185.
- Oparka, K. J., Prior, D. A. M., and Wright, K. M.** (1995). Symplastic communication between primary and developing lateral roots of *Arabidopsis thaliana*. *J. Exper. Bot.* **46**, 187-197.
- Ozcan, S., Dover, J., Rosenwald, A. G., Wolf, S., and Johnston, M.** (1996). Two glucose transporters in *Saccharomyces cerevisiae* are glucose sensors that generate a signal for induction of gene expression. *Proc. Natl. Acad. Sci. USA* **93**, 12428-12432.
- Pao, S. S., Paulsen, I. T., and Saier, M. H.** (1998). Major facilitator superfamily. *Microbiol. Mol. Biol. Rev.* **62**, 1-34.
- Pilgrim, M. L., Caspar, T., Quail, P. H., and McClung, C. R.** (1993). Circadian and light-regulated expression of nitrate reductase in *Arabidopsis*. *Plant Mol. Biol.* **23**, 349-364.
- Quackenbush, J., Cho, J., Lee, D., Liang, F., Holt, I., Karamycheva, S., Parvizi, B., Perte, G., Sultana, R., and White, J.** (2001). The TIGR Gene Indices: analysis of gene transcript sequences in highly sampled eukaryotic species. *Nucl. Acids Res.* **29**, 159-164.
- Quesada, A., Galvan, A., and Fernandez, E.** (1994). Identification of nitrate transporter genes in *Chlamydomonas reinhardtii*. *Plant J.* **5**, 407-419.
- Rastogi, R., Back, E., Schneiderbauer, A., Bowsher, C., Moffatt, B., and Rothstein, S. J.** (1993). A 330 bp region of the spinach nitrite reductase gene promoter directs nitrate-inducible tissue specific expression in transgenic tobacco. *Plant J.* **4**, 317-326.
- Rastogi, R., Bate, N., Sivasankar, S., and Rothstein, S. J.** (1997). Footprinting of the spinach nitrite reductase gene promoter reveals the preservation of nitrate regulatory elements between fungi and higher plants. *Plant Mol. Biol.* **34**, 465-476.
- Rawat, S. R., Hunt, T., and Glass, A. D. M.** (1997). Regulation of ammonium transport in *Arabidopsis thaliana*: physiology and molecular analysis. *Plant Physiol.* **114**, 1283.
- Rawat, S. R., Silim, S. N., Kronzucker, H. J., Siddiqi, M. Y., and Glass, A. D. M.** (1999). AtAMT1 gene expression and NH<sub>4</sub><sup>+</sup> uptake in roots of *Arabidopsis thaliana*: evidence for regulation by root glutamine levels. *Plant J.* **19**, 143-152.
- Redinbaugh, M. G., and Campbell, W. H.** (1991). Higher plant responses to environmental nitrate. *Physiol. Plant.* **82**, 640-650.
- Redinbaugh, M. G., and Campbell, W. H.** (1993). Glutamine synthetase and ferredoxin-dependent glutamate synthase expression in the maize (*Zea mays*) root primary response to nitrate. *Plant Phys.* **101**, 1249-1255.
- Redinbaugh, M. G., and Campbell, W. H.** (1998). Nitrate regulation of the oxidative pentose phosphate pathway in maize (*Zea mays* L.) root plastids: induction of 6-phosphogluconate dehydrogenase activity, protein and transcript levels. *Plant Science* **134**, 129-140.
- Rentsch, D., Laloi, M., Rouhara, I., Schmelzer, E., Delrot, S., and Frommer, W. B.** (1995). *NRT1* encodes a high affinity oligopeptide transporter in *Arabidopsis*. *FEBS Lett.* **370**, 264-268.
- Riechmann, J. L., Heard, J., Martin, G., Reuber, L., Jiang, C. Z., Keddie, J., Adam, L., Pineda, O., Ratcliffe, O. J., Samaha, R. R., and al., e.** (2000). *Arabidopsis* transcription factors: Genome-wide comparative analysis among eukaryotes. *Science* **290**, 2105-2110.

- Ritchie, S. W., Redinbaugh, M. G., Shiraishi, N., Vrba, J. M., and Campbell, W. H.** (1994). Identification of a maize root transcript expressed in the primary response to nitrate - Characterization of a cDNA with homology to Ferredoxin-NADP(+) oxidoreductase. *Plant Molecular Biology* **26**, 679-690.
- Rothstein, S. J., and Sivasankar, S.** (1999). Nitrate inducibility of gene expression using the nitrite reductase gene promoter. In *Inducible Gene Expression in Plants*, P. H. S. Reynolds, ed CAB International), pp.
- Saier, M. H.** (2000). A functional-phylogenetic classification system for transmembrane solute transporters. *Microbiol. Mol. Biol. Rev.* **64**, 354-411.
- Saier, M. H., Eng, B. H., Fard, S., Garg, J., Haggerty, D. A., Hutchinson, W. J., Jack, D. L., Lai, E. C., Liu, H. J., Nusinew, D. P., and al., e.** (1999). Phylogenetic characterization of novel transport protein families revealed by genome analyses. *Biochim. Biophys. Acta-Biomembr.* **1422**, 1-56.
- Sakakibara, H., Kobayashi, K., Deji, A., and Sugiyama, T.** (1997). Partial characterization of the signaling pathway for the nitrate-dependent expression of genes for nitrogen-assimilatory enzymes using detached maize leaves. *Plant and Cell Physiology* **38**, 837-843.
- Sakakibara, H., Takei, K., and Sugiyama, T.** (1996). Isolation and characterization of a cDNA that encodes maize uroporphyrinogen III methyltransferase, an enzyme involved in the synthesis of siroheme, which is a prosthetic group of nitrite reductase. *Plant J.* **10**, 883-892.
- Sakakibara, H., Taniguchi, M., and Sugiyama, T.** (2000). His-Asp phosphorelay signaling: a communication avenue between plants and their environment. *Plant Mol. Biol.* **42**, 273-278.
- Scheible, W. R., GonzalezFontes, A., Lauerer, M., MullerRober, B., Caboche, M., and Stitt, M.** (1997b). Nitrate acts as a signal to induce organic acid metabolism and repress starch metabolism in tobacco. *Plant Cell* **9**, 783-798.
- Scheible, W. R., GonzalezFontes, A., Morcuende, R., Lauerer, M., Geiger, M., Glaab, J., Gojon, A., Schulze, E.-D., and Stitt, M.** (1997a). Tobacco mutants with a decreased number of functional nia genes compensate by modifying the diurnal regulation of transcription, post-translational modification and turnover of nitrate reductase. *Planta* **203**, 304-319.
- Scheible, W. R., Lauerer, M., Schulze, E. D., Caboche, M., and Stitt, M.** (1997c). Accumulation of nitrate in the shoot acts as a signal to regulate shoot-root allocation in tobacco. *Plant J.* **11**, 671-691.
- Schulze, W., Schulze, E. D., Stadler, J., Heilmeier, H., Stitt, M., and Mooney, H. A.** (1994). Growth and reproduction of *Arabidopsis thaliana* in relation to storage of starch and nitrate in the wild-type and in starch- deficient and nitrate-uptake-deficient mutants. *Plant Cell Environ.* **17**, 795-809.
- Schwarz, G., Schulze, J., Bittner, B., Eilers, T., Kuper, J., Bollmann, G., Nerlich, A., Brinkmann, H., and Mendel, R. R.** (2000). The molybdenum cofactor biosynthetic protein Cnx1 complements molybdate-repairable mutants, transfers molybdenum to the metal binding pterin, and is associated with the cytoskeleton. *Plant Cell* **12**, 2455-2471.
- Shelden, M. C., Dong, B., de Bruxelles, G. L., Trevaskis, B., Whelan, J., Ryan, P. R., Howitt, S. M., and Udvardi, M. K.** (2001). Arabidopsis ammonium transporters, AtAMT1;1 and AtAMT1;2, have different biochemical properties and functional roles. *Plant and Soil* **231**, 151-160.
- Siewe, R. M., Weil, B., Burkovski, A., Eikmanns, B. J., Eikmanns, M., and Kramer, R.** (1996). Functional and genetic characterization of the (methyl)ammonium uptake carrier of *Corynebacterium glutamicum*. *J. Biol. Chem.* **271**, 5398-5403.
- Smeekens, S.** (2000). Sugar-induced signal transduction in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **51**, 49-81.
- Sohlenkamp, C., Shelden, M. C., Howitt, S. M., and Udvardi, M. K.** (2000). Characterization of Arabidopsis AtAMT2, a novel ammonium transporter in plants. *FEBS Lett.* **467**, 273-278.
- Song, W., Steiner, H. Y., Zhang, L., Naider, F., Stacey, G., and Becker, J. M.** (1996). Cloning of a second Arabidopsis peptide transport gene. *Plant Physiol.* **110**, 171-178.
- Spalding, E. P., Hirsch, R. E., Lewis, D. R., Qi, Z., Sussman, M. R., and Lewis, B. D.** (1999). Potassium uptake supporting plant growth in the absence of AKT1 channel activity - Inhibition by ammonium and stimulation by sodium. *Journal of General Physiology* **113**, 909-918.
- Stallmeyer, B., Nerlich, A., Schiemann, J., Brinkmann, H., and Mendel, R. R.** (1995). Molybdenum co-factor biosynthesis-the *Arabidopsis thaliana* cDNA *CNX1* encodes a multifunctional two-domain protein homologous to a mammalian neuroproteins, the insect protein *Cinnamon* and three *Eschericia coli* proteins. *Plant J.* **8**, 751-762.
- Steiner, H. Y., Naider, F., and Becker, J. M.** (1995). The PTR family—A new group of peptide transporters. *Mol. Microbiol.* **16**, 825-834.
- Stitt, M.** (1999). Nitrate regulation of metabolism and growth. *Curr. Opin. Plant Biol.* **2**, 178-186.
- Sueyoshi, K., Mitsuyama, T., Sugimoto, T., Kleinhofs, A., Warner, R. L., and Oji, Y.** (1999). Effects of inhibitors for signaling components on the expression of the genes for nitrate reductase and nitrite reductase in excised barley leaves. *Soil Sci. Plant Nutr.* **45**, 1015-1019.
- Tang, P. S., and Wu, H. Y.** (1957). Adaptive formation of nitrate reductase in rice seedlings. *Nature* **179**, 1355-1356.
- Taniguchi, M., Kiba, T., Sakakibara, H., Ueguchi, C., Mizuno, T., and Sugiyama, T.** (1998). Expression of Arabidopsis response regulator homologs is induced by cytokinins and nitrate. *FEBS Letters* **429**, 259-262.
- Thomas, G. H., Mullins, J. G. L., and Merrick, M.** (2000). Membrane topology of the Mep/Amt family of ammonium transporters. *Mol. Microbiol.* **37**, 331-344.
- Thompson, J. D., Higgins, D. G., and Gibson, T. J.** (1994). Clustal-W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl. Acids Res.* **22**, 4673-4680.
- Tischner, R.** (2000). Nitrate uptake and reduction in higher and lower plants. *Plant Cell Envir.* **23**, 1005-1024.
- Tischner, R., Waldeck, B., Goyal, S., and Rains, W. D.** (1993). Effect of nitrate pulses on the nitrate-uptake rate, synthesis of mRNA coding for nitrate reductase, and nitrate reductase activity in the roots of barley seedlings. *Planta* **189**, 533-537.



- Touraine, B., and Glass, A. D. M.** (1997).  $\text{NO}_3^-$  and  $\text{ClO}_3^-$  fluxes in the *chl1-5* mutant of *Arabidopsis thaliana*-Does the *CHL1-5* gene encode a low-affinity  $\text{NO}_3^-$  transporter? *Plant Physiol.* **114**, 137-144.
- Trueman, L. J., Onyeocha, I., and Forde, B. G.** (1996). Recent advances in the molecular biology of a family of eukaryotic high affinity nitrate transporters. *Plant Phys. Biochem.* **34**, 621-627.
- Trueman, L. J., Richardson, A., and Forde, B. G.** (1996). Molecular cloning of higher plant homologues of the high-affinity nitrate transporters of *Chlamydomonas reinhardtii* and *Aspergillus nidulans*. *Gene* **175**, 223-231.
- Tsay, Y.-F., Schroeder, J. I., Feldmann, K. A., and Crawford, N. M.** (1993a). A herbicide sensitivity gene *CHL1* of *Arabidopsis* encodes a nitrate-inducible nitrate transporter. *Cell* **72**, 705-713.
- Unkles, S. E., Hawker, K. L., Grieve, C., Campbell, E. I., Montague, P., and Kinghorn, J. R.** (1991). *crnA* encodes a nitrate transporter in *Aspergillus nidulans*. *Proc. Natl. Acad. Sci. USA* **88**, 204-208.
- Vale, F. R., Jackson, W. A., and Volk, R. J.** (1987). Potassium influx into maize root systems: influence of root potassium concentration and ambient ammonium. *Plant Physiol.* **84**, 1416-1420.
- Von Heijne, G.** (1992). Membrane protein structure prediction: hydrophobicity analysis and the positive-inside rule. *J. Mol. Biol.* **225**, 487-494.
- von Wiren, N., Gazzarrini, S., Gojon, A., and Frommer, W. B.** (2000a). The molecular physiology of ammonium uptake and retrieval. *Curr. Opin. Plant Biol.* **3**, 254-261.
- von Wiren, N., Gojon, A., Chaillou, S., and Raper, D.** (2001). Mechanisms and regulation of ammonium uptake in higher plants. In *Plant Nitrogen*, P. J. Lea and J.-F. Morot-Gaudry, ed (Heidelberg: Springer), pp. 61-77.
- von Wiren, N., Lauter, F. R., Ninnemann, O., Gillissen, B., Walch-Liu, P., Engels, C., Jost, W., and Frommer, W. B.** (2000b). Differential regulation of three functional ammonium transporter genes by nitrogen in root hairs and by light in leaves of tomato. *Plant J.* **21**, 167-175.
- Walch-Liu, P., Neumann, G., Bangerth, F., and Engels, C.** (2000). Rapid effects of nitrogen form on leaf morphogenesis in tobacco. *J. Exper. Bot.* **51**, 227-237.
- Wang, M. Y., Siddiqi, M. Y., Ruth, T. J., and Glass, A. D. M.** (1993). Ammonium uptake by rice roots. II. Kinetics of  $^{13}\text{NH}_4^+$  influx across the plasmalemma. *Plant Physiol.* **125**, 1259-1267.
- Wang, R., and Crawford, N. M.** (1996). Genetic identification of a gene involved in constitutive, high affinity, nitrate transport in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **93**, 9297-9301.
- Wang, R., Guegler, K., LaBrie, S. T., and Crawford, N. M.** (2000). Genomic analysis of a nutrient response in *Arabidopsis* reveals diverse expression patterns and novel metabolic and potential regulatory genes that are induced by nitrate. *Plant Cell* **12**, 1491-1510.
- Wang, R., Liu, D., and Crawford, N. M.** (1998). The *Arabidopsis CHL1* protein plays a major role in high affinity nitrate uptake. *Proc. Natl. Acad. Sci. USA* **95**, 15134-15139.
- Warner, R. L., and Huffaker, R. C.** (1989). Nitrate transport is independent of NADH and NAD(P)H nitrate reductases in barley seedlings. *Plant Physiol.* **91**, 947-953.
- Warner, R. L., and Kleinhofs, A.** (1992). Genetics and molecular biology of nitrate metabolism. *Physiol. Plant.* **85**, 245-252.
- White, P. J.** (1996). The permeation of ammonium through a voltage-independent  $\text{K}^+$  channel in the plasma membrane of rye roots. *J. Membr. Biol.* **152**, 89-99.
- Wilkinson, J. Q., and Crawford, N. M.** (1991). Identification of the *Arabidopsis CHL3* gene as the nitrate reductase structural gene *NIA2*. *Plant Cell* **3**, 461-471.
- Wilkinson, J. Q., and Crawford, N. M.** (1993). Identification and characterization of a chlorate resistant mutant of *Arabidopsis* with mutations in both *NIA1* and *NIA2* nitrate reductase structural genes. *Mol. Gen. Genet.* **239**, 289-297.
- Williams, L. E., and Miller, A. J.** (2001). Transporters responsible for the uptake and partitioning of nitrogenous solutes. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **52**, 659-688.
- Zhang, H., and Forde, B. G.** (2000). Regulation of *Arabidopsis* root development by nitrate availability. *J. Exp. Bot.* **51**, 51-59.
- Zhang, H. M., and Forde, B. G.** (1998). An *Arabidopsis* MADS box gene that controls nutrient-induced changes in root architecture. *Science* **279**, 407-409.
- Zhang, H. M., Jennings, A., Barlow, P. W., and Forde, B. G.** (1999). Dual pathways for regulation of root branching by nitrate. *Proc. Natl. Acad. Sci. USA* **96**, 6529-6534.
- Zhou, J.-J., Fernandez, E., Galvan, A., and Miller, A.** (2000a). A high affinity nitrate transport system from *Chlamydomonas* requires two gene products. *FEBS Lett.* **466**, 225-227.
- Zhou, J. J., Theodoulou, F. L., Muldin, I., Ingemarsson, B., and Miller, A. J.** (1998). Cloning and functional characterization of a *Brassica napus* transporter that is able to transport nitrate and histidine. *J. Biol. Chem.* **273**, 12017-12023.
- Zhou, J. J., Trueman, L. J., Boorer, K. J., Theodoulou, F. L., Forde, B. G., and Miller, A. J.** (2000b). A high affinity fungal nitrate carrier with two transport mechanisms. *J. Biol. Chem.* **275**, 39894-39899.
- Zhuo, D., Okamoto, J., Vidmar, J. J., and Glass, A. D. M.** (1999). Regulation of a putative high-affinity nitrate transporter (*Nrt2;1At*) in roots of *Arabidopsis thaliana*. *Plant J.* **17**, 563-568.
- Zimmer, W., and Mendel, R.** (1999). Molybdenum metabolism in plants. *Plant Biol.* **1**, 160-168.