

RESEARCH ARTICLE

Open Access

# DOF-binding sites additively contribute to guard cell-specificity of *AtMYB60* promoter

Eleonora Cominelli<sup>1,4\*</sup>, Massimo Galbiati<sup>1,2</sup>, Alessandra Albertini<sup>1</sup>, Fabio Fornara<sup>3,5</sup>, Lucio Conti<sup>1,2</sup>, George Coupland<sup>3</sup> and Chiara Tonelli<sup>1\*</sup>

## Abstract

**Background:** We previously demonstrated that the *Arabidopsis thaliana* AtMYB60 protein is an R2R3MYB transcription factor required for stomatal opening. *AtMYB60* is specifically expressed in guard cells and down-regulated at the transcriptional levels by the phytohormone ABA.

**Results:** To investigate the molecular mechanisms governing *AtMYB60* expression, its promoter was dissected through deletion and mutagenesis analyses. By studying different versions of *AtMYB60* promoter::GUS reporter fusions in transgenic plants we were able to demonstrate a modular organization for the *AtMYB60* promoter. Particularly we defined: a minimal promoter sufficient to confer guard cell-specific activity to the reporter gene; the distinct roles of different DOF-binding sites organised in a cluster in the minimal promoter in determining guard cell-specific expression; the promoter regions responsible for the enhancement of activity in guard cells; a promoter region responsible for the negative transcriptional regulation by ABA. Moreover from the analysis of single and multiple mutants we could rule out the involvement of a group of DOF proteins, known as CDFs, already characterised for their involvement in flowering time, in the regulation of *AtMYB60* expression.

**Conclusions:** These findings shed light on the regulation of gene expression in guard cells and provide new promoter modules as useful tools for manipulating gene expression in guard cells, both for physiological studies and future biotechnological applications.

## Background

Land plants uptake carbon dioxide for photosynthesis and lose water vapour by transpiration through stomatal pores, present on the surface of leaves and stems. The opening and closure of the pore is mediated by turgor-driven volume changes of two surrounding guard cells, whose pressure is dynamically adjusted according to environmental and hormonal cues. In response to abiotic stresses, such as drought or high salinity, one of the most rapid responses of plants is the closure of stomata, mediated by the hormone abscisic acid (ABA), to prevent excessive water loss by transpiration (reviewed in [1]).

The genetic manipulation of stomatal activity is emerging as a promising approach to reduce the water requirement of crops, and to enhance productivity under stress conditions [2]. Proper engineering of

stomatal responses requires the use of guard cell-specific promoters, or the identification of guard cell-specific mutants, to avoid undesirable side effects on plant growth and productivity.

Several promoters that confer guard cell-specific gene expression or enhanced gene expression in guard cells have been isolated through different methods: functional characterization of single genes [3-9]; large scale gene- or enhancer-trap screens [10-12]. Moreover transcriptional and proteomic studies have identified additional candidates [13-16]. Nevertheless the majority of these promoters are not guard cell-specific, as they drive the expression of reporter genes in other cell types, including the vascular tissues [6,10,17,18], flower organs [8,9] or starch containing cells [5], significantly reducing the number of true guard cell-specific full size promoters [3,10,14,19,20]. Most importantly, a detailed experimental analysis of guard cell-specific promoters has been performed only in very few cases [11,12,14].

\* Correspondence: cominelli@ibba.cnr.it; chiara.tonelli@unimi.it

<sup>1</sup>Dipartimento di Scienze Biomolecolari e Biotecnologie, Università degli Studi di Milano, Milano, Italy

Full list of author information is available at the end of the article

A true guard cell-specific promoter is driving expression of the Arabidopsis *AtMYB60* (*At1g08810*) gene [10,19,21,22]. We have previously shown that *AtMYB60* is expressed in guard cells [10], and the complete 5' and 3' intergenic genomic regions of this gene, cloned respectively upstream and downstream to reporter genes, were able to drive specific expression in guard cells [10,19]. Guard cell specificity of the *AtMYB60* promoter has been also demonstrated by Nagy *et al.* (2009) and by Meyer *et al.* (2010), who used this promoter to complement the *mrp5-1* mutant phenotype exclusively in guard cells, and to specifically express the AtLMT12 protein at high levels in guard cells, respectively.

Very little information is available concerning promoter *cis*-elements regulating guard cell-specific expression [8,10-12,14,16]. DOF-binding sites have been suggested to have a role in such a regulation [8,10-12]. DOF (DNA binding with One Finger) proteins are plant specific transcription factors involved in light, phytohormones and pathogen signalling and responses as well as seed development (reviewed by [23]). A role for [T/A]AAAG DOF-binding sites in mediating gene expression in guard cells has been experimentally defined only for the potato *KST1* gene [8]. However, in Arabidopsis the role of DOF-motifs in controlling guard cell expression is still controversial [10-12]. The study performed on the potato *KST1* promoter [8] and the bioinformatic analysis performed on several guard-cell specific Arabidopsis promoters [10] suggest that the presence of clusters of DOF *cis*-elements, rather than their absolute number, is important to confer guard cell-specificity to a promoter region [10]. Yet, the role of DOF-binding sites in driving guard cell expression in Arabidopsis and the hypothesis of cluster organization remains to be experimentally investigated.

The guard-cell specific *AtMYB60* promoter presents several DOF clusters, making it an ideal model to test the hypothesis that DOF clusters are important for guard cell-specific expression. Moreover the *AtMYB60* expression is modulated by different environmental cues such as light, dark and drought stress [19], suggesting the presence of different *cis*-elements controlling these transcriptional responses. In this report we aimed to isolate the *cis*-elements responsible for the *AtMYB60* guard cell specific expression. We generated Arabidopsis transgenic lines carrying truncated or mutagenised *AtMYB60* promoter versions fused to the *GUS* reporter gene. Using a combination of histochemical and expression analysis we were able to identify a minimal promoter necessary and sufficient to drive guard cell specific expression. Using the same tools, we were also able to map a region required for ABA-mediated repression.

## Results

### *In-silico* analysis of the *AtMYB60* promoter

In a previous study, we demonstrated that the complete 5' and 3' *AtMYB60* intergenic genomic regions - cloned upstream and downstream of the  $\beta$ -glucuronidase (*GUS*) reporter gene, respectively - could specifically drive strong *GUS* activity in stomata of Arabidopsis seedlings and adult plants [19]. No *GUS* signals were detected in any other cell type or in tissues devoid of stomata [19].

To investigate the possible *cis*-acting elements that regulate *AtMYB60* expression, we surveyed the genomic region upstream of the *AtMYB60* translational start codon for the presence of known transcription factor binding sites using the PLACE software [24]. Our analysis produced a significant enrichment in the [A/T]AAAG motifs in the *AtMYB60* promoter compared to the average distribution of [A/T]AAAG oligos in intergenic regions throughout the Arabidopsis genome ( $P < 0.01$ ) (Figure 1). Interestingly, these [A/T]AAAG motifs, have been shown to be involved in the regulation of guard cell expression of the potato potassium channel *KST1* gene [8]. Also, clusters of [A/T]AAAG motifs, required for the binding of DOF-type transcription factors [25], were over represented in different guard cell-specific promoters [6,10,12]. In particular, Galbiati and colleagues suggested, as guard cell-specific *cis*-element, a cluster of at least three [A/T]AAAG motifs located on the same strand within a region of 100 bp [10]. Using the criteria previously described by Galbiati and collaborators (2008), we found three of these guard cell-specific clusters in the 5' intergenic region of the *AtMYB60* gene (Figure 1), suggesting a conserved mechanism for guard cell specific expression.

### Identification of the *AtMYB60* minimal promoter

To gain more insights into the *cis*-elements that regulate the *AtMYB60* expression in guard cells, we produced a set of Arabidopsis transgenic lines carrying the complete 1,307 bp 5' intergenic region upstream of the translational start codon fused to the reporter *GUS* (construct -1,307::*GUS*, Figure 2A). *GUS* staining analysis of 15 independent T2 lines revealed that this region contains all the *cis*-acting elements required for expression of the reporter in stomata (Figure 2B), while no *GUS* signals were detected in any other cell type or in tissues devoid of stomata (Additional file 1).

Next, we made a series of 5' deletions of the -1,307 bp genomic region to define the minimum sequence length required for the expression in guard cells (Figure 2A). These truncated promoters (fused to the *GUS* gene) were stably transferred to Arabidopsis and 10 to 15 independent T2 transgenic lines were analysed in detail. Deletions of the distal part of the 1,307 bp region to

```
-1307 CACAAGGACACAAGGACATATGGTATGATGATATGCTTTGTTTCTCTGCTTCTCTTACTAATTTGA
-1241 AGCTGTTGGATTGATTTGTCTCTTCTTACGTTCCCTTCTTTTTTTTTCGTTTTCTTTTGTCTGAT
-1175 AGACCAGGCAGGGGCTAGGGCCTAGTGATGGGTATTGGCCAATACTATTGGGTTATTTGCCTGGT
-1109 TTATTATTTTCGATTTTAGGTTAATTCAATTTTAAAGAATACGTAGATTTGTTTGGTTTAGTTGGTT
-1043 TGTTGCACTAAGTTCGGTTTTACATAAATAGAATCTAACACTACTAATTGTTATACGTAAAATAC
-977 AACACAATAACAGATTTTTTCGTTTCAATTTTTTCGTTTAAAGAGGGTAGACATTTTTGGTTTGGTTTGG
-911 TTCATTTTTTTTTTCCCTTTTCAAATTCACATCCTTCACGTAGATGACAAAAATAAAGAAAAACATGA
-845 ATGAAAGTTGTAACCTGTAAGCATCAACATGGAAATCATATCACAAAGAACACAAATCTAACTAAT
-779 GGGTCTTTTACATATTGGTATAATTATAAGTTGTAAGAATATTAGTTAAACAGAGGCAACGAGAG
-713 ATGCGTGATATATGAAAAGTTGAAAACAAAAGACATGGATCTAAAAGAGTCAAGCAAAATGTAATAT
-647 CTTTTTTTCTTCTAAACTTGAGGATGTCCAAGTTGCAGTGAATGATTCCCTTTAATCATGGAGAAA
-581 TTCAATGAAATAATTGTGTTTCTTCCACACTTTATCTTTAATTTATTTTCTTACCACAATTACAAC
-515 TATTATCACAAAAATGTAAGTAACATAGCTTGTGACTCTTCTTCCATTTATGAGTTGATTATCACT
-449 ATATTTATAAGTAATTACCAACGAATGTTCCAAATTAAGCAAAATATTGTAATCGATACACTATGT
-382 ATTCATCTACAATATGTTAACGAGCTCCTTTTATGGAAATATTTTCGATTGAAAAACATTTGATGG
-317 ATCGTTCATAAATAAATAATCCAGTAACGTTTTCTTAAGGGAGATATACATATTCGTGTGGAGAT
-251 CAACATATCTTCGTTAAATTGACTTACGCAAAATAGTTAATGGAAAAGGCAGAGTGACTCGTGAGCTT
-185 GGCAGATCCAAAAGAGGTTGTCAAGAAAAAGCAGATTTAAAAGTTCTTCCCTCTTCTTTAAGTCAC
-119 CCATTAATTTACATATATGTACATACATGTTGCATTTAACTCATATACATACatattctcacatc
-53 tataaagagagcataagactcagagagatctagaggaagagagagagagaaagATG
```

**Figure 1 Nucleotide sequence of the 5'-region of the *AtMYB60* gene.** Nucleotides are numbered on the left with the translational start site designated as +1. The ATG is in bold. The 5' UTR is in lower case letters. The DOF-binding sites are grey boxed, the W-box, considered in the text, is white boxed. Clusters of DOF-binding sites, as defined by Galbiati and colleagues (2008), are underlined. The CAAGTTG motif described as a putative *cis*-element for ABA repression ([16]) is dotted underlined.


position -619 (construct -619::*GUS*), -472 (-472::*GUS*), or -366 (-366::*GUS*) from the ATG codon, did not alter expression of the reporter in guard cells located on both vegetative and floral organs (Figure 2B). Further deletions (to position -262) indicated that the 262 bp proximal region was sufficient to drive expression of the reporter in stomata (Figure 2B). However, the removal of the region between -262 bp and -205 bp (construct -205::*GUS*) completely abolished *GUS* activity in guard cell (Figure 2B). Transgenic lines carrying the -205::*GUS* fusion did not show *GUS* staining in any other cell type, even after prolonged staining (up to 48 h, Figure 2B). This finding suggests that the 57 bp region located between positions -262 and -205 contains *cis*-elements essential for expression in stomatal guard cells. Based on these results, we defined the -262 bp region upstream of the ATG codon as the minimal promoter of the *AtMYB60* gene.

To thoroughly investigate quantitative differences in *GUS* expression among lines carrying different deletion: reporter constructs, we determined the relative amount of *GUS* transcript by quantitative RT-PCR (qRT-PCR). mRNA samples derived from two representative independent lines (A and B) were analysed for each construct (Figure 2C). Lines harbouring the 1,307 bp 5' intergenic region or the -619 deletion fused to the


reporter, did not show any significant differences in their *GUS* transcript accumulation. Conversely, deletions to position -472 and -366 resulted in a two-fold decrease in *GUS* expression compared to the -1,307::*GUS* line, while deletion to position -262 resulted in a five-fold decrease (Figure 2C,  $p < 0.01$ ). These results indicate that one or more sequences with function of enhancer are present in the genomic region between -619 bp and -472 bp and between -472 and -262 from the ATG of *AtMYB60*. In accordance with the results obtained from the histochemical analysis, qRT-PCR experiments did not detect significant *GUS* transcripts accumulation in lines carrying the -205::*GUS* fusion.

#### Site-directed mutagenesis of the *AtMYB60* minimal promoter

Promoter deletion experiments indicate that the *AtMYB60* minimal promoter region (construct -262::*GUS*) contains all the *cis*-acting elements required to sustain expression of a reporter gene in guard cells. This region encompasses the [A/T]AAAAG cluster proximal to the ATG codon, which consists of four AAAAG DOF-binding sites (Figures 1 and 3A). In addition, the PLACE software identified in this region a single W-box, corresponding to the binding site of WRKY transcription factors [26], located upstream of the [A/T]



**Figure 2 Deletion analysis of the *AtMYB60* upstream region.** A, Schematic diagrams of different deletions of the *AtMYB60* upstream region fused to the *GUS* reporter gene. The positions of the different DOF-binding sites and of the W-box, described in the text, are shown. B, Histochemical assay for *GUS* activity in seedlings, rosette leaves and flowers of plants transformed with -1,307::*GUS* (a-d), -619::*GUS* (e-h), -472::*GUS* (i-l), -366::*GUS* (m-p), -262::*GUS* (q-t) and -205::*GUS* (u-x) constructs. The analysis of independent lines harbouring the same construct showed identical patterns of *GUS* staining. Samples were incubated in the staining solution for 16 hours for all the lines, with the exception of line -205::*GUS*, for which the staining was prolonged to 48 hours. Scale bars represent 1 mm. C, Relative expression level of the *GUS* reporter gene in the different transgenic lines harbouring the -1,307::*GUS* (-1,307 A and B), -619::*GUS* (-619 A and B), -472::*GUS*, -366::*GUS* (-366 A and B), -262::*GUS* (-262 A and B) or -205::*GUS* (-205 A and B) constructs. Two lines for each construct were analysed by Real Time RT-PCR. The transcript amount in the line -1,307 A was arbitrarily set to 1 (black column) and used to normalize the relative expression levels in each line. The *ACTIN2* gene (*At3g18780*) was used as a control.



**Figure 3 Role of DOF-binding sites in the minimal promoter in driving GUS activity in guard cells.** A, Schematic diagrams of constructs -262::GUS, -205::GUS and of constructs containing mutagenised version of the minimal promoter in different DOF-binding sites and in the W-box at position -234. B, Percentage of lines for each construct showing strong (column segment in black), intermediate (in dark grey), weak (light grey) or no signal (white). C, A leaf from a line harbouring the -262::GUS construct (a and a particular in e), shown as an example of strong GUS activity. In the following pictures examples of different lines harbouring the *mDOF3*::GUS construct showing respectively an intermediate (b and f), a weak (c and g) and no GUS activity (d and h). Scale bars represent 1 mm (a-d) or 0.1 mm (e-h).



AAAG cluster (Figure 3A). To address the functional significance of the individual *cis*-elements present in the *AtMYB60* minimal promoter, we evaluated the effects of targeted nucleotide substitutions on *GUS* expression (Figure 3A). Mutated versions of the minimal promoter were generated by PCR and fused to *GUS* and at least 30 T2 independent transgenic lines for each mutated promoter::*GUS* combination were visually scored and classified to reflect their relative guard-cell specific *GUS* staining. A representative example of each category is provided in Figure 3C.


We initially tested the role of the single W-box *cis*-element, by replacing the consensus sequence TTGAC, with the non-functional TTGAA motif [27]. Lines carrying the mutated W-box (*mW*::*GUS*) showed similar levels of *GUS* expression to the wild-type promoter, indicating that W-box does not contribute to mediate gene expression in guard cells (Figure 3B). Next, we produced mutant promoters in which single DOF motifs within the [A/T]AAAG cluster were converted to the unrelated CGCGA sequence. Inactivation of the most distal AAAAG site relative to the ATG (hereinafter referred to as *DOF1*) resulted in a dramatic decrease of *GUS* expression (*mDOF1*::*GUS* construct, Figure 4B). 30% of the lines carrying the *mDOF1*::*GUS* construct did not show *GUS* expression, whereas the remaining 70% only showed weak staining, thus indicating a crucial role for *DOF1* in regulating *AtMYB60* expression in guard cells (Figure 3B). Mutations of the second, third or fourth most proximal AAAAG site (hereinafter referred to as *DOF2*, *DOF3* and *DOF4*, respectively), resulted in a reduced *GUS* expression, although to a lesser extent than the one in the *DOF1* (Figure 4B, *mDOF2*::*GUS*, *mDOF3*::*GUS* and *mDOF4*::*GUS* plants). In particular, none of the 30 *mDOF2*::*GUS* transgenic lines displayed strong expression of the reporter, nearly 70% showed intermediate expression, 25% showed weak expression and the remaining 5% did not show any *GUS* staining (Figure 3B). A comparable distribution among strong, intermediate and weak lines was obtained from the analysis of the *mDOF3*::*GUS* and *mDOF4*::*GUS* plants (Figure 3B).

To establish whether DOF-binding sites could exert additive roles in mediating gene expression in stomata we produced a second series of promoters, in which two AAAAG motifs were mutated simultaneously. Mutations of *DOF1* and *DOF2* (*mDOF(1+2)*::*GUS*), *DOF1* and *DOF3* (*mDOF(1+3)*::*GUS*) or *DOF1* and *DOF4* (*mDOF(1+4)*::*GUS*) completely inactivated the minimal promoter, as *GUS* expression was abolished in all the *mDOF(1+2)*::*GUS*, *mDOF(1+3)*::*GUS* and *mDOF(1+4)*::*GUS* lines analysed (Figure 3B). Interestingly, the concurrent mutation of *DOF2* and *DOF3* (*mDOF(2+3)*::*GUS*) resulted in a strong, but yet not complete, inactivation of the

promoter activity in guard cells, as 15% of the *mDOF(2+3)*::*GUS* lines displayed weak expression of the reporter in stomata. Likewise, concomitant inactivation of either *DOF2* and *DOF4*, or *DOF3* and *DOF4* did not completely eliminate *GUS* expression in guard cell (Figure 3B). Taken together, these results indicate that the putative [A/T]AAAG DOF-binding sites located in the *AtMYB60* promoter are necessary to mediate its expression in guard cells.

#### A single DOF cluster is sufficient to drive low expression in guard cell

Our deletion analysis of the *AtMYB60* promoter indicates that the 57 bp region between positions -262 and -205 is essential for gene expression in stomatal guard cells (Figure 2). This region contains the *DOF1* *cis*-element required for guard cell expression as shown by mutagenesis analysis results (Figure 3). To establish whether this 57 bp region was sufficient to activate expression in guard cells, we fused one (1x::*GUS* construct), two (2x::*GUS*) and four tandem copies (4x::*GUS*) of the 57 bp fragment to the minimal *CaMV35S* promoter [28] upstream of the *GUS* reporter gene (Figure 4A), effectively reconstructing an artificial DOF cluster containing one, two or four copies of the *DOF1* element. However, we did not observe *GUS* activity in any of the 30 independent stable transformants produced for each construct, even after prolonged staining (data not shown). These data were confirmed by qRT-PCR analysis of independent lines carrying the 4x::*GUS* fusion (Figure 4B), indicating that the multimerisation of the *DOF1* site *per se* is not sufficient to drive gene expression in guard cell. This might derive from an inappropriate organization and/or spatial distribution of the different DOF elements in the context of the minimal promoter. To test this hypothesis we made two 3' deletions of the *AtMYB60* minimal promoter: the -148-3'::*GUS* and -137-3'::*GUS* constructs containing the first three and four DOF-binding sites respectively of the most proximal cluster fused upstream of the minimal *CaMV35S* promoter (Figure 4B). Our initial histochemical analysis did not reveal any *GUS* positive lines (data not shown). To substantiate this result we also performed a qRT-PCR analysis on fifteen independent lines for each construct. Interestingly, eight lines out of fifteen showed a low but significant *GUS* transcript accumulation compared to the full length minimal promoter (Figure 4B). These results suggest that the presence of the cluster containing three or four DOF-binding sites is sufficient to drive *GUS* activity in guard cells, even though at a very low level. This finding implies that other *cis*-elements present downstream of position -137 are required for the full functionality of the minimal promoter.



### The guard cell-related CDF1, CDF2, CDF3 and CDF5 DOF-type transcription factors do not regulate *AtMYB60* expression in stomata

Target mutagenesis experiments of the *AtMYB60* promoter demonstrated that [A/T]AAAG DNA consensus motifs are essential *cis*-acting elements in the regulation of *AtMYB60* expression in guard cells. Consequently, their cognate DOF proteins represent the most likely candidates as *trans*-acting factors. As the Arabidopsis genome contains 36 DOF-coding genes [23], candidate DOF transcription factors involved in the regulation of *AtMYB60* expression should fulfil two criteria: they should be expressed in guard cells and the loss of their gene function should abolish or significantly down-regulate the expression of *AtMYB60* in this cell type.

The *CYCLING DOF FACTOR 1* (*CDF1*, *At5g62430*) gene, involved in the regulation of photoperiodic flowering, has been shown to be highly expressed in the vascular tissue and guard cells [29]. We thus investigated the expression of the *AtMYB60* gene in the loss-of-function

*cdf1-R* allele. As shown in Additional file 2 we did not detect significant differences in the accumulation of *AtMYB60* transcripts in homozygous *cdf1-R* plants compared with the wild type.

It is important to note that in photoperiodic flowering, CDF1 acts redundantly with three other DOF proteins, namely CDF2 (*At5g39660*), CDF3 (*At3g47500*) and CDF5 (*At1g69570*) [30], belonging to the same phylogenetic group II [31]. Similarly to *CDF1*, promoter::GUS analyses revealed that *CDF2*, *CDF3* and *CDF5* are strongly expressed in guard cells.

We thus analysed the expression of *AtMYB60* in single, double, triple and quadruple *cdf* mutants to determine the possible role of these additional candidate CDF proteins. As for *cdf1-R* mutant, the level of expression of *AtMYB60* was not significantly reduced in the *cdf2-1*, *cdf3-1* and *cdf5-1* single mutants (Additional file 2). Likewise, *AtMYB60* expression was not altered in any of the double, triple or quadruple mutant combinations, indicating that, despite their expression in guard

cells, these four CDF proteins are not *trans*-regulators of *AtMYB60* expression in stomata (Additional file 2).

#### Identification of a promoter region that negatively responds to ABA

We previously reported that transcript accumulation of the *AtMYB60* gene is rapidly down-regulated by exogenous applications of the hormone ABA, which plays a fundamental role in regulating gene expression in response to drought stress [19]. To identify the promoter region responsible for the ABA-mediated *AtMYB60* down-regulation, we applied ABA to the previously described transgenic lines harbouring serial deletions of the *AtMYB60* promoter (Figure 2). Quantitative RT-PCR analysis revealed a similar decrease in *GUS* transcript levels in transgenic lines carrying the full length as well as the -619, -472 and -366::*GUS* fusions (Figure 5). The kinetic of down-regulation of the *GUS* transcript was comparable to the one observed for the endogenous gene *AtMYB60* [19], indicating that -619, -472 and -366 promoters maintain the sequences responsible for transcriptional down-regulation by ABA. Also, these results suggest that the CAAGTTG motif, present in the *AtMYB60* promoter between -619 and -613 (dotted underlined in Figure 1), and recently described as over-represented in ABA-repressed genes [16], does not play a significant role in the ABA-dependent repression of *AtMYB60* expression. Rather, qRT-PCR experiments performed on different independent lines carrying the -246::*GUS* construct showed that the minimal promoter

sequence lacks the region responsible for negative regulation by ABA, as these lines did not show changes in *GUS* expression in response to the hormone as shown in Figure 5.


Taken together these data indicate that, although the minimal promoter maintains the *cis*-elements necessary for guard cell expression, it lacks the motifs that mediate the negative regulation by ABA, becoming ABA-insensitive. We can thus conclude that the region between -366 and -262 contains elements necessary for ABA down-regulation.

#### Discussion

Very few guard cell-specific promoters have been described to date [3,10,14,19,20]. Independent studies demonstrated that the *AtMYB60* promoter can be considered guard cell-specific, being sufficient to drive expression of reporter genes specifically in guard cells [19,21]. Moreover this promoter has also been used to complement a mutant phenotype specifically in guard cells [21], and to investigate subcellular localization exclusively in guard cells [22]. In this study we identified the *AtMYB60* minimal promoter that is necessary and sufficient to drive guard cell-specific expression.

#### DOF-binding sites are required for *AtMYB60* guard-cells expression

Our *in silico* analysis identified three DOF site clusters (Figure 1). Initial deletion studies revealed a prominent role for the most proximal DOF cluster (relative to the



**Figure 5** Expression of the *GUS* gene in different transgenic lines in response to ABA treatment. Two lines for each construct shown in Figure 2 were analysed by Real Time RT-PCR. c represent the control samples. The transcript amount in the sample -1307 A control was arbitrarily set to 1 and used to normalize the relative expression levels in each line. The *ACTIN2* gene was used as a control.



ATG start codon). Site-directed mutagenesis showed that the distal most DOF-binding site (*DOF1* at position -210, Figure 3) plays a major role in driving guard cell expression compared to other DOF motifs of the same cluster (*DOF2* at position -176, *DOF3* at -159 and *DOF4* at -147, Figure 3). These other DOF elements play partially additive roles, as clearly demonstrated by the combined mutagenesis of these sites and *DOF1* site which resulted in a drastically reduced GUS activity (Figure 3). DOF-binding sites are thus key determinants in mediating guard cell expression, in accordance with the DOF cluster hypothesis we previously formulated [10]. A suggestion for a similar involvement of DOF *cis*-elements in Arabidopsis derives from the work of Gardner and colleagues (2009) that identified DOF motifs in a region controlling guard cell expression. Other authors identified a region enriched in DOF-binding sites in the guard cell-specific pGC1 promoter, although the mutation of a single DOF site did not impair promoter activity [14]. Interestingly, a DOF cluster organization is present in the promoter of the grape *VvMYB60* gene, a putative ortholog of *AtMYB60*, indicating a conservation of the cluster structure during the evolution among *AtMYB60* orthologs [32]. The results reported by Plesch and colleagues (2001) on the DOF motif organization in the potato *KST1* promoter highlight a more general evolutionary conservation of this module in the control of guard cell-specific activity of promoters.

Although we cannot rule out the possibility that other unknown transcription factors might interact with those same *cis*-elements, DOF factors represent likely candidates as *AtMYB60* regulators. The most parsimonious hypothesis resulting from combining our results indicates that DOF proteins act as positive regulators of *AtMYB60*. The potato StDOF1 protein has been shown to bind *in vitro* to the guard cell specific promoter of *KST1* [8], while no data are available for any Arabidopsis DOF proteins. Among the Arabidopsis DOF genes, *CDF1*, *CDF2*, *CDF3*, and *CDF5* (CDFs) are expressed in guard cells [29]. However, singles and multiple *cdf* mutants show a wild-type pattern of *AtMYB60* expression, ruling out their involvement in *AtMYB60* regulation (Additional file 2). The majority of Arabidopsis DOF genes are expressed in guard cells [33,34] and may thus act redundantly, as already demonstrated among members of this family [30]. All these aspects do not facilitate the identification of obvious candidates as *AtMYB60* regulators. We are trying to identify the DOF genes involved in the regulation of *AtMYB60* by analysis of its expression in mutants of genes preferentially expressed in the guard cells (<http://bbc.botany.utoronto.ca/efp/cgi-bin/efpWeb.cgi>[33]).

#### Multiple *cis*-elements participate to enhance *AtMYB60* guard-cells expression

Transcriptional *GUS* fusions, harbouring different deletions of the 5' intergenic region to position -262 from the ATG, conferred GUS activity exclusively in guard cells (Figures 2 and Additional file 1). The activity of these promoter regions is in apparent discrepancy with the detection of *AtMYB60* gene expression in seeds, as revealed by available microarray analysis data [33,34] and in roots, as recently reported [35]. One hypothesis to explain this incongruity could be the presence of other regulatory regions present outside the complete 5' and 3' intergenic regions flanking the *AtMYB60* coding sequence. Intron sequences, for example, may be involved in such a regulation, as previously demonstrated for different plant genes ([36] and references herein).

While guard-cell specific expression was invariably maintained by functional *AtMYB60* promoter variants, the levels of expression varied considerably. In addition to DOF-binding sites, other *cis*-elements are required to boost the *AtMYB60* expression. Indeed, an artificial *DOF1* binding site repeated in single or multiple copies could not drive guard cell expression (Figure 4A). The incorporation of the entire proximal DOF cluster (e.g. -137-3'::GUS) resulted in a small but significant guard cell transcriptional activity. Thus, other *cis*-elements downstream of position -137 are required for full activity of the minimal promoter. It is known that *cis*-elements other than DOF-binding sites are involved in the regulation of guard cell expression. In the case of the guard cell-specific *AtPDR3* gene no [A/T]AAAG clusters were identified in a 1000-bp region upstream of the ATG codon, suggesting the presence of other regulatory units [10].

#### Modular organization of the *AtMYB60* promoter

In this study we also investigated the regulation of the *AtMYB60* promoter activity in response to ABA. ABA treatments induce global changes in gene expression in Arabidopsis [16,37-40]. Transcriptomic analyses revealed extensive regulation of gene expression by ABA also in guard cells [13,14,16]. While *cis*-elements that positively regulate the response to ABA have been functionally characterised (for a review, see [41]), those that negatively regulate the response to ABA are largely unknown. A CAA[G/C]TTG motif has been shown to be over-represented in ABA-repressed gene promoters and thus proposed for such a role [16,39]. The *AtMYB60* promoter contains one CAAGTTG motif between -619 and -613 from the ATG, yet our results do not support its proposed role as negative regulator of ABA response. Conversely, a region between positions -366 to -262 contained the entire requirement for the

ABA-mediated repression Figures 5 and 6. It has been proposed that evolution may have favoured the differentiation of mechanisms for ABA down-regulation rather than up-regulation, rendering more difficult for any ABA-repression motif to achieve statistical significance [16]. Our data may provide a valuable model system to clarify the mechanism mediating ABA repression.

Our data suggests a modular organization for the *AtMYB60* promoter as summarised in Figure 6. Through a serial deletion analysis, we defined the *AtMYB60* minimal promoter, sufficient to induce guard cell-specific activity (construct -262::GUS, Figure 2). A 57 bp region, located between position -262 and position -205, is necessary to confer GUS activity in guard cells (Figure 2A). We also identified two regions that enhance the expression of the *GUS* gene between -619 bp and -472 bp and between -472 and -262 (Figure 2B and 2C). Besides providing pieces of evidence for such modular organization, our work indicates that the different portions of the *AtMYB60* promoter may prove useful for manipulating gene expression in guard cells, with the possibility to obtain different level of expression. Moreover, the minimal promoter (whose activity is not influenced by ABA) can be used for ABA-independent expression of target genes in guard cells

Interestingly, both the full length and the minimal promoters maintain their guard cell-specific activity in heterologous systems, such as the crop species tomato and tobacco (Francia, personal communication), thus indicating the conservation of this cell-specific regulatory mechanism among different plant species. Moreover, preliminary results suggest that the *AtMYB60* minimal promoter can be combined with other *cis*-regulatory modules to produce functional guard cell-specific chimeric promoters (Francia, personal communication). As a whole our data demonstrate that both the full length and the minimal *AtMYB60* promoters provide a valuable tool to manipulate gene expression specifically in guard cells, both for

physiological studies and downstream biotechnological applications.

## Conclusions

Our work provides strong evidence for the involvement of [A/T]AAAG elements in the regulation of the *AtMYB60* expression, illustrating their functional cluster organization. Future work will concentrate on the analysis of candidate DOF transcription factors that control this mechanism. Finally we identify a region of the *AtMYB60* promoter required for the negative regulation by ABA, offering the possibility to discover novel *cis*-elements for this kind of regulation.

## Methods


### Plant Material

All plant material described was in the Col-0 accession. The *cdf1-R* line (35S::CDF1-RNAi #23) was kindly provided by Takato Imaizumi [29]. The *cdf2-1*, *cdf3-1* and *cdf5-1* null alleles are T-DNA insertion line. Single, double, triple and quadruple *cdf* mutants have been previously described [30].

### Construction of *AtMYB60* promoter::GUS fusions

5'-deletions of the 5' intergenic genomic region upstream of the *AtMYB60* gene were generated by PCR amplification from plasmid p1.3-2.2::GUS, previously described [19], using different forward primers and a single reverse primer. Forward and reverse primers incorporated a *Hind*III and a *Bam*HI, respectively. The PCR fragments were cloned into the pCR4-TOPO vector (Invitrogen Corporation, Carlsbad, CA), cut with *Hind*III and *Bam*HI and ligated upstream of the *uidA* coding sequence in the pBI101.3 binary vector (Clontech, Palo Alto, CA, USA). The resulting plasmids were renamed -1307::GUS, -619::GUS, -472::GUS, -366::GUS, -262::GUS and -205::GUS (Figure 2).

Chimeric promoters containing different 3'-deleted fragments of the *AtMYB60* minimal promoter and 46-



**Figure 6 Modular organization of the *AtMYB60* promoter.** Different portions of the *AtMYB60* promoter defined through deletion analysis are shown. ABA indicates the region responsible for the negative regulation by ABA treatment.

bp CaMV 35S promoter were produced by amplifying the sequence of the CaMV 35S promoter from -46 to +1 [28] from plasmid pBI121 (Clontech, Palo Alto, CA, USA), using the forward primer 35SXba containing a *Xba*I site and the reverse primer 35SBam with a *Bam*HI site. The PCR product was cloned into the pCR4-TOPO vector and the *Xba*I-*Bam*HI fragment was cloned into the pBI101.3 vector (renamed 35Smin-pBI101.3). The regions from -262 to -137 and from -262 to -148 of the *AtMYB60* minimal promoter were amplified by PCR from plasmid p1.3-2.2::GUS, using the reverse primers p60R6 and p60R7 incorporating a *Xba*I site and a single forward primer p60F3 with the *Hind*III site. The corresponding PCR products were cloned into the pCR4-TOPO vector and the *Hind*III-*Xba*I fragments were cloned into the 35Smin-pBI101.3 vector to give the -137-3'::GUS and -148-3'::GUS vectors, respectively (Figure 3).

Chimeric promoters containing different copies of the region between -262 and -205 of the *AtMYB60* promoter were obtained by synthesising one copy of this sequence, using the forward primer p60F3 with a *Hind*III site and the reverse primer p60R3 with an *Xba*I site. The resulting PCR product was cloned into the pCR4-TOPO vector and the *Hind*III-*Xba*I fragment was ligated into the 35Smin-pBI101.3 vector (construct 1x::GUS). A second copy of this region was generated using the primers p60F3 and p60R5b, both incorporating a *Hind*III site; the fragment *Hind*III-*Hind*III was cloned into the construct 1x::GUS, generating the construct 2x::GUS. This plasmid was used as a template to generate two other copies of the sequence from -262 to -205 using the primers p60F11 and p60R3 incorporating an *Xba*I site. The fragment *Xba*I-*Xba*I was cloned into the plasmid 2x::GUS, to generate the construct 4x::GUS. All the oligonucleotide sequences are reported in Table 1. PCR products were sequenced and the correct orientation of the fragment into the final vector was verified by restriction.

#### Site-directed mutagenesis analysis

Base mutations of the different DOF sites were generated using the megaprimer method [42]. For the mutagenised versions of the *AtMYB60* minimal promoter different megaprimers were PCR amplified from plasmid p1.3-2.2::GUS, using as forward primers mp60DOF1F1, mp60DOF2F1, mp60DOF3F2 and mp60DOF4F2 and the single reverse primer p60R5. The megaprimers were gel purified and used in a second PCR reaction on plasmid p1.3-2.2::GUS with the primer p60F3. The PCR products were cloned into pCR4-TOPO and sequenced before cloning into pBI101.3 vector using the restriction sites *Hind*III and *Bam*HI to generate the following constructs: *mDOF1*::GUS, *mDOF2*::GUS, *mDOF3*::GUS,

**Table 1 Sequence of oligonucleotides used in this study**

Name	Sequence
p60F1	AAGCTTCACAAGGACACAAGGACA
p60 F2b	AAGCTTCAAGTTGCAGTGAATGA
p60F8b	AAGCTTAAACGAGCTCCTTTATGG
p60F9	AAGCTTCCATTTATGAGTTGATTATCA
p60F3	AAGCTTCGTGTGGAGATCAACAT
p60F5	AAGCTTGCAGAGTGACTCGTGA
p60R5	TCTCGGATCCTCTAGATCTCTCTG
p60R6	TCTAGAGAAGAACCTTTTAAATCTGC
p60R7	TCTAGAAAATCTGCTTTTCTTGAC
p60R5b	AAGCTTCTTTCCATTAACATATTTTG
p60F11	TCTAGACGTGTGGAGATCAACAT
p60R3	TCTAGACTTTCCATTAACATATTTTG
35SXba	TCTAGACAAGACCCTTCCTC
35SBam	GGATCCTCCTCTCCAATGA
mp60DOF1F1	AGTTAATGGcgcgaGCAGAGTGACTCGTGA
mp60DOF2F1	TGGCAGATCCcgcgaAGGTGTCAAGAAAA
mp60DOF3F2	TGTC AAGAcgcgaCAGATTTAAAAGTCTT
mp60DOF4F2	CAAGAAAAAGCAGATTTcgcgaTTCTTC
mp60WRKYF1	AAGCTTCGTGTGGAGATCAACATATCTTCGTTAATTGAaTAC GCAAAATA
GUSRTF1	TACGGCAAAGTGTGGGTCAATAATCA
GUSRTR1	CAGGTGTTCGGCGTGGTGTAGAG
ATACT2F	TGCTCTCCATTTGTTTGTTC
ATACT2R	GGCATCAATTCGATCACTCA
qRT-MYB60-F	CATGAAGATGGTGATCATGAGG
qRT-MYB60-R	TTCCATTTGACCCCAAGTAG
PP2a-F	CAGCAACGAATTGTGTTTGG
PP2a-R	AAATACGCCCAACGAACAAA

*Italic and lower case letters indicate restriction and mutagenised sites, respectively*

*mDOF4*::GUS. To generate multiple mutagenised sites the templates for the second PCR amplification were plasmids already carrying a first mutagenised DOF site. In the case of the preparation of the construct *mW*::GUS the megaprimer method was not necessary, as the site to mutagenise is in a position 5' terminal into the minimal promoter and a single PCR reaction was performed with primers mp60WRKYF1 and p60R5, the PCR product was then cloned with the procedure already described.

All the oligonucleotide sequences are reported in Table 1.

#### Arabidopsis transformation and growth conditions

Wild-type Columbia (Col-0) plants were transformed using the *Agrobacterium tumefaciens* strain GV3101 carrying the constructs described above with the floral dip method [43]. Transformed lines were selected on kanamycin and single-insertion lines were selected for further analyses. Analyses of transgenic lines were

performed on T2 or on homozygous T3 plants grown under long-day conditions (16 hr light; 8 hr dark at 100  $\mu\text{mol m}^{-2} \text{sec}^{-1}$ ) at 22°C in a growth chamber. Seeds were germinated in Petri dishes containing Murashige and Skoog medium, 1% w/v sucrose and 0.8% w/v agar for seedling analysis or directly on soil for adult plant organ analysis. The ABA treatment was performed as previously described [19].

#### GUS activity assays and histochemical staining

For detection of GUS activity, tissues were fixed for 2 h in 90% (v/v) acetone at -20°C, incubated for 16-48 hours, at 37°C, in 0.05% (w/v) X-glucuronic acid, 0.1% (v/v) Triton X-100, and 0.5mM ferrocyanidine in 50 mM phosphate buffer (pH 7) and subsequently cleared in 70% (v/v) ethanol. Seedlings and flowers were cleared with a chloral hydrate:glycerol:water solution (8:1:2, v/v). Samples were examined using a Leica M205 FA stereomicroscope (Leica Microsystems GmbH Wetzlar, Germany) and a Zeiss Axiophot D1 microscope (Carl Zeiss MicroImaging, LLC Thornwood, New York, USA). Stereomicroscope images were recorded using the Leica LAS software version 2.8.1. Microscope images were recorded with an AxioCam MRc5 camera (Zeiss) using the AxioVision program (version 5.0).

#### Quantification of mRNA expression

RNA isolation, reverse transcription, qRT-PCR reactions and data analysis were performed as previously described [30]. *GUS* expression was analysed using primers GUSRT-F1 and GUSRT-R1, *ACTIN2* gene (primers ATACT2F, ATACT2R) was used as a reference for normalization. *AtMYB60* expression in different *cdf* mutants was analysed using primers qRT-MYB60-F and qRT-MYB60-R. *PP2A* gene, corresponding to *At1g13320* (primers PP2a-F and PP2a-R) was used as a reference for normalization [44]. All primer sequences are reported in Table 1.

#### Additional material

**Additional file 1: Analysis of GUS activity in seeds at different developmental stages in 1,307::GUS line.** A: open silique showing signal only in stomata and not in developing seeds. B: mature-green-stage seed (13 DAP). C: a 24 h imbibed seed. D: embryo isolated from a 24 h imbibed seed. The same results were obtained in all transgenic lines described in Figure 2. Scale bars represent 0.1 mm.

**Additional file 2: Relative expression of the *AtMYB60* gene in the different *cdf* mutants.** *cdf1-R* is an RNAi line ([29]). The other single and multiple mutants have been previously described ([30]). The *PP2a* (*At1g13320*) gene was used as a control [44].

#### Acknowledgements

This work was supported by the Italian "Progetto AGER, bando Viticoltura da Vino" (SERRES 2010-2015), by the BIOGESTECA 15083/RCC project, funded by

Regione Lombardia and by the AGRISOST project, funded by Fondazione Umberto Veronesi per il Progresso delle Scienze, Milano, Italy.

#### Author details

<sup>1</sup>Dipartimento di Scienze Biomolecolari e Biotecnologie, Università degli Studi di Milano, Milano, Italy. <sup>2</sup>Fondazione Filarete, Milano, Italy. <sup>3</sup>Max Planck Institute for Plant Breeding Research, Cologne, Germany. <sup>4</sup>Istituto di Biologia e Biotecnologia Agraria, CNR, Milano, Italy. <sup>5</sup>Dipartimento di Biologia, Università degli Studi di Milano, Milano, Italy.

#### Authors' contributions

EC carried out the construction of promoter-reporter plasmids, plant transformation and drafted the manuscript. EC, AA and LC did transgenic Arabidopsis analysis. FF carried out *cdf* mutant analysis. CT, MG, and GC conceived the study, participated in its design and coordination. MG, LC and FF helped to draft the manuscript. All authors read and approved the final manuscript.

Received: 14 September 2011 Accepted: 16 November 2011

Published: 16 November 2011

#### References

1. Kim TH, Böhmer M, Hu H, Nishimura N, Schroeder JI: Guard cell signal transduction network: advances in understanding abscisic acid, CO<sub>2</sub>, and Ca<sup>2+</sup> signaling. *Annu Rev Plant Biol* 2010, **61**:13.11-13.31.
2. Schroeder JI, Kwak JM, Allen GJ: Guard cell abscisic acid signalling and engineering drought hardiness in plants. *Nature* 2001, **410**:327-330.
3. Gray JE, Holroyd GH, van der Lee FM, Bahrami AR, Sijmons PC, Woodward FI, Schuch W, Hetherington AM: The HIC signalling pathway links CO<sub>2</sub> perception to stomatal development. *Nature* 2000, **408**:713-716.
4. Kopka J, Provart NJ, Muller-Rober B: Potato guard cells respond to drying soil by a complex change in the expression of genes related to carbon metabolism and turgor regulation. *Plant J* 1997, **11**:871-882.
5. Muller-Rober B, La Cognata U, Sonnewald U, Willmitzer L: A truncated version of an ADP-glucose pyrophosphorylase promoter from potato specifies guard cell-selective expression in transgenic plants. *Plant Cell* 1994, **6**:601-612.
6. Nakamura RL, McKendree WL, Hirsch RE, Sedbrook JC, Gaber RF, Sussman MR: Expression of an Arabidopsis potassium channel gene in guard cells. *Plant Physiol* 1995, **109**:371-374.
7. Nylander M, Svensson J, Palva ET, Welin BV: Stress-induced accumulation and tissue-specific localization of dehydrins in Arabidopsis thaliana. *Plant Mol Biol* 2001, **45**:263-279.
8. Plesch G, Ehrhardt T, Mueller-Roeber B: Involvement of TAAAG elements suggests a role for Dof transcription factors in guard cell-specific gene expression. *Plant J* 2001, **28**:455-464.
9. Terry N, Arias MB, Engler G, Tire C, Villarreal R, Van Montagu M, Inze D: rha1, a gene encoding a small GTP binding protein from Arabidopsis, is expressed primarily in developing guard cells. *Plant Cell* 1993, **5**:1761-1769.
10. Galbiati M, Simoni L, Pavesi G, Cominelli E, Francia P, Vavasseur A, Nelson T, Bevan M, Tonelli C: Gene trap lines identify Arabidopsis genes expressed in stomatal guard cells. *Plant J* 2008, **53**:750-762.
11. Gardner MJ, Baker AJ, Assie JM, Poethig RS, Haseloff JP, Webb AA: GAL4 GFP enhancer trap lines for analysis of stomatal guard cell development and gene expression. *J Exp Bot* 2009, **60**:213-226.
12. Plesch G, Kamann E, Mueller-Roeber B: Cloning of regulatory sequences mediating guard-cell-specific gene expression. *Gene* 2000, **249**:83-89.
13. Leonhardt N, Kwak JM, Robert N, Waner D, Leonhardt G, Schroeder JI: Microarray expression analyses of Arabidopsis guard cells and isolation of a recessive abscisic acid hypersensitive protein phosphatase 2C mutant. *Plant Cell* 2004, **16**:596-615.
14. Yang Y, Costa A, Leonhardt N, Siegel RS, Schroeder JI: Isolation of a strong Arabidopsis guard cell promoter and its potential as a research tool. *Plant Methods* 2008, **4**:6.
15. Zhao Z, Zhang W, Stanley BA, Assmann SM: Functional proteomics of Arabidopsis thaliana guard cells uncovers new stomatal signaling pathways. *Plant Cell* 2008, **20**:3210-3226.
16. Wang RS, Pandey S, Li S, Gookin TE, Zhao Z, Albert R, Assmann SM: Common and unique elements of the ABA-regulated transcriptome of Arabidopsis guard cells. *BMC Genomics* 2011, **12**:216.



17. Husebye H, Chadchawan S, Winge P, Thangstad OP, Bones AM: **Guard cell- and phloem idioblast-specific expression of thioglucoside glucohydrolase 1 (myrosinase) in Arabidopsis.** *Plant Physiol* 2002, **128**:1180-1188.
18. Liang YK, Dubos C, Dodd IC, Holroyd GH, Hetherington AM, Campbell MM: **AtMYB61, an R2R3-MYB transcription factor controlling stomatal aperture in Arabidopsis thaliana.** *Curr Biol* 2005, **15**:1201-1206.
19. Cominelli E, Galbiati M, Vavasseur A, Conti L, Sala T, Vuylsteke M, Leonhardt N, Dellaporta SL, Tonelli C: **A guard-cell-specific MYB transcription factor regulates stomatal movements and plant drought tolerance.** *Curr Biol* 2005, **15**:1196-1200.
20. Francia P, Simoni L, Cominelli E, Tonelli C, Galbiati M: **Gene trap-based identification of a guard cell promoter in Arabidopsis.** *Plant Signal Behav* 2008, **3**:684-686.
21. Nagy R, Grob H, Weder B, Green P, Klein M, Frelet-Barrand A, Schjoerring JK, Brearley C, Martinoia E: **The Arabidopsis ATP-binding cassette protein AtMRP5/AtABCC5 is a high affinity inositol hexakisphosphate transporter involved in guard cell signaling and phytate storage.** *J Biol Chem* 2009, **284**:33614-33622.
22. Meyer S, Mumm P, Imes D, Endler A, Weder B, Al-Rasheid KA, Geiger D, Marten I, Martinoia E, Hedrich R: **AtALMT12 represents an R-type anion channel required for stomatal movement in Arabidopsis guard cells.** *Plant J* 2010, **63**:1054-1062.
23. Yanagisawa S: **Dof domain proteins: plant-specific transcription factors associated with diverse phenomena unique to plants.** *Plant Cell Physiol* 2004, **45**:386-391.
24. PLACE. **A Database of Plant Cis-acting Regulatory DNA Elements.** [http://www.dna.affrc.go.jp/PLACE/].
25. Yanagisawa S, Schmidt RJ: **Diversity and similarity among recognition sequences of Dof transcription factors.** *Plant J* 1999, **17**:209-214.
26. Eulgem T, Rushton PJ, Robatzek S, Somssich IE: **The WRKY superfamily of plant transcription factors.** *Trends Plant Sci* 2000, **5**:199-206.
27. Yu D, Chen C, Chen Z: **Evidence for an important role of WRKY DNA binding proteins in the regulation of NPR1 gene expression.** *Plant Cell* 2001, **13**:1527-1540.
28. Velten J, Morey KJ, Cazzonelli CI: **Plant viral intergenic DNA sequence repeats with transcription enhancing activity.** *Virology* 2005, **2**:16.
29. Imaizumi T, Schultz TF, Harmon FG, Ho LA, Kay SA: **FKF1 F-box protein mediates cyclic degradation of a repressor of CONSTANS in Arabidopsis.** *Science* 2005, **309**:293-297.
30. Fornara F, Panigrahi KC, Gissot L, Sauerbrunn N, Ruhl M, Jarillo JA, Coupland G: **Arabidopsis DOF transcription factors act redundantly to reduce CONSTANS expression and are essential for a photoperiodic flowering response.** *Dev Cell* 2009, **17**:75-86.
31. Yanagisawa S: **The Dof family of plant transcription factors.** *Trends Plant Sci* 2002, **7**:555-560.
32. Galbiati M, Matus JT, Francia P, Rusconi F, Canon P, Medina C, Conti L, Cominelli E, Tonelli C, Arce-Johnson P: **The grapevine guard cell-related VvMYB60 transcription factor is involved in the regulation of stomatal activity and is differentially expressed in response to ABA and osmotic stress.** *BMC Plant Biol* 2011, **11**:142.
33. Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ: **An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets.** *PLoS One* 2007, **2**:e718.
34. **Arabidopsis eFP Browser.** [http://bbc.botany.utoronto.ca/efp/cgi-bin/efpWeb.cgi].
35. Oh JE, Kwon Y, Kim JH, Noh H, Hong SW, Lee H: **A dual role for MYB60 in stomatal regulation and root growth of Arabidopsis thaliana under drought stress.** *Plant Mol Biol* 2011, **77**:91-103.
36. Schauer SE, Schluter PM, Baskar R, Gheyselinck J, Bolanos A, Curtis MD, Grossniklaus U: **Intronic regulatory elements determine the divergent expression patterns of AGAMOUS-LIKE6 subfamily members in Arabidopsis.** *Plant J* 2009, **59**:987-1000.
37. Seki M, Ishida J, Narusaka M, Fujita M, Nanjo T, Umezawa T, Kamiya A, Nakajima M, Enju A, Sakurai T, et al: **Monitoring the expression pattern of around 7,000 Arabidopsis genes under ABA treatments using a full-length cDNA microarray.** *Funct Integr Genomics* 2002, **2**:282-291.
38. Zeller G, Henz SR, Widmer CK, Sachsenberg T, Ratsch G, Weigel D, Laubinger S: **Stress-induced changes in the Arabidopsis thaliana transcriptome analyzed using whole-genome tiling arrays.** *Plant J* 2009, **58**:1068-1082.
39. Choudhury A, Lahiri A: **Comparative analysis of abscisic acid-regulated transcriptomes in Arabidopsis.** *Plant Biol* 2011, **13**:28-35.
40. Matsui A, Ishida J, Morosawa T, Mochizuki Y, Kaminuma E, Endo TA, Okamoto M, Nambara E, Nakajima M, Kawashima M, et al: **Arabidopsis transcriptome analysis under drought, cold, high-salinity and ABA treatment conditions using a tiling array.** *Plant Cell Physiol* 2008, **49**:1135-1149.
41. Yamaguchi-Shinozaki K, Shinozaki K: **Organization of cis-acting regulatory elements in osmotic- and cold-stress-responsive promoters.** *Trends Plant Sci* 2005, **10**:88-94.
42. Kammann M, Laufs J, Schell J, Gronenborn B: **Rapid insertional mutagenesis of DNA by polymerase chain reaction (PCR).** *Nucleic Acids Res* 1989, **17**:5404.
43. Clough SJ, Bent AF: **Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana.** *Plant J* 1998, **16**:735-743.
44. Hong SM, Bahn SC, Lyu A, Jung HS, Ahn JH: **Identification and testing of superior reference genes for a starting pool of transcript normalization in Arabidopsis.** *Plant Cell Physiol* 2010, **51**:1694-1706.

doi:10.1186/1471-2229-11-162

**Cite this article as:** Cominelli et al.: DOF-binding sites additively contribute to guard cell-specificity of AtMYB60 promoter. *BMC Plant Biology* 2011 **11**:162.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

