

Review

Algal evolution in relation to atmospheric CO₂: carboxylases, carbon-concentrating mechanisms and carbon oxidation cycles

John A. Raven^{1,2,*}, Mario Giordano³, John Beardall⁴
and Stephen C. Maberly⁵

¹*Division of Plant Sciences, The James Hutton Institute, University of Dundee at TŷHI, Invergowrie, Dundee DD2 5DA, UK*

²*School of Plant Biology, University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia*

³*Department of Marine Sciences, Università Politecnica delle Marche, 60131 Ancona, Italy*

⁴*School of Biological Sciences, Monash University, Clayton Victoria 3800, Australia*

⁵*Centre for Ecology and Hydrology, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster LA1 4AP, UK*

Oxygenic photosynthesis evolved at least 2.4 Ga; all oxygenic organisms use the ribulose biphosphate carboxylase-oxygenase (Rubisco)–photosynthetic carbon reduction cycle (PCRC) rather than one of the five other known pathways of autotrophic CO₂ assimilation. The high CO₂ and (initially) O₂-free conditions permitted the use of a Rubisco with a high maximum specific reaction rate. As CO₂ decreased and O₂ increased, Rubisco oxygenase activity increased and 2-phosphoglycolate was produced, with the evolution of pathways recycling this inhibitory product to sugar phosphates. Changed atmospheric composition also selected for Rubiscos with higher CO₂ affinity and CO₂/O₂ selectivity correlated with decreased CO₂-saturated catalytic capacity and/or for CO₂-concentrating mechanisms (CCMs). These changes increase the energy, nitrogen, phosphorus, iron, zinc and manganese cost of producing and operating Rubisco–PCRC, while biosphere oxygenation decreased the availability of nitrogen, phosphorus and iron. The majority of algae today have CCMs; the timing of their origins is unclear. If CCMs evolved in a low-CO₂ episode followed by one or more lengthy high-CO₂ episodes, CCM retention could involve a combination of environmental factors known to favour CCM retention in extant organisms that also occur in a warmer high-CO₂ ocean. More investigations, including studies of genetic adaptation, are needed.

Keywords: inorganic carbon; mixing depth; photosynthetically active radiation; Rubisco; temperature; nutrients; UV radiation

1. INTRODUCTION

Algae are oxygenic photosynthetic organisms other than embryophytic plants, and by this definition include the cyanobacteria as well as a wide range of eukaryotic lineages. Cyanobacteria, as indicated by the occurrence of oxygenic photosynthesis, evolved at least 2.4 Ga, although fossil (including chemical biomarker) evidence for cyanobacteria does not go back beyond 2.1 Ga [1]. Eukaryotic algae have occurred since at least 1.2 Ga [2–4] and from freshwaters, and possibly lake margins, since 1.1 Ga [5] (table 1). Since 2.4 Ga, the biosphere has become increasingly oxygenated [1], reflecting the colonization of the oceans by cyanobacteria after their origin in freshwater habitats, with a corresponding

increase in the capacity of these organisms to have global biogeochemical influence [15,16] (table 1). A significant increase in oxygen, with oxygenation of the deep ocean, occurred in the Neoproterozoic 0.54–1 Ga [32], with variations in the Phanerozoic including the highest known level in the Permo-Carboniferous glaciation [33]. Carbon dioxide varied with a general downward trend, with minima generally related to glaciations [10,34]. CO₂ was relatively constant at about 23 (range of estimates 10–100) times the present level between 2.5 and 1.8 Ga, with a very significant decrease between 1.8 and 1.1 Ga [35], further variations in the Neoproterozoic [36–38] and relatively well-established changes in the Phanerozoic with minima in the Permo-Carboniferous and in the Pleistocene glaciations [33,39]. In this article, we consider how these environmental changes have influenced algal evolution, both through direct effects of the concentrations of CO₂ and O₂ on photosynthesis and related metabolism, and through indirect

* Author for correspondence (j.a.raven@dundee.ac.uk).

One contribution of 12 to a Theme Issue ‘Atmospheric CO₂ and the evolution of photosynthetic eukaryotes: from enzymes to ecosystems’.

Table 1. Inorganic carbon acquisition characteristics of cyanobacteria and algae related to earliest known occurrence of the taxon. General references on earliest known occurrence of algae: [6–8]. References on presence or absence of CO₂-concentrating mechanisms (CCMs): [9–14].

taxon	occurrence of CCM in extant organisms	oldest known fossil	references
Cyanobacteria	CCM ubiquitous	2.15 Ga (biomarkers); 2.45 Ga (O ₂)	[1,15–17]
Chlorophyta: Prasinophyceae	CCM ubiquitous	1.3 Ga	[18,19]
Chlorophyta: Chlorophyceae	CCM present in all?	(450 Ma) ³²	[19,20]
Chlorophyta: Trebouxiophyceae	CCM present or absent	450 Ma	[19–23]
Chlorophyta: Ulvophyceae	CCM usually present; absent in some; C ₄ in one	540 Ma	[6,19,20]
Streptophyta: Charophyceae	CCM present in all?	450 Ma	[6,20]
Streptophyta: Embryophytes	CCM usually absent; pyrenoid-based CCM in some anthocerophytes, C ₄ or CAM in some freshwater tracheophytes, CCMs not involving C ₄ and CAM in some freshwater and all marine tracheophytes	475 Ma	[6,24,25]
Rhodophyta: Bangiophyceae	CCM in all?	1.2 Ga	[2]
Rhodophyta: Florideophyceae	CCM in many, absent from some marine, many freshwater	600 Ma	[26,27]
Ochista: Bacillariophyceae	CCM in all?	120 Ma	[28,29]
Ochista: Fucophyceae	CCM in all?	(570 Ma?)	[27,30]
Ochista: Tribophyceae	CCM in all?	600 Ma	[3,31]
Ochista: Chrysophyceae and Synurophyceae	CCM absent in all	?	[9]

effects of changes in temperature (CO₂) and the availability of combined nitrogen, phosphorus and iron (CO₂ and O₂).

2. AUTOTROPHIC CARBOXYLASES

Six pathways of autotrophic CO₂ fixation are known in extant organisms, including ribulose biphosphate carboxylase-oxygenase carboxylase activity in the photosynthetic carbon reduction cycle (Rubisco–PCRC), using CO₂ as the inorganic carbon species assimilated, which is at the core of inorganic carbon assimilation in extant oxygenic photosynthetic organisms [40–45]. These pathways are summarized in table 2 with respect to their stoichiometric requirement for reductant and ATP, their affinities for inorganic carbon expressed in terms of the half-saturation value for CO₂ and the influence of oxygen on their functioning.

Converting one CO₂ to the oxidation–reduction level of carbohydrates (CH₂O) requires four reducing equivalents at, or lower than, the midpoint redox potential of the NADPH.NADP⁺ couple. The values given in table 2 indicate the minimal stoichiometry, assuming no redox side reactions or futile cycles. An example of a side reaction for the Rubisco–PCRC pathway is the Rubisco oxygenase activity in the photosynthetic carbon oxidation cycles(s) (Rubisco–PCOC), which occurs at a relatively low [CO₂]/[O₂] ratio. All of the autotrophic CO₂ fixation pathways also require ATP (table 2), with a range of stoichiometries from one to three ATPs for CO₂ converted to carbohydrate. The Rubisco–PCRC pathway has the equal highest energy cost of converting CO₂ to carbohydrate, with the further

energy cost of the side reaction of the Rubisco–PCOC at low [CO₂]/[O₂] ratios.

The Rubisco–PCRC pathway seems more appropriate to CO₂ fixation at the present atmospheric CO₂ level when the half-saturation value for CO₂ is considered. The forms of Rubisco with the highest affinities for CO₂ (Form IB from some algae and vascular plants relying on CO₂ diffusion to Rubisco; Form ID from other algae) have half-saturation values for CO₂ almost as low as those of two other pathways, while the values for the other three pathways are considerably higher (table 2). The final criterion in table 1 is the effect of oxygen on the pathways. While the Rubisco–PCRC pathway is competitively (with CO₂) inhibited by O₂, the enzymes of this pathway are not damaged by O₂: some of the other pathways have one or more enzymes that are subject to irreversible inhibition by O₂ (table 2). For lack of accurate information, the resource cost of synthesizing the enzymic machinery needed to fix one mole of CO₂ per second from the present atmospheric CO₂ concentration is not shown in table 2. This value is a function of the stoichiometry of the enzymic protein components in the pathway, with that of the carboxylase(s) set by their CO₂ affinity, and the *M_r* (relative molecular mass) values of the enzymes [40,41]. Here, the relatively low-specific reaction rate and high *M_r* of Rubisco would tend to make the Rubisco–PCRC pathway expensive at present CO₂ concentrations, even without the O₂ effect, though some other pathways are probably at least as expensive as a result of the low CO₂ affinity of their carboxylases and the consequent large quantity of carboxylase needed to fix one mole of CO₂ per second from the present atmospheric CO₂ concentration.

Table 2. Energy (NADPH and ATP) stoichiometry, affinity for inorganic carbon expressed as the half-saturation concentration for CO₂, competitive inhibition by O₂ and damage by O₂ for six autotrophic inorganic carbon assimilation pathways. Based on [40–45].

pathway from inorganic carbon to carbohydrate	NAD(P)H per CO ₂	ATP per CO ₂	K _{1/2} CO ₂ mmol m ⁻³	O ₂ competitive inhibition	O ₂ damage to one or more enzymes
Rubisco–PCRC, saturating CO ₂ no O ₂	2	3	≥10	yes	no
reverse TCAC	2	1.67	>1500	no	yes ^a
3-hydroxypropionate	2	2	10	no	no ^b
3-hydroxypropionate–4-hydroxybutyrate	2	3	>2000	no	O ₂ -insensitive pathway in some organisms living in microaerobic habitats
dicarboxylate–4-hydroxybutyrate	2	2.67	>2000	no	no; some organisms live in microaerophilic habitats
Wood–Ljungdahl pathway	2	1	40 000 ^c	no	yes

^aThe reverse tricarboxylic acid cycle (TCAC) can occur in thermophilic aerobic chemolithotrophs as a result of low O₂ solubility and high respiratory rates maintaining a low O₂ concentration inside the cells, and expression of an O₂-insensitive version of the 2-oxoglutarate–ferredoxin oxidoreductase which has at least a fivefold lower specific activity than the O₂-sensitive enzyme [43].

^bThe most oxygen-sensitive enzyme, methylmalonyl-CoA mutase, can be assayed and even purified at atmospheric equilibrium O₂ concentrations; it may not be sufficiently O₂-tolerant to function in illuminated cells of oxygenic photosynthetic organisms [43].

^cAlthough acetogens live in habitats with higher CO₂ concentrations than correspond to atmospheric equilibrium [46], the *in vitro* K_{1/2} value cited does seem very high.

The occurrence of Rubisco–PCRC as the core carboxylase in oxygenic photosynthetic organisms can be related to opportunity and functionality. By opportunity is meant the occurrence of the pathway at the time (about 2.4 Ga) at which the earliest evidence for oxygenic photosynthetic organisms is known. The Rubisco–PCRC pathway originated before oxygenic photosynthesis evolved (see below). By functionality is meant the carboxylase CO₂ affinity and carboxylase *M_r* values as determinants of the quantity of carboxylase (mass of protein) needed to fix one mole of CO₂ per second from the current atmosphere, the extent to which O₂ competitively inhibits or damages the enzymes of the pathway, and the ATP cost of the pathway per CO₂ assimilated (table 2). The CO₂ affinity criterion apparently rules out three of the five pathways (two of which are also very oxygen-sensitive), leaving the 3-hydroxypropionate pathway and the Rubisco–PCRC pathway. While not oxygen inhibited in the manner found for Rubisco, the 3-hydroxypropionate pathway may be sufficiently O₂-sensitive to restrict its functionality in oxygenic organisms once O₂ had begun to accumulate in the part of the biosphere occupied by oxygenic photosynthetic organisms (tables 1 and 2). In such ways can the role of the Rubisco–PCRC pathway in all known oxygenic photosynthetic organisms be rationalized.

3. RUBISCO CARBOXYLASE ACTIVITY AND THE PHOTOSYNTHETIC CARBON REDUCTION CYCLE

Rubisco evolved before the origin of oxygenic photosynthesis [47–49]. The Rubisco gene family not only contains the forms I, II and III Rubiscos that catalyse the Rubisco carboxylase and Rubisco oxygenase reactions, but also the form IV Rubisco-like protein (RLP) which does not catalyse the typical Rubisco reactions and which is involved in methionine salvage in some bacteria [47–49]. Molecular phylogenetic analysis [47–49] suggests that an ancestral form III Rubisco

arose in a methanogen (i.e. a member of the Archaea) and gave rise, by two vertical transmissions, to all other form III Rubiscos and to form IV. Horizontal gene transfer then moved form III and form IV Rubiscos to an ancestral bacterium, where form III gave rise to the ancestors of forms I and II, and hence by vertical transmission to the form IV RLP of bacilli and to forms I, II and IV of an ancestral proteobacterium.

Vertical descent and further diversification gave rise to the forms IA, IC, ID and II, and the form IV RLP in extant Proteobacteria. Horizontal gene transfer from the Proteobacteria transferred form IA to cyanobacteria, where form IB evolved from form IA [47–50]. However, there is also evidence that extant α -cyanobacteria with form 1A Rubisco (α -cyanobacteria) acquired their form IA Rubisco (and other α -carboxysomal proteins) from a proteobacterium, displacing the form IB Rubisco and associated β -carboxysomal proteins [51–56]. Endosymbiosis of a heterocystous β -cyanobacterium [57] gave rise to the plastids of the Archaeplastida (i.e. Plantae), where the form IB Rubisco of the endosymbiont was retained by glaucocystophyte and green algae and hence by vertical transfer to embryophytic ('higher') plants, and was moved by secondary endosymbiosis to the (rhizarian) chlorarachniophytes and the (excavate-discicrinate) euglenoids. Horizontal gene transfer from a proteobacterium accounts for the presence of form ID Rubisco in red algae and in the chromistan algae (ochrista or heterokontophytes, cryptophytes and haptophytes) whose plastids arose by secondary endosymbiosis of a red alga. The peridinin-containing dinoflagellate and basal apicomplexans (represented today by the photosynthetically competent *Chromera velia* and an as yet un-named close relative) also obtained their plastids by secondary endosymbiosis from a red alga, but horizontal transfer of form II Rubisco from a proteobacterium replaced the form ID Rubisco [58]. Other dinoflagellates have other forms of Rubisco as a result of tertiary endosymbioses. Finally, a second

primary endosymbiosis involving an α -cyanobacterium with form IA Rubisco accounts for the presence of plastids and form IA Rubisco in the (rhizarian) euglyphid amoeba *Paulinella* [54,55].

Some of the enzymes of the PCRC of vascular plants are derived from cyanobacteria in the primary endosymbiosis leading to the Archaeplastida (i.e. Plantae), while others are of host origin [59]. This work has now been extended to include the glaucocystophyte and red algal members of the Archaeplastida [60] and the diatoms whose plastids arose from secondary endosymbiosis of a red alga [61] following an earlier secondary endosymbiosis of a green alga [62]. There are also differences among algae in the regulation of enzymes of the PCRC [63]. Further work is needed to establish the relevance, if any, of atmospheric changes to these differences in the regulation of enzymes of the PCRC [63,64], and also the absence of Rubisco activase from algae with the form ID Rubisco in the few cases examined [61].

It is likely that the earliest Rubiscos in chemolithotrophic and anoxygenic photosynthetic Archaea and Bacteria were operating in a high-CO₂ environment and, because there was no oxygenic photosynthesis, in the essential absence of O₂. Such an environment would have provided little or no selective pressure for a high CO₂ affinity or a high CO₂/O₂ selectivity relative to a high CO₂-saturated maximum catalytic rate, using the mechanistic arguments of Tcherkez *et al.* [65] that a high maximum catalytic rate is incompatible with high CO₂ affinity and higher CO₂/O₂ selectivity.

The early organisms using Rubisco as a carboxylase would, then, be able to use diffusive entry of CO₂, with none of the resource costs associated with Rubisco operating at below the CO₂-saturated rate in the presence of O₂ when Rubisco oxygenase activity is expressed. The additional resource costs can be considered as capital (synthetic) costs and running costs. The capital costs are those of synthesizing additional Rubisco enzyme per cell if the per cell rate of CO₂ assimilation is to be retained, as well as the costs of making the enzymes and transporters related to the operation of a PCOC and/or a CO₂-concentrating mechanism (CCM), and include energy, carbon and nitrogen as well as phosphorus for any additional ribosomes that are required [66]. The running costs are in terms of energy, both for synthesizing 2-phosphoglycolate and metabolizing it back to sugar through a PCOC and for operating a CCM.

The early Rubisco-containing organisms could still permit CO₂ saturation of a high specific reaction rate Rubisco, with corresponding savings of resources in constructing the enzymic machinery able to assimilate one mole of CO₂ per second. In other words, the conditions described would give the lowest possible energy, carbon and nitrogen costs of producing the amount of Rubisco capable of fixing one mole of CO₂ per second, and also give the lowest possible energy cost of operating Rubisco, i.e. two NADPH and three ATP per CO₂ converted to carbohydrate. A correlated saving associated with this minimal requirement for NADPH and ATP in autotrophic CO₂ assimilation is in the quantity of redox and ATP synthesis machinery needed for NADPH and ATP production. Minimizing

the protein requirements for autotrophy also decreases the phosphorus requirement for mRNA in ribosomes and in tRNA, and mRNA needed to produce the Rubisco and associated enzymes. These characteristics would minimize the nitrogen needed to produce the machinery associated with a given rate of CO₂ fixation, and thus the phosphorus in RNA needed to synthesize the proteins [66]. Costs in energy, nitrogen, phosphorus, iron and manganese will be considered below in the context of decreasing CO₂ and increasing O₂ [66–74].

The mechanistically constrained covariation in Rubisco kinetics mentioned above, i.e. a high CO₂-saturated specific reaction rate correlated with low CO₂ affinity and CO₂/O₂ selectivity and vice versa [65] has parallels with the growth strategies of vascular plants [75] and of algae [76] according to the competitive–stress-tolerant–ruderal (CSR) paradigm, for which there are also mechanistic bases in terms of having high rates of metabolism and growth in ruderals, and lower rates but perhaps a more effective use of resources in stress-tolerators [77–79].

4. OXYGEN ACCUMULATION, RUBISCO OXYGENASE AND THE METABOLISM OF PHOSPHOGLYCOLATE

The build-up of O₂ has had many effects on algal evolution, permitting respiration and, via the occurrence of stratospheric ozone, a decreased UV-B flux, and production of reactive oxygen species from O₂ rather than UV-B action on cell constituents [79,80]. The presence of O₂ does not, however, inhibit O₂ production by oxygenic photosynthetic organisms [81,82]. The accumulation of O₂ in the habitats of oxygenic photosynthetic organisms using Rubisco as their autotrophic carboxylase permits Rubisco oxygenase activity to occur, provided the CO₂ concentration is below saturation for Rubisco. Such decreased CO₂ concentrations, combined with at least local O₂ accumulation, probably occurred about 2 Ga, when there is evidence of an ice age extending to low palaeolatitudes [35,83].

The product of the Rubisco oxygenase activity is, as well as one 3-phosphoglycerate per O₂ consumed, one 2-phosphoglycolate. In addition to sequestering the often limiting resource phosphorus if 2-phosphoglycolate continues to accumulate, 2-phosphoglycolate is also an inhibitor of some reactions involving phosphate esters, including some in the PCRC [84]. Accordingly, all organisms using Rubisco in the presence of O₂, i.e. oxygenic photolithotrophs and some chemolithotrophs, have 2-phosphoglycolate phosphatase [84,85]. This enzyme would not have been needed to deal with 2-phosphoglycolate from Rubisco oxygenase activity in anoxygenic photosynthetic proteobacteria, or in any anaerobic chemolithotrophs, using the Rubisco–PCR pathway before build-up of O₂ in the biosphere. 2-Phosphoglycolate phosphatase also occurs in some non-autotrophic bacteria that have no Rubisco, where it is thought to be involved in some forms of DNA repair [86,87]. Since DNA damage and its repair must have occurred before oxygenic photosynthesis, the 2-phosphoglycolate phosphatase in oxygenic photosynthetic organisms could have been recruited from bacteria

lacking autotrophic 2-phosphoglycolate synthesis. However, the 2-phosphoglycolate phosphatase from eukaryotes does not seem to have been derived from the 2-phospho-glycolate phosphatase of cyanobacteria [88].

The organic carbon product of 2-phosphoglycolate phosphatase is glycolate. This can be excreted, with loss from the organism of the energy and carbon used in its synthesis. Alternatively, glycolate can be salvaged by metabolism to 3-phosphoglycerate, and hence triose phosphate, which occur in the PCRC, albeit with the input of energy and the release of CO₂ [85]. The cyanobacteria have two variants on pathways converting glycolate to 3-phosphoglycerate.

One means of converting glycolate to 3-phosphoglycerate is the pathway via glycine and serine, with recycling of ammonia, as in the classic PCOC of embryophytic plants and at least some eukaryotic algae [85,88–92]. The pathway through glycine and serine seems to have been gained by eukaryotic photosynthetic organisms during the primary endosymbiosis yielding the plastids of the Archaeplastida (i.e. Plantae), although some of the genes in eukaryotes came from α -proteobacteria rather than cyanobacteria [88]. The β -cyanobacterial plastid ancestor gave rise to the glycolate oxidase, glycerate kinase and hydroxypyruvate reductase of algae and embryophytes, while serine hydroxymethyl transferase and the L, P and T subunits of glycine decarboxylase came from α -proteobacteria by horizontal gene transfer [88]. The origin of the other eukaryotic PCOC genes, i.e. those encoding the H subunit of glycine decarboxylase, glutamate–glyoxylate aminotransferase and serine–glyoxylate aminotransferase, has not yet been established [88].

The other pathway from glycolate to 3-phosphoglycerate involves tartronic semialdehyde, and is called the tartronic semialdehyde pathway by phycologists; bacteriologists call it the glycerate pathway, even though glycerate is also an intermediate of the PCOC [85,88–90]. Parts of the PCOC (glycerate dehydrogenase, serine transaminases and serine hydroxymethyltransferase) could have been recruited from core metabolism synthesizing serine and glycine from glycolytic intermediates, while others (glycolate dehydrogenase/oxidase, glycine decarboxylase and glycerate kinase) have no known roles other than in the metabolism of glycolate to PCRC intermediates [93,94].

It seems likely that at least one of the metabolic pathways from glycolate to 3-phosphoglycerate evolved in oxygenic photosynthetic organisms relying on diffusive CO₂ entry before CCMs evolved. Not only is there at least a minimal flux through Rubisco oxygenase and thence to intermediates of a glycolate metabolism pathway despite high levels of expression of CCMs [85,89,90,92,95], but elimination of the pathways of glycolate metabolism is fatal to the organism [85,90]. Previous misgivings [91,92] about the occurrence of the complete PCOC in diatoms have now been largely overcome, although there are still doubts as to the glycerate kinase step [61,88].

The changes to CO₂ fixation in oxygenic photolithotrophs in relation to decreasing CO₂ and, especially, increasing O₂ is part of a wider range of resource cost increases as the biosphere becomes

oxygenated. Falkowski & Godfrey [72] point out that not only is Rubisco impacted by increasing O₂ with decreasing CO₂, but that the potential for oxygen damage to nitrogenase becomes manifest, and the very source of the O₂, the reaction centre of photosystem II, is itself subject to photodamage both directly through excitation energy transfer to the reaction centre, but also indirectly through the accumulated O₂ forming reactive O₂ species ([66]). As Raven [66,68–70] points out, the effects on Rubisco demand additional nitrogen in the enzyme itself and in related enzymes, more iron and manganese in additional thylakoid redox agents, and more phosphorus in the RNA needed to make the additional protein, if the rate of photosynthesis is to be maintained. More energy input as NADPH and ATP is also needed to run CO₂ assimilation [66,69,70]. For oxygen damage to nitrogenase, there is generally synthesis of ‘reserve’ nitrogenase in addition to what is needed in the absence of oxygen damage to satisfy the combined nitrogen requirements of cell growth. Synthesis of the ‘reserve’ nitrogenase requires the nitrogen and energy needed for the synthesis of any protein, but also the iron and (in almost all cases) molybdenum used in the nitrogenase cofactors. When oxygen damage does occur and reserve nitrogenase is used catalytically, more energy (but not nitrogen, iron and molybdenum) is needed to synthesize nitrogenase to replace what is damaged. In both cases, the production of more nitrogenase than would be required in the absence of oxygen involves the use of more phosphorus for the RNA required for the additional protein synthesis. In the case of photoinhibition, more nitrogen and energy are needed to synthesize reserve photosystem II reaction centres, more energy is needed to synthesize replacement photosystem II reaction centres, and more phosphorus is needed for the RNAs needed for the extra protein synthesis. A further aspect of damage to proteins by O₂ concerns the absence of any core autotrophic CO₂ assimilation pathway other than Rubisco–PCRC from oxygenic photosynthetic organisms. In addition to the CO₂ affinity problems outlined for some of the alternative pathways, there would also be the requirement for additional resources (photosynthetically active radiation (PAR), nitrogen and phosphorus) to make and use additional RNA needed for the additional resynthesis of O₂-damaged protein (see [66] and above, for other cases).

Compounding this need for additional nitrogen and phosphorus per unit CO₂ or N₂ assimilated and photons used in photochemistry is the effect of increased O₂ on the availability of nitrogen and phosphorus. Falkowski & Godfrey [72] point out that oxygenation of the biosphere not only decreases the potential for diazotrophy, but allows nitrifying microbes to convert NH₄⁺ to NO₃⁻, which in hypoxic or anoxic micro-habitats can be denitrified to produce N₂O and N₂. This nitrification–denitrification sequence decreases the availability of combined nitrogen to non-diazotrophic primary producers. In the case of phosphorus, the availability of O₂ converts Fe(II) to oxidized iron (Fe(III)), which binds phosphate and thus decreases global phosphorus availability [96,97].

Phosphorus is one of the biogeochemical regulators of the O₂ content of the atmosphere [96,97]. This topic will be returned to below in the context of ocean deoxygenation as a function of increases in CO₂ and temperature.

5. INTRODUCTION TO CO₂-CONCENTRATING MECHANISMS

Diffusive entry of CO₂ to Rubisco was presumably the ancestral mechanism of autotrophic CO₂ assimilation in oxygenic photosynthetic organisms. Entry of CO₂ to Rubisco by diffusion is found today in the majority, by species number and contribution to global primary productivity, of terrestrial oxygenic photosynthetic organisms, but in a minority of oxygenic photosynthetic organisms in aquatic environments where photolithotrophs with CCMs predominate [11–13,34,67,79, 98–101] (table 1). The references just cited show that CCMs are very widely distributed among algae, both phylogenetically and geographically, although they seem to be absent from chrysophycean and synurophycean algae [9]. The mechanistic, including molecular, details of the CCMs of cyanobacteria are now known [51–53,56,102]. The CCMs of eukaryotic algae are less clearly understood at both the molecular and the mechanistic levels, although they are clearly polyphyletic [10,13,34,67,79,103,104].

As to the evolutionary origin of CCMs, the selective factors were presumably decreasing CO₂ and increasing O₂. The variability of other gases over the last 2.4 Gyr suggests that there were several periods at which CCMs could have been resource-effective (energy, nitrogen, phosphorus, iron, zinc and manganese) alternatives to diffusive CO₂ entry to Rubisco with attendant high activity of Rubisco oxygenase and expression of high levels of enzymes converting 2-phosphoglycolate to sugar phosphate [10,67–71,74,105,106]. We shall return to the question of the timing(s) of the origin(s) of CCMs and how, if they originated early, they were retained through intervening high-CO₂ episodes. First, we consider what the transition from diffusive CO₂ entry to CCM-based delivery of CO₂ to Rubisco involves.

6. THE FUNCTIONING OF CO₂-CONCENTRATING MECHANISMS IN COMPARISON WITH THE DIFFUSIVE ENTRY OF CO₂

One factor in the evolution of a CCM in an organism previously relying on diffusive CO₂ movement from the medium to Rubisco as CO₂ availability decreases is the kinetics of the Rubisco used by the organism. A Rubisco with a high CO₂-saturated catalytic capacity, and a correspondingly low CO₂ affinity and CO₂/O₂ selectivity, such as occurs in the form IA and form IB Rubiscos of extant cyanobacteria, would be expected to present a stronger evolutionary case for a CCM at a given CO₂ concentration and CO₂/O₂ ratio than would eukaryotic form IB and form ID Rubisco with higher CO₂ affinity and CO₂/O₂ selectivity but a lower CO₂-saturated maximum catalytic rate [65]. Young *et al.* [107] have used molecular phylogenetic evidence to show that there was positive selection of the form

ID Rubiscos in some eukaryotes that correspond to low-CO₂ and low-CO₂/O₂ episodes in the geological record, and that these episodes of positive selection could have corresponded to the time of evolution of CCMs. To be effective, the CCM must maintain a higher CO₂ concentration at the site of Rubisco than would be possible by CO₂ diffusion alone [10].

The essential component of a CCM is the accumulation of CO₂ in the compartment containing Rubisco to a higher steady-state concentration than occurs in the growth medium, and hence even higher than the steady-state concentration near Rubisco which could occur with diffusive CO₂ entry. For algae, one mechanism of accumulation could involve C₄-like photosynthetic metabolism with an ATP-dependent (C₃ + C₁) carboxylation in the cytosol, using HCO₃⁻ obtained directly or indirectly (via CO₂ entry from the medium followed by carbonic anhydrase catalysis) from the medium, followed by a (C₄–C₁) decarboxylation in the chloroplast [104]. The alternative mechanisms do not involve the inorganic carbon transferred from the medium to Rubisco forming organic intermediates. These alternative mechanisms involve transmembrane-active transport mechanisms which move an inorganic carbon species (CO₂ or HCO₃⁻), or H⁺, against a free-energy gradient. Such transporters could be (and have been) derived from transporter gene families by change of specificity of the transported substrate (to CO₂ or HCO₃⁻), with changes in regulation and, perhaps, changes in intracellular targeting [51–53,56]. An exception to the need for active transport across a membrane is CO₂ use in the cyanobacterial CCMs, where diffusive CO₂ entry across the plasmalemma is followed by energized conversion to HCO₃⁻ by the NAD(P)H–PQ oxidoreductase of the thylakoid membrane [51–53,56]. Here, a high CO₂ permeability of the plasmalemma is required. Such a high membrane permeability to CO₂ is needed for diffusive CO₂ entry all the way to Rubisco in organisms lacking a CCM, and (as noted above) in CO₂ ‘active transport’ in cyanobacteria. A high CO₂ permeability is also necessary in organisms with a CCM mechanism involving CO₂ production from HCO₃⁻ in a compartment acidified by a H⁺ pump followed by transmembrane movement of CO₂ into the compartment containing Rubisco [13]. The energized conversion of CO₂ to HCO₃⁻ in the cyanobacterial cytosol could increase the chance of any CO₂ that leaks out of the carboxysomes being trapped as HCO₃⁻ in the cytosol. In all other cases, a CCM is most energetically efficient with minimal CO₂ flux from the compartment in which it is accumulated back to the medium, i.e. with very low membrane permeabilities to CO₂.

The energetic savings that could result from a low CO₂ permeability of the plasmalemma, and of the inner plastid envelope (if that is the membrane involved in active transport of inorganic carbon), in eukaryotes with CCMs based on active transport of an inorganic carbon species would, if verified, suggest a phylogenetic, and in many cases, an acclimatory (changes from growth in high to low CO₂) decrease in CO₂ permeability. This question has been addressed by a number of workers [103,108], who found relatively high CO₂ permeability in the eukaryotic algal membranes examined, regardless of

whether the algae had been cultured in high (CCM repressed) or low (CCM expressed) inorganic carbon concentrations. The permeability values for CO₂ of the plasmalemma of high-CO₂ and low-CO₂-grown cells of *Chlamydomonas reinhardtii* range from 0.76 to 1.8.10⁻³ cm s⁻¹ [108], while for four species of diatom the range is 15–56.10⁻³ cm s⁻¹ [103], with the range of values probably related to methodological as well as phylogenetic differences. While models for CCMs in diatoms consistent with the available data show relatively modest energy losses during CCM operation, they do involve constraints such as the membrane(s) at which active inorganic carbon transport occurs, and the chemical species involved in this active transport [103]. The same argument applies to the protein shell of the carboxysome, for which a restriction on CO₂ diffusion, but not on the diffusion of anionic substrates (HCO₃⁻ and ribulose-1,5-bisphosphate) and product (3-phosphoglycerate) has been demonstrated [109].

Most CCMs also involve one or more carbonic anhydrase enzymes: an exception is a C₄ mechanism which involves, as indicated above, HCO₃⁻ entry to the cytosol, (C₃ + C₁) carboxylation using HCO₃⁻, and (C₄ - C₁) decarboxylation in the plastid stroma with CO₂ as the inorganic carbon product [13,92,95,104,110]. All other well-investigated CCMs seem to involve 'normal' carbonic anhydrases, i.e. those catalysing the equilibration of CO₂ and HCO₃⁻ [11,67,111]: this is the case for 'active CO₂ influx' in cyanobacteria, which involves a carbonic anhydrase in the carboxysome as well as the energized conversion of CO₂ to HCO₃⁻ at the thylakoid membrane, which is effectively a unidirectional carbonic anhydrase [56].

The functioning of CCMs is influenced by a number of factors other than the availability of inorganic carbon (and, in some cases, O₂), e.g. photosynthetically active radiation, UV-B radiation, the form and the concentration of combined nitrogen, the phosphate concentration and the iron concentration [66,67,74,105]. The influence of these factors on the expression and functioning of CCMs is presented in table 3 [66,67,74]. There are also predicted effects of expression of CCMs rather than reliance on diffusive entry of CO₂ on the resource costs of synthesis of the photosynthetic apparatus, and of its operation; these are discussed below. We next discuss the possible influence of these interactions on the evolution of CCMs, their retention through any high-CO₂ episodes between their origin in a low-CO₂ habitat and today, and their fate in a future higher CO₂ and warmer world.

7. THE ORIGINS OF CO₂-CONCENTRATING MECHANISMS

The 'why' of the origin of CCMs presumably concerns the occurrence of low CO₂, both in absolute terms and in relation to O₂, which was indicated above as requiring additional protein (hence RNA and phosphorus) in more Rubisco and in enzymes metabolizing 2-phosphoglycolate to sugar phosphate, as well as additional energy input per net CO₂ assimilated by diffusive CO₂ entry of CO₂ to Rubisco and metabolism of 2-phosphoglycolate. Depending on the circumstances, e.g. the form of Rubisco present and the environmental

conditions as well as the mechanism of the particular CCM [10,34,68–71,74,79,91,103,105], a CCM could require less energy, nitrogen, phosphorus, zinc and iron for its synthesis, and less energy for its operation, than diffusive CO₂ entry with metabolism of 2-phosphoglycolate (table 2). Other factors that could have influenced the evolution of CCMs include the decreasing UV-B flux with the increased stratospheric O₃ resulting from the build-up of O₂, which is itself an influence on the evolution of CCMs [79]. UV-B radiation causes damage to Rubisco and to photosystem II, but has less effect on photosystem I. The limited data available suggest that UV-B has little effect on CCM activity in the green alga *Dunaliella tertiolecta* [122] but elevated CO₂ can increase the sensitivity of microalgae to UV-B [123,124]. There is no information about the possibility of a differential impact of UV-B on the various forms of Rubisco, though it would be interesting to know if changes in UV-B in the past relate to the evolution of different Rubiscos.

The 'how' of the origin of CCMs concerns the ancestry of the various components of the pathway. For the active transport components, H⁺ pumps are ubiquitous and anion transporters/pumps (hence HCO₃⁻ transporters/pumps) are also widespread, as could be the ancestors of CO₂ pumps [10,13,34,51–53,56,79,102,104]. Facilitators of downhill transmembrane CO₂ transport, yielding permeabilities in excess of those owing to the lipid phase alone, are required for cyanobacterial active transporters and for the mechanism involving an acidified compartment generating CO₂ from HCO₃⁻ with subsequent transmembrane CO₂ diffusion to Rubisco. Such facilitators would presumably have originally been components of the diffusive pathway for CO₂ from the medium to Rubisco. Carbonic anhydrases could have had a number of roles prior to their co-option into CCMs, including that of facilitating diffusion of CO₂ (as HCO₃⁻) in the diffusive entry of CO₂ from the medium to Rubisco. Cyanobacterial carboxysomes are part of a larger family of bacterial micro-compartments [125]. This brief view may be over-optimistic as to the ease of co-opting existing mechanisms into CCMs [34], and does not address the origin of the eukaryotic pyrenoid [14]; however, it does indicate some possibilities.

There are a number of options as to the 'when' of the evolution of CCMs. We initially consider times of relatively low CO₂, based on low palaeotemperatures (with the requirement that greenhouse gases corrected for the faint young Sun) or on biogeochemical or biological proxies. Glacial/low CO₂ episodes occurred 2.4–2.1 Ga and at 750, 650 and 320–270 Ma, as well as the Pleistocene (last 2.4 Myr) [10]. All but the earliest of these times would have been relevant to at least some of the eukaryotic as well as cyanobacterial oxygenic photosynthetic organisms (table 1). There is no direct fossil evidence as to the origin of CCMs, and little help from molecular clocks [10,67,79], although recent work by Young *et al.* [107] shows episodes of positive selection of form ID Rubisco in diatoms and haptophytes which correspond to low-CO₂ episodes and hence possibly relate to the origin of CCMs. Assuming, as seems very likely, that cyanobacterial and at least some algal CCMs evolved before the

Table 3. Influence of environmental conditions on the expression of CCMs and the resource cost of CCM operation versus diffusive entry of CO₂. Effects on CCMs of environmental factors, and the direction of change of these environmental factors in algal and aquatic plant habitats with global environmental change between icehouse episodes. Modified from [10]. Further details and references are given in the text and in [10–12,101,105,106,110,112–120]. Predicted resource costs of synthesizing and operating a photosynthetic apparatus using a CCM relative to one relying on entry of CO₂ by diffusion [66,68–71,73,91,105,121]. DOC, dissolved organic carbon; n.a., not applicable.

factor	change to algal environment caused by variation in the factor	effects on expression of CCMs and on their affinity for CO ₂	predicted effect of CCM expression on resource costs of synthesis and (for PAR) operation of the photosynthetic apparatus
CO ₂	increase in CO ₂ in essentially all environments, although less predictable effect in freshwaters which can be out of equilibrium with the atmosphere	decreased inorganic carbon affinity with growth at high CO ₂ ; can be a switch to diffusive CO ₂ entry in some eukaryotes	no general effect on carbon cost of synthesis of the photosynthetic apparatus
temperature	increase in temperature in all environments	prediction of increased CCM expression, or increased fraction of organisms with CCMs, at higher temperatures as a result of lower CO ₂ solubility, is not uniformly supported by the available data.	inapplicable: temperature (Boltzmann energy) is not resource consumed in the synthesis of the photosynthetic apparatus, in its operation or in its maintenance
PAR	increase in PAR in pelagic planktonic environments	decreased inorganic carbon affinity with growth at low PAR	possible decreased energy needed in synthesis of photosynthetic apparatus which uses a CCM. Energy saving in operation of a CCM if there is low CO ₂ leakage and a low CO ₂ affinity and low CO ₂ :O ₂ selectivity
nitrogen	decrease in combined nitrogen in upper mixed layer of lotic environments	generally increased inorganic carbon affinity with growth at low NO ₃ ⁻ . One example each of decreased carbon affinity with growth at lowest NO ₃ ⁻ concentration tested, and with growth over entire NH ₄ ⁺ range tested.	decreased nitrogen cost of synthesis of the photosynthetic apparatus incorporating a CCM if the savings in the synthesis of smaller amounts per cell of Rubisco and of the PCOC enzymes are not offset by the nitrogen cost of the synthesis of CCM components. No consumption of nitrogen in the operation of the CCM
phosphorus	decrease in phosphate in upper mixed layer of lotic environments	two examples of increased inorganic carbon affinity, two examples of decreased inorganic carbon affinity, with growth at low phosphate	where there is a decreased protein content in the photosynthetic apparatus there could be a corresponding decrease in the need for phosphorus needed to synthesize the photosynthetic apparatus as a result of decreased requirement for RNA
iron	probable decrease in iron in upper mixed layer of lotic environments	one example of increased inorganic carbon affinity with growth at low iron, one example of no effect	decreased Fe content of the photosynthetic apparatus if the decreased requirement for NADPH in the near-absence of Rubisco oxygenase activity, and the PCOC, with correspondingly lower content of non-cyclic electron transport components, is not offset by the additional thylakoid components needed for the additional ATP requirement for CCM operation, especially if this additional ATP is made using cyclic photophosphorylation using only photosystem I with its high Fe content
zinc	as for iron	no direct measurements	variable predictions of relative zinc requirements depending on CCM mechanism

(Continued.)

Table 3. (Continued.)

factor	change to algal environment caused by variation in the factor	effects on expression of CCMs and on their affinity for CO ₂	predicted effect of CCM expression on resource costs of synthesis and (for PAR) operation of the photosynthetic apparatus
UV-A	increase in UV-A in lotic planktonic environments, but decrease with higher concentration of DOC	no data	n.a.
UV-B	increase in UV-B in lotic planktonic environments, but decrease with higher concentration of DOC	variable responses of CCMs with increased UV-B flux for growth	n.a.

Pleistocene, they would have had to have survived intervening period(s) of higher CO₂ and higher temperatures. The mechanisms of retention of CCMs in these apparently unfavourable environments are now considered in the context of the response of present day CCMs to such environments.

8. RETENTION OF CO₂-CONCENTRATING MECHANISMS IN HIGH-CO₂ EPISODES

It is possible to argue for long-lasting low-CO₂ microhabitats, e.g. in benthic microbial mats, including stromatolites, where inorganic carbon diffusion from the bulk medium into the cells is restricted by thick diffusion boundary layers [79]. Biogeochemically more important are planktonic habitats where such low CO₂ refuges are less plausible. CCM retention is considered in the context of present work on the response of phytoplankton to current and expected environmental change, and especially increasing CO₂ and the associated warming.

More widespread (planktonic) retention of CCMs in relatively high CO₂ concentrations has been argued by Raven *et al.* [10] in the context of what might happen with increasing CO₂ and temperature over the next several decades. The argument here is that the increased buoyancy of warmer surface waters will lead to a shoaling of the thermocline, which will decrease fluxes of combined nitrogen and of phosphorus from the deeper ocean where mineralization of sinking particulate organic matter occurs, which have been modelled as decreasing global marine primary productivity [126–128]. This decreased nutrient supply, with the increased mean flux of photosynthetically active radiation (and UV-B) incident on the cells in the shallower upper mixed layer will, on the basis of observations on extant phytoplankton (table 3), favour retention of CCMs despite higher CO₂ concentrations [10].

A further relevant consideration which was not mentioned by Raven *et al.* [10] is that a warmer upper mixed layer has decreased oxygen solubility. For the same rate of net oxygen production at two temperatures, there will be greater degree of oxygen supersaturation and hence a greater loss of oxygen to the atmosphere at the higher temperature. Together with the decreased solute transfer between the upper mixed layer and deeper ocean, there is less transfer of oxygen below the

thermocline [129–131]. The widespread, but not universal, decrease in calcification by calcified plankton [132–134], and the much smaller effect on silicification by diatoms [135] in a higher CO₂, warmer ocean means less ballasting of sinking particulate organic matter, hence slower sinking and more microbial mineralization just below the thermocline ([136], cf. [137]). While this higher nutrient concentration just below the thermocline might be expected to partly offset the lower rate constant for nutrient transfer to the upper mixed layer, another factor must be considered. The combination of more microbial respiration and increased oxygen supply can lead to hypoxia and even anoxia in certain sub-surface waters, with implications for loss of the nitrate and nitrite forms of combined nitrogen produced from organic matter by mineralization and nitrification in less deoxygenated zones followed by denitrification or the anammox reaction in more deoxygenated places [131], with a decrease in the nitrogen:phosphorus ratio in the nutrients reaching the upper mixed layer. Although, in the long term (thousands of years and more), a warmer world would heat the ocean interior as well as the upper mixed layer and potentially decrease the extent to which the thermocline shoals, there is a well-established correlation of widespread deep-ocean anoxia (and even euxinia) with warmer, high-CO₂ episodes in Earth history [138], so at least a decreased upward flux of combined nitrogen across the thermocline would continue, favouring retention of CCMs.

These arguments are based on the response of extant algae to the changes occurring, and predicted, in their environment as a result of increased CO₂ and temperature, with downstream effects on the marine and inland water inorganic system, the mixed layer depth and water body oxygenation. This can be used to inform us of how CCMs were retained in past episodes of high CO₂. Dealing first with the organisms studied, almost all of the work has been carried out with organisms which have only been exposed to the increased CO₂ and associated changes in temperature and the availability of other resources (table 3) for time periods of days to weeks. This time period allows 1–100 generation, meaning that only regulatory (altering the existing proteome by post-translational modification, and changes in the metabolome) and acclimatory (altering the expressed proteome based on the existing genome) [139] responses of extant

algae can be expressed. In very few cases has relevant evolutionary evidence been sought in laboratory experiments for increased CO₂ [139–146] and, using different methods, higher temperatures [147]. An example of where evolution has been unable to cope with natural CO₂ enrichment present for several decades concerns calcified red algae growing on seagrass leaves near an underwater vent in the Mediterranean [148]. Even for studies of regulation and acclimation there can be problems with the length of time that an alga has been in culture and frequently exposed to CO₂, nitrogen, phosphorus, PAR, UV and temperature which has little relevance to their natural environment [149]. Furthermore, there is a relative lack of studies in which changes in CO₂ have been combined with other relevant environmental changes: table 3 and Raven *et al.* [10] analyse and give references to the important work in which such interactions have been studied. Finally, there are some relatively poorly constrained factors of ocean chemistry in the past [80,150–154], and until recently little consensus as to the appropriate methodology [134,155–158] for laboratory and mesocosm experimentation on increased CO₂.

Despite these reservations, which also apply to other models of past and future atmosphere–ocean–organism interactions, the suggestions of Raven *et al.* [27] provide a lead into further studies in how CCMs could be retained in lengthy episodes between shorter low-CO₂ episodes. Any retention of CCMs in high-CO₂ episodes would be a further complication in the use of stable carbon isotope ratios of phytoplankton-derived organic carbon from marine sediments as a palaeobarometer for CO₂, since most marine phytoplankton today have CCMs [67,98,159,160]. Organisms lacking CCMs, e.g. terrestrial liverworts and mosses, do not suffer from this problem when used in palaeobarometry of CO₂ [161].

9. CONCLUSIONS

The changing CO₂ and O₂ concentration over the last 2.4 Gyr have had significant effects on the physiology and ecology of cyanobacteria and algae. From the presumed ancestral diffusive CO₂ entry to Rubisco, all extant cyanobacteria have CCMs, Rubiscos with high CO₂-saturated catalytic activity and low CO₂ affinity and CO₂/O₂ selectivity, and an essential role for the capacity to convert the 2-phosphoglycolate formed as a very small fraction of the total carbon flux into triose phosphates. Most eukaryotic algae have CCMs: a greater fraction has CCMs in the sea than in freshwaters, and there is no strong relationship to water temperatures. The evolution of CCMs can apparently be related to decreased CO₂ availability and to the presence of oxygen, modulated by the kinetics of the form of Rubisco in the organisms, with some components of the CCMs adapted in evolution from the roles in other pathways. The retention of CCMs during the high-CO₂ episodes predominant through Earth history could have been related in part to the interaction of CCM expression with other environmental factors which change in high-CO₂ water bodies.

The University of Dundee is a registered Scottish Charity, No SC10596. J.A.R.'s research on inorganic carbon acquisition was funded by BBSRC UK and NERC UK. J.A.R. thanks David Beerling for the invitation to lecture at the meeting at the Kavli Royal Society International Centre from which this paper, and the Theme Issue, arose, and also Anthony Cohen for the invitation to the Ocean Deoxygenation meeting at the Kavli Royal Society International Centre which expanded his reading on ocean deoxygenation. J.B. is grateful to the Australian Research Council for their support of his work on inorganic carbon acquisition in relation to climate change. M.G.'s research on CCMs was partially funded by the Fondazione Cariverona, Italy, the Southeast Wisconsin Energy Technology Research Center, USA and the Ministry for Agriculture and Forestry (MIPAF), Italy (project Bioforme). We thank Ian McCulloch for his help in converting the references from Harvard to Vancouver.

REFERENCES

- 1 Rasmussen, B., Fletcher, I. R., Brocks, J. J. & Kilburn, M. R. 2008 Reassessing the first appearance of eukaryotes and cyanobacteria. *Nature* **455**, 1101–1104. (doi:10.1038/nature07381)
- 2 Butterfield, N. J. 2000 *Bangiomorpha pubescens* n. gen., n. sp.: implications for the evolution of sex, multicellularity, and the Mesoproterozoic/Neoproterozoic radiation of eukaryotes. *Paleobiology* **26**, 386–404. (doi:10.1666/0094-8373(2000)026<0386:bpngns<2.0.co;2)
- 3 Butterfield, N. J. 2004 A vaucheriacean alga from the middle Neoproterozoic of Spitsbergen: implications for the evolution of Proterozoic eukaryotes and the Cambrian explosion. *Paleobiology* **30**, 231–252. (doi:10.1666/0094-8373(2004)030<0231:avafm<2.0.co;2)
- 4 Butterfield, N. J., Knoll, A. H. & Swett, K. 1988 Exceptional preservation of fossils in an upper Proterozoic shale. *Nature* **334**, 424–427. (doi:10.1038/334424a0)
- 5 Strother, P. K., Battison, L., Brasier, M. D. & Wellman, C. H. 2011 Earth's earliest non-marine eukaryotes. *Nature* **473**, 505–509. (doi:10.1038/nature09943)
- 6 Taylor, T. N., Taylor, E. L. & Krings, M. 2009 *Paleobotany: the biology and evolution of fossil plants*, 2nd edn. London, UK: Academic.
- 7 Graham, L. E. & Wilcox, L. W. 2000 *Algae*. Upper Saddle River, NJ: Prentice Hall.
- 8 Van den Hoek, C., Mann, D. G. & Jahns, H. M. 1995 *Algae: an introduction to phycology*. Cambridge, UK: Cambridge University Press.
- 9 Maberly, S. C., Ball, L. A., Raven, J. A. & Suetemeyer, D. 2009 Inorganic carbon acquisition by chrysophytes. *J. Phycol.* **45**, 1052–1061. (doi:10.1111/j.1529-8817.2009.00734.x)
- 10 Raven, J. A., Giordano, M., Beardall, J. & Maberly, S. C. 2011 Algal and aquatic plant carbon concentrating mechanisms in relation to environmental change. *Photosynth. Res.* **109**, 1–16. (doi:10.1007/s11120-011-9632-6)
- 11 Raven, J. A. *et al.* 2002 Seaweeds in cold seas: evolution and carbon acquisition. *Ann. Bot.* **90**, 525–536. (doi:10.1093/aob/mcf171)
- 12 Raven, J. A. *et al.* 2002 Mechanistic interpretation of carbon isotope discrimination by marine macroalgae and seagrasses. *Funct. Plant Biol.* **29**, 355–378. (doi:10.1071/pp01201)
- 13 Raven, J. A. 2010 Inorganic carbon acquisition by eukaryotic algae: four current questions. *Photosynth. Res.* **106**, 123–134. (doi:10.1007/s11120-010-9563-7)
- 14 Badger, M. R., Andrews, T. J., Whitney, S. M., Ludwig, M., Yellowlees, D. C., Leggat, W. & Price, G. D. 1998

- The diversity and coevolution of Rubisco, plastids, pyrenoids, and chloroplast-based CO₂-concentrating mechanisms in algae. *Can. J. Bot. Rev. Can. Bot.* **76**, 1052–1071. (doi:10.1139/cjb-76-6-1052)
- 15 Sanchez-Baracaldo, P., Hayes, P. K. & Blank, C. E. 2005 Morphological and habitat evolution in the Cyanobacteria using a compartmentalization approach. *Geobiology* **3**, 145–165. (doi:10.1111/j.1472-4669.2005.00050.x)
- 16 Blank, C. E. & Sanchez-Baracaldo, P. 2010 Timing of morphological and ecological innovations in the cyanobacteria—a key to understanding the rise in atmospheric oxygen. *Geobiology* **8**, 1–23. (doi:10.1111/j.1472-4669.2009.00220.x)
- 17 Schirmermeister, B. E., Antonelli, A. & Bagheri, H. C. 2011 The origin of multicellularity in cyanobacteria. *BMC Evol. Biol.* **11**, 45. (doi:10.1186/1471-2148-11-45)
- 18 Teyssedre, B. 2006 Are the green algae (phylum Viridiplantae) two billion years old? *Carnets Geol.* **3**, 1–15.
- 19 Becker, B. & Marin, B. 2009 Streptophyte algae and the origin of embryophytes. *Ann. Bot.* **103**, 999–1004. (doi:10.1093/aob/mcp044)
- 20 Lewis, L. A. & McCourt, R. M. 2004 Green algae and the origin of land plants. *Am. J. Bot.* **91**, 1535–1556. (doi:10.3732/ajb.91.10.1535)
- 21 Derenne, S., Metzger, P., Largeau, C., Vanbergen, P. F., Gatellier, J. P., Damste, J. S. S., Deleuw, J. W. & Berkaloﬀ, C. 1992 Similar morphological and chemical variations of *Gloeocapsomorpha prisca* in Ordovician sediments and cultured *Botryococcus braunii* as a response to changes in salinity. *Org. Geochem.* **19**, 299–313. (doi:10.1016/0146-6380(92)90001-e)
- 22 Senousy, H. H., Beakes, G. W. & Hack, E. 2004 Phylogenetic placement of *Botryococcus braunii* (Trebouxiophyceae) and *Botryococcus sudeticus* isolate UTEX 2629 (Chlorophyceae). *J. Phycol.* **40**, 412–423. (doi:10.1046/j.1529-8817.2004.03173.x)
- 23 Zhang, Z., Metzger, P. & Sachs, J. P. 2007 Biomarker evidence for the co-occurrence of three races (A, B and L) of *Botryococcus braunii* in El Junco Lake, Galápagos. *Org. Geochem.* **38**, 1459–1478. (doi:10.1016/j.orggeochem.2007.05.015)
- 24 Rubinstein, C. V., Gerrienne, P., de la Puente, G. S., Astini, R. A. & Steemans, P. 2010 Early Middle Ordovician evidence for land plants in Argentina (Eastern Gondwana). *New Phytol.* **188**, 365–369. (doi:10.1111/j.1469-8137.2010.03433.x)
- 25 Wellman, C. H. 2010 The invasion of the land by plants: when and where? *New Phytol.* **188**, 306–309. (doi:10.1111/j.1469-8137.2010.03471.x)
- 26 Xiao, S. H., Knoll, A. H., Yuan, X. L. & Poeschel, C. M. 2004 Phosphatized multicellular algae in the Neoproterozoic Doushantuo Formation, China, and the early evolution of florideophyte red algae. *Am. J. Bot.* **91**, 214–227. (doi:10.3732/ajb.91.2.214)
- 27 Yuan, X., Chen, Z., Xiao, S., Zhou, C. & Hua, H. 2011 An early Ediacaran assemblage of macroscopic and morphologically differentiated eukaryotes. *Nature* **470**, 389–392. (doi:10.1038/nature09810)
- 28 Raven, J. A. & Waite, A. M. 2004 The evolution of silicification in diatoms: inescapable sinking and sinking as escape? *New Phytol.* **162**, 45–61. (doi:10.1111/j.1469-8137.2004.01022.x)
- 29 Sims, P. A., Mann, D. G. & Medlin, L. K. 2006 Evolution of the diatoms: insights from fossil, biological and molecular data. *Phycologia* **45**, 361–402. (doi:10.2216/05-22.1)
- 30 Xu, Z. L. 2001 New discoveries of phaeophycean fossils in the Early Cambrian, Haikou, Kunming, Yunnan, southwest China. *Acta Bot. Sin.* **43**, 1072–1076.
- 31 Javaux, E. J., Knoll, A. H. & Walter, M. 2003 Recognizing and interpreting the fossils of early eukaryotes. *Orig. Life Evol. Biosphere* **33**, 75–94. (doi:10.1023/a:1023992712071)
- 32 Shields-Zhou, G. & Och, L. 2011 The case for a Neoproterozoic oxygenation event: geochemical evidence and biological consequences. *GSA Today* **21**, 4–11. (doi:10.1130/GSATG102A.1)
- 33 Berner, R. A. 2006 GEOCARBSULF: a combined model for Phanerozoic atmospheric O₂ and CO₂. *Geochim. Cosmochim. Acta* **70**, 5653–5664. (doi:10.1016/j.gca.2005.11.032)
- 34 Raven, J. A., Giordano, M. & Beardall, J. 2008 Insights into the evolution of CCMs from comparisons with other resource acquisition and assimilation processes. *Physiol. Plant* **133**, 4–14. (doi:10.1111/j.1399-3054.2007.01024.x)
- 35 Sheldon, N. D. 2006 Precambrian paleosols and atmospheric CO₂ levels. *Precambrian Res.* **147**, 148–155. (doi:10.1016/j.precamres.2006.02.004)
- 36 Riding, R. 2006 Cyanobacterial calcification, carbon dioxide concentrating mechanisms, and Proterozoic–Cambrian changes in atmospheric composition. *Geobiology* **4**, 299–316. (doi:10.1111/j.1472-4669.2006.00087.x)
- 37 Bao, H., Lyons, J. R. & Zhou, C. 2008 Triple oxygen isotope evidence for elevated CO₂ levels after a Neoproterozoic glaciation. *Nature* **453**, 504–506. (doi:10.1038/nature06959)
- 38 Bjerrum, C. J. & Canfield, D. E. 2011 Towards a quantitative understanding of the late Neoproterozoic carbon cycle. *Proc. Natl Acad. Sci. USA* **108**, 5542–5547. (doi:10.1073/pnas.1101755108)
- 39 Riding, R. 2009 An atmospheric stimulus for cyanobacterial-bioinduced calcification ca. 350 million years ago? *Palaio* **24**, 685–696. (doi:10.2110/palo.2009.p09-033r)
- 40 Raven, J. A. 2009 Contributions of anoxygenic and oxygenic phototrophy and chemolithotrophy to carbon and oxygen fluxes in aquatic environments. *Aquat. Microb. Ecol.* **56**, 177–192. (doi:10.3354/ame01315)
- 41 Bar-Even, A., Noor, E., Lewis, N. E. & Milo, R. 2010 Design and analysis of synthetic carbon fixation pathways. *Proc. Natl Acad. Sci. USA* **107**, 8889–8894. (doi:10.1073/pnas.0907176107)
- 42 Berg, I. A., Kockelkorn, D., Ramos-Vera, W. H., Say, R. F., Zarzycki, J., Huegler, M., Alber, B. E. & Fuchs, G. 2010 Autotrophic carbon fixation in archaea. *Nat. Rev. Microbiol.* **8**, 447–460. (doi:10.1038/nrmicro2365)
- 43 Berg, I. A. 2011 Ecological aspects of the distribution of different autotrophic CO₂ fixation pathways. *Appl. Environ. Microbiol.* **77**, 1925–1936. (doi:10.1128/aem.02473-10)
- 44 Boyle, N. R. & Morgan, J. A. 2011 Computation of metabolic fluxes and efficiencies for biological carbon dioxide fixation. *Metabol. Eng.* **13**, 150–158. (doi:10.1016/j.ymben.2011.01.005)
- 45 Huegler, M. & Sievert, S. M. 2011 Beyond the Calvin cycle: autotrophic carbon fixation in the ocean. *Annu. Rev. Mar. Sci.* **3**, 261–289. (doi:10.1146/annurev-marine-120709-142712)
- 46 Ljungdhal, L. G. 1986 The autotrophic pathway of acetate synthesis in acetogenic bacteria. *Annu. Rev. Microbiol.* **40**, 415–450. (doi:10.1146/annurev.mi.40.100186.002215)
- 47 Tabita, F. R., Hanson, T. E., Li, H., Satagopan, S., Singh, J. & Chan, S. 2007 Function, structure, and evolution of the RubisCO-like proteins and their RubisCO homologs. *Microbiol. Mol. Biol. Rev.* **71**, 576. (doi:10.1128/mmmbr.00015-07)
- 48 Tabita, F. R., Hanson, T. E., Satagopan, S., Witte, B. H. & Kreeel, N. E. 2008 Phylogenetic and evolutionary

- relationships of RubisCO and the RubisCO-like proteins and the functional lessons provided by diverse molecular forms. *Phil. Trans. R. Soc. B* **363**, 2629–2640. (doi:10.1098/rstb.2008.0023)
- 49 Tabita, F. R., Satagopan, S., Hanson, T. E., Kreel, N. E. & Scott, S. S. 2008 Distinct form I, II, III, and IV Rubisco proteins from the three kingdoms of life provide clues about Rubisco evolution and structure/function relationships. *J. Exp. Bot.* **59**, 1515–1524. (doi:10.1093/jxb/erm361)
- 50 Marin, B., Nowack, E., Glockner, G. & Melkonian, M. 2007 The ancestor of the *Paulinella* chromatophore obtained a carboxysomal operon by horizontal gene transfer from a *Nitrococcus*-like γ -proteobacterium. *BMC Evol. Biol.* **7**, 85. (doi:10.1186/1471-2148-7-85)
- 51 Badger, M. R., Hanson, D. & Price, G. D. 2002 Evolution and diversity of CO₂ concentrating mechanisms in cyanobacteria. *Funct. Plant Biol.* **29**, 161–173. (doi:10.1071/pp01213)
- 52 Badger, M. R. & Price, G. D. 2003 CO₂ concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution. *J. Exp. Bot.* **54**, 609–622. (doi:10.1093/jxb/erg076)
- 53 Badger, M. R., Price, G. D., Long, B. M. & Woodger, F. J. 2006 The environmental plasticity and ecological genomics of the cyanobacterial CO₂ concentrating mechanism. *J. Exp. Bot.* **57**, 249–265. (doi:10.1093/jxb/eri286)
- 54 Marin, B., Nowack, E. C. M. & Melkonian, M. 2005 A plastid in the making: evidence for a second primary endosymbiosis. *Protist* **156**, 425–432. (doi:10.1016/j.protis.2005.09.001)
- 55 Nowack, E. C. M., Melkonian, M. & Gloeckner, G. 2008 Chromatophore genome sequence of *Paulinella* sheds light on acquisition of photosynthesis by eukaryotes. *Curr. Biol.* **18**, 410–418. (doi:10.1016/j.cub.2008.02.051)
- 56 Price, G. D., Badger, M. R., Woodger, F. J. & Long, B. M. 2008 Advances in understanding the cyanobacterial CO₂-concentrating mechanism (CCM): functional components, C_i transporters, diversity, genetic regulation and prospects for engineering into plants. *J. Exp. Bot.* **59**, 1441–1461. (doi:10.1093/jxb/erm112)
- 57 Deusch, O., Landan, G., Roettger, M., Gruenheit, N., Kowallik, K. V., Allen, J. F., Martin, W. & Dagan, T. 2008 Genes of cyanobacterial origin in plant nuclear genomes point to a heterocyst-forming plastid ancestor. *Mol. Biol. Evol.* **25**, 748–761. (doi:10.1093/molbev/msn022)
- 58 Janouskovec, J., Horak, A., Obornik, M., Lukes, J. & Keeling, P. J. 2010 A common red algal origin of the apicomplexan, dinoflagellate, and heterokont plastids. *Proc. Natl Acad. Sci. USA* **107**, 10 949–10 954. (doi:10.1073/pnas.1003335107)
- 59 Martin, W. & Schnarrenberger, C. 1997 The evolution of the Calvin cycle from prokaryotic to eukaryotic chromosomes: a case study of functional redundancy in ancient pathways through endosymbiosis. *Curr. Genet.* **32**, 1–18. (doi:10.1007/s002940050241)
- 60 Reyes-Prieto, A. & Bhattacharya, D. 2007 Phylogeny of Calvin cycle enzymes supports Plantae monophyly. *Mol. Phylogenet. Evol.* **45**, 384–391. (doi:10.1016/j.ympev.2007.02.026)
- 61 Kroth, P. G. *et al.* 2008 A model for carbohydrate metabolism in the diatom *Phaeodactylum tricornutum* deduced from comparative whole genome analysis. *PLoS ONE* **3**, e1426. (doi:10.1371/journal.pone.0001426)
- 62 Moustafa, A., Beszteri, B., Maier, U. G., Bowler, C., Valentin, K. & Bhattacharya, D. 2009 Genomic footprints of a cryptic plastid endosymbiosis in diatoms. *Science* **324**, 1724–1726. (doi:10.1126/science.1172983)
- 63 Maberly, S. C., Courcelle, C., Groben, R. & Gontero, B. 2010 Phylogenetically-based variation in the regulation of the Calvin cycle enzymes, phosphoribulokinase and glyceraldehyde-3-phosphate dehydrogenase, in algae. *J. Exp. Bot.* **61**, 735–745. (doi:10.1093/jxb/erp337)
- 64 Groben, R., Kaloudas, D., Raines, C. A., Offmann, B., Maberly, S. C. & Gontero, B. 2010 Comparative sequence analysis of CP12, a small protein involved in the formation of a Calvin cycle complex in photosynthetic organisms. *Photosynth. Res.* **103**, 183–194. (doi:10.1007/s11120-010-9542-z)
- 65 Tcherkez, G. G. B., Farquhar, G. D. & Andrews, T. J. 2006 Despite slow catalysis and confused substrate specificity, all ribulose biphosphate carboxylases may be nearly perfectly optimized. *Proc. Natl Acad. Sci. USA* **103**, 7246–7251. (doi:10.1073/pnas.0600605103)
- 66 Raven, J. A. 2011 The cost of photoinhibition. *Physiol. Plant* **142**, 87–104. (doi:10.1111/j.1399-3054.2011.01465.x)
- 67 Giordano, M., Beardall, J. & Raven, J. A. 2005 CO₂ concentrating mechanisms in algae: mechanisms, environmental modulation, and evolution. *Annu. Rev. Plant Biol.* **56**, 99–131. (doi:10.1146/annurev.arplant.56.032604.144052)
- 68 Raven, J. A. 1990 Predictions of Mn and Fe use efficiencies of phototrophic growth as a function of light availability for growth and of C assimilation pathway. *New Phytol.* **116**, 1–18. (doi:10.1111/j.1469-8137.1990.tb00505.x)
- 69 Raven, J. A. 1991 Physiology of inorganic C acquisition and implications for resource use efficiency by marine-phytoplankton—relation to increased CO₂ and temperature. *Plant Cell Environ.* **14**, 779–794. (doi:10.1111/j.1365-3040.1991.tb01442.x)
- 70 Raven, J. A. 1991 Implications of inorganic carbon utilization—ecology, evolution, and geochemistry. *Can. J. Bot. Rev. Can. Bot.* **69**, 908–924. (doi:10.1139/b91-118)
- 71 Raven, J. A. & Johnston, A. M. 1991 Mechanisms of inorganic carbon acquisition in marine phytoplankton and their implications for the use of other resources. *Limnol. Oceanogr.* **36**, 1701–1714. (doi:10.4319/lo.1991.36.8.1701)
- 72 Falkowski, P. G. & Godfrey, L. V. 2008 Electrons, life and the evolution of Earth's oxygen cycle. *Phil. Trans. R. Soc. B* **363**, 2705–2716. (doi:10.1098/rstb.2008.0054)
- 73 Raven, J. A., Andrews, M. & Quigg, A. 2005 The evolution of oligotrophy: implications for the breeding of crop plants for low input agricultural systems. *Ann. Appl. Biol.* **146**, 261–280. (doi:10.1111/j.1744-7348.2005.040138.x)
- 74 Raven, J. A., Brown, K., Mackay, M., Beardall, J., Giordano, M., Granum, E., Leegood, R. C., Kilminster, K. & Walker, D. I. 2005 Iron, nitrogen, phosphorus and zinc cycling and consequences for primary productivity in the oceans. In *Micro-organisms and earth systems—advances in geomicrobiology* (eds G. M. Gadd, K. T. Semple & H. M. Lappin-Scott), pp. 247–272. Cambridge, UK: Cambridge University Press.
- 75 Grime, J. P. 2001 *Plant strategies, vegetation processes, and ecosystem properties*, 2nd edn. Chichester, UK: John Wiley.
- 76 Reynolds, C. S. 2006 *Ecology of phytoplankton*. Cambridge, UK: Cambridge University Press.
- 77 Flynn, K. J. 2009 Going for the slow burn: why should possession of a low maximum growth rate be advantageous for microalgae? *Plant Ecol. Diversity* **2**, 179–189. (doi:10.1080/17550870903207268)

- 78 Lambers, H., Raven, J. A., Shaver, G. R. & Smith, S. E. 2008 Plant nutrient-acquisition strategies change with soil age. *Trends Ecol. Evol.* **23**, 95–103. (doi:10.1016/j.tree.2007.10.008)
- 79 Raven, J. A., Cockell, C. S. & De La Rocha, C. L. 2008 The evolution of inorganic carbon concentrating mechanisms in photosynthesis. *Phil. Trans. R. Soc. B* **363**, 2641–2650. (doi:10.1098/rstb.2008.0020)
- 80 Falkowski, P. G. & Raven, J. A. 2007 *Aquatic photosynthesis*, 2nd edn. Princeton, NJ: Princeton University Press.
- 81 Raven, J. A. & Larkum, A. W. D. 2007 Are there ecological implications for the proposed energetic restrictions on photosynthetic oxygen evolution at high oxygen concentrations? *Photosynth. Res.* **94**, 31–42. (doi:10.1007/s11120-007-9211-z)
- 82 Shevela, D., Beckmann, K., Clausen, J., Junge, W. & Messinger, J. 2011 Membrane-inlet mass spectrometry reveals a high driving force for oxygen production by photosystem II. *Proc. Natl Acad. Sci. USA* **108**, 3602–3607. (doi:10.1073/pnas.1014249108)
- 83 Evans, D. A., Beukes, N. J. & Kirschvink, J. L. 1997 Low-latitude glaciation in the Palaeoproterozoic era. *Nature* **386**, 262–266. (doi:10.1038/386262a0)
- 84 Schwarte, S. & Bauwe, H. 2007 Identification of the photorespiratory 2-phosphoglycolate phosphatase, PGLP1, in *Arabidopsis*. *Plant Physiol.* **144**, 1580–1586. (doi:10.1104/pp.107.099192)
- 85 Bauwe, H., Hagemann, M. & Fernie, A. R. 2010 Photorespiration: players, partners and origin. *Trends Plant Sci.* **15**, 330–336. (doi:10.1016/j.tplants.2010.03.006)
- 86 Kim, Y. C. et al. 2004 Structure- and function-based characterization of a new phosphoglycolate phosphatase from *Thermoplasma acidophilum*. *J. Biol. Chem.* **279**, 517–526. (doi:10.1074/jbc.M306054200)
- 87 Pellicer, M. T., Nunez, M. F., Aguilar, J., Badia, J. & Baldoma, L. 2003 Role of 2-phosphoglycolate phosphatase of *Escherichia coli* in metabolism of the 2-phosphoglycolate formed in DNA repair. *J. Bacteriol.* **185**, 5815–5821. (doi:10.1128/jb.185.19.5815-5821.2003)
- 88 Kern, R., Bauwe, H. & Hagemann, M. 2011 Evolution of enzymes involved in the photorespiratory 2-phosphoglycolate cycle from cyanobacteria via algae toward plants. *Photosynth. Res.* **109**, 1–12. (doi:10.1007/s11120-010-9615-z)
- 89 Eisenhut, M. et al. 2006 The plant-like C₂ glycolate cycle and the bacterial-like glycerate pathway cooperate in phosphoglycolate metabolism in cyanobacteria. *Plant Physiol.* **142**, 333–342. (doi:10.1104/pp.106.082982)
- 90 Eisenhut, M., Ruth, W., Haimovich, M., Bauwe, H., Kaplan, A. & Hagemann, M. 2008 The photorespiratory glycolate metabolism is essential for cyanobacteria and might have been conveyed endosymbiotically to plants. *Proc. Natl. Acad. Sci. USA* **105**, 17 199–17 204. (doi:10.1073/pnas.0807043105)
- 91 Raven, J. A., Kubler, J. E. & Beardall, J. 2000 Put out the light, and then put out the light. *J. Mar. Biol. Assoc. UK* **80**, 1–25. (doi:10.1017/s0025315499 001526)
- 92 Roberts, K., Granum, E., Leegood, R. & Raven, J. 2007 Carbon acquisition by diatoms. *Photosynth. Res.* **93**, 79–88. (doi:10.1007/s11120-007-9172-2)
- 93 Bartsch, O., Hagemann, M. & Bauwe, H. 2008 Only plant-type (GLYK) glycerate kinases produce D-glycerate-3-phosphate. *FEBS Lett.* **582**, 3025–3028. (doi:10.1016/j.febslet.2008.07.038)
- 94 Boldt, R., Edner, C., Kolkusaoglu, U., Hagemann, M., Weckwerth, W., Wienkoop, S., Morgenthal, K. & Bauwe, H. 2005 D-Glycerate-3-kinase, the last unknown enzyme in the photorespiratory cycle in *Arabidopsis*, belongs to a novel kinase family. *Plant Cell* **17**, 2413–2420. (doi:10.1105/tpc.105.033993)
- 95 Roberts, K., Granum, E., Leegood, R. C. & Raven, J. A. 2007 C₃ and C₄ pathways of photosynthetic carbon assimilation in marine diatoms are under genetic, not environmental, control. *Plant Physiol.* **145**, 230–235. (doi:10.1104/pp.107.102616)
- 96 Holland, H. D. 1978 *The chemistry of the atmosphere and oceans*. New York, NY: Wiley.
- 97 Lenton, T. M. & Watson, A. J. 2000 Redfield revisited: 2 What regulates the oxygen content of the atmosphere? *Global Biogeochem. Cycles* **14**, 249–268. (doi:10.1029/1999gb900076)
- 98 Johnston, A. M. & Kennedy, H. 1998 Carbon isotope fractionation in marine systems: open ocean studies and laboratory studies. In *Stable isotopes: integration of biological, ecological and geochemical processes* (ed. H. Griffiths), pp. 239–256. Oxford, UK: BIOS Scientific.
- 99 Maberly, S. C. 1990 Exogenous sources of inorganic carbon for photosynthesis by marine macroalgae. *J. Phycol.* **26**, 439–449. (doi:10.1111/j.0022-3646.1990.00439.x)
- 100 Hepburn, C. D., Pritchard, D. W., Cornwall, C. E., McLeod, R. J., Beardall, J., Raven, J. A. & Hurd, C. L. 2011 Diversity of carbon use strategies in a kelp forest community: implications for a high CO₂ ocean. *Global Change Biol.* **17**, 2488–2497. (doi:10.1111/j.1365-2486.2011.02411.x)
- 101 Marconi, M., Giordano, M. & Raven, J. A. 2011 Impact of taxonomy, geography and depth of on δ¹³C and δ¹⁵N variation in a large collection of macroalgae. *J. Phycol.* **47**, 1023–1035. (doi:10.1111/j.1529-88172011.01045.x)
- 102 Raven, J. A. 2011 Carbon. In *Ecology of cyanobacteria* (ed. B. A. Whitton), 2nd edn. Berlin, Germany: Springer.
- 103 Hopkinson, B. M., Dupont, C. L., Allen, A. E. & Morel, F. M. M. 2011 Efficiency of the CO₂-concentrating mechanism of diatoms. *Proc. Natl Acad. Sci. USA* **108**, 3830–3837. (doi:10.1073/pnas.1018062108)
- 104 Reinfelder, J. R. 2011 Carbon concentrating mechanisms in eukaryotic marine phytoplankton. *Annu. Rev. Mar. Sci.* **3**, 291–315. (doi:10.1146/annurev-marine-120709-142720)
- 105 Raven, J. A., Ball, L. A., Beardall, J., Giordano, M. & Maberly, S. C. 2005 Algae lacking carbon-concentrating mechanisms. *Can. J. Bot.-Rev. Can. Bot.* **83**, 879–890. (doi:10.1139/b05-074)
- 106 Spijkerman, E. 2011 The expression of a carbon concentrating mechanism in *Chlamydomonas acidophila* under variable phosphorus, iron, and CO₂ concentrations. *Photosynth. Res.* **109**, 1–11. (doi:10.1007/s11120-010-9607-z)
- 107 Young, J. N., Rickaby, R. E. M., Kapralov, M. V. & Filatov, D. A. 2012 Adaptive signals in algal Rubisco reveal a history of ancient atmospheric carbon dioxide. *Phil. Trans. R. Soc. B* **367**, 483–492. (doi:10.1098/rstb.2011.0145)
- 108 Sultemeyer, D. & Rinast, K. A. 1996 The CO₂ permeability of the plasma membrane of *Chlamydomonas reinhardtii*: mass-spectrometric ¹⁸O exchange measurements from CO₂ ¹³C-¹⁸O in suspensions of carbonic anhydrase-loaded plasma-membrane vesicles. *Planta* **200**, 358–368. (doi:10.1007/BF00200304)
- 109 Dou, Z., Heinhorst, S., Williams, E. B., Murin, C. D., Shively, J. M. & Cannon, G. C. 2008 CO₂ fixation kinetics of *Halothiobacillus neapolitanus* mutant carboxysomes lacking carbonic anhydrase suggest the shell acts as a diffusional barrier for CO₂. *J. Biol. Chem.* **283**, 10 377–10 384. (doi:10.1074/jbc.M709285200)
- 110 Reinfelder, J. R., Kraepiel, A. M. L. & Morel, F. M. M. 2000 Unicellular C₄ photosynthesis in a marine diatom. *Nature* **407**, 996–999. (doi:10.1038/35039612)

- 111 Giordano, M., Norici, A., Forssen, M., Eriksson, M. & Raven, J. A. 2003 An anaplerotic role for mitochondrial carbonic anhydrase in *Chlamydomonas reinhardtii*. *Plant Physiol.* **132**, 2126–2134. (doi:10.1104/pp.103.023424)
- 112 Beardall, J. & Giordano, M. 2002 Ecological implications of microalgal and cyanobacterial CO₂ concentrating mechanisms, and their regulation. *Funct. Plant Biol.* **29**, 335–347. (doi:10.1071/pp01195)
- 113 Beardall, J., Stojkovic, S. & Larsen, S. 2009 Living in a high CO₂ world: impacts of global climate change on marine phytoplankton. *Plant Ecol. Diversity* **2**, 191–205. (doi:10.1080/17550870903271363)
- 114 Finkel, Z. V., Beardall, J., Flynn, K. J., Quigg, A., Rees, T. A. V. & Raven, J. A. 2010 Phytoplankton in a changing world: cell size and elemental stoichiometry. *J. Plankton Res.* **32**, 119–137. (doi:10.1093/plankt/fbp098)
- 115 Gao, G., Gao, K. & Giordano, M. 2009 Responses to solar UV radiation of the diatom *Skeletonema costatum* (Bacillariophyceae) grown at different Zn²⁺ concentrations. *J. Phycol.* **45**, 119–129. (doi:10.1111/j.1529-8817.2008.00616.x)
- 116 Gao, K. & Zheng, Y. 2010 Combined effects of ocean acidification and solar UV radiation on photosynthesis, growth, pigmentation and calcification of the coralline alga *Corallina sessilis* (Rhodophyta). *Global Change Biol.* **16**, 2388–2398. (doi:10.1111/j.1365-2486.2009.02113.x)
- 117 Ihnken, S., Roberts, S. & Beardall, J. 2011 Differential responses of growth and photosynthesis in the marine diatom *Chaetoceros muelleri* to CO₂ and light availability. *Phycologia* **50**, 182–193. (doi:10.2216/10-11.1)
- 118 Kohinata, T., Nishino, H. & Fukuzawa, H. 2008 Significance of zinc in a regulatory protein, CCM1, which regulates the carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Plant Cell. Physiol.* **49**, 273–283. (doi:10.1093/pcp/pcn003)
- 119 Morel, F. M. M., Cox, E. H., Kraepiel, A. M. L., Lane, T. W., Milligan, A. J., Schaperdorth, I., Reinfelder, J. R. & Tortell, P. D. 2002 Acquisition of inorganic carbon by the marine diatom *Thalassiosira weissflogii*. *Funct. Plant Biol.* **29**, 301–308. (doi:10.1071/pp01199)
- 120 Raven, J. A. & Geider, R. J. 1988 Temperature and algal growth. *New Phytol.* **110**, 441–461. (doi:10.1111/j.1469-8137.1988.tb00282.x)
- 121 Raven, J. A., Evans, M. C. W. & Korb, R. E. 1999 The role of trace metals in photosynthetic electron transport in O₂-evolving organisms. *Photosynth. Res.* **60**, 111–149. (doi:10.1023/a:1006282714942)
- 122 Shelly, K., Héraud, P. & Beardall, J. 2002 Nitrogen limitation in *Dunaliella tertiolecta* (Chlorophyceae) leads to increased susceptibility to damage by ultraviolet-B radiation but also increased repair capacity. *J. Phycol.* **38**, 713–720. (doi:10.1046/j.1529-8817.2002.01147.x)
- 123 Sobrino, C., Neale, P. J., Phillips-Kress, J. D., Moeller, R. E. & Porter, J. A. 2009 Elevated CO₂ increases sensitivity to ultraviolet radiation in lacustrine phytoplankton assemblages. *Limnol. Oceanogr.* **54**, 2448–2459. (doi:10.4319/lo.2009.54.6_part_2.2448)
- 124 Sobrino, C., Ward, M. L. & Neale, P. J. 2008 Acclimation to elevated carbon dioxide and ultraviolet radiation in the diatom *Thalassiosira pseudonana*: effects on growth, photosynthesis, and spectral sensitivity of photoinhibition. *Limnol. Oceanogr.* **53**, 494–505. (doi:10.4319/lo.2008.53.2.0494)
- 125 Bobik, T. A. 2006 Polyhedral organelles compartmenting bacterial metabolic processes. *Appl. Microbiol. Biotechnol.* **70**, 517–525. (doi:10.1007/s00253-005-0295-0)
- 126 Boyce, D. G., Lewis, M. R. & Worm, B. 2010 Global phytoplankton decline over the past century. *Nature* **466**, 591–596. (doi:10.1038/nature09268)
- 127 Richardson, A. J. 2008 In hot water: zooplankton and climate change. *ICES J. Mar. Sci.* **65**, 279–295. (doi:10.1093/icesjms/fsn028)
- 128 Steinacher, M. *et al.* 2010 Projected 21st century decrease in marine productivity: a multi-model analysis. *Biogeosciences* **7**, 979–1005. (doi:10.5194/bg-7-979-2010)
- 129 Froelicher, T. L., Joos, F., Plattner, G. K., Steinacher, M. & Doney, S. C. 2009 Natural variability and anthropogenic trends in oceanic oxygen in a coupled carbon cycle-climate model ensemble. *Global Biogeochem. Cycles* **23**, GB1003. (doi:10.1029/2008gb003316)
- 130 Keeling, R. F., Körtzinger, A. & Gruber, N. 2010 Ocean deoxygenation in a warming world. *Annu. Rev. Mar. Sci.* **2**, 199–229. (doi:10.1146/annurev.marine.010908.163855)
- 131 Tyrrell, T. 2011 Anthropogenic modification of the oceans. *Phil. Trans. R. Soc. A* **369**, 887–908. (doi:10.1098/rsta.2010.0334)
- 132 Doney, S. C., Fabry, V. J., Feely, R. A. & Kleypas, J. A. 2009 Ocean acidification: the other CO₂ problem. *Annu. Rev. Mar. Sci.* **1**, 169–192. (doi:10.1146/annurev.marine.010908.163834)
- 133 Raven, J. A. 2011 Effects on marine algae of changed seawater chemistry with increasing atmospheric CO₂. *Biol. Environ. Proc. R. Ir. Acad.* **111B**, 1–17. (doi:10.3318/bioe.2011.01)
- 134 Riebesell, U., Koertzing, A. & Oschlies, A. 2009 Sensitivities of marine carbon fluxes to ocean change. *Proc. Natl Acad. Sci. USA* **106**, 20 602–20 609. (doi:10.1073/pnas.0813291106)
- 135 Milligan, A. J., Varela, D. E., Brzezinski, M. A. & Morel, F. M. M. 2004 Dynamics of silicon metabolism and silicon isotopic discrimination in a marine diatom as a function of pCO₂. *Limnol. Oceanogr.* **49**, 322–329. (doi:10.4319/lo.2004.49.2.0322)
- 136 Hofmann, M. & Schellnhuber, J. 2009 Oceanic acidification affects marine carbon pump and triggers extended marine oxygen holes. *Proc. Natl Acad. Sci. USA* **106**, 3017–3022. (doi:10.1073/pnas.0813384106)
- 137 Devol, A. H. & Hartnett, H. E. 2001 Role of the oxygen-deficient zone in transfer of organic carbon to the deep ocean. *Limnol. Oceanogr.* **46**, 1684–1690. (doi:10.4319/lo.2001.46.7.1684)
- 138 Jenkyns, H. C. 2010 Geochemistry of oceanic anoxic events. *Geochem. Geophys. Geosyst.* **11**, 1–30. (doi:10.1029/2009gc002788)
- 139 Raven, J. A. & Geider, R. D. 2003 Adaptation, acclimation and regulation of photosynthesis in algae. In *Photosynthesis in algae* (eds A. W. D. Larkum, S. E. Douglas & J. A. Raven), pp. 385–412. Dordrecht, The Netherlands: Kluwer Academic.
- 140 Bell, G. & Collins, S. 2008 Adaptation, extinction and global change. *Evol. Appl.* **1**, 3–16. (doi:10.1111/j.1752-4571.2007.00011.x)
- 141 Collins, S. 2011 Competition limits adaptation and productivity in a photosynthetic alga at elevated CO₂. *Proc. R. Soc. B* **278**, 247–255. (doi:10.1098/rspb.2010.1173)
- 142 Collins, S. & Bell, G. 2004 Phenotypic consequences of 1,000 generations of selection at elevated CO₂ in a green alga. *Nature* **431**, 566–569. (doi:10.1038/nature02945)
- 143 Collins, S. & Bell, G. 2006 Rewinding the tape: selection of algae adapted to high CO₂ at current and pleistocene levels of CO₂. *Evolution* **60**, 1392–1401. (doi:10.1111/j.0014-3820.2006.tb01218.x)

- 144 Collins, S. & Bell, G. 2006 Evolution of natural algal populations at elevated CO₂. *Ecol. Lett.* **9**, 129–135. (doi:10.1111/j.1461-0248.2005.00854.x)
- 145 Collins, S. & Gardner, A. 2009 Integrating physiological, ecological and evolutionary change: a Price equation approach. *Ecol. Lett.* **12**, 744–757. (doi:10.1111/j.1461-0248.2009.01340.x)
- 146 Collins, S., Suetemeyer, D. & Bell, G. 2006 Changes in C uptake in populations of *Chlamydomonas reinhardtii* selected at high CO₂. *Plant Cell Environ.* **29**, 1812–1819. (doi:10.1111/j.1365-3040.2006.01559.x)
- 147 Huertas, I. E., Rouco, M., López-Rodas, V. & Costas, E. 2011 Warming will affect phytoplankton differently: evidence through a mechanistic approach. *Proc. R. Soc. B* **278**, 3534–3543. (doi:10.1098/rspb.2011.0160)
- 148 Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M., Rowley, S. J., Tedesco, D. & Buia, C. 2008 Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* **454**, 96–99. (doi:10.1038/nature07051)
- 149 Mercado, J. & Gordillo, F. 2011 Inorganic carbon acquisition in algal communities: are the laboratory data relevant to the natural ecosystems? *Photosynth. Res.* **109**, 1–11. (doi:10.1007/s11120-011-9646-0)
- 150 Bots, P., Benning, L. G., Rickaby, R. E. M. & Shaw, S. 2011 The role of SO₄ in the switch from calcite to aragonite seas. *Geology* **39**, 331–334. (doi:10.1130/g31619.1)
- 151 Lee, J. & Morse, J. W. 2010 Influences of alkalinity and pCO₂ on CaCO₃ nucleation from estimated Cretaceous composition seawater representative of ‘calcite seas’. *Geology* **38**, 115–118. (doi:10.1130/g30537.1)
- 152 Ries, J. B. 2009 Effects of secular variation in seawater Mg/Ca ratio (calcite-aragonite seas) on CaCO₃ sediment production by the calcareous algae *Halimeda*, *Penicillus* and *Udotea*—evidence from recent experiments and the geological record. *Tera Nov.* **21**, 323–339. (doi:10.1111/j.1365-3121.2009.00899.x)
- 153 Zachos, J. C. et al. 2005 Rapid acidification of the ocean during the Paleocene–Eocene thermal maximum. *Science* **308**, 1611–1615. (doi:10.1126/science.1109004)
- 154 Zeebe, R. E. & Wolf-Gladrow, D. A. 2001 *CO₂ in Seawater: equilibrium, kinetics, isotopes*. Amsterdam, The Netherlands: Elsevier.
- 155 Hurd, C. L., Hepburn, C. D., Currie, K. I., Raven, J. A. & Hunter, K. A. 2009 Testing the effects of ocean acidification on algal metabolism: considerations for experimental designs. *J. Phycol.* **45**, 1236–1251. (doi:10.1111/j.1529-8817.2009.00768.x)
- 156 Iglesias-Rodriguez, M. D. et al. 2008 Phytoplankton calcification in a high-CO₂ world. *Science* **320**, 336–340. (doi:10.1126/science.1154122)
- 157 Riebesell, U., Fabry, V. J., Hansson, L. & Gattuso, J.-P. (eds) 2010 *Guide to best practices for ocean acidification research and data reporting*. Luxembourg: Publications Office of the European Union.
- 158 Shi, D., Xu, Y. & Morel, F. M. M. 2009 Effects of the pH/pCO₂ control method on medium chemistry and phytoplankton growth. *Biogeosciences* **6**, 1199–1207. (doi:10.5194/bg-6-1199-2009)
- 159 Laws, E. A., Popp, B. N., Cassar, N. & Tanimoto, J. 2002 ¹³C discrimination patterns in oceanic phytoplankton: likely influence of CO₂ concentrating mechanisms, and implications for palaeoreconstructions. *Funct. Plant Biol.* **29**, 323–333. (doi:10.1071/pp01183)
- 160 Raven, J. A., Johnston, A. M. & Turpin, D. H. 1993 Influence of changes in CO₂ concentration and temperature on marine phytoplankton ¹³C/¹²C ratios—an analysis of possible mechanisms. *Glob. Planet Change* **8**, 1–12. (doi:10.1016/0921-8181(93)90058-v)
- 161 Fletcher, B. J., Beerling, D. J., Brentnall, S. J. & Royer, D. L. 2005 Fossil bryophytes as recorders of ancient CO₂ levels: experimental evidence and a Cretaceous case study. *Global Biogeochem. Cycles* **19**, GB3012. (doi:10.1029/2005gb002495)