



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Minireview

Drosophila as a genetic model for studying pathogenic human viruses

Tamara T. Hughes, Amanda L. Allen, Joseph E. Bardin, Megan N. Christian, Kansei Daimon, Kelsey D. Dozier, Caom L. Hansen, Lisa M. Holcomb, Joseph Ahlander*

Department of Natural Sciences, Northeastern State University, Tahlequah, OK 74464, USA

ARTICLE INFO

Article history:

Received 11 September 2011
 Returned to author for revision
 22 September 2011
 Accepted 21 November 2011
 Available online 15 December 2011

Keywords:

Drosophila
 Viruses
 GAL4/UAS system
 Transgenic expression
 Model organism
 Human pathogens

ABSTRACT

Viruses are infectious particles whose viability is dependent on the cells of living organisms, such as bacteria, plants, and animals. It is of great interest to discover how viruses function inside host cells in order to develop therapies to treat virally infected organisms. The fruit fly *Drosophila melanogaster* is an excellent model system for studying the molecular mechanisms of replication, amplification, and cellular consequences of human viruses. In this review, we describe the advantages of using *Drosophila* as a model system to study human viruses, and highlight how *Drosophila* has been used to provide unique insight into the gene function of several pathogenic viruses. We also propose possible directions for future research in this area.

© 2011 Elsevier Inc. All rights reserved.

Introduction

Viral infection is associated with a number of diseases ranging from the common cold to cancer. It is therefore of great interest to understand the molecular basis of viral infection and propagation to minimize the threat of these viruses to human health.

During infection, viruses release their genetic material into the host cell. These foreign genes are expressed, producing viral proteins which hijack the molecular machinery of the host cell through interactions with endogenous cellular proteins. One strategy for minimizing the damaging effects of a virus is to identify and inhibit the molecular mechanisms by which viruses replicate in cells.

The fruit fly *Drosophila melanogaster* is currently being used as a genetic system to model many human diseases, such as Parkinson's disease (Feany and Bender, 2000), heritable cancer syndromes such as multiple endocrine neoplasia (Read et al., 2005), and metabolic disorders like obesity and diabetes (Musselman et al., 2011). *Drosophila* has been used for decades to study the molecular and genetic functions of a range of viruses, as well as giving important insight into the mechanisms of host antiviral immunity (Sabin et al., 2010). Of particular note are a number of human viruses and their gene products that have been studied using *Drosophila* (Table 1). For example, *Drosophila* cells have been used in genome-wide RNA interference screens to rapidly identify cellular factors required for replication of influenza

and dengue viruses (Hao et al., 2008; Sessions et al., 2009). The discovery of host factors involved in viral pathogenesis may lead to the development of novel treatments.

In this article, we discuss how *D. melanogaster* can be used to study viral gene function. We also review some of the published research that has used *Drosophila* to study important human viral pathogens. Finally, we suggest opportunities for future studies using this approach.

D. melanogaster as a model to study gene function

D. melanogaster has already proven to be a powerful tool for understanding the molecular function of viral proteins (Table 1). The conserved genetic pathways between fly and human combined with the availability of numerous genetic resources to study gene function makes *D. melanogaster* a natural model system to study molecular mechanisms related to human biology (Reiter et al., 2001).

Drosophila possesses many characteristics desired in a model organism that allow rapid, meaningful analysis of viral gene function. First, the genetics of the *Drosophila* are relatively simple. *Drosophila* contains fewer genes than humans, indicating less overall genetic redundancy. This allows for a simpler analysis when studying the effects of genes on biological processes (Dimova and Dyson, 2005; Zhang et al., 2007). Second, these model organisms can be genetically modified and propagated quickly. The developmental time of *Drosophila* ranges from about one to three weeks and is dependent on temperature and other environmental conditions. The entire lifespan is approximately one month in length. In addition, female flies can produce hundreds of offspring within a couple of weeks, and those

* Corresponding author at: Northeastern State University, Department of Natural Sciences, 600 North Grand Avenue, Tahlequah, OK 74464, USA. Fax: +1 918 458 2325.
 E-mail address: ahlander@nsuok.edu (J. Ahlander).

Table 1
Human viruses studied using *Drosophila melanogaster*.

Virus	Involvement in human disease	Experimental systems used	References
Dengue virus (DENV)	Dengue fever, hemorrhagic fever	<i>Drosophila</i> cell culture (infected)	Mukherjee and Hanley (2010); Sessions et al. (2009)
Epstein–Barr virus (EBV)	Infectious mononucleosis, various cancers, autoimmune disease	Live <i>Drosophila</i> (transgenic)	Adamson et al. (2005)
Hepatitis B virus (HBV) Human immunodeficiency virus 1 (HIV-1)	Hepatitis B, liver cancer Acquired immune deficiency syndrome (AIDS)	<i>Drosophila</i> cell culture (transfected) <i>Drosophila</i> cell culture (transfected), live <i>Drosophila</i> (transgenic)	Wang et al. (1998) Battaglia et al. (2001); Brighty and Rosenberg (1994); Chan et al. (2002); Chaudhuri et al. (2007); Ivey-Hoyle and Rosenberg (1990); Lee et al. (2005); Leulier et al. (2003); Ponti et al. (2008) Steinberg et al. (2008)
Human cytomegalovirus (HCMV)	Birth defects, mononucleosis, severe complications in immunocompromised individuals	Live <i>Drosophila</i> (transgenic)	
Influenza A virus	Flu pandemics, pneumonia, respiratory failure	<i>Drosophila</i> cell culture (infected), live <i>Drosophila</i> (transgenic)	Adamson et al. (2011); Chou et al. (2007); Hao et al. (2008); Lam et al. (2010); Li et al. (2004)
SARS coronavirus (SARS-CoV)	Severe acute respiratory syndrome (SARS)	Live <i>Drosophila</i> (transgenic)	Chan et al. (2007, 2009); Wong et al. (2005)
Simian vacuolating virus 40 (SV40) Sindbis virus (SINV)	Possibly oncogenic Sindbis fever, Pogosta disease	Live <i>Drosophila</i> (transgenic) <i>Drosophila</i> cell culture (infected), live <i>Drosophila</i> (infected)	Kotadia et al. (2008) Avadhanula et al. (2009); Brackney et al. (2010); Galiana-Arnoux et al. (2006); Mudiganti et al. (2006, 2010); Rose et al. (2011); Sabin et al. (2009); Saleh et al. (2009)
Vaccinia virus (VACV)	Fever, rash, used as vaccine to protect against smallpox	<i>Drosophila</i> cell culture (infected), live <i>Drosophila</i> (infected, transgenic)	Chou et al. (2007); Li et al. (2004); Moser et al. (2010); Sabin et al. (2009)
Vesicular stomatitis virus (VSV)	Flu-like symptoms in humans, usually infects livestock	<i>Drosophila</i> cell culture (infected), live <i>Drosophila</i> (infected)	Cherry (2009); Mueller et al. (2010); Sabin et al. (2009); Shelly et al. (2009)
West Nile virus (WNV)	West Nile fever, encephalitis	<i>Drosophila</i> cell culture (infected), live <i>Drosophila</i> (infected)	Brackney et al. (2010); Chotkowski et al. (2008); Glaser and Meola (2010)

offspring become sexually mature within the first day of adulthood, enabling the life cycle to start over once again. They are also convenient to grow in the lab due to their small size and simple diet. Hence, large numbers of flies can be maintained inexpensively in the laboratory. Third, the action of viral genes can be studied in the context of whole *Drosophila* tissues *in vivo*, which more closely models the cellular environment of viral infection. This is particularly valuable because the effects of the virus at different stages of development can be explored.

Despite its many advantages, there are some limitations of using *Drosophila* as a system to model human biology. First, there are physiological differences between *Drosophila* and humans. For example, the optimal temperature for culturing fruit flies is between 18 °C and 27 °C. On the other hand, the average body temperature of a human is approximately 37 °C. Human viruses may be adapted for optimal function at 37 °C, so some viral proteins may be unable to function properly at temperatures lower than normal body human temperature. Second, genetic differences between *Drosophila* and humans may pose a challenge to using flies for studying some human viruses. Since genomic conservation is not comprehensive between humans and fruit flies some genes found in humans are absent in the *Drosophila* genome, which could make it difficult to study some viruses that may require host factors that are not found in *Drosophila* cells. Finally, there are also biochemical differences between *Drosophila* and humans. For example, the influenza virus binds to a sialic acid residue on the surface of human cells during infection. One study required that the influenza virus be modified using a different viral coat protein to aid it in infecting *Drosophila* cells, since these insect cells lacked the sialic acid necessary for viral entry (Hao et al., 2008). However, once inside the cell these viral genes are expressed and appear to function similarly to when they are inside of human cells.

Undoubtedly, the differences between human and insect cells do not necessarily need to become a permanent obstacle to the use of *Drosophila* as a model system, as many limitations have been and can be overcome through modification of either the virus or the host cells (Chaudhuri et al., 2007; Hao et al., 2008). Furthermore, an alternative to altering viral coat proteins to allow infection would be

to introduce viral genes into *Drosophila* cells through transfection or transgenesis. However, rather than remaining a standalone system to study viral mechanisms, *Drosophila* may be most beneficial as a tool to rapidly screen the *in vivo* function of viral genes followed by complementary studies with mammalian cells.

The GAL4/UAS system for *in vivo* expression of viral transgenes

A strategy often used to express viral genes in *Drosophila* is the binary GAL4/UAS gene expression system. In this system a gene of interest is constructed so that its expression is under the control of the upstream activating sequence (UAS), which is activated by binding of the GAL4 transcription factor (Fig. 1). *Drosophila* expression vectors are available to insert any gene of interest for the generation of transgenic flies and can efficiently accommodate genes greater than 5 kb in size. In addition, there are publicly available fly stocks for hundreds of different inducible or tissue-specific GAL4 transgenes, which permit precise control over transgene expression. Adult flies carrying a UAS-linked transgene are mated to flies carrying a GAL4 driver, producing progeny containing both elements of the system. The GAL4 gene can then induce expression of the gene of interest in a predictable pattern in the organism. Transcription of the target gene requires the presence of GAL4, so in its absence the gene of interest remains silent in cells that do not express GAL4. One advantage of this system is the ability to study toxic or lethal gene products by restricting transgene expression to cells in non-essential tissues like the eye or wing (Duffy, 2002).

Mutant phenotypes generated by transgenic expression of gene products such as viral proteins can be used to study the molecular and genetic mechanisms that underpin the function of those genes. Such gain-of-function phenotypes are particularly amenable to genetic screens to uncover the cellular host factors involved in the regulation of viral pathogenesis. In addition, a library of drug compounds may be fed to developing flies to discover inhibitors of the mutant phenotypes caused by the viral proteins being studied. Such pharmacological screens have the potential to discover new candidate drugs for the treatment of viral infections.




Fig. 1. The GAL4/UAS system can be used to express foreign viral proteins *in vivo* in *Drosophila melanogaster*. A gene of interest is placed next to the upstream activating sequence (UAS), which allows activation of gene expression by the GAL4 transcription factor. Flies with a UAS-responsive transgene are crossed with other flies with a specific GAL4 driver gene. In this illustrated example, the offspring has a GAL4 gene under control of the actin promoter, which expresses GAL4 ubiquitously throughout development. GAL4 then binds to UAS and turns on the viral gene in all cells that express GAL4. If expression had to be targeted to a specific subset of cells, then a different GAL4 line would be used to restrict viral gene expression to specific tissues. For example, GAL4 expression could be targeted to the eye where the viral gene would only be expressed in those cells but not in the other tissues of the organism.

Several strategies can be employed when designing viral transgenes to ensure robust gene expression in *Drosophila*. First, it is important that these genes are expressed at sufficient levels in *Drosophila* cells, since the level of transgene expression can have a significant effect on phenotypes. A single transgene is sometimes not sufficient to produce a phenotype or produces a weak phenotype with low penetrance. However, GAL4 activity is temperature sensitive, so raising the rearing temperature of the flies a few degrees can enhance expression (Duffy, 2002). Such a shift in temperature may also have a positive effect on the activity of human viral proteins, since these typically function under the higher physiological temperature of the human body. Furthermore, stronger expression through an increase of the transgene copy number can also help to generate a phenotype (Asano and Wharton, 1999; Hong et al., 2008). However, improved expression vectors for making transgenic flies have been generated that increase transgene expression several fold over previous constructs, which in some cases will eliminate the need to combine multiple copies of transgenes to boost expression (Pfeiffer et al., 2010).

Second, codon optimization of viral gene sequences should also be considered when constructing transgenes for expression in heterologous hosts (Welch et al., 2011). The *D. melanogaster* genome shows a preference for particular codons compared to other organisms, a phenomenon called codon bias, and it is presumed that genes that encode proteins using rare codons will be translated at a slower rate. In other words, a gene that expresses well in one host species may express poorly in a different species, so using codon optimization for the target species may improve translation efficiency. Indeed, codon optimization of enhanced green fluorescent protein (EGFP) for expression in *Drosophila* resulted in a 50% increase in EGFP production compared to the standard coding sequence (Pfeiffer et al., 2010). Codon optimized gene sequences can be generated in-house through site-directed mutagenesis, or alternatively they can be commercially synthesized *de novo*.

Third, like the Kozak consensus sequence used for efficient translation initiation in mammalian genes, highly expressed *Drosophila* genes also have the Kozak-like sequence CAAAAUG (Cavener, 1987). Inclusion of this sequence in viral gene constructs may enhance translation initiation and increase protein expression levels. Together these genetic engineering strategies may enhance expression of viral transgenes in *Drosophila* to greatly advance functional study of these genes.

Investigators should be aware that special approval may be required by governmental agencies before making transgenic organisms to study gene sequences from some viruses, particularly those viruses that are regarded as potential bioterrorism threats. Examples of these would likely include some viruses that cause hemorrhagic fever or encephalitis. The genes of many viruses, however, may be freely permitted for use in making transgenic insects. For example, in the United States, the National Institutes of Health allows the generation of transgenic invertebrates with DNA derived from most eukaryotic viruses, as long as it contains less than two-thirds of the viral genome and cannot lead to the production of infectious viruses (NIH, 2011). These issues should be carefully considered during the design of experiments to study pathogenic viruses using insect systems.

Studies of human viruses using *D. melanogaster*

Numerous studies have shown that *D. melanogaster* is a valuable system for studying human viruses (Table 1). Here we review a few of these studies that highlight the efficacy of this approach. Specifically, we summarize important findings that helped to advance understanding of the SARS and HIV viruses.

SARS

Severe Acute Respiratory Syndrome corona virus (SARS-CoV) was the cause of a worldwide pneumonia outbreak in 2003 (Rota et al., 2003). SARS is an enveloped, single-stranded RNA virus that infects tissues of the intestines and lungs via air-borne transmission (Chen et al., 2011). An effective drug to treat SARS is still being pursued, since most pharmaceutical treatment of SARS patients so far have proven ineffective (Stockman et al., 2006). Research using *D. melanogaster* has elucidated how SARS-CoV proteins function within the cellular environment.

In vivo expression of the SARS-CoV 3a protein using transgenic *Drosophila* caused an increase in apoptosis in the developing eye (Wong et al., 2005). Genetic interaction studies with these flies further showed that apoptosis caused by 3a expression occurred through the mitochondrial pathway via cytochrome c, and this result was later validated using human cells (Padhan et al., 2008). Through the use of genetic modifier screens, the function of 3a was also linked to other cellular processes, including calcium regulation, ubiquitination, and transcription (Wong et al., 2005). A subsequent report studied structure–function relationships of the 3a protein using a combination of experiments with human cell culture and transgenic *Drosophila* (Chan et al., 2009). Importantly, pharmaceutical blockage of the 3a ion channel activity prevents its ability to induce apoptosis both *in vitro* (human cells) and *in vivo* (transgenic *Drosophila*). Another study using transgenic flies showed that the SARS-CoV membrane (M) protein induces apoptosis in the eye by suppressing survival signaling pathways (Chan et al., 2007). Thus, research in *Drosophila* has identified novel cellular targets that may be useful for future research to discover drugs that control the activity of these SARS-CoV proteins, leading to treatments that could alleviate symptoms and limit the spread of this disease.

HIV

More than 30 million individuals are infected with the human immunodeficiency virus (HIV) worldwide, resulting in about 2 million

deaths annually (Kilmarx, 2009). HIV-1 is an enveloped retrovirus that uses its own reverse transcriptase to replicate its genomic single-stranded RNA through a DNA intermediate. During its life cycle, this viral DNA can become permanently integrated into the host cell DNA where its genes are expressed (Cherepanov et al., 2011). The virus is generally spread through sexual contact or contact with blood products. Although antiviral drugs can suppress the infection for many years, there is currently no cure for HIV. In an effort to better understand this virus, *D. melanogaster* has been used to study the function of genes from HIV-1. Described below are examples of three different HIV-1 genes whose functions were further clarified using *Drosophila*. These studies highlight the strength and versatility of this genetic model system.

HIV-1 Nef is a membrane-associated protein involved in the downregulation of the cell surface receptor CD4 through endocytosis (Garcia and Miller, 1991). Human CD4 and HIV-1 Nef proteins were co-expressed in cultured *Drosophila* S2 cells, where Nef was shown to downregulate CD4 (Chaudhuri et al., 2007). Using RNA interference to target cellular factors involved in protein trafficking, it was revealed that Nef-dependent CD4 downregulation required a specific interaction with AP2, a complex involved in clathrin-mediated endocytosis, but not other AP complexes. This discovery was followed up using HeLa cells where it was shown that the Nef-AP2 interaction is functionally conserved in humans. Another study used transgenic *Drosophila* to show that Nef expression in larval wing discs also caused apoptosis through activation of the conserved JNK signaling pathway (Lee et al., 2005). In addition, Nef expression negatively affected the *Drosophila* immune system by inhibiting NF- κ B signaling in fat body cells. These findings may help to explain how Nef expression during HIV infection contributes to the decline of T-cell immune function that is characteristic of AIDS progression.

Tat is an HIV-1 protein required for viral gene expression and is essential for viral replication. Tat was expressed in transgenic *Drosophila*, where it disrupted microtubule polymerization and kinetochore dynamics via a direct interaction with tubulin (Battaglia et al., 2001). Ensuing research in human cells validated the importance of this finding that helped to advance the understanding of the mechanisms of HIV pathogenesis (Butler et al., 2011; Chen et al., 2002). Tat was previously shown to localize to nucleoli in human cells, but the function of Tat in the nucleolus was unclear. Another study demonstrated that expression of Tat protein in the *Drosophila* ovary showed nucleolar localization (Ponti et al., 2008). In these transgenic females, Tat was shown to affect the maturation of ribosomes through the inhibition of rRNA processing, which resulted in a reduced number of ribosomes in the cytoplasm. Many viruses regulate protein production to facilitate viral replication and to modulate the apoptotic response of the host cell, so this research suggests a mechanism by which Tat may play a role in HIV-1 pathogenesis.

HIV-1 Rev is a protein that has been shown to regulate expression of HIV proteins, for example by facilitating export and translation of viral *env* mRNA. Rev was studied in cultures of *Drosophila* S2 cells through the use of a *Rev* gene co-transfected with a plasmid containing a copy of the viral *env* gene (Ivey-Hoyle and Rosenberg, 1990). It was found that Rev acts in *Drosophila* cells as it does in mammalian cells by promoting the transport of *env* mRNA from the nucleus to the cytoplasm. This suggests that the Rev protein functions by interacting with host cellular factors that are conserved between humans and insects (Brighty and Rosenberg, 1994). Future research will benefit from using *Drosophila* to study HIV protein function due to the high conservation between insect and human cellular pathways.

Future directions

D. melanogaster has proven to be an excellent system for studying the pathogenic mechanisms of human viruses. However, we believe that this tool remains underutilized and holds great potential for

the study of other human viruses. Viruses that would make good candidates for future study in *Drosophila* would include those that have a large impact on human populations. A small viral genome would allow for a simpler selection of candidate genes for further study. In addition, viruses that have known strains of different pathogenic characteristics (HIV, HPV, etc.) may also be good candidates for study. For example, a comparison of the functional differences between genes of the different strains could help to uncover what makes one strain more pathogenic than another. Based on these criteria, we have identified three candidates – the human papillomavirus, the hepatitis C virus, and the yellow fever virus – which could potentially benefit from studies using *Drosophila* as a model. We anticipate that in the future *D. melanogaster* will prove to be a productive system for uncovering the molecular mechanisms of these and other pathogenic human viruses.

Acknowledgments

This publication was made possible by Grant Number DUE-0856396 from the National Science Foundation and by Grant Number P2ORR016478 from the National Center for Research Resources (NCRR), a component of the National Institutes of Health (NIH). We are also grateful to Dr. Brian Ceresa for helpful comments during the preparation of the manuscript.

References

- Adamson, A.L., Wright, N., Lajeunesse, D.R., 2005. Modeling early Epstein–Barr virus infection in *Drosophila melanogaster*: the BZLF1 protein. *Genetics* 171, 1125–1135.
- Adamson, A.L., Chohan, K., Swenson, J., Lajeunesse, D.R., 2011. A *Drosophila* model for genetic analysis of influenza viral/host interactions. *Genetics* 189, 495–506.
- Asano, M., Wharton, R.P., 1999. E2F mediates developmental and cell cycle regulation of ORC1 in *Drosophila*. *EMBO J.* 18, 2435–2448.
- Avadhanula, V., Weasner, B.P., Hardy, G.G., Kumar, J.P., Hardy, R.W., 2009. A novel system for the launch of alphavirus RNA synthesis reveals a role for the Imd pathway in arthropod antiviral response. *PLoS Pathog.* 5, e1000582.
- Battaglia, P.A., Zito, S., Macchini, A., Gigliani, F., 2001. A *Drosophila* model of HIV-Tat-related pathogenicity. *J. Cell Sci.* 114, 2787–2794.
- Brackney, D.E., Scott, J.C., Sagawa, F., Woodward, J.E., Miller, N.A., Schilkey, F.D., Mudge, J., Wilusz, J., Olson, K.E., Blair, C.D., Ebel, G.D., 2010. C6/36 *Aedes albopictus* cells have a dysfunctional antiviral RNA interference response. *PLoS Negl. Trop. Dis.* 4, e856.
- Brighty, D.W., Rosenberg, M., 1994. A cis-acting repressive sequence that overlaps the Rev-responsive element of human immunodeficiency virus type 1 regulates nuclear retention of *env* mRNAs independently of known splice signals. *Proc. Natl. Acad. Sci. U. S. A.* 91, 8314–8318.
- Butler, T.R., Smith, K.J., Self, R.L., Braden, B.B., Prendergast, M.A., 2011. Neurodegenerative effects of recombinant HIV-1 Tat(1–86) are associated with inhibition of microtubule formation and oxidative stress-related reductions in microtubule-associated protein-2(a, b). *Neurochem. Res.* 36, 819–828.
- Cavener, D.R., 1987. Comparison of the consensus sequence flanking translational start sites in *Drosophila* and vertebrates. *Nucleic Acids Res.* 15, 1353–1361.
- Chan, H.Y., Tung, C.K., O’Kane, C.J., 2002. Rev-GFP transgenic lines for studies of nucleocytoplasmic transport in *Drosophila*. *Genesis* 34, 139–141.
- Chan, C.M., Ma, C.W., Chan, W.Y., Chan, H.Y., 2007. The SARS-coronavirus membrane protein induces apoptosis through modulating the Akt survival pathway. *Arch. Biochem. Biophys.* 459, 197–207.
- Chan, C.M., Tsoi, H., Chan, W.M., Zhai, S., Wong, C.O., Yao, X., Chan, W.Y., Tsui, S.K., Chan, H.Y., 2009. The ion channel activity of the SARS-coronavirus 3a protein is linked to its pro-apoptotic function. *Int. J. Biochem. Cell Biol.* 41, 2232–2239.
- Chaudhuri, R., Lindwasser, O.W., Smith, W.J., Hurley, J.H., Bonifacino, J.S., 2007. Downregulation of CD4 by human immunodeficiency virus type 1 Nef is dependent on clathrin and involves direct interaction of Nef with the AP2 clathrin adaptor. *J. Virol.* 81, 3877–3890.
- Chen, D., Wang, M., Zhou, S., Zhou, Q., 2002. HIV-1 Tat targets microtubules to induce apoptosis, a process promoted by the pro-apoptotic Bcl-2 relative Bim. *EMBO J.* 21, 6801–6810.
- Chen, C., Zhao, B., Yang, X., Li, Y., 2011. Role of two-way airflow owing to temperature difference in severe acute respiratory syndrome transmission: revisiting the largest nosocomial severe acute respiratory syndrome outbreak in Hong Kong. *J. R. Soc. Interface* 8, 699–710.
- Cherepanov, P., Maertens, G.N., Hare, S., 2011. Structural insights into the retroviral DNA integration apparatus. *Curr. Opin. Struct. Biol.* 21, 249–256.
- Cherry, S., 2009. VSV infection is sensed by *Drosophila*, attenuates nutrient signaling, and thereby activates antiviral autophagy. *Autophagy* 5, 1062–1063.
- Chotkowski, H.L., Ciota, A.T., Jia, Y., Puig-Basagoiti, F., Kramer, L.D., Shi, P.Y., Glaser, R.L., 2008. West Nile virus infection of *Drosophila melanogaster* induces a protective RNAi response. *Virology* 377, 197–206.

- Chou, Y.T., Tam, B., Linay, F., Lai, E.C., 2007. Transgenic inhibitors of RNA interference in *Drosophila*. *Fly (Austin)* 1, 311–316.
- Dimova, D.K., Dyson, N.J., 2005. The E2F transcriptional network: old acquaintances with new faces. *Oncogene* 24, 2810–2826.
- Duffy, J.B., 2002. GAL4 system in *Drosophila*: a fly geneticist's Swiss army knife. *Genesis* 34, 1–15.
- Feany, M.B., Bender, W.W., 2000. A *Drosophila* model of Parkinson's disease. *Nature* 404, 394–398.
- Galiana-Arnoux, D., Dostert, C., Schneemann, A., Hoffmann, J.A., Imler, J.L., 2006. Essential function *in vivo* for Dicer-2 in host defense against RNA viruses in *Drosophila*. *Nat. Immunol.* 7, 590–597.
- Garcia, J.V., Miller, A.D., 1991. Serine phosphorylation-independent downregulation of cell-surface CD4 by nef. *Nature* 350, 508–511.
- Glaser, R.L., Meola, M.A., 2010. The native Wolbachia endosymbionts of *Drosophila melanogaster* and *Culex quinquefasciatus* increase host resistance to West Nile virus infection. *PLoS One* 5, e11977.
- Hao, L., Sakurai, A., Watanabe, T., Sorensen, E., Nidom, C.A., Newton, M.A., Ahlquist, P., Kawaoka, Y., 2008. *Drosophila* RNAi screen identifies host genes important for influenza virus replication. *Nature* 454, 890–893.
- Hong, S.W., Qi, W., Brabant, M., Bosco, G., Martinez, J.D., 2008. Human 14-3-3 gamma protein results in abnormal cell proliferation in the developing eye of *Drosophila melanogaster*. *Cell Div* 3, 2.
- Ivey-Hoyle, M., Rosenberg, M., 1990. Rev-dependent expression of human immunodeficiency virus type 1 gp160 in *Drosophila melanogaster* cells. *Mol. Cell. Biol.* 10, 6152–6159.
- Kilmarx, P.H., 2009. Global epidemiology of HIV. *Curr. Opin. HIV AIDS* 4, 240–246.
- Kotadia, S., Kao, L.R., Comerford, S.A., Jones, R.T., Hammer, R.E., Megraw, T.L., 2008. PP2A-dependent disruption of centrosome replication and cytoskeleton organization in *Drosophila* by SV40 small tumor antigen. *Oncogene* 27, 6334–6346.
- Lam, V.K., Tokusumi, T., Cerabona, D., Schulz, R.A., 2010. Specific cell ablation in *Drosophila* using the toxic viral protein M2(H37A). *Fly (Austin)* 4, 338–343.
- Lee, S.B., Park, J., Jung, J.U., Chung, J., 2005. Nef induces apoptosis by activating JNK signaling pathway and inhibits NF-kappaB-dependent immune responses in *Drosophila*. *J. Cell Sci.* 118, 1851–1859.
- Leulier, F., Marchal, C., Miletich, I., Limbourg-Bouchon, B., Benarous, R., Lemaitre, B., 2003. Directed expression of the HIV-1 accessory protein Vpu in *Drosophila* fat-body cells inhibits Toll-dependent immune responses. *EMBO Rep.* 4, 976–981.
- Li, W.X., Li, H., Lu, R., Li, F., Dus, M., Atkinson, P., Brydon, E.W., Johnson, K.L., Garcia-Sastre, A., Ball, L.A., Palese, P., Ding, S.W., 2004. Interferon antagonist proteins of influenza and vaccinia viruses are suppressors of RNA silencing. *Proc. Natl. Acad. Sci. U. S. A.* 101, 1350–1355.
- Moser, T.S., Jones, R.G., Thompson, C.B., Coyne, C.B., Cherry, S., 2010. A kinome RNAi screen identified AMPK as promoting poxvirus entry through the control of actin dynamics. *PLoS Pathog.* 6, e1000954.
- Mudiganti, U., Hernandez, R., Ferreira, D., Brown, D.T., 2006. Sindbis virus infection of two model insect cell systems—a comparative study. *Virus Res.* 122, 28–34.
- Mudiganti, U., Hernandez, R., Brown, D.T., 2010. Insect response to alphavirus infection—establishment of alphavirus persistence in insect cells involves inhibition of viral polyprotein cleavage. *Virus Res.* 150, 73–84.
- Mueller, S., Gausson, V., Vodovar, N., Deddouche, S., Troxler, L., Perot, J., Pfeffer, S., Hoffmann, J.A., Saleh, M.C., Imler, J.L., 2010. RNAi-mediated immunity provides strong protection against the negative-strand RNA vesicular stomatitis virus in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.* 107, 19390–19395.
- Mukherjee, S., Hanley, K.A., 2010. RNA interference modulates replication of dengue virus in *Drosophila melanogaster* cells. *BMC Microbiol.* 10, 127.
- Musselman, L.P., Fink, J.L., Narzinski, K., Ramachandran, P.V., Hathiramani, S.S., Cagan, R.L., Baranski, T.J., 2011. A high-sugar diet produces obesity and insulin resistance in wild-type *Drosophila*. *Dis. Model. Mech.* 4, 842–849.
- NIH, 2011. NIH Guidelines for Research Involving Recombinant DNA Molecules, May ed, pp. III-D-4-a.
- Padhan, K., Minakshi, R., Towheed, M.A., Jameel, S., 2008. Severe acute respiratory syndrome coronavirus 3a protein activates the mitochondrial death pathway through p38 MAP kinase activation. *J. Gen. Virol.* 89, 1960–1969.
- Pfeiffer, B.D., Ngo, T.T., Hibbard, K.L., Murphy, C., Jenett, A., Truman, J.W., Rubin, G.M., 2010. Refinement of tools for targeted gene expression in *Drosophila*. *Genetics* 186, 735–755.
- Ponti, D., Troiano, M., Bellenchi, G.C., Battaglia, P.A., Gliani, F., 2008. The HIV Tat protein affects processing of ribosomal RNA precursor. *BMC Cell Biol.* 9, 32.
- Read, R.D., Goodfellow, P.J., Mardis, E.R., Novak, N., Armstrong, J.R., Cagan, R.L., 2005. A *Drosophila* model of multiple endocrine neoplasia type 2. *Genetics* 171, 1057–1081.
- Reiter, L.T., Potocki, L., Chien, S., Gribskov, M., Bier, E., 2001. A systematic analysis of human disease-associated gene sequences in *Drosophila melanogaster*. *Genome Res.* 11, 1114–1125.
- Rose, P.P., Hanna, S.L., Spiridigliozzi, A., Wannissorn, N., Beiting, D.P., Ross, S.R., Hardy, R.W., Bambina, S.A., Heise, M.T., Cherry, S., 2011. Natural resistance-associated macrophage protein is a cellular receptor for Sindbis virus in both insect and mammalian hosts. *Cell Host Microbe* 10, 97–104.
- Rota, P.A., Oberste, M.S., Monroe, S.S., Nix, W.A., Campagnoli, R., Icenogle, J.P., Penaranda, S., Bankamp, B., Maher, K., Chen, M.H., Tong, S., Tamin, A., Lowe, L., Frace, M., DeRisi, J.L., Chen, Q., Wang, D., Erdman, D.D., Peret, T.C., Burns, C., Ksiazek, T.G., Rollin, P.E., Sanchez, A., Liffick, S., Holloway, B., Limor, J., McCaustland, K., Olsen-Rasmussen, M., Fouchier, R., Gunther, S., Osterhaus, A.D., Drosten, C., Pallansch, M.A., Anderson, L.J., Bellini, W.J., 2003. Characterization of a novel coronavirus associated with severe acute respiratory syndrome. *Science* 300, 1394–1399.
- Sabin, L.R., Zhou, R., Gruber, J.J., Lukinova, N., Bambina, S., Berman, A., Lau, C.K., Thompson, C.B., Cherry, S., 2009. Ars2 regulates both miRNA- and siRNA-dependent silencing and suppresses RNA virus infection in *Drosophila*. *Cell* 138, 340–351.
- Sabin, L.R., Hanna, S.L., Cherry, S., 2010. Innate antiviral immunity in *Drosophila*. *Curr. Opin. Immunol.* 22, 4–9.
- Saleh, M.C., Tassetto, M., van Rij, R.P., Goic, B., Gausson, V., Berry, B., Jacquier, C., Antoniewski, C., Andino, R., 2009. Antiviral immunity in *Drosophila* requires systemic RNA interference spread. *Nature* 458, 346–350.
- Sessions, O.M., Barrows, N.J., Souza-Neto, J.A., Robinson, T.J., Hershey, C.L., Rodgers, M.A., Ramirez, J.L., Dimopoulos, G., Yang, P.L., Pearson, J.L., Garcia-Blanco, M.A., 2009. Discovery of insect and human dengue virus host factors. *Nature* 458, 1047–1050.
- Shelly, S., Lukinova, N., Bambina, S., Berman, A., Cherry, S., 2009. Autophagy is an essential component of *Drosophila* immunity against vesicular stomatitis virus. *Immunity* 30, 588–598.
- Steinberg, R., Shemer-Avni, Y., Adler, N., Neuman-Silberberg, S., 2008. Human cytomegalovirus immediate-early-gene expression disrupts embryogenesis in transgenic *Drosophila*. *Transgenic Res.* 17, 105–119.
- Stockman, L.J., Bellamy, R., Garner, P., 2006. SARS: systematic review of treatment effects. *PLoS Med.* 3, e343.
- Wang, H.D., Trivedi, A., Johnson, D.L., 1998. Regulation of RNA polymerase I-dependent promoters by the hepatitis B virus X protein via activated Ras and TATA-binding protein. *Mol. Cell. Biol.* 18, 7086–7094.
- Welch, M., Villalobos, A., Gustafsson, C., Minshull, J., 2011. Designing genes for successful protein expression. *Methods Enzymol.* 498, 43–66.
- Wong, S.L., Chen, Y., Chan, C.M., Chan, C.S., Chan, P.K., Chui, Y.L., Fung, K.P., Waye, M.M., Tsui, S.K., Chan, H.Y., 2005. *In vivo* functional characterization of the SARS-coronavirus 3a protein in *Drosophila*. *Biochem. Biophys. Res. Commun.* 337, 720–729.
- Zhang, J., Schulze, K.L., Hiesinger, P.R., Suyama, K., Wang, S., Fish, M., Acar, M., Hoskins, R.A., Bellen, H.J., Scott, M.P., 2007. Thirty-one flavors of *Drosophila* rab proteins. *Genetics* 176, 1307–1322.