

A Manganese–Porphyrin Complex Decomposes H₂O₂, Inhibits Apoptosis, and Acts as a Radiation Mitigator in Vivo


Detcho A. Stoyanovsky,^{*,†} Zhentai Huang,[†] Jianfei Jiang,[†] Natalia A. Belikova,[†] Vladimir Tyurin,[†] Michael W. Epperly,[§] Joel S. Greenberger,[§] Hülya Bayir,[‡] and Valerian E. Kagan^{*,†}

Departments of [†]Environmental and Occupational Health, [‡]Critical Care Medicine, and [§]Radiation Oncology, University of Pittsburgh, Pittsburgh, Pennsylvania 15260

S Supporting Information

ABSTRACT: Ionizing radiation triggers mitochondrial overproduction of H₂O₂ with concomitant induction of intrinsic apoptosis, whereby clearance of H₂O₂ upon overexpression of mitochondrial catalase increases radioresistance in vitro and in vivo. As an alternative to gene therapy, we tested the potential of Mn^(III)–porphyrin complexes to clear mitochondrial H₂O₂. We report that triphenyl-[(2*E*)-2-[4-[(1*Z*,4*Z*,9*Z*,15*Z*)-10,15,20-tris(4-aminophenyl)-21,23-dihydroporphyrin-5-yl]phenyl]iminoethyl]phosphonium-Mn^(III) compartmentalizes preferentially into mitochondria of mouse embryonic cells, reacts with H₂O₂, impedes γ -ray-induced mitochondrial apoptosis, and increases the survival of mice exposed to whole body irradiation with γ -rays.

KEYWORDS: Manganese–porphyrin complex, H₂O₂, apoptosis, radiation mitigator




By radiolyzing water and damaging biomolecules via generation of O-, C-, S-, and N-centered radicals, ionizing radiation (IR) triggers cell death via three general mechanisms: mitochondria-mediated apoptosis, mitosis-linked death or loss of clonogenicity, and necrosis. The completion of these cell death pathways includes a cascade of reactions whose development may take from hours to days,¹ thus affording time for radiomitigation. In contrast, the initial radiolytic events happen instantaneously; therefore, protection of cells can be attained by radioprotectors (IRP) present at the time of radiation exposure. In search for IRP against IR, over 4000 scavengers of free radicals have been synthesized and tested in mice.² Currently, however, only 2-butylaminoethylsulfanylphosphonic acid (amifostine; WR-2721) has found clinical applications as an IRP.³

Apoptosis causes massive cell loss in radiosensitive tissues.⁴ Recent studies suggest that inhibition of apoptosis and induction of regeneration-promoting cytokines are viable strategies in the discovery of mechanism-based radiomitigators (IRM).^{4,5}

It has been shown that mitochondria exposed to IR overproduce H₂O₂.⁶ Mitochondrial production of H₂O₂ can trigger inner-sphere oxidation of cardiolipin (CL) in its complex with cytochrome c ([CL-cyt c]; Scheme 1), which is an early apoptotic event.⁷ Notably, clearance of H₂O₂ upon overexpression of mitochondrial catalase increases radioresistance in vitro and in vivo.⁸

As an alternative to gene therapy, we tested the potential of the porphyrin complexes **1a** and **1b** to act as IRM via clearance of mitochondrial H₂O₂ in cells exposed to IR (Scheme 2). The rationale for selection of **1a,b** was based both on the ability of

Scheme 1



amphiphilic organic cations to preferentially compartmentalize into mitochondria⁹ and on the potential of Mn^(III)–porphyrin complexes to react with H₂O₂.^{10,11}

We speculated that reduction of the porphyrin–Mn^(V)=O complex **2** by ascorbic acid (Asc) and/or glutathione (GSH) would close a redox cycle that clears mitochondrial H₂O₂.

Figure 1A depicts the reaction of **1a** with H₂O₂ at pH 8.0, a value characteristic of intramitochondrial environment.¹² Consecutive additions of H₂O₂ to solution of **1a** resulted in a linear response of the H₂O₂-sensitive electrode, whose background readings were restored via decomposition of H₂O₂ with catalase (trace 1). The rates of the reactions of **1a** and **1b** with H₂O₂ markedly decreased at pH 7.0, which suggested the involvement of HOO⁻, rather than H₂O₂ (Figure 1B,C); unexpectedly, at pH 7.0–7.4, H₂O₂ did not react with **1a** to any significant extent

Received: June 13, 2011

Accepted: September 13, 2011

Published: September 13, 2011

Scheme 2





Figure 1. Complexes **1a** and **1b** react with H₂O₂. Experiments were carried out at 25 °C in 0.1 M phosphate buffer. Catalase, 400 units/mL (A); H₂O₂, 3.3 (A) and 200 μM (D); **1a**: 10, 20, 40, and 60 (A, traces 2–5) and 10 μM (B); **1b**: 10 (C) and 30 μM (D). The results in B–D represent the mean ± SE (*n* = 3).

(Figure 1B). Under these conditions, neither **1a** nor **1b** reacted with superoxide anion radical (O₂^{•-}); Supporting Information, Figure 3SI). Both **1a** and **1b** (5–20 μM) were relatively stable in the presence of equimolar concentrations of H₂O₂. At concentrations exceeding 0.2 mM, H₂O₂ “bleached” the chromophores (*t*_{1/2} ~ 60 min), most likely due to oxidative release of Mn^(III). In control experiments, we found that H₂O₂ (up to 0.2 mM) did not react with the porphyrin lacking Mn^(III). Hence, the tentative mechanism presented in Scheme 1 presumes the addition of HOO⁻ to Mn^(III) with concomitant heterolytic fission of the –O–O– bond (HO–Mn^(III)–OOH → HO–Mn^(V)=O + HO⁻). Similar reaction mechanisms have been proposed for the breakdown of peroxynitrite and peroxyacids by Mn^(III)-TMPyP.^{13,14} In a reaction system consisting of **1a,b** (20 μM), H₂O₂ (100 μM), and phenol (10 mM), we did

not observe formation of benzene-1,2-diol and benzene-1,4-diol, which attests for the lack of homolytic fission of HO–Mn^(III)–OOH to HO[•] and Mn^(III)–O[•] ↔ Mn^(IV)=O (C₆H₅OH + HO[•] + O₂ → HO₂[•] + C₆H₄(OH)₂;¹⁵ data not shown). Alternatively, release of H₂O from HO–Mn^(III)–OOH following a “push–pull” mechanism would yield the short-leaved dioxo species O=Mn^(V)=O. This pathway has been reported by Jin et al., who have shown that the rates of formation of O=Mn^(V)=O from Mn^(III)-TMPyP in the presence of HSO₅⁻, H₂O₂, and *t*-BuOOH follow bell-shaped curves with maxima at pH 10.¹⁶

The reaction of **1b** with H₂O₂ yielded **2** (or O=Mn^(V)=O), as suggested by the enhanced one-electron oxidation of Asc (Figure 1D) to semidehydroascorbyl radical (Scheme 2, [Asc]••; Figure 1D, inset) and by the increased oxidation of GSH by H₂O₂ in the presence of **1b** in both buffered aqueous solutions and in mouse embryonic cells (MEC; Supporting Information, Figures 5SI and 6SI). “Bleaching” of **1a,b** by H₂O₂ was fully inhibited by ascorbate (1 mM) and GSH (1 mM; data not shown), suggesting that the intracellular oxidation of **1b** to secondary porphyrin radicals will be hindered. In model experiments with CL-containing liposomes, **1b** also impeded oxidative modification of CL by Cyt c with H₂O₂ as a source of oxidizing equivalents (Supporting Information, Figure 7SI).

Next, we assessed the compartmentalization and IRM properties of **1b** in MEC. Upon incubation with MEC for 30 min, **1b** accumulated exclusively into mitochondria. Assuming that the volume of a single MEC is ~2.75 μm³ and its mitochondrial compartment occupies ~10% of the volume of the cytoplasm,¹⁷ we estimated the mitochondrial and cytosolic concentrations of **1b** to be ~3.5 and 0.005 mM, respectively (Figure 2A). These data indicate a considerable accumulation of **1b** at the sites of H₂O₂ production and execution of mitochondrial stages of apoptosis in cells exposed to IR.

Treatment of MEC with **1b** led to decreased levels of mitochondrial H₂O₂ (Figure 2B–D). Experiments with MEC were performed after mitochondria-targeted expression of HyPer, a fluorescent protein sensor for H₂O₂.¹⁸ HyPer has been engineered via insertion of yellow fluorescent protein (cpYFP) into the regulatory domain of the *E. coli* protein OxyR.^{18,19} Notably, changes in the fluorescence spectrum of HyPer are observed at submicromolar concentrations of H₂O₂ without any significant interference from other cellular oxidants. Flow cytometric experiments indicate that **1b** decreased the fluorescence of HyPer in control MEC and cells exposed to either H₂O₂ or IR (Figure 2B,C, respectively), suggesting that it competed with HyPer for mitochondrial H₂O₂. The effects of irradiation, H₂O₂, and **1b** on the fluorescence of HyPer were visualized by confocal microscopy (Figure 2D). Altogether, these data supported the hypothesis that **1b** can act as a radiation mitigator via clearance of mitochondrial H₂O₂. To further verify this hypothesis, we explored whether **1b** protects MEC against IR-induced apoptosis. MEC were irradiated at a dose of 10 Gy; 15 min thereafter, the cells were incubated with **1b**. Several end points of apoptotic cell death were investigated 48 h post-IR. Compound **1b** inhibited in a dose-dependent manner the activities of caspases 3/7 (Figure 2E), the externalization of phosphatidylserine (Anexin V positive cells; Figure 2F), and the release of cyt c from mitochondria (Figure 2F, inset). Similar protection was found with **1a**, although at higher concentrations (Supporting Information, Figure 4SI).




Figure 2. Cellular compartmentalization and radiomitigative properties of **1b**. (A) Compound **1b** preferentially compartmentalizes into mitochondria. (B and D) Effects of γ -rays (10 Gy), H₂O₂ (100 μ M), and **1b** (10 μ M) on the fluorescence of HyPer. (E and F) Effects of **1b** on caspases 3/7 activities and PS externalization in control and irradiated MEC; (inset E) Western bolt analysis of cytosolic Cyt c in control and irradiated MEC. (F) Radiomitigative effects of **1b** on mice exposed to γ -rays (9.25 Gy). The results in G are from two independent experiments and in A–C and E and F represent the mean \pm SE ($n = 3$).

We further assessed whether the in vitro radiomitigative effects of **1b** was translated into an in vivo effect in the mouse. Groups of 20 mice received **1b** (5 mg/kg body weight) 15 min after whole body irradiation (9.25 Gy). As shown in Figure 2G, \sim 80% of irradiated control mice were dead by day 21 (open circles), while nonirradiated mice receiving only ethanol as a vehicle (open triangles) survived the treatment. Irradiated mice post-treated with **1b** showed a significant increase in survival, with over 75% of mice alive at day 35 (closed circles).

In conclusion, the data presented herein indicate that **1b** acts as a potent radiation mitigator. It has been reported that the polycationic porphyrin complex Mn^(III)–TMPyP⁵⁺, when pre-administered for 14 days, significantly increased the survival of mice exposed to whole-body irradiation.²⁰ The pharmacological action of MnTMPyP⁵⁺ has been associated with its superoxide dismutase (SOD)-like activity. Cytotoxic, IR-induced mitochondrial production of (O₂^{•-}) precedes that of H₂O₂ (2O₂^{•-} + 2H⁺ \rightarrow H₂O₂),²¹ and it has been associated with inactivation of a

number of mitochondrial enzymes.²² While the reduction potentials of monocationic Mn^(III)–porphyrin complexes preclude reactions with (O₂^{•-}), targeted synthesis of SOD mimetics has been achieved via conjugation of the porphyrin ring with charged groups (reviewed in ref 23). This structural modification leads to decreased electron density near Mn^(III), thus increasing its ability to undergo one-electron reduction. However, the charge of these complexes reduces their ability to cross cell membranes (and thereby their bioavailability) and increases their toxicity due to intercalation into nucleic acids.²³ Our data complement previous studies on the medicinal chemistry of Mn^(III)–porphyrin complexes within the context of clearance of mitochondrial H₂O₂¹¹ rather than O₂^{•-}. The radiomitigative properties of porphyrin–Mn^(III) complexes may also reflect detoxification of peroxynitrite,²⁴ a possibility that has not been explored thus far. It is noteworthy that **1b** is a lipophilic compound whose bioavailability is not restricted by its molecular charge; we found millimolar mitochondrial concentrations of **1b** vs micromolar concentrations of MnTMPyP⁵⁺, reported

in ref 25. Furthermore, **1b** acted as a radiation mitigator when administered to mice after exposure to IR. This is an important pharmacological advantage, as IR damage to organisms is often the result of accidents, whereby post-IR treatment with free radical scavengers is not a viable option. It is possible that derivatives of **1b** containing functions with SOD-like activity may further enhance its pharmacological effectiveness and radiomitigative potential.

■ ASSOCIATED CONTENT

S Supporting Information. Additional details for biological assays and the synthesis and characterization of **1a,b**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Tel: +1-412-383-6910. E-mail: stoanovskyd@upmc.edu (D.A.S.). Tel: +1-412-624-9479. E-mail: Kagan@pitt.edu (V.E.K.).

Funding Sources

This work was supported by NIH Grant 2U19AI068021-06.

■ ABBREVIATIONS

GSH, glutathione; Asc, ascorbate; CL, cardiolipin; MEC, mouse embryonic cells; IR, ionizing radiation; IRP, radioprotectors; IRM, radiomitigators

■ REFERENCES

- (1) Shinomiya, N.; Kuno, Y.; Yamamoto, F.; Fukasawa, M.; Okumura, A.; Uefuji, M.; Rokutanda, M. Different mechanisms between premitotic apoptosis and postmitotic apoptosis in X-irradiated U937 cells. *Int. J. Radiat. Oncol., Biol., Phys.* **2000**, *47*, 767–777.
- (2) Maisin, J. R. Bacq and Alexander Award lecture—Chemical radioprotection: past, present, and future prospects. *Int. J. Radiat. Biol.* **1998**, *73*, 443–450.
- (3) Kouvaris, J. R.; Kouloulis, V. E.; Vlahos, L. J. Amifostine: The first selective-target and broad-spectrum radioprotector. *Oncologist* **2007**, *12*, 738–747.
- (4) Gudkov, A. V.; Komarova, E. A. The role of p53 in determining sensitivity to radiotherapy. *Nat. Rev. Cancer* **2003**, *3*, 117–129.
- (5) Burdelya, L. G.; Krivokrysenko, V. I.; Tallant, T. C.; Strom, E.; Gleiberman, A. S.; Gupta, D.; Kurnasov, O. V.; Fort, F. L.; Osterman, A. L.; Didonato, J. A.; Feinstein, E.; Gudkov, A. V. An agonist of toll-like receptor 5 has radioprotective activity in mouse and primate models. *Science* **2008**, *320*, 226–230.
- (6) Sohal, R. S.; Dubey, A. Mitochondrial oxidative damage, hydrogen peroxide release, and aging. *Free Radical Biol. Med.* **1994**, *16*, 621–626.
- (7) Kagan, V. E.; Tyurin, V. A.; Jiang, J.; Tyurina, Y. Y.; Ritov, V. B.; Amoscato, A. A.; Osipov, A. N.; Belikova, N. A.; Kapralov, A. A.; Kini, V.; Vlasova, I. I.; Zhao, Q.; Zou, M.; Di, P.; Svistunenko, D. A.; Kurnikov, I. V.; Borisenko, G. G. Cytochrome c acts as a cardiolipin oxygenase required for release of proapoptotic factors. *Nat. Chem. Biol.* **2005**, *1*, 223–232.
- (8) Epperly, M. W.; Melendez, J. A.; Zhang, X.; Nie, S.; Pearce, L.; Peterson, J.; Franicola, D.; Dixon, T.; Greenberger, B. A.; Komanduri, P.; Wang, H.; Greenberger, J. S. Mitochondrial targeting of a catalase transgene product by plasmid liposomes increases radioresistance in vitro and in vivo. *Radiat. Res.* **2009**, *171*, 588–595.
- (9) Murphy, M. P.; Smith, R. A. Targeting antioxidants to mitochondria by conjugation to lipophilic cations. *Annu. Rev. Pharmacol. Toxicol.* **2007**, *47*, 629–656.
- (10) Meunier, B. Metalloporphyrins as versatile catalysts for oxidation reactions and oxidative DNA cleavage. *Chem. Rev.* **1992**, *92*, 1411–1456.
- (11) Day, B. J. Catalytic antioxidants: A radical approach to new therapeutics. *Drug Discovery Today* **2004**, *9*, 557–566.
- (12) Abad, M. F.; Di Benedetto, G.; Magalhaes, P. J.; Filippin, L.; Pozzan, T. Mitochondrial pH monitored by a new engineered green fluorescent protein mutant. *J. Biol. Chem.* **2004**, *279*, 11521–11529.
- (13) Groves, J. T.; Lee, J.; Marla, S. S. Detection and Characterization of an Oxomanganese(V) Porphyrin Complex by Rapid-Mixing Stopped-Flow Spectrophotometry. *J. Am. Chem. Soc.* **1997**, *119*, 6269–6273.
- (14) Lee, J.; Hunt, J. A.; Groves, J. T. Manganese Porphyrins as Redox-Coupled Peroxynitrite Reductases. *J. Am. Chem. Soc.* **1998**, *120*, 6053–6061.
- (15) Ivanova, J.; Salama, G.; Clancy, R. M.; Schor, N. F.; Nylander, K. D.; Stoyanovsky, D. A. Formation of nitroxyl and hydroxyl radical in solutions of sodium trioxodinitrate: effects of pH and cytotoxicity. *J. Biol. Chem.* **2003**, *278*, 42761–42768.
- (16) Jin, N.; Lahaye, D. E.; Groves, J. T. A “Push-Pull” Mechanism for Heterolytic O–O Bond Cleavage in Hydroperoxo Manganese Porphyrins. *Inorg. Chem.* **2010**, *49*, 11516–11524.
- (17) Posakony, J. W.; England, J. M.; Attardi, G. Mitochondrial growth and division during the cell cycle in HeLa cells. *J. Cell Biol.* **1977**, *74*, 468–491.
- (18) Belousov, V. V.; Fradkov, A. F.; Lukyanov, K. A.; Staroverov, D. B.; Shakhbazov, K. S.; Tersikh, A. V.; Lukyanov, S. Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat. Methods* **2006**, *3*, 281–286.
- (19) Christman, M. F.; Storz, G.; Ames, B. N. OxyR, a positive regulator of hydrogen peroxide-inducible genes in *Escherichia coli* and *Salmonella typhimurium*, is homologous to a family of bacterial regulatory proteins. *Proc. Natl. Acad. Sci. U.S.A.* **1989**, *86*, 3484–3488.
- (20) Lee, J. H.; Park, J. W. A manganese porphyrin complex is a novel radiation protector. *Free Radical Biol. Med.* **2004**, *37*, 272–283.
- (21) Epperly, M. W.; Sikora, C. A.; DeFilippi, S. J.; Gretton, J. A.; Zhan, Q.; Kufe, D. W.; Greenberger, J. S. Manganese superoxide dismutase (SOD2) inhibits radiation-induced apoptosis by stabilization of the mitochondrial membrane. *Radiat. Res.* **2002**, *157*, 568–577.
- (22) Zhang, Y.; Marcillat, O.; Giulivi, C.; Ernster, L.; Davies, K. J. The oxidative inactivation of mitochondrial electron transport chain components and ATPase. *J. Biol. Chem.* **1990**, *265*, 16330–16336.
- (23) Batinic-Haberle, I.; Reboucas, J. S.; Spasojevic, I. Superoxide dismutase mimics: Chemistry, pharmacology, and therapeutic potential. *Antioxid. Redox Signaling* **2010**, *13*, 877–918.
- (24) Groves, J. T.; Marla, S. S. Peroxynitrite-Induced DNA Strand Scission Mediated by a Manganese Porphyrin. *J. Am. Chem. Soc.* **1995**, *117*, 9578–9579.
- (25) Spasojevic, I.; Chen, Y.; Noel, T. J.; Yu, Y.; Cole, M. P.; Zhang, L.; Zhao, Y.; St. Clair, D. K.; Batinic-Haberle, I. Mn porphyrin-based superoxide dismutase (SOD) mimic, MnIIIITE-2-PyP5+, targets mouse heart mitochondria. *Free Radical Biol. Med.* **2007**, *42*, 1193–1200.