

## 4-(2,2-Difluoro-1,3-benzodioxol-4-yl)-1*H*-pyrrole-3-carbonitrile

Fan-Wei Meng, Guang-Feng Hou, Ying-Hui Yu and Jin-Sheng Gao\*

Engineering Research Center of Pesticides of Heilongjiang University, Heilongjiang University, Harbin 150050, People's Republic of China, and College of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China

Correspondence e-mail: hgf1000@163.com


Received 12 December 2011; accepted 19 December 2011

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.043;  $wR$  factor = 0.118; data-to-parameter ratio = 14.1.

In the title compound,  $\text{C}_{12}\text{H}_6\text{F}_2\text{N}_2\text{O}_2$ , the 2,2-difluoro-1,3-benzodioxole ring system is approximately planar [maximum deviation = 0.012 (2)  $\text{\AA}$ ] and its mean plane is twisted with respect to the pyrrole ring, making a dihedral angle of 2.51 (9) $^\circ$ . In the crystal, N—H $\cdots$ N hydrogen bonds link the molecules into chains running along the  $a$  axis.  $\pi$ — $\pi$  stacking is also observed between parallel benzene rings of adjacent molecules, the centroid–centroid distance being 3.7527 (13)  $\text{\AA}$ .


### Related literature

For background to the title compound, see: Li *et al.* (2009); Pfluger *et al.* (1990). For the synthesis, see: Nyfeler & Ehrenfreund (1986).



### Experimental

#### Crystal data



$M_r = 248.19$

Triclinic, $P\bar{1}$	$V = 523.42 (18)\text{ \AA}^3$
$a = 7.5726 (15)\text{ \AA}$	$Z = 2$
$b = 7.8114 (16)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 8.9785 (18)\text{ \AA}$	$\mu = 0.13\text{ mm}^{-1}$
$\alpha = 93.58 (3)^\circ$	$T = 293\text{ K}$
$\beta = 94.65 (3)^\circ$	$0.39 \times 0.32 \times 0.15\text{ mm}$
$\gamma = 97.47 (3)^\circ$	

#### Data collection

Rigaku R-AXIS RAPID diffractometer	5120 measured reflections
Absorption correction: multi-scan ( <i>ABSCOR</i> ; Higashi, 1995)	2359 independent reflections
$T_{\min} = 0.950$ , $T_{\max} = 0.980$	1485 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.026$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.118$	$\Delta\rho_{\max} = 0.19\text{ e \AA}^{-3}$
$S = 1.04$	$\Delta\rho_{\min} = -0.15\text{ e \AA}^{-3}$
2359 reflections	
167 parameters	
1 restraint	

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N}1-\text{H}1\text{O} \cdots \text{N}2^i$	0.89 (1)	2.15 (1)	3.034 (2)	169 (2)

Symmetry code: (i)  $x + 1, y, z$ .

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors thank the Project of Innovation Service Platform of Heilongjiang Province (PG09J001) and Heilongjiang University, China, for supporting the work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5407).

### References

- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.  
Li, C., Miu, H.-D., Zeng, Z.-W., Wang, M.-J., Wu, Z.-X., Yang, F. & Shi, W.-J. (2009). *Modern Agrochem.* **8**, 19–24.  
Nyfeler, R. & Ehrenfreund, J. (1986). Switzerland Patent No. EP0206999.  
Pfluger, R. W., Indermühle, J. & Felix, F. (1990). Switzerland Patent No. EP0378046.  
Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.  
Rigaku/MSC (2002). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

## **supplementary materials**

*Acta Cryst.* (2012), E68, o222 [doi:10.1107/S1600536811054523]

## 4-(2,2-Difluoro-1,3-benzodioxol-4-yl)-1*H*-pyrrole-3-carbonitrile

F.-W. Meng, G.-F. Hou, Y.-H. Yu and J.-S. Gao

### Comment

Fludioxonil also know as Maxim, which is kind of fungicide developed and produced by Novartis (Li *et al.*, 2009; Pfluger *et al.*, 1990). Herein we report its structure.

In the title compound, phenyl and pyrrole ring are almost coplanar with a small dihedral angle of 2.51 (9) $^{\circ}$  (Figure 1). Intermolecular N—H···N hydrogen bonds link molecules into chains along [100] (Figure 2, Table 1).

### Experimental

The title compound was prepared by the reaction of 2-cyano-3-(2,2-difluoro-1,3-benzodioxol-4-yl)-2-propenamide and tosylmethyl isocyanide under alkaline condition (Robert & Josef, 1986). Colorless block crystals suitable for singl crystal X-ray diffraction were obtained by the recrystallization of title compound from a dichloromethane solution.

### Refinement

N-bound H atom was located in a differece Fourier map and positional parameters were refined,  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{N})$ . Other H atoms were placed in calculated positions with C—H = 0.93 Å, and refined in riding mode with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. The molecular structure of the title compound, showing displacement ellipsoids at the 50% probability level for non-H atoms.



Fig. 2. A partial packing view, showing the hydrogen-bonding chain structure along [100].

## 4-(2,2-Difluoro-1,3-benzodioxol-4-yl)-1*H*-pyrrole-3-carbonitrile

### Crystal data

$\text{C}_{12}\text{H}_6\text{F}_2\text{N}_2\text{O}_2$

$Z = 2$

$M_r = 248.19$

$F(000) = 252$

Triclinic,  $P\bar{1}$

$D_x = 1.575 \text{ Mg m}^{-3}$

Hall symbol: -P 1

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

# supplementary materials

---

$a = 7.5726 (15)$ Å	Cell parameters from 3390 reflections
$b = 7.8114 (16)$ Å	$\theta = 3.4\text{--}27.5^\circ$
$c = 8.9785 (18)$ Å	$\mu = 0.13 \text{ mm}^{-1}$
$\alpha = 93.58 (3)^\circ$	$T = 293$ K
$\beta = 94.65 (3)^\circ$	Block, colorless
$\gamma = 97.47 (3)^\circ$	$0.39 \times 0.32 \times 0.15$ mm
$V = 523.42 (18)$ Å <sup>3</sup>	

## Data collection

Rigaku R-AXIS RAPID diffractometer	2359 independent reflections
Radiation source: fine-focus sealed tube graphite	1485 reflections with $I > 2\sigma(I)$
$\omega$ scan	$R_{\text{int}} = 0.026$
Absorption correction: multi-scan ( <i>ABSCOR</i> ; Higashi, 1995)	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.4^\circ$
$T_{\text{min}} = 0.950, T_{\text{max}} = 0.980$	$h = -9 \rightarrow 9$
5120 measured reflections	$k = -10 \rightarrow 10$
	$l = -11 \rightarrow 10$

## Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.043$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.118$	$w = 1/[\sigma^2(F_o^2) + (0.0612P)^2 + 0.0143P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2359 reflections	$\Delta\rho_{\text{max}} = 0.19 \text{ e \AA}^{-3}$
167 parameters	$\Delta\rho_{\text{min}} = -0.15 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: <i>SHELXTL</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.020 (6)

## Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0680 (2)	0.6305 (2)	0.7663 (2)	0.0492 (4)
C2	-0.1137 (2)	0.6131 (3)	0.7677 (2)	0.0580 (5)
H2	-0.1723	0.5712	0.8478	0.070*
C3	-0.2036 (2)	0.6626 (3)	0.6407 (2)	0.0604 (5)
H3	-0.3277	0.6534	0.6348	0.072*
C4	-0.1158 (2)	0.7254 (2)	0.5222 (2)	0.0533 (5)
H4	-0.1833	0.7566	0.4395	0.064*
C5	0.0715 (2)	0.7442 (2)	0.52139 (18)	0.0412 (4)
C6	0.1551 (2)	0.6918 (2)	0.64884 (19)	0.0425 (4)
C7	0.3564 (2)	0.6277 (3)	0.8215 (2)	0.0580 (5)
C8	0.1679 (2)	0.8093 (2)	0.39662 (18)	0.0406 (4)
C9	0.0963 (2)	0.8705 (2)	0.25982 (19)	0.0424 (4)
C10	0.2365 (2)	0.9151 (2)	0.1754 (2)	0.0515 (5)
H10	0.2281	0.9581	0.0812	0.062*
C11	0.3488 (2)	0.8223 (3)	0.3852 (2)	0.0523 (5)
H11	0.4323	0.7921	0.4572	0.063*
C12	-0.0831 (2)	0.8924 (2)	0.2144 (2)	0.0474 (4)
F1	0.46357 (16)	0.74161 (18)	0.91577 (13)	0.0804 (4)
F2	0.43750 (16)	0.48669 (18)	0.81288 (16)	0.0803 (4)
N1	0.38690 (19)	0.8860 (2)	0.25235 (18)	0.0575 (5)
H101	0.4958 (17)	0.901 (3)	0.220 (3)	0.086*
N2	-0.2273 (2)	0.9115 (2)	0.17807 (19)	0.0631 (5)
O1	0.19296 (17)	0.5902 (2)	0.87638 (15)	0.0650 (4)
O2	0.33825 (15)	0.69183 (17)	0.68299 (13)	0.0540 (4)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0499 (10)	0.0565 (11)	0.0440 (10)	0.0091 (8)	0.0105 (8)	0.0133 (8)
C2	0.0507 (10)	0.0722 (13)	0.0562 (12)	0.0089 (9)	0.0222 (9)	0.0210 (10)
C3	0.0379 (9)	0.0835 (14)	0.0652 (13)	0.0130 (9)	0.0160 (8)	0.0240 (11)
C4	0.0403 (9)	0.0718 (12)	0.0515 (11)	0.0115 (8)	0.0092 (8)	0.0185 (10)
C5	0.0381 (8)	0.0465 (9)	0.0405 (9)	0.0070 (7)	0.0077 (7)	0.0078 (7)
C6	0.0354 (8)	0.0514 (9)	0.0422 (9)	0.0057 (7)	0.0092 (7)	0.0078 (8)
C7	0.0467 (10)	0.0838 (14)	0.0474 (11)	0.0123 (10)	0.0066 (8)	0.0252 (10)
C8	0.0374 (8)	0.0470 (9)	0.0386 (9)	0.0068 (7)	0.0065 (7)	0.0072 (7)
C9	0.0382 (8)	0.0512 (10)	0.0396 (9)	0.0084 (7)	0.0061 (7)	0.0081 (7)
C10	0.0451 (9)	0.0727 (12)	0.0402 (10)	0.0117 (8)	0.0082 (7)	0.0193 (9)
C11	0.0388 (9)	0.0759 (12)	0.0461 (11)	0.0121 (8)	0.0066 (7)	0.0224 (9)
C12	0.0442 (10)	0.0602 (11)	0.0398 (10)	0.0085 (8)	0.0064 (7)	0.0134 (8)
F1	0.0666 (8)	0.1180 (11)	0.0512 (7)	-0.0057 (7)	-0.0045 (6)	0.0148 (7)
F2	0.0734 (8)	0.0941 (9)	0.0855 (10)	0.0318 (7)	0.0215 (6)	0.0415 (8)
N1	0.0377 (8)	0.0870 (12)	0.0525 (10)	0.0100 (8)	0.0134 (7)	0.0254 (8)
N2	0.0437 (9)	0.0914 (13)	0.0580 (11)	0.0144 (8)	0.0042 (7)	0.0246 (9)

## supplementary materials

---

O1	0.0525 (8)	0.1003 (11)	0.0473 (8)	0.0114 (7)	0.0113 (6)	0.0335 (7)
O2	0.0382 (6)	0.0824 (9)	0.0444 (7)	0.0082 (6)	0.0065 (5)	0.0242 (6)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )


C1—C2	1.366 (3)	C7—F1	1.331 (2)
C1—C6	1.368 (2)	C7—O1	1.372 (2)
C1—O1	1.391 (2)	C7—O2	1.373 (2)
C2—C3	1.383 (3)	C8—C11	1.373 (2)
C2—H2	0.9300	C8—C9	1.437 (2)
C3—C4	1.382 (2)	C9—C10	1.375 (2)
C3—H3	0.9300	C9—C12	1.421 (2)
C4—C5	1.407 (2)	C10—N1	1.336 (2)
C4—H4	0.9300	C10—H10	0.9300
C5—C6	1.373 (2)	C11—N1	1.358 (2)
C5—C8	1.468 (2)	C11—H11	0.9300
C6—O2	1.3960 (19)	C12—N2	1.145 (2)
C7—F2	1.330 (2)	N1—H101	0.891 (10)
C2—C1—C6	122.93 (17)	F2—C7—O2	109.90 (18)
C2—C1—O1	127.94 (16)	F1—C7—O2	109.89 (16)
C6—C1—O1	109.12 (15)	O1—C7—O2	110.86 (15)
C1—C2—C3	114.73 (17)	C11—C8—C9	104.79 (14)
C1—C2—H2	122.6	C11—C8—C5	126.80 (15)
C3—C2—H2	122.6	C9—C8—C5	128.40 (14)
C4—C3—C2	122.42 (16)	C10—C9—C12	123.08 (16)
C4—C3—H3	118.8	C10—C9—C8	107.74 (15)
C2—C3—H3	118.8	C12—C9—C8	129.12 (15)
C3—C4—C5	122.74 (17)	N1—C10—C9	108.11 (15)
C3—C4—H4	118.6	N1—C10—H10	125.9
C5—C4—H4	118.6	C9—C10—H10	125.9
C6—C5—C4	112.87 (15)	N1—C11—C8	109.45 (15)
C6—C5—C8	123.29 (14)	N1—C11—H11	125.3
C4—C5—C8	123.83 (15)	C8—C11—H11	125.3
C1—C6—C5	124.29 (15)	N2—C12—C9	179.4 (2)
C1—C6—O2	108.24 (15)	C10—N1—C11	109.90 (14)
C5—C6—O2	127.46 (14)	C10—N1—H101	125.5 (15)
F2—C7—F1	105.63 (16)	C11—N1—H101	124.5 (15)
F2—C7—O1	110.18 (16)	C7—O1—C1	105.74 (14)
F1—C7—O1	110.26 (18)	C7—O2—C6	106.02 (13)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N1—H101 $\cdots$ N2 <sup>i</sup>	0.89 (1)	2.15 (1)	3.034 (2)	169 (2)

Symmetry codes: (i)  $x+1, y, z$ .

Fig. 1



## **supplementary materials**

---

**Fig. 2**

