

Published in final edited form as:

J Neurochem. 2012 January ; 120(Suppl 1): 9–21. doi:10.1111/j.1471-4159.2011.07519.x.

Proteolytic processing of Alzheimer's β -amyloid precursor protein

Han Zhang^{*,†,1}, Qilin Ma^{‡,1}, Yun-wu Zhang^{*}, and Huaxi Xu^{*,†}

^{*}Fujian Provincial Key Laboratory of Neurodegenerative Disease and Aging Research, College of Medicine, Xiamen University, Xiamen, Fujian 361005, China

[†]Neurodegenerative Disease Research Program, Sanford-Burnham Medical Research Institute, 10901 North Torrey Pines Road, La Jolla 92037, CA, USA

[‡]Department of Neurology, the First Affiliated Hospital of Xiamen University, Xiamen, Fujian 361005, China

Abstract

β -amyloid precursor protein (APP) is a critical factor in the pathogenesis of Alzheimer's disease (AD). APP undergoes posttranslational proteolysis/processing to generate the hydrophobic β -amyloid (A β) peptides. Deposition of A β in the brain, forming oligomeric A β and plaques, is identified as one of the key pathological hallmarks of AD. The processing of APP to generate A β is executed by β - and γ -secretase and is highly regulated. A β toxicity can lead to synaptic dysfunction, neuronal cell death, impaired learning/memory and abnormal behaviors in AD models *in vitro* and *in vivo*. Aside from A β , proteolytic cleavages of APP can also give rise to the APP intracellular domain (AICD), reportedly involved in multiple types of cellular events such as gene transcription and apoptotic cell death. In addition to amyloidogenic processing, APP can also be cleaved by α -secretase to form a soluble or secreted APP ectodomain (sAPP- α) that has been shown to be mostly neuro-protective. In this review, we describe the mechanisms involved in APP metabolism and the likely functions of its various proteolytic products to give a better understanding of the patho/physiological functions of APP.

Keywords

β -amyloid precursor protein; α -secretase; β -secretase; γ -secretase; caspase

Alzheimer's disease (AD), first officially described by the German psychiatrist and neuropathologist Alois Alzheimer in 1906, is the most common form of dementia. Characterized by progressive cognitive impairment, loss of memory and abnormal behavior, AD generally affects people over the age of 65. However, around 5% of AD patients develop the disease phenotype at a much younger age (~40 to 50 years old) and are classified as early-onset, most of which are also familial cases. Plaques consisting of β -amyloid (A β) peptide (Selkoe 1998), neurofibrillary tangles (NFTs) consisting largely of hyperphosphorylated microtubule-associated tau protein (Buee *et al.* 2000; Gendron and Petrucelli 2009) and neuron loss in the hippocampus and cortex regions are the major pathological hallmarks of Alzheimer's disease. As A β is believed to play a crucial role in the

Address correspondence and reprint requests to Dr. Huaxi Xu, Sanford-Burnham Medical Research Institute, 10901 N. Torrey Pines Rd., La Jolla, CA 92037, USA. Telephone: +1 858 7955246; Fax: +1 858 7955273. xuh@sanfordburnham.org.

¹These authors contributed equally to this study.

The authors declare no conflict of interest with respect to the research reported herein.

pathogenesis of AD, its precursor protein APP has become one of the most studied proteins in the field of AD research.

Basic knowledge of APP

The amyloid plaques associated with AD were first purified and found to consist of multimeric aggregates of A β polypeptide containing about 40 amino acid residues in the mid-1980s (Glenner and Wong 1984; Masters *et al.* 1985). Cloning of the complementary DNA (cDNA) of A β revealed that A β is derived from a larger precursor protein (Tanzi *et al.* 1987). The full-length cDNA of the amyloid precursor protein (APP) was later isolated and sequenced and APP was predicted to be a glycosylated integral membrane cell surface receptor protein with 695 amino acids (Kang *et al.* 1987). The A β peptide was found to be a cleavage product derived from the transmembrane domain of this large precursor protein.

The *APP* gene is located on chromosome 21 and contains 18 exons. Although alternative splicing of transcripts from the single *APP* gene results in several isoforms of the gene product, APP695, whose encoding cDNA lacks the gene sequence from exons 7 and 8, is preferentially expressed in neurons (Sandbrink *et al.* 1994). APP751, lacking exon 8, and APP770, encoded with all 18 exons, are predominant variants elsewhere (Yoshikai *et al.* 1990). Two homologues of APP were also identified and named APP-like protein 1 and 2 (APLP1 and APLP2) (Wasco *et al.* 1992; Wasco *et al.* 1993; Coulson *et al.* 2000). APLP2, similarly to APP, is expressed ubiquitously while APLP1 is only expressed in the brain and is only found in mammals.

The APP protein is a type I integral membrane protein with a large extracellular portion, a hydrophobic transmembrane domain, and a short C-terminus designated the APP intracellular domain (AICD). The extracellular portion of APP contains E1 and E2 domains and a Kunitz protease inhibitor (KPI) domain that is missing in APP695 (Kang and Muller-Hill 1990; Rohan de Silva *et al.* 1997). The E1 domain is reported to function as the major interaction interface for dimerization of cellular APP/APLPs (Soba *et al.* 2005). Although trans-dimerization of the E2 domain is also observed in X-ray structures (Wang and Ha 2004), biochemical assays have failed to confirm such a trans-dimerization (Soba *et al.* 2005). The levels of APP isoforms with a KPI domain seem to be elevated in patients with AD (Menendez-Gonzalez *et al.* 2005) and a splicing shift in neurons from APP695 to KPI-containing APP isoforms, along with increased A β generation, is observed when the NMDA receptor is activated (Bordji *et al.* 2010). These studies suggest that an alteration in APP splicing may contribute to disease pathogenesis.

APLP1 and APLP2 are both type I integral membrane proteins and share conserved structures with APP, including a large extracellular motif containing the E1 and E2 domains and a short intracellular domain. However, the transmembrane domain is not conserved among APP and APLPs and the A β sequence is unique to APP. In addition, APLP1 does not possess the KPI domain due to a lack of the encoding exon in the *APLP1* gene.

Manipulations of the APP/APLPs genes suggest that they are partially functionally redundant. Upon individual deletion of any of these three genes, mice are viable with only relatively subtle abnormalities (Zheng *et al.* 1995; Dawson *et al.* 1999). However, APP/APLP2, APLP1/APLP2 double knockout or APP/APLP1/APLP2 triple knockout mice show early postnatal lethality. Interestingly, APP/APLP1 double knockout mice are viable, indicating a crucial function of APLP2 in the absence of either APP or APLP1 (von Koch *et al.* 1997; Heber *et al.* 2000; Herms *et al.* 2004). Although it was reported that neurons generated from APP/APLP1/APLP2 triple knockout embryonic stem cells behave normally *in vitro* and *in vivo* (Bergmans *et al.* 2010), abnormal developments in the peripheral and central nervous system were repeatedly observed in both APP/APLP2 double knockout mice

and APP/APLP1/APLP2 triple knockout mice (Wang *et al.* 2005; Yang *et al.* 2005; Wang *et al.* 2007; Wang *et al.* 2009). Studies reported that the APP/APLP2 double knockout results in impaired neuromuscular junctions, indicating that the trans-synaptic interaction of APP is necessary for the proper development of motor neurons. Cortical dysplasia due to defective neuronal migration was also shown in APP/APLP1/APLP2 triple knockout mice (Herms *et al.* 2004). Although the physiological functions of APP/APLPs have not been well characterized, these loss-of-function phenotypes suggest the comprehensive involvement of these proteins during development.

APP processing

APP undergoes posttranslational processing, involving several different secretases and proteases, via two major pathways. In the non-amyloidogenic pathway, APP is sequentially cleaved by α -secretase and γ -secretase. α -cleavage, which cuts APP at the 17th amino acid inside the A β peptide sequence (Fig. 1), releases a large secreted extracellular domain (sAPP- α) and a membrane-associated C-terminal fragment consisting of 83 amino acids (C83). APP C83 is further cleaved by γ -secretase to release a P3 peptide and the APP intracellular domain (AICD), both of which are degraded rapidly. In the amyloidogenic pathway, APP is primarily processed by β -secretase at the 1st residue or at the 11th residue (so called β' site) of the A β peptide sequence (Fig. 1), shedding sAPP- β and generating a membrane associated C terminal fragment consisting of 99 amino acids (C99) (Sarah and Robert 2007). γ -secretase further cleaves C99 to release AICD and the amyloidogenic A β peptide which aggregates and fibrillates to form amyloid plaques in the brain.

α -secretase and α -cleavage

Since APP was found to be constitutively cleaved at the α -site to yield sAPP- α (Esch *et al.* 1990), three members of the ADAMs (a disintegrin and metalloproteinase), ADAM-10, ADAM-17 and ADAM-9 have been proposed as the α -secretase (Buxbaum *et al.* 1998; Koike *et al.* 1999; Lammich *et al.* 1999).

ADAMs are type I integral membrane proteins that belong to the zinc protease super family and have been implicated in the control of cytokine and growth factor shedding. ADAM10 is widely expressed in the brain and in other tissues (Chantry and Glynn 1990; Howard *et al.* 1996) and a several fold increase in sAPP- α levels in cell lines overexpressing ADAM10 can be observed (Lammich *et al.* 1999). Moderate neuronal overexpression of human ADAM10 increases sAPP- α production while reducing A β generation/plaque formation in mice carrying the human APP V717I mutation, while expression of a catalytically-inactive form of the ADAM10 mutation increases the size and number of amyloid plaques in mouse brains (Lammich *et al.* 1999). These findings suggest that ADAM10 may be responsible for constitutive α -cleavage activity. On the other hand, although sAPP- α generation is not affected in ADAM9/17 knock-down cell lines nor in mice carrying deficient ADAM9/17 genes (Weskamp *et al.* 2002; Kuhn *et al.* 2010), overexpression of ADAM9/17 does increase the level of sAPP- α under some conditions, suggesting that ADAM9 and ADAM17 are more likely involved in the regulated α -cleavage of APP rather than in constitutive α -cleavage.

In addition to APP, ADAMs have many other substrates (Qiang *et al.* 2009; Altmepfen *et al.* 2011). For example, ADAM17 has been identified as the protease responsible for shedding of the transmembrane form of tumor necrosis factor- α at its physiological processing site (Black *et al.* 1997; Moss *et al.* 1997). ADAM17 also cleaves the epidermal growth factor (EGF) family members (Peschon *et al.* 1998; Lee *et al.* 2003) and ADAM17-deficient mice possess the phenotype of mice with an EGF signaling defect (Buxbaum *et al.* 1998). ADAM10 can cleave Notch and ADAM10-deficient mice develop a loss of Notch

signaling phenotype (Hartmann *et al.* 2002). In addition, ADAM10 and ADAM17 were found to be directly responsible for the constitutive and protein kinase c-regulated processing of prions, respectively (Gouras *et al.* 2000; Vincent *et al.* 2001), whereas another ADAM member, ADAM9, acts as an upstream activator of ADAM10 during this process (Cisse *et al.* 2005).

β -secretase and β -cleavage

A β generation is initiated by β -cleavage at the ectodomain of APP, resulting in the generation of an sAPP- β domain and the membrane associated APP C-terminal fragment C99. The putative β -secretase, β -site APP cleaving enzyme 1 (BACE1), was first identified and characterized in 1999 (Sinha *et al.* 1999; Vassar *et al.* 1999; Yan *et al.* 1999; Hussain *et al.* 2000; Lin *et al.* 2000). BACE1 is a type I transmembrane aspartyl protease with its active site on the luminal side of the membrane. The originally identified full-length BACE1 has 501 amino acids (BACE1–501) and is predominantly expressed in perinuclear post-Golgi membranes, vesicular structures throughout the cytoplasm, as well as on the cell surface (Ehehalt *et al.* 2002). Several other minor transcripts of BACE1 (BACE1–476, 457 and 432) derived from alternative splicing were found later (Tanahashi and Tabira 2001), however their β -cleavage activity and subcellular localization are different from those of BACE1–501 (Tanahashi and Tabira 2001; Ehehalt *et al.* 2002). Knocking out the BACE1 gene prevents A β generation and completely abolishes A β pathology in mice expressing the Swedish mutation of human APP (Farzan *et al.* 2000; Cai *et al.* 2001; Luo *et al.* 2001; Roberds *et al.* 2001; Ohno *et al.* 2004; Laird *et al.* 2005). The expression level and activity of BACE1 were also found to be elevated in AD patients (Holsinger *et al.* 2002; Yang *et al.* 2003). However, there is no evidence linking BACE1 gene variants with familial AD (FAD) and the BACE1 gene variants do not affect BACE1's activity or APP processing/A β generation (Sjolander *et al.* 2010). More recently, BACE1 knockout mice have been found to exhibit hypomyelination and altered neurological behaviors, such as reduced grip strength and elevated pain sensitivity, probably because the physiological functions of other BACE1 substrates, such as neuregulin 1 and the voltage-gated sodium channel (Nav1) β 2 subunit, are negatively affected in these mice (Laird *et al.* 2005; Hu *et al.* 2006; Willem *et al.* 2006; Gersbacher *et al.* 2010; Hitt *et al.* 2010).

BACE2 is a homolog of BACE1 (Acquati *et al.* 2000; Xin *et al.* 2000) and human BACE2 shares 71% homology and 45% identity with human BACE1. Similar to BACE1, BACE2 is predominantly localized in post-Golgi structures and on the cell surface (Ehehalt *et al.* 2002). Notably, several splice variants of BACE2 were also identified and found to be expressed in several central nervous system subregions (Solans *et al.* 2000). However, the expression level of BACE2 is much lower in the brain than BACE1 and is mostly expressed in glial cells (Laird *et al.* 2005). In addition, BACE2 cleaves APP within the A β domain in a manner more similar to α -secretase than β -secretase (Farzan *et al.* 2000).

Cathepsin B was once suspected as a β -secretase candidate (Hook *et al.* 2005; Schechter and Ziv 2008). However, recent studies suggest that Cathepsin B can degrade A β into harmless fragments. It is therefore thought that Cathepsin B plays a role in the body's natural defense against AD (Mueller-Steiner *et al.* 2006).

γ -secretase and γ -cleavage

Both α -cleavage and β -cleavage generate short APP C-terminal fragments that are further processed by γ -secretase. Distinct from α -/ β -secretases, γ -activity involves a large proteinase complex consisting of at least four major protein components (Presenilin1 or Presenilin2, PEN2, APH1 and Nicastrin) (Vetrivel *et al.* 2006).

Presenilins (PSs) were identified and cloned in the mid-1990s (Levy-Lahad *et al.* 1995; Sherrington *et al.* 1995). Genetic mutations of PSs are found in a large portion of familiar AD (FAD) cases, indicating its crucial role in AD. Although other proteins are required for the γ -secretase complex, PSs are believed to contain the actual protease activity (Wolfe *et al.* 1999; Wen *et al.* 2008; Ahn *et al.* 2010a). PSs are multi-transmembrane proteins and can be cleaved at the cytoplasmic loop between the 6th and 7th transmembrane regions to generate an N terminal and a C terminal fragment during post-translational maturation (Thinakaran *et al.* 1996). The two fragments interact with each other after the cleavage and they are both necessary for γ -secretase activity. Transgenic mice overexpressing PSs with FAD mutations show significantly increased A β 42 levels, suggesting that PS mutations probably induce AD by producing more of the hydrophobic A β 42 form (Duff *et al.* 1996; Xia *et al.* 2001).

Nicastrin, identified as a protein that interacts with PS in 2000 (Yu *et al.* 2000), is a type I membrane glycoprotein with a large ectodomain (Perrin *et al.* 2009). Nicastrin undergoes a glycosylation/maturation process that causes a conformation change in its ectodomain, which is crucial for the assembly and maturation of the γ -secretase complex and γ -activity (Shirotani *et al.* 2003; Chavez-Gutierrez *et al.* 2008). Mature nicastrin can bind to the ectodomain of APP CTFs derived through α -/ β -secretase cleavage and may act as a substrate receptor of γ -secretase (Shirotani *et al.* 2003; Kaether *et al.* 2004; Shah *et al.* 2005).

PEN2 and APH1 are another two γ -secretase complex components that were originally identified as the enhancers of PSs (Francis *et al.* 2002). APH1 is a multiple transmembrane protein with seven transmembrane domains and a cytosolic C terminus (Fortna *et al.* 2004). APH1 interacts with immature nicastrin and PS to form a relatively stable pre-complex which is then translocated to the trans-Golgi from the ER/cis-Golgi for further maturation (Lee *et al.* 2002; Kimberly *et al.* 2003; Takasugi *et al.* 2003; Jankowsky *et al.* 2004). PEN2 is a hairpin-like protein with two transmembrane domains and with both ends in the lumen (Crystal *et al.* 2003; Takasugi *et al.* 2003). PEN2 is found to mediate the endoproteolysis of PS (Luo *et al.* 2003). Enhanced γ -secretase activity is also observed when PEN2 is exogenously expressed (Shiraishi *et al.* 2004).

The γ -secretase complex is assembled in sequential steps. Nicastrin and APH1 initially form a subcomplex (Shirotani *et al.* 2004) and then PS binds to the Nicastrin-APH1 subcomplex (Takasugi *et al.* 2003). The joining of PEN2 results in a conformation-dependent activation of γ -secretase (Kimberly *et al.* 2003; Takasugi *et al.* 2003). Nicastrin, PEN2, APH1 and PS interact with each other and also mutually modulate each other (Lee *et al.* 2002; Steiner *et al.* 2002; Pasternak *et al.* 2003; Kaether *et al.* 2004). Nicastrin ablation leads to decreased levels of APH1, PEN2 and PS1 fragments, accompanied by increased levels of immature full-length PS1 (Zhang *et al.* 2005). Downregulation of APH1, PEN2 or nicastrin also reduces the processing of PS and results in impaired γ -cleavage of APP and Notch (Francis *et al.* 2002). PS deficiency also results in decreased levels of PEN2 and APH1, as well as impaired glycosylation/maturation of Nicastrin (Zhang *et al.* 2005).

γ -secretase cleaves APP at multiple sites and in sequential steps to generate A β peptides of different lengths (Fig. 1). The majority of A β peptides produced are 40 amino acids long, however, peptides ranging from 38 to 43 amino acids are found *in vivo*. Besides the dominant γ -cleavage site at 40 and 42 residues, ζ -cleavage at 46 and ϵ -cleavage at 49 residues are also thought to be mediated by γ -secretase (Weidemann *et al.* 2002; Zhao *et al.* 2004; Raben *et al.* 2005; Sato *et al.* 2005). Accordingly, various AICDs (C50, C53, C57 and C59) can be generated during these multi-site cleavages executed by γ -secretase. However, all of the endogenous AICD forms are rarely detected, probably due to their very rapid degradation (Ag 2000; Lu *et al.* 2000; Sastre *et al.* 2001; Yu *et al.* 2001; Sato *et al.* 2003). Interestingly, as the substrate of γ -secretase, APP itself can regulate the intracellular

trafficking and cell surface delivery of PS1 (Kuzuya *et al.* 2007; Liu *et al.* 2009). In addition, APP has been found to possess a domain that negatively modulates γ -secretase activity in A β production by binding to an allosteric site within the γ -secretase complex (Kuzuya *et al.* 2007; Ahn *et al.* 2010b; Zhang and Xu 2010). These results reveal a novel mutual regulation between γ -secretase and its substrate.

In addition to cleaving APP, γ -secretase cleaves many other single transmembrane proteins within the transmembrane domain (Lee *et al.* 2002). One of the most important γ -secretase substrates, Notch, can release its intracellular domain (NICD) upon γ -cleavage. NICD is well-known as a signal molecule that transactivates a number of genes critical to development (Kopan *et al.* 1996; Schroeter *et al.* 1998). Mice with ablation of PS1, Nicastrin, or APH1A show Notch-deficient-like lethal phenotypes and neuronal tube formation defects (Shen *et al.* 1997; Donoviel *et al.* 1999; Li *et al.* 2003; Ma *et al.* 2005). Postnatal forebrain-specific inactivation of PS1 in APP transgenic mice, although preventing A β accumulation, failed to rescue contextual memory long-term. Conditional inactivation of γ -secretase components in the forebrain resulted in progressive memory impairment and age-related neurodegeneration (Dominguez *et al.* 2005; Saura *et al.* 2005; Serneels *et al.* 2009; Tabuchi *et al.* 2009).

Caspase-cleavage

In addition to cleavages involving secretases, APP can be cleaved by caspases independently at its C terminus (Asp664 of APP695), releasing a short tail containing the last 31 amino acids (C31) of APP and a fragment (Jcasp) from between the γ - and caspase-cleavage sites (Lu *et al.* 2000). Caspase-cleavages of APP are thought to be harmful since both C31 and the Jcasp fragment generated were found to be cytotoxic (Lu *et al.* 2003a). Transgenic mice with Swedish and Indiana mutations of APP show increased susceptibility to seizures, which can be abolished by a D664A mutation that disrupts caspase-cleavages (Ghosal *et al.* 2009). These results indicate that caspase-cleavages of APP contribute, at least in part, to the neurotoxicity of APP processing products.

Function of APP and its fragments

Full-length APP

Due to its highly similar structure to Notch, APP has been proposed to function as a cell surface receptor [reviewed in (Zheng and Koo 2011)]. Several studies have reported that certain ligands, including A β , F-spondin, Nogo-66, netrin-1 and BRI2, bind to the extracellular domain of APP, resulting in modulated APP processing and sequential downstream signals (Lorenzo *et al.* 2000; Lu *et al.* 2003b; Ho and Sudhof 2004; Park *et al.* 2006; Lourenco *et al.* 2009; Matsuda *et al.* 2009; Zheng and Koo 2011). However, the physiological functions of these interactions remain to be determined. Nevertheless, APP is more widely accepted as a protein contributing to cell adhesion via its extracellular domain. Studies have demonstrated that the E1 and E2 regions of APP can interact with extracellular matrix proteins (Small *et al.* 1999). Furthermore, the E1 and E2 regions of APP were found to interact with themselves, in parallel or anti-parallel, forming homo- (with APP) or heterodimers (with APLPs) (Wang and Ha 2004; Soba *et al.* 2005; Dahms *et al.* 2010). Recent studies also suggest APP/APLPs as synaptic adhesion molecules as silencing of APP led to defects in neuronal migration (Young-Pearse *et al.* 2007; Wang *et al.* 2009; Norstrom *et al.* 2010).

sAPP- α

The constitutively secreted sAPP- α has been found to be neuro-protective (Mattson *et al.* 1993; Furukawa *et al.* 1996; Han *et al.* 2005; Ma *et al.* 2009). sAPP- α is thought to promote

neurite outgrowth and synaptogenesis as well as cell adhesion (Mattson 1997; Gakhar Koppole 2008). Studies have found that sAPP- α is a growth factor (Herzog *et al.* 2004; Siemes *et al.* 2006) that regulates the proliferation of embryonic and adult neural stem cells (Ohsawa *et al.* 1999; Caille *et al.* 2004). *In vivo* studies have also shown that sAPP- α promotes learning and memory in animal models (Meziane *et al.* 1998; Taylor *et al.* 2008). sAPP- α alone is able to rescue most of the abnormalities of APP deficient mice (Ring *et al.* 2007), implying that most of the physiological functions of APP are conducted by its extracellular domain.

sAPP- β

Although there are only 17 amino acids difference between sAPP- β and sAPP- α , sAPP- β reportedly lacks most of the neuroprotective effects of sAPP- α (Furukawa *et al.* 1996). A recent study suggested that sAPP- β can be cleaved to generate an N-terminal fragment that is a ligand for death receptor 6, activating caspase-6 which further stimulates axonal pruning and neuronal cell death (Nikolaev *et al.* 2009).

A β

The physiological and pathological functions of A β have been extensively investigated due to its central role in AD. Studies have demonstrated that A β overproduction leads to neurotoxicity, neuronal tangle formation, synaptic damage and eventually neuron loss in the pathologically affected brain regions (Selkoe 1998; Shankar and Walsh 2009). Among the various A β peptides generated by the multiple-site cleavages of secretases, A β 42 has proved to be more hydrophobic and amyloidogenic than others (Burdick *et al.* 1992). Most mutations related to hereditary familial AD either increase A β generation or the ratio of A β 42/ A β 40 (Borchelt *et al.* 1996; Scheuner *et al.* 1996). Studies also suggest that increased A β 42 levels probably provide the core for oligomerization, fibrillation and amyloid plaque generation (Jarrett *et al.* 1993; Iwatsubo *et al.* 1994).

Although excessive A β causes neurotoxicity, some studies have shown that A β 40 protects neurons against A β 42-induced neuronal damage and is required for the viability of central neurons (Plant *et al.* 2003; Zou *et al.* 2003). Moreover, two groups recently reported that low doses (picomolar) of A β can positively modulate synaptic plasticity and memory by increasing long-term potentiation (Morley *et al.* 2008; Puzzo *et al.* 2008), revealing a novel physiological function of A β under normal conditions.

AICD

Depending on the exact cleavage site of γ -secretase, AICD may have 59, 57, 53 or 50 residues. However, because AICD is quickly degraded after γ -cleavage, the biochemical features and physiological functions of AICD *in vivo* are difficult to study. So far, most of the information on AICD is deduced based on exogenous systems.

There are several conserved regions located on AICD: a YTSI (653–656 residues according to APP695) motif near the cell membrane, a VTPEER (667–762 residues) helix capping box and a YENPTY (682–687 residues) domain which mediates the interaction between APP/AICD and phosphotyrosine binding (PTB) domain containing proteins. Many AICD binding proteins have been identified. Some of the proteins, including KLC, Fe65, Shc, JIP, Numb, X11, Clathrin and mDab1, were found to share one or several common PTB domains that specifically interact with the Asn-Pro-X-Tyr amino acid sequence present in the YENPTY motif of AICD (Borg *et al.* 1996; Brassler *et al.* 1996; M. McLoughlin and Cj Miller 1996; Howell *et al.* 1999; Salcini *et al.* 1999; Kamal *et al.* 2000; Matsuda *et al.* 2001; Weggen *et al.* 2001; Roncarati *et al.* 2002; Tarr *et al.* 2002b; Tarr *et al.* 2002a; Inomata *et al.* 2003; Matsuda *et al.* 2003). Other proteins, such as PAT1, SET and 14-3-3 γ , are believed to bind

to the YTSI or VTPEER motif of AICD (Zheng *et al.* 1998; Madeira *et al.* 2005; Sumioka *et al.* 2005). AICD, therefore, may have different functions when interacting with its' various binding partners.

AICD also contains three phosphorylation sites, including two threonine residues at 654 and 668 and a serine residue at 665. AICD has been found to be phosphorylated by PKC, calcium-calmodulin dependent-kinase II, GSK3- β , Cdk5 and JNK at the Ser/Thr sites mentioned above. Such phosphorylation may affect APP processing or the binding of AICD-interacting proteins, thus affecting the function of AICD (Gandy *et al.* 1988; Iwatsubo *et al.* 1994; Iijima *et al.* 2000; Inomata *et al.* 2003).

Since the generation of AICD is very similar to that of many other signaling molecules, such as NICD, which is also generated by γ -cleavage and can mediate the transcription of genes important for development, AICD may function in a similar fashion. Indeed, a role for AICD in gene transactivation is supported by several studies. The most widely accepted mechanism is that AICD, together with Fe65 and Tip60, forms a transcriptionally-active complex. Fe65 is one of the most well studied proteins that bind to the YENPTY motif of AICD. Tip60, a histone acetyltransferase, is a component of a larger nuclear complex with DNA binding, ATPase and DNA helicase activities (Ikura *et al.* 2000). Although Tip60 does not bind to AICD directly, an indirect interaction between AICD and Tip60 is mediated by Fe65. Upon forming this complex, AICD is stabilized and can be translocated into the nucleus to regulate expression of genes such as KAI1, Neprilysin, LRP1, p53, GSK-3 β and EGFR (Baek *et al.* 2002; Kim *et al.* 2003; Cao and Sudhof 2004; Pardossi-Piquard *et al.* 2005; Alves da Costa *et al.* 2006; Liu *et al.* 2007; Zhang *et al.* 2007). Another transactivating complex consisting of AICD, Fe65 and CP2/LSF/LBP1 has also been reported to induce the expression of GSK3- β (Kim *et al.* 2003). Although the transactivation effect of AICD has been observed by different research groups, the model of how AICD functions is still controversial (Cao and Sudhof 2004; Hass and Yankner 2005; Nakaya and Suzuki 2006). It was originally widely accepted that translocating into the nucleus was necessary for AICD to activate gene expression. However, some studies have suggested that an AICD mediated conformational change in Fe65 is sufficient and that the nuclear translocation of AICD is not required (Cao and Sudhof 2004; Hass and Yankner 2005). Another study has suggested that the nuclear translocation of AICD is independent of Fe65 and is a result of its phosphorylation at T668 (Nakaya and Suzuki 2006). More investigation is required to further elucidate the mechanism.

As an adaptor protein involved in protein sorting and trafficking, X11 has been suggested as affecting APP trafficking/metabolism by interacting with AICD, leading to reduced A β production. X11 has also been found to suppress the transactivation of AICD, possibly by competing with AICD for the recruitment of Fe65, as they share the same binding motif (Biederer *et al.* 2002).

Many studies have documented that AICD is cytotoxic and that overexpressing different AICDs (C31, C57, C59) in HeLa, H4, N2a or PC12 cell lines, as well as neuronal cell lines, induces cell death (Lu *et al.* 2000). AICDs lacking the NPTY motif that mediates the interaction between AICD and other proteins failed to induce apoptosis, suggesting the functional involvement of an AICD-binding protein (Xu *et al.* 2006). A Tip60 loss-of-function mutation makes HeLa cells more resistant to apoptosis (Ikura *et al.* 2000). These results indicate that the cytotoxicity of AICD may be mediated by its interacting proteins. For example, JIP is the scaffolding protein of the JNK pathway kinase and is involved in various cell events including neuronal apoptosis and axonal transporting. By binding to AICD, JIP mediates APP/AICD phosphorylation at Thr668, thus modulating APP

trafficking, maturation and processing. Additionally, there is evidence suggesting that the cell death triggered by AICD is partially mediated by JIP (Taru *et al.* 2002).

In addition, AICD-induced cytotoxicity may be mediated by its regulation targets (Kim *et al.* 2003; Alves da Costa *et al.* 2006; Vetrivel *et al.* 2006). For example, P53 expression, as well as p53-mediated apoptosis, can be enhanced by AICD (Alves da Costa *et al.* 2006; Ozaki *et al.* 2006). Another AICD target gene, GSK3- β , may also contribute to AICD-related cytotoxicity by upregulating tau hyperphosphorylation. GSK-3 β activation and CRMP-2 phosphorylation, along with downstream tau hyper-phosphorylation/aggregation, neurodegeneration and memory loss, are observed in an AICD C59 transgenic mouse line in which Fe65 is co-expressed (Ryan and Pimplikar 2005; Ghosal *et al.* 2009). However, another group found only increased neuronal sensitivity to toxic and apoptotic stimuli in mice overexpressing AICD either alone or with Fe65 (Giliberto *et al.* 2008).

On the other hand, C31, a short form of AICD generated by caspase cleavage, has been reported to directly activate caspase-3 in the tumor cell death process (Lu *et al.* 2000; Bertrand *et al.* 2001; Nishimura *et al.* 2002; Madeira *et al.* 2005). C31 also appears to induce a caspase-independent toxicity by selectively increasing A β 42 (Dumanchin-Njock *et al.* 2001). A D664A mutation in AICD, blocking caspase cleavage, abolished the synaptic loss, axonal defects and behavioral changes found in PDAPP(D664A) mice (Galvan *et al.* 2006). APP binding protein 1 (APBP1) reportedly interacts with AICD and activates the neddylation pathway (Chen 2004), further downregulating the level of β -catenin and potentially resulting in apoptosis. In addition, cellular Ca²⁺ homeostasis appears to be modulated by AICD (Hamid *et al.* 2007).

Conclusion

As a crucial protein in AD, APP is multifunctional and can be post-translationally processed via different pathways. The physiological and pathological functions of APP are closely related to the processing of APP, which may suggest therapeutic treatments for AD. In this review, we have summarized the current knowledge of APP metabolism. Further understanding of APP processing and the patho/physiological functions of various APP metabolites will be important for the development of an AD treatment.

Abbreviations used

Aβ	β -amyloid
AD	Alzheimer's disease
ADAM	a disintegrin and metalloproteinase
AICD	APP intracellular domain
APLP1	APP-like protein 1
APLP2	APP-like protein 2
APP	β -amyloid precursor protein
BACE1	β -site APP cleaving enzyme 1
cDNA	complementary DNA
EGF	epidermal growth factor
FAD	familiar Alzheimer's disease
KPI	Kunitz protease inhibitor

NICD	Notch intracellular domain
PS	presenilin
PTB	phosphotyrosine binding

Acknowledgments

This work was supported in part by National Institutes of Health grants (R01AG021173, R01NS046673, R01AG030197 and R03AG034366 to H.X.), and grants from the Alzheimer's Association (to H.X. and Y.-w.Z.), the American Health Assistance Foundation (to H.X.), National Natural Science Foundation of China (30973150 to Y.-w.Z.), the 973 Prophase Project (2010CB535004 to Y.-w.Z.), and Natural Science Funds for Distinguished Young Scholar of Fujian Province (2009J06022 to Y.-w.Z.). Y.-w.Z. is supported by the Program for New Century Excellent Talents in Universities (NCET), the Fundamental Research Funds for the Central Universities, and Fok Ying Tung Education Foundation.

References

- Acquati F, Accarino M, Nucci C, Fumagalli P, Jovine L, Ottolenghi S, Taramelli R. The gene encoding DRAP (BACE2), a glycosylated transmembrane protein of the aspartic protease family, maps to the down critical region. *FEBS Lett.* 2000; 468:59–64. [PubMed: 10683441]
- Ag CB. Generation of an Apoptotic Intracellular Peptide by gamma-Secretase Cleavage of Alzheimer's Amyloid Protein Precursor. *J. Alzheimer's Dis.* 2000; 2:289–301. [PubMed: 12214090]
- Ahn K, Shelton CC, Tian Y, Zhang X, Gilchrist ML, Sisodia SS, Li YM. Activation and intrinsic gamma-secretase activity of presenilin 1. *Proc. Natl. Acad. Sci. USA.* 2010; 107:21435. [PubMed: 21115843]
- Altmeppen HC, Prox J, Puig B, Kluth MA, Bernreuther C, Thurm D, Jorissen E, Petrowitz B, Bartsch U, De Strooper B, Saftig P, Glatzel M. Lack of a-disintegrin-and-metalloproteinase ADAM10 leads to intracellular accumulation and loss of shedding of the cellular prion protein in vivo. *Mol. Neurodegener.* 2011; 6:36. [PubMed: 21619641]
- Alves, da Costa C.; Sunyach, C.; Pardossi-Piquard, R.; Sevalle, J.; Vincent, B.; Boyer, N.; Kawarai, T.; Girardot, N.; St, George-Hyslop P.; Checler, F. Presenilin-dependent gamma-secretase-mediated control of p53-associated cell death in Alzheimer's disease. *J. Neurosci.* 2006; 26:6377–6385. [PubMed: 16763046]
- Baek SH, Ohgi KA, Rose DW, Koo EH, Glass CK, Rosenfeld MG. Exchange of N-CoR corepressor and Tip60 coactivator complexes links gene expression by NF-kappaB and beta-amyloid precursor protein. *Cell.* 2002; 110:55–67. [PubMed: 12150997]
- Bergmans BA, Shariati SA, Habets RL, Verstreken P, Schoonjans L, Muller U, Dotti CG, De Strooper B. Neurons generated from APP/APLP1/APLP2 triple knockout embryonic stem cells behave normally in vitro and in vivo: lack of evidence for a cell autonomous role of the amyloid precursor protein in neuronal differentiation. *Stem Cells.* 2010; 28:399–406. [PubMed: 20049903]
- Bertrand E, Brouillet E, Caille I, Bouillot C, Cole GM, Prochiantz A, Allinquant B. A short cytoplasmic domain of the amyloid precursor protein induces apoptosis in vitro and in vivo. *Mol. Cell. Neurosci.* 2001; 18:503–511. [PubMed: 11922141]
- Biederer T, Cao X, Sudhof TC, Liu X. Regulation of APP-dependent transcription complexes by Mint/X11s: differential functions of Mint isoforms. *J. Neurosci.* 2002; 22:7340–7351. [PubMed: 12196555]
- Black RA, Rauch CT, Kozlosky CJ, Peschon JJ, Slack JL, Wolfson MF, Castner BJ, Stocking KL, Reddy P, Srinivasan S, Nelson N, Boiani N, Schooley KA, Gerhart M, Davis R, Fitzner JN, Johnson RS, Paxton RJ, March CJ, Cerretti DP. A metalloproteinase disintegrin that releases tumour-necrosis factor-alpha from cells. *Nature.* 1997; 385:729–733. [PubMed: 9034190]
- Borchelt DR, Thinakaran G, Eckman CB, Lee MK, Davenport F, Ratovitsky T, Prada CM, Kim G, Seekins S, Yager D. Familial Alzheimer's disease-linked presenilin 1 variants elevate A-beta 1–42/1–40 ratio in vitro and in vivo. *Neuron.* 1996; 17:1005–1013. [PubMed: 8938131]

- Bordji K, Becerril-Ortega J, Nicole O, Buisson A. Activation of extrasynaptic, but not synaptic, NMDA receptors modifies amyloid precursor protein expression pattern and increases amyloid-ss production. *J. Neurosci.* 2010; 30:15927–15942. [PubMed: 21106831]
- Borg JP, Ooi J, Levy E, Margolis B. The phosphotyrosine interaction domains of X11 and FE65 bind to distinct sites on the YENPTY motif of amyloid precursor protein. *Mol. Cell. Biol.* 1996; 16:6229–6241. [PubMed: 8887653]
- Brassler SL, Gray MD, Sopher BL, Hu Q, Hearn MG, Pham DG, Dinulos MB, Fukuchi KI, Sisodia SS, Miller MA. cDNA cloning and chromosome mapping of the human Fe65 gene: interaction of the conserved cytoplasmic domains of the human -amyloid precursor protein and its homologues with the mouse Fe65 protein. *Hum. Mol. Genet.* 1996; 5:1589. [PubMed: 8894693]
- Buee L, Bussiere T, Buee-Scherrer V, Delacourte A, Hof PR. Tau protein isoforms, phosphorylation and role in neurodegenerative disorders. *Brain Res. Brain Res. Rev.* 2000; 33:95–130. [PubMed: 10967355]
- Burdick D, Soreghan B, Kwon M, Kosmoski J, Knauer M, Henschen A, Yates J, Cotman C, Glabe C. Assembly and aggregation properties of synthetic Alzheimer's A4/beta amyloid peptide analogs. *J. Biol. Chem.* 1992; 267:546–554. [PubMed: 1730616]
- Buxbaum JD, Liu KN, Luo Y, Slack JL, Stocking KL, Peschon JJ, Johnson RS, Castner BJ, Cerretti DP, Black RA. Evidence that tumor necrosis factor alpha converting enzyme is involved in regulated alpha-secretase cleavage of the Alzheimer amyloid protein precursor. *J. Biol. Chem.* 1998; 273:27765–27767. [PubMed: 9774383]
- Cai H, Wang Y, McCarthy D, Wen H, Borchelt DR, Price DL, Wong PC. BACE1 is the major beta-secretase for generation of Abeta peptides by neurons. *Nat. Neurosci.* 2001; 4:233–234. [PubMed: 11224536]
- Caille I, Allinquant B, Dupont E, Bouillot C, Langer A, Muller U, Prochiantz A. Soluble form of amyloid precursor protein regulates proliferation of progenitors in the adult subventricular zone. *Development.* 2004; 131:2173–2181. [PubMed: 15073156]
- Cao X, Sudhof TC. Dissection of amyloid-beta precursor protein-dependent transcriptional transactivation. *J. Biol. Chem.* 2004; 279:24601–24611. [PubMed: 15044485]
- Chantry A, Glynn P. A novel metalloproteinase originally isolated from brain myelin membranes is present in many tissues. *Biochem. J.* 1990; 268:245–248. [PubMed: 1693075]
- Chavez-Gutierrez L, Tolia A, Maes E, Li T, Wong PC, de Strooper B. Glu(332) in the Nicastrin ectodomain is essential for gamma-secretase complex maturation but not for its activity. *J. Biol. Chem.* 2008; 283:20096–20105. [PubMed: 18502756]
- Chen YZ. APP induces neuronal apoptosis through APP-BP1-mediated downregulation of beta-catenin. *Apoptosis.* 2004; 9:415–422. [PubMed: 15192323]
- Cisse MA, Sunyach C, Lefranc-Jullien S, Postina R, Vincent B, Checler F. The disintegrin ADAM9 indirectly contributes to the physiological processing of cellular prion by modulating ADAM10 activity. *J. Biol. Chem.* 2005; 280:40624–40631. [PubMed: 16236709]
- Coulson EJ, Paliga K, Beyreuther K, Masters CL. What the evolution of the amyloid protein precursor supergene family tells us about its function. *Neurochem. Int.* 2000; 36:175–184. [PubMed: 10676850]
- Crystal AS, Morais VA, Pierson TC, Pijak DS, Carlin D, Lee VM, Doms RW. Membrane topology of gamma-secretase component PEN-2. *J. Biol. Chem.* 2003; 278:20117–20123. [PubMed: 12639958]
- Dahms SO, Hoefgen S, Roeser D, Schlott B, Guhrs KH, Than ME. Structure and biochemical analysis of the heparin-induced E1 dimer of the amyloid precursor protein. *Proc. Natl. Acad. Sci. USA.* 2010; 107:5381–5386. [PubMed: 20212142]
- Dawson GR, Seabrook GR, Zheng H, Smith DW, Graham S, O'Dowd G, Bowery BJ, Boyce S, Trumbauer ME, Chen HY, Van der Ploeg L. H, Sirinathsinghji DJ. Age-related cognitive deficits, impaired long-term potentiation and reduction in synaptic marker density in mice lacking the beta-amyloid precursor protein. *Neuroscience.* 1999; 90:1–13. [PubMed: 10188929]
- Donoviel DB, Hadjantonakis AK, Ikeda M, Zheng H, Hyslop PS, Bernstein A. Mice lacking both presenilin genes exhibit early embryonic patterning defects. *Genes Dev.* 1999; 13:2801–2810. [PubMed: 10557208]

- Duff K, Eckman C, Zehr C, Yu X, Prada CM, Perez-tur J, Hutton M, Buee L, Harigaya Y, Yager D, Morgan D, Gordon MN, Holcomb L, Refolo L, Zenk B, Hardy J, Younkin S. Increased amyloid-beta42(43) in brains of mice expressing mutant presenilin 1. *Nature*. 1996; 383:710–713. [PubMed: 8878479]
- Dumanchin-Njock C, Alves da Costa C. A, Mercken L, Pradier L, Checler F. The caspase-derived C-terminal fragment of betaAPP induces caspase-independent toxicity and triggers selective increase of Abeta42 in mammalian cells. *J. Neurochem*. 2001; 78:1153–1161. [PubMed: 11553689]
- Eehalt R, Michel B, De Pietri Tonelli D, Zacchetti D, Simons K, Keller P. Splice variants of the beta-site APP-cleaving enzyme BACE1 in human brain and pancreas. *Biochem. Biophys. Res. Commun*. 2002; 293:30–37. [PubMed: 12054559]
- Esch FS, Keim PS, Beattie EC, Blacher RW, Culwell AR, Oltersdorf T, McClure D, Ward PJ. Cleavage of amyloid beta peptide during constitutive processing of its precursor. *Science*. 1990; 248:1122–1124. [PubMed: 2111583]
- Fagan R, Swindells M, Overington J, Weir M. Nicastrin, a presenilin-interacting protein, contains an aminopeptidase/transferrin receptor superfamily domain. *Trends. Biochem. Sci*. 2001; 26:213–214. [PubMed: 11295540]
- Farzan M, Schnitzler CE, Vasilieva N, Leung D, Choe H. BACE2, a beta -secretase homolog, cleaves at the beta site and within the amyloid-beta region of the amyloid-beta precursor protein. *Proc. Natl. Acad. Sci. USA*. 2000; 97:9712–9717. [PubMed: 10931940]
- Figuerola DJ, Morris JA, Ma L, Kandpal G, Chen E, Li YM, Austin CP. Presenilin-dependent gamma-secretase activity modulates neurite outgrowth. *Neurobiol. Dis*. 2002; 9:49–60. [PubMed: 11848684]
- Fiore F, Zambrano N, Minopoli G, Donini V, Duilio A, Russo T. The regions of the Fe65 protein homologous to the phosphotyrosine interaction/phosphotyrosine binding domain of Shc bind the intracellular domain of the Alzheimer's amyloid precursor protein. *J. Biol. Chem*. 1995; 270:30853–30856. [PubMed: 8537337]
- Fortna RR, Crystal AS, Morais VA, Pijak DS, Lee VM, Doms RW. Membrane topology and nicastrin-enhanced endoproteolysis of APH-1, a component of the gamma-secretase complex. *J. Biol. Chem*. 2004; 279:3685–3693. [PubMed: 14593096]
- Francis R, McGrath G, Zhang J, Ruddy DA, Sym M, Apfeld J, Nicoll M, Maxwell M, Hai B, Ellis MC, Parks AL, Xu W, Li J, Gurney M, Myers RL, Himes CS, Hiebsch R, Ruble C, Nye JS, Curtis D. *aph-1* and *pen-2* are required for Notch pathway signaling, gamma-secretase cleavage of betaAPP, and presenilin protein accumulation. *Dev. Cell*. 2002; 3:85–97. [PubMed: 12110170]
- Furukawa K, Barger SW, Blalock EM, Mattson MP. Activation of K⁺ channels and suppression of neuronal activity by secreted beta-amyloid-precursor protein. *Nature*. 1996a; 379:74–78. [PubMed: 8538744]
- Furukawa K, Sopher BL, Rydel RE, Begley JG, Pham DG, Martin GM, Fox M, Mattson MP. Increased Activity Regulating and Neuroprotective Efficacy of Secretase Derived Secreted Amyloid Precursor Protein Conferred by a C Terminal Heparin Binding Domain. *J. Neurochem*. 1996b; 67:1882–1896. [PubMed: 8863493]
- Gakhar Koppole N, Hundeshagen P, Mandl C, Weyer SW, Allinquant B, Muller U, Ciccolini F. Activity requires soluble amyloid precursor protein to promote neurite outgrowth in neural stem cell derived neurons via activation of the MAPK pathway. *Eur. J. Neurosci*. 2008; 28:871–882. [PubMed: 18717733]
- Galvan V, Gorostiza OF, Banwait S, Ataie M, Logvinova AV, Sitaraman S, Carlson E, Sagi SA, Chevallier N, Jin K, Greenberg DA, Bredesen DE. Reversal of Alzheimer's-like pathology and behavior in human APP transgenic mice by mutation of Asp664. *Proc. Natl. Acad. Sci. USA*. 2006; 103:7130–7135. [PubMed: 16641106]
- Gandy S, Czernik AJ, Greengard P. Phosphorylation of Alzheimer disease amyloid precursor peptide by protein kinase C and Ca²⁺/calmodulin-dependent protein kinase II. *Proc. Natl. Acad. Sci. USA*. 1988; 85:6218–6221. [PubMed: 3137567]
- Gendron TF, Petrucelli L. The role of tau in neurodegeneration. *Mol. Neurodegener*. 2009; 4:13. [PubMed: 19284597]

- Gersbacher MT, Kim DY, Bhattacharyya R, Kovacs DM. Identification of BACE1 cleavage sites in human voltage-gated sodium channel beta 2 subunit. *Mol. Neurodegener.* 2010; 5:61. [PubMed: 21182789]
- Ghosal K, Vogt DL, Liang M, Shen Y, Lamb BT, Pimplikar SW. Alzheimer's disease-like pathological features in transgenic mice expressing the APP intracellular domain. *Proc. Natl. Acad. Sci. USA.* 2009; 106:18367–18372. [PubMed: 19837693]
- Gilberto L, Acker CM, Davies P, D'Adamio L. Transgenic Expression of the Amyloid- Precursor Protein-Intracellular Domain Does Not Induce Alzheimer's Disease-like Traits In Vivo. *PLoS One.* 2010; 5:e11609. [PubMed: 20661273]
- Gilberto L, Zhou D, Weldon R, Tamagno E, De Luca P, Tabaton M, D'Adamio L. Evidence that the Amyloid beta Precursor Protein-intracellular domain lowers the stress threshold of neurons and has a "regulated" transcriptional role. *Mol. Neurodegener.* 2008; 3:12. [PubMed: 18764939]
- Glennner GG, Wong CW. Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem. Biophys. Res. Commun.* 1984; 120:885–890. [PubMed: 6375662]
- Hamid R, Kilger E, Willem M, Vassallo N, Kostka M, Bornhovd C, Reichert AS, Kretschmar HA, Haass C, Herms J. Amyloid precursor protein intracellular domain modulates cellular calcium homeostasis and ATP content. *J. Neurochem.* 2007; 102:1264–1275. [PubMed: 17763532]
- Han P, Dou F, Li F, Zhang X, Zhang YW, Zheng H, Lipton SA, Xu H, Liao FF. Suppression of cyclin-dependent kinase 5 activation by amyloid precursor protein: a novel excitoprotective mechanism involving modulation of tau phosphorylation. *J. Neurosci.* 2005; 25:11542. [PubMed: 16354912]
- Hartmann D, de Strooper B, Serneels L, Craessaerts K, Herreman A, Annaert W, Umans L, Lubke T, Lena Illert A, von Figura K, Saftig P. The disintegrin/metalloprotease ADAM 10 is essential for Notch signalling but not for alpha-secretase activity in fibroblasts. *Hum. Mol. Genet.* 2002; 11:2615–2624. [PubMed: 12354787]
- Hass MR, Yankner BA. A {gamma}-secretase-independent mechanism of signal transduction by the amyloid precursor protein. *J. Biol. Chem.* 2005; 280:36895–36904. [PubMed: 16103124]
- Heber S, Herms J, Gajic V, Hainfellner J, Aguzzi A, Rulicke T, von Kretschmar H, von Koch C, Sisodia S, Tremml P, Lipp HP, Wolfer DP, Muller U. Mice with combined gene knock-outs reveal essential and partially redundant functions of amyloid precursor protein family members. *J. Neurosci.* 2000; 20:7951–7963. [PubMed: 11050115]
- Herms J, Anliker B, Heber S, Ring S, Fuhrmann M, Kretschmar H, Sisodia S, Muller U. Cortical dysplasia resembling human type 2 lissencephaly in mice lacking all three APP family members. *EMBO J.* 2004; 23:4106–4115. [PubMed: 15385965]
- Herzog V, Kirfel G, Siemes C, Schmitz A. Biological roles of APP in the epidermis. *Eur. J. Cell. Biol.* 2004; 83:613–624. [PubMed: 15679106]
- Hitt DB, Jaramillo CT, Chetkoyich MD, Vassar R. BACE1 mice exhibit seizure activity that does not correlate with sodium channel level or axonal localization. *Mol. Neurodegener.* 2010; 5:31. [PubMed: 20731874]
- Ho A, Sudhof TC. Binding of F-spondin to amyloid-beta precursor protein: a candidate amyloid-beta precursor protein ligand that modulates amyloid-beta precursor protein cleavage. *Proc. Natl. Acad. Sci. USA.* 2004; 101:2548. [PubMed: 14983046]
- Holsinger RM, McLean CA, Beyreuther K, Masters CL, Evin G. Increased expression of the amyloid precursor beta-secretase in Alzheimer's disease. *Ann. Neurol.* 2002; 51:783–786. [PubMed: 12112088]
- Hook V, Toneff T, Bogoy M, Greenbaum D, Medzihradzky KF, Neveu J, Lane W, Hook G, Reisine T. Inhibition of cathepsin B reduces beta-amyloid production in regulated secretory vesicles of neuronal chromaffin cells: evidence for cathepsin B as a candidate beta-secretase of Alzheimer's disease. *Biol. Chem.* 2005; 386:931–940. [PubMed: 16164418]
- Howard L, Lu X, Mitchell S, Griffiths S, Glynn P. Molecular cloning of MADM: a catalytically active mammalian disintegrin-metalloprotease expressed in various cell types. *Biochem. J.* 1996; 317(Pt 1):45–50. [PubMed: 8694785]

- Howell BW, Lanier LM, Frank R, Gertler FB, Cooper JA. The disabled 1 phosphotyrosine-binding domain binds to the internalization signals of transmembrane glycoproteins and to phospholipids. *Mol. Cell. Biol.* 1999; 19:5179–5188. [PubMed: 10373567]
- Hu X, Hicks CW, He W, Wong P, Macklin WB, Trapp BD, Yan R. Bace1 modulates myelination in the central and peripheral nervous system. *Nat. Neurosci.* 2006; 9:1520–1525. [PubMed: 17099708]
- Hussain I, Powell DJ, Howlett DR, Chapman GA, Gilmour L, Murdock PR, Tew DG, Meek TD, Chapman C, Schneider K, Ratcliffe SJ, Tattersall D, Testa TT, Southan C, Ryan DM, Simmons DL, Walsh FS, Dingwall C, Christie G. ASP1 (BACE2) cleaves the amyloid precursor protein at the beta-secretase site. *Mol. Cell. Neurosci.* 2000; 16:609–619. [PubMed: 11083922]
- Iijima K, Ando K, Takeda S, Satoh Y, Seki T, Itohara S, Greengard P, Kirino Y, Nairn AC, Suzuki T. Neuron-specific phosphorylation of Alzheimer's beta-amyloid precursor protein by cyclin-dependent kinase 5. *J. Neurochem.* 2000; 75:1085–1091. [PubMed: 10936190]
- Ikura T, Ogryzko VV, Grigoriev M, Groisman R, Wang J, Horikoshi M, Scully R, Qin J, Nakatani Y. Involvement of the TIP60 histone acetylase complex in DNA repair and apoptosis. *Cell.* 2000; 102:463–473. [PubMed: 10966108]
- Inomata H, Nakamura Y, Hayakawa A, Takata H, Suzuki T, Miyazawa K, Kitamura N. A scaffold protein JIP-1b enhances amyloid precursor protein phosphorylation by JNK and its association with kinesin light chain 1. *J. Biol. Chem.* 2003; 278:22946–22955. [PubMed: 12665528]
- Iwatsubo T, Odaka A, Suzuki N, Mizusawa H, Nukina N, Ihara Y. Visualization of A beta 42(43) and A beta 40 in senile plaques with end-specific A beta monoclonals: evidence that an initially deposited species is A beta 42(43). *Neuron.* 1994; 13:45–53. [PubMed: 8043280]
- Jarrett JT, Berger EP, Lansbury PT Jr. The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry.* 1993; 32:4693–4697. [PubMed: 8490014]
- Kaether C, Capell A, Edbauer D, Winkler E, Novak B, Steiner H, Haass C. The presenilin C-terminus is required for ER-retention, nicastrin-binding and gamma-secretase activity. *EMBO J.* 2004; 23:4738–4748. [PubMed: 15549135]
- Kamal A, Stokin GB, Yang Z, Xia CH, Goldstein LS. Axonal transport of amyloid precursor protein is mediated by direct binding to the kinesin light chain subunit of kinesin-I. *Neuron.* 2000; 28:449–459. [PubMed: 11144355]
- Kang J, Muller-Hill B. Differential splicing of Alzheimer's disease amyloid A4 precursor RNA in rat tissues: PreA4(695) mRNA is predominantly produced in rat and human brain. *Biochem. Biophys. Res. Commun.* 1990; 166:1192–1200. [PubMed: 1689572]
- Kang J, Lemaire HG, Unterbeck A, Salbaum JM, Masters CL, Grzeschik KH, Multhaup G, Beyreuther K, Muller-Hill B. The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature.* 1987; 325:733–736. [PubMed: 2881207]
- Kim HS, Kim EM, Lee JP, Park CH, Kim S, Seo JH, Chang KA, Yu E, Jeong SJ, Chong YH. C-terminal fragments of amyloid precursor protein exert neurotoxicity by inducing glycogen synthase kinase-3 beta expression. *FASEB J.* 2003; 17:1951. [PubMed: 12923068]
- Koike H, Tomioka S, Sorimachi H, Saido TC, Maruyama K, Okuyama A, Fujisawa-Sehara A, Ohno S, Suzuki K, Ishiura S. Membrane-anchored metalloprotease MDC9 has an alpha-secretase activity responsible for processing the amyloid precursor protein. *Biochem. J.* 1999; 343(Pt 2):371–375. [PubMed: 10510302]
- Kojro E, Gimpl G, Lammich S, Marz W, Fahrenholz F. Low cholesterol stimulates the nonamyloidogenic pathway by its effect on the alpha-secretase ADAM 10. *Proc. Natl. Acad. Sci. USA.* 2001; 98:5815–5820. [PubMed: 11309494]
- Kopan R, Schroeter EH, Weintraub H, Nye JS. Signal transduction by activated mNotch: importance of proteolytic processing and its regulation by the extracellular domain. *Proc. Natl. Acad. Sci. USA.* 1996; 93:1683–1688. [PubMed: 8643690]
- Kuhn PH, Wang H, Dislich B, Colombo A, Zeitschel U, Ellwart JW, Kremmer E, Rossner S, Lichtenthaler SF. ADAM10 is the physiologically relevant, constitutive alpha-secretase of the amyloid precursor protein in primary neurons. *EMBO J.* 2010; 29:3020–3032. [PubMed: 20676056]

- Laird FM, Cai H, Savonenko AV, Farah MH, He K, Melnikova T, Wen H, Chiang HC, Xu G, Koliatsos VE, Borchelt DR, Price DL, Lee HK, Wong PC. BACE1, a major determinant of selective vulnerability of the brain to amyloid-beta amyloidogenesis, is essential for cognitive, emotional, and synaptic functions. *J. Neurosci.* 2005; 25:11693–11709. [PubMed: 16354928]
- Lammich S, Kojro E, Postina R, Gilbert S, Pfeiffer R, Jasionowski M, Haass C, Fahrenholz F. Constitutive and regulated alpha-secretase cleavage of Alzheimer's amyloid precursor protein by a disintegrin metalloprotease. *Proc. Natl. Acad. Sci. USA.* 1999; 96:3922–3927. [PubMed: 10097139]
- LaVoie MJ, Fraering PC, Ostaszewski BL, Ye W, Kimberly WT, Wolfe MS, Selkoe DJ. Assembly of the gamma-secretase complex involves early formation of an intermediate subcomplex of Aph-1 and nicastrin. *J. Biol. Chem.* 2003; 278:37213–37222. [PubMed: 12857757]
- Lee DC, Sunnarborg SW, Hinkle CL, Myers TJ, Stevenson MY, Russell WE, Castner BJ, Gerhart MJ, Paxton RJ, Black RA, Chang A, Jackson LF. TACE/ADAM17 processing of EGFR ligands indicates a role as a physiological convertase. *Ann. N. Y. Acad. Sci.* 2003; 995:22–38. [PubMed: 12814936]
- Lee SF, Shah S, Li H, Yu C, Han W, Yu G. Mammalian APH-1 interacts with presenilin and nicastrin and is required for intramembrane proteolysis of amyloid-beta precursor protein and Notch. *J. Biol. Chem.* 2002; 277:45013–45019. [PubMed: 12297508]
- Levy-Lahad E, Wasco W, Poorkaj P, Romano DM, Oshima J, Pettingell WH, Yu CE, Jondro PD, Schmidt SD, Wang K, et al. Candidate gene for the chromosome 1 familial Alzheimer's disease locus. *Science.* 1995; 269:973–977. [PubMed: 7638622]
- Li T, Ma G, Cai H, Price DL, Wong PC. Nicastrin is required for assembly of presenilin/gamma-secretase complexes to mediate Notch signaling and for processing and trafficking of beta-amyloid precursor protein in mammals. *J. Neurosci.* 2003; 23:3272–3277. [PubMed: 12716934]
- Lin X, Koelsch G, Wu S, Downs D, Dashti A, Tang J. Human aspartic protease memapsin 2 cleaves the beta-secretase site of beta-amyloid precursor protein. *Proc. Natl. Acad. Sci. USA.* 2000; 97:1456–1460. [PubMed: 10677483]
- Liu Q, Zerbinatti CV, Zhang J, Hoe HS, Wang B, Cole SL, Herz J, Muglia L, Bu G. Amyloid precursor protein regulates brain apolipoprotein E and cholesterol metabolism through lipoprotein receptor LRP1. *Neuron.* 2007; 56:66–78. [PubMed: 17920016]
- Liu Y, Zhang Y, Wang X, Zhang H, You X, Liao FF, Xu H. Intracellular trafficking of presenilin 1 is regulated by -amyloid precursor protein and phospholipase D1. *J. Biol. Chem.* 2009; 284:12145. [PubMed: 19276086]
- Lorenzo A, Yuan M, Zhang Z, Paganetti PA, Sturchler-Pierrat C, Staufenbiel M, Mautino J, Vigo FS, Sommer B, Yankner BA. Amyloid beta interacts with the amyloid precursor protein: a potential toxic mechanism in Alzheimer's disease. *Nat. Neurosci.* 2000; 3:460–464. [PubMed: 10769385]
- Lourenco FC, Galvan V, Fombonne J, Corset V, Llambi F, Muller U, Bredesen DE, Mehlen P. Netrin-1 interacts with amyloid precursor protein and regulates amyloid-beta production. *Cell Death Differ.* 2009; 16:655–663. [PubMed: 19148186]
- Lu DC, Soriano S, Bredesen DE, Koo EH. Caspase cleavage of the amyloid precursor protein modulates amyloid protein toxicity. *J. Neurochem.* 2003a; 87:733–741. [PubMed: 14535955]
- Lu DC, Shaked GM, Masliah E, Bredesen DE, Koo EH. Amyloid protein toxicity mediated by the formation of amyloid protein precursor complexes. *Ann. Neurol.* 2003b; 54:781–789. [PubMed: 14681887]
- Lu DC, Rabizadeh S, Chandra S, Shayya RE, Ellerby LM, Ye X, Salvesen GS, Koo EH, Bredesen DE. A second cytotoxic proteolytic peptide derived from amyloid beta-protein precursor. *Nat. Med.* 2000; 6:397–404. [PubMed: 10742146]
- Luo WJ, Wang H, Li H, Kim BS, Shah S, Lee HJ, Thinakaran G, Kim TW, Yu G, Xu H. PEN-2 and APH-1 coordinately regulate proteolytic processing of presenilin 1. *J. Biol. Chem.* 2003; 278:7850–7854. [PubMed: 12522139]
- Luo Y, Bolon B, Kahn S, Bennett BD, Babu-Khan S, Denis P, Fan W, Kha H, Zhang J, Gong Y, Martin L, Louis JC, Yan Q, Richards WG, Citron M, Vassar R. Mice deficient in BACE1, the Alzheimer's beta-secretase, have normal phenotype and abolished beta-amyloid generation. *Nat. Neurosci.* 2001; 4:231–232. [PubMed: 11224535]

- McLoughlin D, Cj Miller C. The intracellular cytoplasmic domain of the Alzheimer's disease amyloid precursor protein interacts with phosphotyrosine-binding domain proteins in the yeast two-hybrid system. *FEBS Lett.* 1996; 397:197–200. [PubMed: 8955346]
- Ma G, Li T, Price DL, Wong PC. APH-1a is the principal mammalian APH-1 isoform present in gamma-secretase complexes during embryonic development. *J. Neurosci.* 2005; 25:192–198. [PubMed: 15634781]
- Ma T, Zhao YB, Kwak YD, Yang Z, Thompson R, Luo Z, Xu H, Liao FF. Statin's excitoprotection is mediated by sAPP and the subsequent attenuation of calpain-induced truncation events, likely via rho-ROCK signaling. *J. Neurosci.* 2009; 29:11226. [PubMed: 19741129]
- Madeira A, Pommet JM, Prochiantz A, Allinquant B. SET protein (TAF1beta, I2PP2A) is involved in neuronal apoptosis induced by an amyloid precursor protein cytoplasmic subdomain. *FEBS J.* 2005; 19:1905–1907.
- Marquez-Sterling NR, Lo AC, Sisodia SS, Koo EH. Trafficking of cell-surface beta-amyloid precursor protein: evidence that a sorting intermediate participates in synaptic vesicle recycling. *J. Neurosci.* 1997; 17:140–151. [PubMed: 8987743]
- Masters CL, Simms G, Weinman NA, Multhaup G, McDonald BL, Beyreuther K. Amyloid plaque core protein in Alzheimer disease and Down syndrome. *Proc. Natl. Acad. Sci. USA.* 1985; 82:4245–4249. [PubMed: 3159021]
- Matsuda S, Matsuda Y, D'Adamio L. Amyloid beta protein precursor (AbetaPP), but not AbetaPP-like protein 2, is bridged to the kinesin light chain by the scaffold protein JNK-interacting protein 1. *J. Biol. Chem.* 2003; 278:38601–38606. [PubMed: 12893827]
- Matsuda S, Matsuda Y, Snapp EL, D'Adamio L. Maturation of BRI2 generates a specific inhibitor that reduces APP processing at the plasma membrane and in endocytic vesicles. *Neurobiol. Aging.* 2009; 32:1400–1408. [PubMed: 19748705]
- Matsuda S, Yasukawa T, Homma Y, Ito Y, Niikura T, Hiraki T, Hirai S, Ohno S, Kita Y, Kawasumi M, Kouyama K, Yamamoto T, Kyriakis JM, Nishimoto I. c-Jun N-terminal kinase (JNK)-interacting protein-1b/islet-brain-1 scaffolds Alzheimer's amyloid precursor protein with JNK. *J. Neurosci.* 2001; 21:6597–6607. [PubMed: 11517249]
- Mattson MP. Cellular actions of beta-amyloid precursor protein and its soluble and fibrillogenic derivatives. *Physiol. Rev.* 1997; 77:1081–1132. [PubMed: 9354812]
- Mattson MP, Cheng B, Culwell AR, Esch FS, Lieberburg I, Rydel RE. Evidence for excitoprotective and intraneuronal calcium-regulating roles for secreted forms of the beta-amyloid precursor protein. *Neuron.* 1993; 10:243–254. [PubMed: 8094963]
- Menendez-Gonzalez M, Perez-Pinera P, Martinez-Rivera M, Calatayud MT, Blazquez Menes B. APP processing and the APP-KPI domain involvement in the amyloid cascade. *Neurodegener. Dis.* 2005; 2:277–283. [PubMed: 16909010]
- Meziane H, Dodart JC, Mathis C, Little S, Clemens J, Paul SM, Ungerer A. Memory-enhancing effects of secreted forms of the beta-amyloid precursor protein in normal and amnesic mice. *Proc. Natl. Acad. Sci. USA.* 1998; 95:12683–12688. [PubMed: 9770546]
- Morley JE, Farr SA, Banks WA, Johnson SN, Yamada KA, Xu L. A physiological role for amyloid-beta protein: enhancement of learning and memory. *J. Alzheimer's Dis.* 2008; 19:441–449. [PubMed: 19749407]
- Moss ML, Jin SL, Milla ME, Bickett DM, Burkhart W, Carter HL, Chen WJ, Clay WC, Didsbury JR, Hassler D, Hoffman CR, Kost TA, Lambert MH, Leesnitzer MA, McCauley P, McGeehan G, Mitchell J, Moyer M, Pahel G, Rocque W, Overton LK, Schoenen F, Seaton T, Su JL, Becherer JD, et al. Cloning of a disintegrin metalloproteinase that processes precursor tumour-necrosis factor-alpha. *Nature.* 1997; 385:733–736. [PubMed: 9034191]
- Mueller-Steiener S, Zhou Y, Arai H, Roberson ED, Sun B, Chen J, Wang X, Yu G, Esposito L, Mucke L, Gan L. Anti-amyloidogenic and neuroprotective functions of cathepsin B: implications for Alzheimer's disease. *Neuron.* 2006; 51:703–714. [PubMed: 16982417]
- Nakaya T, Suzuki T. Role of APP phosphorylation in FE65-dependent gene transactivation mediated by AICD. *Genes. Cells.* 2006; 11:633–645. [PubMed: 16716194]
- Niimura M, Isoo N, Takasugi N, Tsuruoka M, Ui-Tei K, Saigo K, Morohashi Y, Tomita T, Iwatsubo T. Aph-1 contributes to the stabilization and trafficking of the gamma-secretase complex through

- mechanisms involving intermolecular and intramolecular interactions. *J. Biol. Chem.* 2005; 280:12967–12975. [PubMed: 15644323]
- Nikolaev A, McLaughlin T, O'Leary DD, Tessier-Lavigne M. APP binds DR6 to trigger axon pruning and neuron death via distinct caspases. *Nature.* 2009; 457:981–989. [PubMed: 19225519]
- Nishimura I, Uetsuki T, Kuwako K, Hara T, Kawakami T, Aimoto S, Yoshikawa K. Cell death induced by a caspase-cleaved transmembrane fragment of the Alzheimer amyloid precursor protein. *Cell Death Differ.* 2002; 9:199–208. [PubMed: 11840170]
- Norstrom EM, Zhang C, Tanzi R, Sisodia SS. Identification of NEEP21 as a ss-amyloid precursor protein-interacting protein in vivo that modulates amyloidogenic processing in vitro. *J. Neurosci.* 2010; 30:15677–15685. [PubMed: 21084623]
- Ohno M, Sametsky EA, Younkin LH, Oakley H, Younkin SG, Citron M, Vassar R, Disterhoft JF. BACE1 deficiency rescues memory deficits and cholinergic dysfunction in a mouse model of Alzheimer's disease. *Neuron.* 2004; 41:27–33. [PubMed: 14715132]
- Ohsawa I, Takamura C, Morimoto T, Ishiguro M, Kohsaka S. Amino-terminal region of secreted form of amyloid precursor protein stimulates proliferation of neural stem cells. *Eur. J. Neurosci.* 1999; 11:1907–1913. [PubMed: 10336659]
- Ozaki T, Li Y, Kikuchi H, Tomita T, Iwatsubo T, Nakagawara A. The intracellular domain of the amyloid precursor protein (AICD) enhances the p53-mediated apoptosis. *Biochem. Biophys. Res. Commun.* 2006; 351:57–63. [PubMed: 17054906]
- Pardossi-Piquard R, Petit A, Kawarai T, Sunyach C, Alves da Costa C, Vincent B, Ring S, D'Adamio L, Shen J, Muller U, St George Hyslop P, Checler F. Presenilin-dependent transcriptional control of the Abeta-degrading enzyme neprilysin by intracellular domains of betaAPP and APLP. *Neuron.* 2005; 46:541–554. [PubMed: 15944124]
- Park JH, Gimbel DA, GrandPre T, Lee JK, Kim JE, Li W, Lee DH, Strittmatter SM. Alzheimer precursor protein interaction with the Nogo-66 receptor reduces amyloid-beta plaque deposition. *J. Neurosci.* 2006; 26:1386–1395. [PubMed: 16452662]
- Pasternak SH, Bagshaw RD, Guiral M, Zhang S, Ackerley CA, Pak BJ, Callahan JW, Mahuran DJ. Presenilin-1, nicastrin, amyloid precursor protein, and gamma-secretase activity are co-localized in the lysosomal membrane. *J. Biol. Chem.* 2003; 278:26687–26694. [PubMed: 12736250]
- Peschon JJ, Slack JL, Reddy P, Stocking KL, Sunnarborg SW, Lee DC, Russell WE, Castner BJ, Johnson RS, Fitzner JN, Boyce RW, Nelson N, Kozlosky CJ, Wolfson MF, Rauch CT, Cerretti DP, Paxton RJ, March CJ, Black RA. An essential role for ectodomain shedding in mammalian development. *Science.* 1998; 282:1281–1284. [PubMed: 9812885]
- Plant LD, Boyle JP, Smith IF, Peers C, Pearson HA. The production of amyloid beta peptide is a critical requirement for the viability of central neurons. *J. Neurosci.* 2003; 23:5531–5535. [PubMed: 12843253]
- Postina R, Schroeder A, Dewachter I, Bohl J, Schmitt U, Kojro E, Prinzen C, Endres K, Hiemke C, Blessing M, Flamez P, Dequenne A, Godaux E, van Leuven F, Fahrenholz F. A disintegrin-metalloproteinase prevents amyloid plaque formation and hippocampal defects in an Alzheimer disease mouse model. *J. Clin. Invest.* 2004; 113:1456–1464. [PubMed: 15146243]
- Qian S, Jiang P, Guan XM, Singh G, Trumbauer ME, Yu H, Chen HY, Van de Ploeg L. H, Zheng H. Mutant human presenilin 1 protects presenilin 1 null mouse against embryonic lethality and elevates Abeta1–42/43 expression. *Neuron.* 1998; 20:611–617. [PubMed: 9539133]
- Qiang L, Juan Z, Hien T, Marcel V, Karina R, Guojun B. LRP1 shedding in human brain: roles of ADAM10 and ADAM17. *Mol. Neurodegener.* 2009; 4
- Ring S, Weyer SW, Kilian SB, Waldron E, Pietrzik CU, Filippov MA, Herms J, Buchholz C, Eckman CB, Korte M, Wolfer DP, Muller UC. The secreted beta-amyloid precursor protein ectodomain APPs alpha is sufficient to rescue the anatomical, behavioral, and electrophysiological abnormalities of APP-deficient mice. *J. Neurosci.* 2007; 27:7817–7826. [PubMed: 17634375]
- Roberds SL, Anderson J, Basi G, Bienkowski MJ, Branstetter DG, Chen KS, Freedman SB, Frigon NL, Games D, Hu K, Johnson-Wood K, Kappenman KE, Kawabe TT, Kola I, Kuehn R, Lee M, Liu W, Motter R, Nichols NF, Power M, Robertson DW, Schenk D, Schoor M, Shopp GM, Shuck ME, Sinha S, Svensson KA, Tatsuno G, Tintrup H, Wijsman J, Wright S, McConlogue L. BACE knockout mice are healthy despite lacking the primary beta-secretase activity in brain:

- implications for Alzheimer's disease therapeutics. *Hum. Mol. Genet.* 2001; 10:1317–1324. [PubMed: 11406613]
- Rohan de Silva HA, Jen A, Wickenden C, Jen LS, Wilkinson SL, Patel AJ. Cell-specific expression of beta-amyloid precursor protein isoform mRNAs and proteins in neurons and astrocytes. *Brain Res. Mol. Brain Res.* 1997; 47:147–156. [PubMed: 9221912]
- Roncarati R, Sestan N, Scheinfeld MH, Berechid BE, Lopez PA, Meucci O, McGlade JC, Rakic P, D'Adamio L. The gamma-secretase-generated intracellular domain of -amyloid precursor protein binds Numb and inhibits Notch signaling. *Proc. Natl. Acad. Sci. USA.* 2002; 99:7102. [PubMed: 12011466]
- Ryan KA, Pimplikar SW. Activation of GSK-3 and phosphorylation of CRMP2 in transgenic mice expressing APP intracellular domain. *J. Cell Biol.* 2005; 171:327–335. [PubMed: 16230462]
- Sandbrink R, Masters CL, Beyreuther K. Beta A4-amyloid protein precursor mRNA isoforms without exon 15 are ubiquitously expressed in rat tissues including brain, but not in neurons. *J. Biol. Chem.* 1994; 269:1510–1517. [PubMed: 8288617]
- Sarah C, Robert V. The Alzheimer's disease beta-secretase enzyme, BACE1. *Mol. Neurodegener.* 2007; 2
- Sastre M, Steiner H, Fuchs K, Capell A, Multhaup G, Condron MM, Teplow DB, Haass C. Presenilin-dependent gamma-secretase processing of beta-amyloid precursor protein at a site corresponding to the S3 cleavage of Notch. *EMBO Rep.* 2001; 2:835–841. [PubMed: 11520861]
- Sato T, Tanimura Y, Hirofumi N, Saido TC, Morishima-Kawashima M, Ihara Y. Blocking the cleavage at midportion between gamma- and epsilon-sites remarkably suppresses the generation of amyloid beta-protein. *FEBS Lett.* 2005; 579:2907–2912. [PubMed: 15890346]
- Sato T, Dohmae N, Qi Y, Kakuda N, Misonou H, Mitsumori R, Maruyama H, Koo EH, Haass C, Takio K, Morishima-Kawashima M, Ishiura S, Ihara Y. Potential link between amyloid beta-protein 42 and C-terminal fragment gamma 49–99 of beta-amyloid precursor protein. *J. Biol. Chem.* 2003; 278:24294–24301. [PubMed: 12707272]
- Saura CA, Chen G, Malkani S, Choi SY, Takahashi RH, Zhang D, Gouras GK, Kirkwood A, Morris RG, Shen J. Conditional inactivation of presenilin 1 prevents amyloid accumulation and temporarily rescues contextual and spatial working memory impairments in amyloid precursor protein transgenic mice. *J. Neurosci.* 2005; 25:6755–6764. [PubMed: 16033885]
- Schechter I, Ziv E. Kinetic properties of cathepsin D and BACE 1 indicate the need to search for additional beta-secretase candidate(s). *Biol. Chem.* 2008; 389:313–320. [PubMed: 18177262]
- Scheinfeld MH, Roncarati R, Vito P, Lopez PA, Abdallah M, D'Adamio L. Jun NH2-terminal kinase (JNK) interacting protein 1 (JIP1) binds the cytoplasmic domain of the Alzheimer's -amyloid precursor protein (APP). *J. Biol. Chem.* 2002; 277:3767. [PubMed: 11724784]
- Scheuner D, Eckman C, Jensen M, Song X, Citron M, Suzuki N, Bird TD, Hardy J, Hutton M, Kukull W, Larson E, Levy-Lahad E, Viitanen M, Peskind E, Poorkaj P, Schellenberg G, Tanzi R, Wasco W, Lannfelt L, Selkoe D, Younkin S. Secreted amyloid beta-protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease. *Nat. Med.* 1996; 2:864–870. [PubMed: 8705854]
- Schroeter EH, Kisslinger JA, Kopan R. Notch-1 signalling requires ligand-induced proteolytic release of intracellular domain. *Nature.* 1998; 393:382–386. [PubMed: 9620803]
- Selkoe DJ. The cell biology of beta-amyloid precursor protein and presenilin in Alzheimer's disease. *Trends Cell Biol.* 1998; 8:447–453. [PubMed: 9854312]
- Serneels L, DeJaegere T, Craessaerts K, Horre K, Jorissen E, Tousseyn T, Hebert S, Coolen M, Martens G, Zwijsen A, Annaert W, Hartmann D, De Strooper B. Differential contribution of the three Aph1 genes to gamma-secretase activity in vivo. *Proc. Natl. Acad. Sci. USA.* 2005; 102:1719–1724. [PubMed: 15665098]
- Serneels L, Van Biervliet J, Craessaerts K, DeJaegere T, Horre K, Van Houtvin T, Esselmann H, Paul S, Schafer MK, Berezovska O, Hyman BT, Sprangers B, Sciot R, Moons L, Jucker M, Yang Z, May PC, Karran E, Wiltfang J, D'Hooge R, De Strooper B. gamma-Secretase heterogeneity in the Aph1 subunit: relevance for Alzheimer's disease. *Science.* 2009; 324:639–642. [PubMed: 19299585]

- Shah S, Lee SF, Tabuchi K, Hao YH, Yu C, LaPlant Q, Ball H, Dann CE, 3rd Sudhof T, Yu G. Nicastrin functions as a gamma-secretase-substrate receptor. *Cell*. 2005; 122:435–447. [PubMed: 16096062]
- Shankar GM, Walsh DM. Alzheimer's disease: synaptic dysfunction and A β . *Mol. Neurodegener*. 2009; 4:48. [PubMed: 19930651]
- Shen J, Bronson RT, Chen DF, Xia W, Selkoe DJ, Tonegawa S. Skeletal and CNS defects in Presenilin-1-deficient mice. *Cell*. 1997; 89:629–639. [PubMed: 9160754]
- Sherrington R, Rogaev EI, Liang Y, Rogaeva EA, Levesque G, Ikeda M, Chi H, Lin C, Li G, Holman K, Tsuda T, Mar L, Foncin JF, Bruni AC, Montesi MP, Sorbi S, Rainero I, Pinessi L, Nee L, Chumakov I, Pollen D, Brookes A, Sanseau P, Polinsky RJ, Wasco W, Da Silva H. A, Haines JL, Pericak-Vance MA, Tanzi RE, Roses AD, Fraser PE, Rommens JM, St George-Hyslop P. H. Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature*. 1995; 375:754–760. [PubMed: 7596406]
- Shiraishi H, Sai X, Wang HQ, Maeda Y, Kurono Y, Nishimura M, Yanagisawa K, Komano H. PEN-2 enhances gamma-cleavage after presenilin heterodimer formation. *J. Neurochem*. 2004; 90:1402–1413. [PubMed: 15341524]
- Shirovani K, Edbauer D, Kostka M, Steiner H, Haass C. Immature nicastrin stabilizes APH 1 independent of PEN 2 and presenilin: identification of nicastrin mutants that selectively interact with APH 1. *J. Neurochem*. 2004; 89:1520–1527. [PubMed: 15189355]
- Shirovani K, Edbauer D, Capell A, Schmitz J, Steiner H, Haass C. Gamma-Secretase activity is associated with a conformational change of nicastrin. *J. Biol. Chem*. 2003; 278:16474. [PubMed: 12644462]
- Siemes C, Quast T, Kummer C, Wehner S, Kirfel G, Muller U, Herzog V. Keratinocytes from APP/APLP2-deficient mice are impaired in proliferation, adhesion and migration in vitro. *Exp. Cell Res*. 2006; 312:1939–1949. [PubMed: 16584729]
- Sinha S, Anderson JP, Barbour R, Basi GS, Caccavello R, Davis D, Doan M, Dovey HF, Frigon N, Hong J, Jacobson-Croak K, Jewett N, Keim P, Knops J, Lieberburg I, Power M, Tan H, Tatsuno G, Tung J, Schenk D, Seubert P, Suomensaaari SM, Wang S, Walker D, Zhao J, McConlogue L, John V. Purification and cloning of amyloid precursor protein beta-secretase from human brain. *Nature*. 1999; 402:537–540. [PubMed: 10591214]
- Sjolander A, Zetterberg H, Andreasson U, Minthon L, Blennow K. Gene variants do not influence BACE1 activity, levels of APP or A β isoforms in CSF in Alzheimer's disease. *Mol. Neurodegener*. 2010; 5:37. [PubMed: 20849576]
- Small DH, Clarris HL, Williamson TG, Reed G, Key B, Mok SS, Beyreuther K, Masters CL, Nurcombe V. Neurite-outgrowth regulating functions of the amyloid protein precursor of Alzheimer's disease. *J. Alzheimers. Dis*. 1999; 1:275–285. [PubMed: 12214125]
- Soba P, Eggert S, Wagner K, Zentgraf H, Siehl K, Kreger S, Lower A, Langer A, Merdes G, Paro R, Masters CL, Muller U, Kins S, Beyreuther K. Homo- and heterodimerization of APP family members promotes intercellular adhesion. *EMBO J*. 2005; 24:3624–3634. [PubMed: 16193067]
- Solans A, Estivill X, de La Luna S. A new aspartyl protease on 21q22.3, BACE2, is highly similar to Alzheimer's amyloid precursor protein beta-secretase. *Cytogenet. Cell Genet*. 2000; 89:177–184. [PubMed: 10965118]
- Steiner H, Winkler E, Edbauer D, Prokop S, Basset G, Yamasaki A, Kostka M, Haass C. PEN-2 is an integral component of the gamma-secretase complex required for coordinated expression of presenilin and nicastrin. *J. Biol. Chem*. 2002; 277:39062–39065. [PubMed: 12198112]
- Sumioka A, Nagaishi S, Yoshida T, Lin A, Miura M, Suzuki T. Role of 14-3-3gamma in FE65-dependent gene transactivation mediated by the amyloid beta-protein precursor cytoplasmic fragment. *J. Biol. Chem*. 2005; 280:42364–42374. [PubMed: 16223726]
- Suzuki T, Oishi M, Marshak DR, Czernik AJ, Nairn AC, Greengard P. Cell cycle-dependent regulation of the phosphorylation and metabolism of the Alzheimer amyloid precursor protein. *EMBO J*. 1994; 13:1114. [PubMed: 8131745]
- Tabuchi K, Chen G, Sudhof TC, Shen J. Conditional forebrain inactivation of nicastrin causes progressive memory impairment and age-related neurodegeneration. *J. Neurosci*. 2009; 29:7290–7301. [PubMed: 19494151]

- Takasugi N, Tomita T, Hayashi I, Tsuruoka M, Niimura M, Takahashi Y, Thinakaran G, Iwatsubo T. The role of presenilin cofactors in the gamma-secretase complex. *Nature*. 2003; 422:438–441. [PubMed: 12660785]
- Tanahashi H, Tabira T. Three novel alternatively spliced isoforms of the human beta-site amyloid precursor protein cleaving enzyme (BACE) and their effect on amyloid beta-peptide production. *Neurosci. Lett*. 2001; 307:9–12. [PubMed: 11516562]
- Tanzi RE, Gusella JF, Watkins PC, Bruns GA, St George-Hyslop P, Van Keuren ML, Patterson D, Pagan S, Kurnit DM, Neve RL. Amyloid beta protein gene: cDNA, mRNA distribution, and genetic linkage near the Alzheimer locus. *Science*. 1987; 235:880–884. [PubMed: 2949367]
- Tarr PE, Roncarati R, Pelicci G, Pelicci PG, D'Adamio L. Tyrosine phosphorylation of the beta-amyloid precursor protein cytoplasmic tail promotes interaction with Shc. *J. Biol. Chem*. 2002; 277:16798–16804. [PubMed: 11877420]
- Taru H, Kirino Y, Suzuki T. Differential roles of JIP scaffold proteins in the modulation of amyloid precursor protein metabolism. *J. Biol. Chem*. 2002; 277:27567–27574. [PubMed: 12023290]
- Taylor CJ, Ireland DR, Ballagh I, Bourne K, Marechal NM, Turner PR, Bilkey DK, Tate WP, Abraham WC. Endogenous secreted amyloid precursor protein-alpha regulates hippocampal NMDA receptor function, long-term potentiation and spatial memory. *Neurobiol. Dis*. 2008; 31:250–260. [PubMed: 18585048]
- Thinakaran G, Borchelt DR, Lee MK, Slunt HH, Spitzer L, Kim G, Ratovitsky T, Davenport F, Nordstedt C, Seeger M. Endoproteolysis of presenilin 1 and accumulation of processed derivatives in vivo. *Neuron*. 1996; 17:181–190. [PubMed: 8755489]
- Tian Y, Bassit B, Chau D, Li YM. An APP inhibitory domain containing the Flemish mutation residue modulates gamma-secretase activity for Aβ production. *Nat. Struct. Mol. Biol*. 2010; 17:151–158. [PubMed: 20062056]
- Uemura K, Farmer KC, Nasser-Ghods N, Jones P, Berezovska O. Reciprocal relationship between APP positioning relative to the membrane and PS1 conformation. *Mol. Neurodegener*. 2011; 6:15. [PubMed: 21310068]
- Vassar R, Bennett BD, Babu-Khan S, Kahn S, Mendiola EA, Denis P, Teplow DB, Ross S, Amarante P, Loeloff R, Luo Y, Fisher S, Fuller J, Edenson S, Lile J, Jarosinski MA, Biere AL, Curran E, Burgess T, Louis JC, Collins F, Treanor J, Rogers G, Citron M. Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science*. 1999; 286:735–741. [PubMed: 10531052]
- Vetrivel KS, Zhang YW, Xu H, Thinakaran G. Pathological and physiological functions of presenilins. *Mol. Neurodegener*. 2006; 1:4. [PubMed: 16930451]
- Vincent B, Paitel E, Frobert Y, Lehmann S, Grassi J, Checler F. Phorbol ester-regulated cleavage of normal prion protein in HEK293 human cells and murine neurons. *J. Biol. Chem*. 2000; 275:35612–35616. [PubMed: 10952979]
- Vincent B, Paitel E, Saftig P, Frobert Y, Hartmann D, De Strooper B, Grassi J, Lopez-Perez E, Checler F. The disintegrins ADAM10 and TACE contribute to the constitutive and phorbol ester-regulated normal cleavage of the cellular prion protein. *J. Biol. Chem*. 2001; 276:37743–37746. [PubMed: 11477090]
- Von Koch CS, Zheng H, Chen H, Trumbauer M, Thinakaran G, van der Ploeg L. H, Price DL, Sisodia SS. Generation of APLP2 KO mice and early postnatal lethality in APLP2/APP double KO mice. *Neurobiol. Aging*. 1997; 18:661–669. [PubMed: 9461064]
- Wang B, Yang L, Wang Z, Zheng H. Amyloid precursor protein mediates presynaptic localization and activity of the high-affinity choline transporter. *Proc. Natl. Acad. Sci. USA*. 2007; 104:14140. [PubMed: 17709753]
- Wang P, Yang G, Mosier DR, Chang P, Zaidi T, Gong YD, Zhao NM, Dominguez B, Lee KF, Gan WB. Defective neuromuscular synapses in mice lacking amyloid precursor protein (APP) and APP-Like protein 2. *J. Neurosci*. 2005; 25:1219. [PubMed: 15689559]
- Wang Y, Ha Y. The X-ray structure of an antiparallel dimer of the human amyloid precursor protein E2 domain. *Mol. Cell*. 2004; 15:343–353. [PubMed: 15304215]

- Wang Z, Wang B, Yang L, Guo Q, Aithmitti N, Songyang Z, Zheng H. Presynaptic and postsynaptic interaction of the amyloid precursor protein promotes peripheral and central synaptogenesis. *J. Neurosci.* 2009; 29:10788. [PubMed: 19726636]
- Wasco W, Bupp K, Magendantz M, Gusella JF, Tanzi RE, Solomon F. Identification of a mouse brain cDNA that encodes a protein related to the Alzheimer disease-associated amyloid beta protein precursor. *Proc. Natl. Acad. Sci. USA.* 1992; 89:10758–10762. [PubMed: 1279693]
- Wasco W, Gurubhagavatula S, Paradis MD, Romano DM, Sisodia SS, Hyman BT, Neve RL, Tanzi RE. Isolation and characterization of APLP2 encoding a homologue of the Alzheimer's associated amyloid beta protein precursor. *Nat. Genet.* 1993; 5:95–100. [PubMed: 8220435]
- Weidemann A, Eggert S, Reinhard FB, Vogel M, Paliga K, Baier G, Masters CL, Beyreuther K, Evin G. A novel epsilon-cleavage within the transmembrane domain of the Alzheimer amyloid precursor protein demonstrates homology with Notch processing. *Biochemistry.* 2002; 41:2825–2835. [PubMed: 11851430]
- Weskamp G, Cai H, Brodie TA, Higashiyama S, Manova K, Ludwig T, Blobel CP. Mice lacking the metalloprotease-disintegrin MDC9 (ADAM9) have no evident major abnormalities during development or adult life. *Mol. Cell. Biol.* 2002; 22:1537–1544. [PubMed: 11839819]
- Willem M, Garratt AN, Novak B, Citron M, Kaufmann S, Rittger A, DeStrooper B, Saftig P, Birchmeier C, Haass C. Control of peripheral nerve myelination by the beta-secretase BACE1. *Science.* 2006; 314:664–666. [PubMed: 16990514]
- Wolfe MS, Xia W, Ostaszewski BL, Diehl TS, Kimberly WT, Selkoe DJ. Two transmembrane aspartates in presenilin-1 required for presenilin endoproteolysis and gamma-secretase activity. *Nature.* 1999; 398:513–517. [PubMed: 10206644]
- Xin H, Stephans JC, Duan X, Harrowe G, Kim E, Grieshammer U, Kingsley C, Giese K. Identification of a novel aspartic-like protease differentially expressed in human breast cancer cell lines. *Biochim. Biophys. Acta.* 2000; 1501:125–137. [PubMed: 10838186]
- Xu Y, Kim HS, Joo Y, Choi Y, Chang KA, Park CH, Shin KY, Kim S, Cheon YH, Baik TK. Intracellular domains of amyloid precursor-like protein 2 interact with CP2 transcription factor in the nucleus and induce glycogen synthase kinase-3 expression. *Cell Death Differ.* 2006; 14:79–91. [PubMed: 16645641]
- Yan R, Bienkowski MJ, Shuck ME, Miao H, Tory MC, Pauley AM, Brashier JR, Stratman NC, Mathews WR, Buhl AE, Carter DB, Tomasselli AG, Parodi LA, Heinrikson RL, Gurney ME. Membrane-anchored aspartyl protease with Alzheimer's disease beta-secretase activity. *Nature.* 1999; 402:533–537. [PubMed: 10591213]
- Yang G, Gong YD, Gong K, Jiang WL, Kwon E, Wang P, Zheng H, Zhang XF, Gan WB, Zhao NM. Reduced synaptic vesicle density and active zone size in mice lacking amyloid precursor protein (APP) and APP-like protein 2. *Neurosci. Lett.* 2005; 384:66–71. [PubMed: 15919150]
- Yang LB, Lindholm K, Yan R, Citron M, Xia W, Yang XL, Beach T, Sue L, Wong P, Price D, Li R, Shen Y. Elevated beta-secretase expression and enzymatic activity detected in sporadic Alzheimer disease. *Nat. Med.* 2003; 9:3–4. [PubMed: 12514700]
- Yoshikai S, Sasaki H, Doh-ura K, Furuya H, Sakaki Y. Genomic organization of the human amyloid beta-protein precursor gene. *Gene.* 1990; 87:257–263. [PubMed: 2110105]
- Young-Pearse TL, Bai J, Chang R, Zheng JB, LoTurco JJ, Selkoe DJ. A critical function for beta-amyloid precursor protein in neuronal migration revealed by in utero RNA interference. *J. Neurosci.* 2007; 27:14459–14469. [PubMed: 18160654]
- Yu C, Kim SH, Ikeuchi T, Xu H, Gasparini L, Wang R, Sisodia SS. Characterization of a presenilin-mediated amyloid precursor protein carboxyl-terminal fragment gamma. Evidence for distinct mechanisms involved in gamma-secretase processing of the APP and Notch1 transmembrane domains. *J. Biol. Chem.* 2001; 276:43756–43760. [PubMed: 11583985]
- Yu G, Nishimura M, Arawaka S, Levitan D, Zhang L, Tandon A, Song YQ, Rogaeva E, Chen F, Kawarai T. Nicastrin modulates presenilin-mediated notch/glp-1 signal transduction and beta-APP processing. *Nature.* 2000; 407:48–54. [PubMed: 10993067]
- Zhang YW, Xu H. Substrate check of gamma-secretase. *Nat. Struct. Mol. Biol.* 2010; 17:140–141. [PubMed: 20130572]

- Zhang YW, Wang R, Liu Q, Zhang H, Liao FF, Xu H. Presenilin/gamma-secretase-dependent processing of beta-amyloid precursor protein regulates EGF receptor expression. *Proc. Natl. Acad. Sci. USA.* 2007; 104:10613–10618. [PubMed: 17556541]
- Zhang YW, Luo WJ, Wang H, Lin P, Vetrivel KS, Liao F, Li F, Wong PC, Farquhar MG, Thinakaran G, Xu H. Nicastrin is critical for stability and trafficking but not association of other presenilin/gamma-secretase components. *J. Biol. Chem.* 2005; 280:17020–17026. [PubMed: 15711015]
- Zhao G, Mao G, Tan J, Dong Y, Cui MZ, Kim SH, Xu X. Identification of a new presenilin-dependent zeta-cleavage site within the transmembrane domain of amyloid precursor protein. *J. Biol. Chem.* 2004; 279:50647–50650. [PubMed: 15485850]
- Zhao G, Cui MZ, Mao G, Dong Y, Tan J, Sun L, Xu X. gamma-Cleavage is dependent on zeta-cleavage during the proteolytic processing of amyloid precursor protein within its transmembrane domain. *J. Biol. Chem.* 2005; 280:37689. [PubMed: 16157587]
- Zheng H, Koo EH. Biology and pathophysiology of the amyloid precursor protein. *Mol. Neurodegener.* 2011; 6:27. [PubMed: 21527012]
- Zheng H, Jiang M, Trumbauer ME, Sirinathsingji DJ, Hopkins R, Smith DW, Heavens RP, Dawson GR, Boyce S, Conner MW, Stevens KA, Slunt HH, Sisoda SS, Chen HY, Van der Ploeg L. H. beta-Amyloid precursor protein-deficient mice show reactive gliosis and decreased locomotor activity. *Cell.* 1995; 81:525–531. [PubMed: 7758106]
- Zheng P, Eastman J, Vande Pol S, Pimplikar SW. PAT1, a microtubule-interacting protein, recognizes the basolateral sorting signal of amyloid precursor protein. *Proc. Natl. Acad. Sci. USA.* 1998; 95:14745. [PubMed: 9843960]
- Zou K, Kim D, Kakio A, Byun K, Gong JS, Kim J, Kim M, Sawamura N, Nishimoto S, Matsuzaki K, Lee B, Yanagisawa K, Michikawa M. Amyloid beta-protein (Abeta)1–40 protects neurons from damage induced by Abeta1–42 in culture and in rat brain. *J. Neurochem.* 2003; 87:609–619. [PubMed: 14535944]




Fig. 1.
Proteolytic processing of APP.