

α 1,3 Glucans Are Dispensable in *Aspergillus fumigatus*

Christine Henry, Jean-Paul Latgé, and Anne Beauvais

Unité des Aspergillus, Institut Pasteur, Paris, France

A triple α 1,3 glucan synthase mutant of *Aspergillus fumigatus* obtained by successive deletions of the three α 1,3 glucan synthase genes (*AGS1*, *AGS2*, and *AGS3*) has a cell wall devoid of α 1,3 glucans. The lack of α 1,3 glucans affects neither conidial germination nor mycelial vegetative growth and is compensated by an increase in β 1,3 glucan and/or chitin content.

In *Aspergillus fumigatus*, α 1,3 glucans are a major amorphous cell wall polysaccharide, accounting for 35 to 40% of the mycelial cell wall and 20 to 25% of the conidial cell wall (21). α 1,3 glucans are also a major cell wall component of the yeast form of the human pathogens *Paracoccidioides brasiliensis*, *Histoplasma capsulatum*, *Blastomyces dermatitidis*, and *Cryptococcus neoformans* and of the nonpathogenic model yeast *Schizosaccharomyces pombe* (14, 15, 24, 25, 27). In these species, the concentrations reported vary from 28% in *S. pombe* to 35 to 46% in the virulent yeast forms (13). In all pathogenic species studied to date, α 1,3 glucans are important for virulence. In addition, in *S. pombe*, α 1,3 glucans are essential for fungal viability. For *A. fumigatus*, it was shown that α 1,3 glucans have a major adhesive role in the interactions between hyphae or germinating conidia (3, 8).

α 1,3 glucans are synthesized by α 1,3 glucan synthases, which are transmembrane enzymes with very high molecular masses (>200 kDa). The number of genes coding for Ags proteins varies between fungi. Only one *AGS* gene was found in *H. capsulatum* and *C. neoformans*, whereas five *AGS* genes were identified in *S. pombe* (14, 24, 25). In *A. fumigatus*, α 1,3 glucans are synthesized by three α 1,3 glucan synthases (Ags1p [AFUA_3G00910], Ags2p [AFUA_2G11270], and Ags3p [AFUA_1G15440]) (21). All three *A. fumigatus* *AGS* genes are expressed constitutively (21). Three single *AGS* mutants have been constructed in *A. fumigatus*. The *ags1* Δ mutant had a 50% reduction in the cell wall α 1,3 glucan content of the mycelium (2). In spite of this cell wall defect, deletion of *AGS1* did not reduce the virulence of the strain in an experimental murine model of invasive aspergillosis (2). In contrast to the case for the *ags1* Δ mutant, the mycelial cell walls of *ags2* Δ and *ags3* Δ mutants had α 1,3 glucan levels similar to that of the parental strain. In addition, no modification of the α 1,3 glucan content of the conidial cell wall was seen in all single *ags* Δ mutants. Compensatory expression of the other members of the *AGS* family, which has been seen in all single *ags* Δ mutants, could explain the lack of a significant phenotype for each single *ags* Δ mutant (2, 21). For example, *AGS3* and *AGS2* showed increased expression in the *ags1* Δ mutant, and *AGS1* expression compensated for the lack of *AGS3* in the *ags3* Δ mutant (21).

Understanding the functional role of α 1,3 glucans in *A. fumigatus* required the construction of a triple *ags1* Δ *ags2* Δ *ags3* Δ mutant devoid of α 1,3 glucans. We report here the construction and growth phenotype of this triple mutant. As expected for the triple *AGS* deletion, the cell wall of the *ags1* Δ *ags2* Δ *ags3* Δ mutant did not contain any α 1,3 glucans, but surprisingly, the mutant did not show any reduction in fungal viability and growth *in vitro*.

MATERIALS AND METHODS

Strains, plasmids, and culture conditions. The *AkuB*^{ku80} *pyrG* Δ strain was used as the parental strain (6). The *AkuB*^{ku80} *pyrG* Δ strain and mutants were maintained on 2% malt agar slants. Conidia were produced on 2% malt agar slants for 5 days at 37°C and recovered by vortexing with 0.05% (vol/vol) Tween 20 aqueous solution. For DNA extraction, mycelium was grown for 16 h at 37°C in a Sabouraud liquid medium supplemented with 10 mM uracil. For transformation experiments, complete medium supplemented with 100 μ g/ml hygromycin (Sigma) was used for screening of the single mutant strain (*ags1* Δ ::*HPH*), with 5 μ g/ml phleomycin (Invitrogen) added for the double deletion mutant strain (*ags1* Δ ::*HPH* *ags2* Δ ::*BLE*), and minimal medium was used for the selection of the triple deletion mutant strain (*ags1* Δ ::*HPH* *ags2* Δ ::*BLE* *ags3* Δ ::*PYRG*; named the *ags1* Δ *ags2* Δ *ags3* Δ strain in this study). For complementation, minimal medium supplemented with 50 μ g/ml sulfonylurea (chlorimuron-ethyl; Sigma) or 10 μ g/ml pyriithiamine (sigma) was used for screening of the transformants (10, 17, 29). For cell wall analysis, the mycelium was grown for 24 h at 37°C in Brian's medium (4). For nikkomycin Z (Sigma) susceptibility testing, 1% yeast extract medium was used.

Deletion of the *AGS* genes and construction of a triple *AGS* mutant. Genomic DNA was extracted as described by Girardin et al. (12). For Southern blot analysis, 10 μ g of digested genomic DNA was size fractionated in 0.7% agarose and blotted onto a positively charged nylon membrane (Hybond-N+; GE Healthcare).

Deletion cassettes were constructed by joining both 5'- and 3'-flanking sequences of each gene to be deleted with the positively selectable marker *HPH*, *BLE*, or *PYRG*, using the overlap method and the primers described in Table S1 in the supplemental material (18). Upstream and downstream *AGS1*, *AGS2*, or *AGS3* sequences were amplified from *AkuB*^{ku80} genomic DNA. Hygromycin, phleomycin, and *PYRG* resistance cassettes were amplified from pAN 7.1, pAN 8.1, and pAB4-1, respectively (7, 20, 23). The *ags1* Δ ::*HPH* (*ags1* Δ) single mutant was constructed by replacing *AGS1* with the *HPH* gene (conferring resistance to hygromycin). The *ags1* Δ ::*HPH* *ags2* Δ ::*BLE* (*ags1* Δ *ags2* Δ) double deletion mutant was constructed by replacing the *AGS2* open reading frame (ORF) with the *BLE* gene (conferring resistance to phleomycin) in the *ags1* Δ background. In the *ags1* Δ ::*HPH* *ags2* Δ ::*BLE* *ags3* Δ ::*PYRG* (*ags1* Δ *ags2* Δ *ags3* Δ) triple deletion mutant, the *AGS3* ORF was replaced with the *PYRG* gene from *Aspergillus niger* (conferring the ability to grow on medium without uracil and uridine) (28) in the double deletion mutant (*ags1* Δ *ags2* Δ) background (see Fig. S1). Transformations were achieved by following an

Received 20 October 2011 Accepted 24 October 2011

Published ahead of print 4 November 2011

Address correspondence to Anne Beauvais, anne.beauvais@pasteur.fr.

Supplemental material for this article may be found at <http://ec.asm.org/>.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/EC.05270-11




FIG 1 Cell wall compositions of the *ags1Δ ags2Δ ags3Δ* mutant and the *AKuB^{ku80}* parental strain. The monosaccharide compositions of the alkali-soluble (AS) and alkali-insoluble (AI) fractions are shown. Glc, glucose; Man, mannose; Gal, galactose; GlcNAc, *N*-acetylglucosamine; GalNAc, *N*-acetylgalactosamine.

electroporation or protoplast protocol (2, 19). The following day, drugs were added to the plates. The plates were incubated for 1 week at 20°C for the *ags1Δ* and *ags1Δ ags2Δ* strains and at 37°C for the *ags1Δ ags2Δ ags3Δ* strain.

For each transformation, transformants obtained under appropriate selective conditions were screened by PCR amplification using three pairs of primers (see Table S1 in the supplemental material). The first pair consisted of one primer designed to target outside the 5' end of the deletion cassette coupled with one primer designed to target inside the resistance gene. The second pair of primers consisted of one primer designed to target outside the 3' end of the deletion cassette coupled with one primer designed to target inside the resistance gene. The third pair of primers was designed to target inside the *AGS* gene. Positive amplicons obtained with only the first two pairs of primers indicated that the full-length deletion cassette was incorporated at the appropriate locus and that the *AGS* gene was deleted in these transformants. To confirm the results obtained by PCR, genomic DNA was digested by appropriate enzymes and analyzed by Southern blotting (see Fig. S1).

Complementation of the triple mutant. Strategies to complement the triple *AGS* mutant are available in the supplemental material.

Phenotypic analysis of mutants. The carbohydrate compositions of the cell walls of conidia and mycelia were determined as described previously, using three different batches of culture (21, 22). The alkali-insoluble (AI) and alkali-soluble (AS) fractions were extracted from the lyophilized cell wall (22). Monosaccharide composition was analyzed by gas chromatography after hydrolysis with 4 N trifluoroacetic acid for hexoses and 8 N HCl for hexosamines for 4 h at 100°C, followed by reduction and peracetylation of the alkali-insoluble and alkali-soluble fractions (22). α - and β 1,3 glucans were determined by measuring the reducing sugar released by recombinant α - and β 1,3 glucanases (2). An aggregation assay of germinating conidia was done as previously described (8). Nikkomycin Z susceptibility was tested following an adaptation of the resazurin method described by Clavaud et al. (5).

RESULTS AND DISCUSSION

Construction of a triple *AGS* mutant. A triple *AGS* mutant was generated through successive deletions of the three *AGS* genes (see Fig. S1 in the supplemental material). Deleting *AGS* genes in *A. fumigatus* has always been difficult. Originally, single mutants were obtained in a CBS144-89 wild-type background, with a transformation efficiency of <1% (2, 21). In spite of many attempts, no triple mutants were obtained using this parental strain.

In order to improve the transformation efficiency, an *AKuB^{ku80}* derivative of the same strain, deficient for nonhomologous end joining and favoring homologous recombination, was used (6). Even though the transformation efficiency was improved, the total number of transformants per transformation remained very low (<6/transformation). Four *ags1Δ* mutants were obtained after 3 transformation experiments; 2 double mutant *ags1Δ ags2Δ* transformants were selected after 5 transformation experiments. It was verified that the phenotype of the *ags1Δ* mutant was the same in *AKuB^{ku80}* and CBS144-89. The *ags1Δ ags2Δ* mutant had the same phenotype (growth, sporulation, and cell wall composition) as the *ags1Δ* mutant (data not shown). Finally, after 3 transformations, a unique clone was isolated subsequent to the integration of the *ags3Δ* cassette at the *AGS3* position in the *ags1Δ ags2Δ* strain to create the triple *ags1Δ ags2Δ ags3Δ* mutant. The correct integration of the resistance marker at the right locus was verified by Southern blotting.

The triple *AGS* mutant of *A. fumigatus* is totally deficient in α 1,3 glucans. The chemical composition of the *A. fumigatus* cell wall was investigated in the parental and triple *AGS* mutant strains. The ratios of AI concentration to AS concentration in the mycelial and conidial cell walls of the *ags1Δ ags2Δ ags3Δ* mutant were increased in comparison to those of the *AKuB^{ku80}* parental strain. Ratios of 10 and 2.6 were seen for the *ags1Δ ags2Δ ags3Δ* and parental strains, respectively, for the mycelium, with ratios of 8 and 2.2, respectively, for the conidia. The high alkali-insoluble/alkali-soluble hexose ratio was due to a reduction in the glucose content of the alkali-soluble fraction of the *ags1Δ ags2Δ ags3Δ* mutant: there was 4% glucose in the *ags1Δ ags2Δ ags3Δ* mycelium and 38% glucose in the *AKuB^{ku80}* mycelium, while there was 5% glucose in the *ags1Δ ags2Δ ags3Δ* conidia and 24% glucose in the *AKuB^{ku80}* conidia (Fig. 1). Unexpectedly, a small amount of glucan was found in the AS fraction of the *ags1Δ ags2Δ ags3Δ* mutant. However, after α 1,3 glucanase treatment, the mycelial AS fraction of the parental strain also contained 4% glucan and the conidial AS fraction contained 7% glucan, similar to the amounts encountered in the triple *AGS* mutant. The remaining glucan in the α 1,3 glucanase-treated mycelial and conidial AS fractions of the *ags1Δ*




FIG 2 Colonies of the triple AGS mutant (*ags1*Δ *ags2*Δ *ags3*Δ) and the *AkuB*^{ku80} parental strain grown for 48 h at 37°C on Sabouraud agar medium.

*ags2*Δ *ags3*Δ mutant and parental strains was totally degraded by β 1,3 glucanase (data not shown). This result showed that the remaining glucan was a β 1,3 glucan. The origin and structure of this small amount of β 1,3 glucan in the alkali-soluble fraction remain unknown. This result confirmed, however, that the triple AGS mutant was totally devoid of α 1,3 glucans.

The decrease in the α 1,3 glucan in the alkali-soluble fraction was compensated by an increase in the galactosamine present in the alkali-soluble and alkali-insoluble fractions (Fig. 1). Polylactosamine constituted the galactosaminogalactan, which is also an amorphous polysaccharide of the *A. fumigatus* cell wall (9). β 1,3 glucans and chitin were also increased in the mycelial alkali-insoluble fraction, whereas only chitin was increased in the conidial alkali-insoluble fraction (Fig. 1). Similarly, analysis of the cell wall of a *C. neoformans* *ags1*Δ mutant demonstrated that the loss of α 1,3 glucans was accompanied by a compensatory increase in the chitin/chitosan concentration and a redistribution of β 1,3 glucans between the cell fractions (26). The loss of α 1,3 glucans in the cell wall of the *ags1*Δ *ags2*Δ *ags3*Δ mutant was compensated by an increase in the alkali-insoluble polysaccharide content (Fig. 1). The compensatory increase of the alkali-insoluble polysaccharide amount showed that *A. fumigatus* needed to reinforce the rigidity of the cell wall affected by the loss of α 1,3 glucans. These compensatory reactions explain why the thicknesses of the cell walls of the triple mutant and parental strains observed by transmission electron microscopy were similar (data not shown). The increase in the amount of chitin in the cell wall of the *ags1*Δ *ags2*Δ *ags3*Δ mutant was correlated with a decrease in the susceptibility of the mutant to the chitin synthase inhibitor nikkomycin Z. MIC values for nikkomycin were 25 μ g/ml and >200 μ g/ml for *AkuB*^{ku80} and the *ags1*Δ *ags2*Δ *ags3*Δ mutant, respectively. Seven chitin synthases were found in the *A. fumigatus* genome (1).

α 1,3 glucans are not essential in *A. fumigatus*. Unexpectedly, the growth of the triple mutant was similar to that of the *AkuB*^{ku80} strain in Sabouraud's, Brian's, complete, and minimal media at all temperatures and pHs tested (Fig. 2; data not shown). This result showed that α 1,3 glucans were not essential in *A. fumigatus* and suggested that alkali-soluble polysaccharides of *A. fumigatus* did not play any structural role in the cell wall of this fungal species. Similarly, a strain of *H. capsulatum* that did not contain any α 1,3 glucans in the cell wall grew like wild-type strains that contained more than 35% α 1,3 glucans in the cell wall (16). In other fungi, such as *S. pombe* and *C. neoformans*, α 1,3 glucans are essential for yeast morphogenesis (14, 26). Similar results were also obtained for other dimorphic pathogenic fungi in which the switch between β 1,3 glucans and α 1,3 glucans is associated with the changes between mycelium and yeast forms (15, 27). In *S. pombe*, the content




FIG 3 Conidial aggregation after 2 h of incubation at 37°C with shaking (150 rpm). Note the lack of conidial aggregation for the triple *ags1*Δ *ags2*Δ *ags3*Δ mutant.

of α 1,3 glucans is 28%. Five AGS genes are present in *S. pombe*, but only *AGS1* of *S. pombe* is an essential gene (14). At a semipermissive temperature, the thermosensitive *ags1.1* mutant contains only 7% α 1,3 glucans in its cell wall (14). The cells are rounded and pear-shaped, and the cell wall becomes looser and thicker than that of the parental strain.

The conidiogenesis of the triple mutant was slightly decreased, as found for the single *ags1*Δ and *ags2*Δ mutants, compared to that of the parental strain (2). The viability of the conidia was not affected, and the germination level of the triple AGS mutant was similar to that of the parental strain (data not shown). However, the analysis of the aggregation phenotype of the germinating conidia also confirmed that the triple AGS mutant did not contain α 1,3 glucans in the cell wall. As reported earlier, conidia of the parental strain started to aggregate after 90 min of incubation in a culture medium at 37°C, as soon as α 1,3 glucans emerged on the cell wall surface. After 3 h, large aggregates containing more than 95% of germinating conidia were seen for the parental *AkuB*^{ku80} strain (Fig. 3). Previous studies (8) have demonstrated that this aggregation depends exclusively on α 1,3 glucan- α 1,3 glucan interactions. In contrast, and in agreement with the lack of α 1,3 glucans in the cell wall, no aggregation was observed with the germinating conidia of the *ags1*Δ *ags2*Δ *ags3*Δ mutant (Fig. 3).

Unfortunately, in spite of many transformation experiments using the various strategies summarized above, it was impossible to complement the triple AGS mutant. Reasons for these repeated failures remain unknown, but they could be due to the large size of the genes (>8 kb), their *in vivo* three-dimensional conformation, or the presence of one of these AGS genes (*AGS1*) in the subtelomeric region of a chromosome (chromosome 3), which is always difficult to manipulate genetically (11). Nevertheless, the lack of a growth phenotype in a triple AGS mutant totally devoid of α 1,3 glucans showed definitively that the α 1,3 glucans that are the major cell wall component are fully dispensable for *A. fumigatus* vegetative growth.

REFERENCES

- Alcazar-Fuoli L, et al. 2011. Functional analysis of the fungal/plant class chitinase family in *Aspergillus fumigatus*. *Fungal Genet. Biol.* 48: 418–429.
- Beauvais A, et al. 2005. Two alpha(1-3) glucan synthases with different functions in *Aspergillus fumigatus*. *Appl. Environ. Microbiol.* 71: 1531–1538.
- Beauvais A, et al. 2007. An extracellular matrix glues together the aerial-grown hyphae of *Aspergillus fumigatus*. *Cell. Microbiol.* 9:1588–1600.
- Brian PW, et al. 1961. Phytotoxic compounds produced by *Fusarium equiseti*. *J. Exp. Bot.* 12:1–12.
- Clavaud C, Beauvais A, Barbin L, Munier-Lehman H, Latgé JP. The composition of the culture medium influences the β 1,3 glucan metabo-

- lism of *Aspergillus fumigatus* and the antifungal activity of inhibitors of β 1,3 glucan synthesis tested alone or in combination. *Antimicrob. Agents Chemother.*, in press.
6. da Silva Ferreira ME, et al. 2006. The akuB(KU80) mutant deficient for nonhomologous end joining is a powerful tool for analyzing pathogenicity in *Aspergillus fumigatus*. *Eukaryot. Cell* 5:207–211.
 7. d'Enfert C. 1996. Selection of multiple disruption events in *Aspergillus fumigatus* using the orotidine-5'-decarboxylase gene, pyrG, as a unique transformation marker. *Curr. Genet.* 30:76–82.
 8. Fontaine T, et al. 2010. Cell wall alpha-1,3 glucans induce the aggregation of germinating conidia of *Aspergillus fumigatus*. *Fungal Genet. Biol.* 47:707–712.
 9. Fontaine T, et al. 2011. Galactosaminogalactan, a new immunosuppressive polysaccharide of *Aspergillus fumigatus*. *PLoS Pathog.* 7:e1002372.
 10. Fudal I, Collemare J, Bohnert HU, Melayah D, Lebrun MH. 2007. Expression of Magnaporthe grisea avirulence gene ACE1 is connected to the initiation of appressorium-mediated penetration. *Eukaryot. Cell* 6:546–554.
 11. Gibbons JG, et al. 2012. Global transcriptome changes underlying colony growth in the opportunistic human pathogen *Aspergillus fumigatus*. *Eukaryot. Cell* 11:68–78.
 12. Girardin H, Latge JP, Srikantha T, Morrow B, Soll DR. 1993. Development of DNA probes for fingerprinting *Aspergillus fumigatus*. *J. Clin. Microbiol.* 31:1547–1554.
 13. Grun CH, et al. 2005. The structure of cell wall alpha-glucan from fission yeast. *Glycobiology* 15:245–257.
 14. Hochstenbach F, et al. 1998. Identification of a putative alpha-glucan synthase essential for cell wall construction and morphogenesis in fission yeast. *Proc. Natl. Acad. Sci. U. S. A.* 95:9161–9166.
 15. Hogan LH, Klein BS. 1994. Altered expression of surface alpha-1,3-glucan in genetically related strains of *Blastomyces dermatitidis* that differ in virulence. *Infect. Immun.* 62:3543–3546.
 16. Klimpel KR, Goldman WE. 1988. Cell walls from avirulent variants of *Histoplasma capsulatum* lack alpha-(1,3)-glucan. *Infect. Immun.* 56:2997–3000.
 17. Kubodera T, Yamashita N, Nishimura A. 2002. Transformation of *Aspergillus* sp. and *Trichoderma reesei* using the pyrithiamine resistance gene (ptrA) of *Aspergillus oryzae*. *Biosci. Biotechnol. Biochem.* 66:404–406.
 18. Lamarre C, Ibrahim-Granet O, Du C, Calderone R, Latge JP. 2007. Characterization of the SKN7 ortholog of *Aspergillus fumigatus*. *Fungal Genet. Biol.* 44:682–690.
 19. Lambou K, Perkhofe S, Fontaine T, Latge J-P. 2010. Comparative functional analysis of the OCH1 mannosyltransferase families in *Aspergillus fumigatus* and *Saccharomyces cerevisiae*. *Yeast* 75:910–923.
 20. Mattern IE, Punt PJ, Van den Hondel CAMJJ. 1988. A vector of *Aspergillus* transformation conferring phleomycin resistance. *Fungal Genet. Newsl.* 35:25.
 21. Maubon D, et al. 2006. AGS3, an alpha(1-3)glucan synthase gene family member of *Aspergillus fumigatus*, modulates mycelium growth in the lung of experimentally infected mice. *Fungal Genet. Biol.* 43:366–375.
 22. Mouyna I, et al. 2010. Members of PMT family in *Aspergillus fumigatus* differentially affect growth, morphogenesis, and viability. *Mol. Microbiol.* 76:1205–1221.
 23. Punt PJ, Oliver RP, Dingemans MA, Pouwels PH, van den Hondel CA. 1987. Transformation of *Aspergillus* based on the hygromycin B resistance marker from *Escherichia coli*. *Gene* 56:117–124.
 24. Rappleye CA, Engle JT, Goldman WE. 2004. RNA interference in *Histoplasma capsulatum* demonstrates a role for alpha-(1,3)-glucan in virulence. *Mol. Microbiol.* 53:153–165.
 25. Reese AJ, Doering TL. 2003. Cell wall alpha-1,3-glucan is required to anchor the *Cryptococcus neoformans* capsule. *Mol. Microbiol.* 50:1401–1409.
 26. Reese AJ, et al. 2007. Loss of cell wall alpha(1-3) glucan affects *Cryptococcus neoformans* from ultrastructure to virulence. *Mol. Microbiol.* 63:1385–1398.
 27. San-Blas F, San-Blas G, Cova LJ. 1976. A morphological mutant of *Paracoccidioides brasiliensis* strain IVIC Pb9. Isolation and wall characterization. *J. Gen. Microbiol.* 93:209–218.
 28. Weidner G, d'Enfert C, Koch A, Mol PC, Brakhage AA. 1998. Development of a homologous transformation system for the human pathogenic fungus *Aspergillus fumigatus* based on the pyrG gene encoding orotidine 5'-monophosphate decarboxylase. *Curr. Genet.* 33:378–385.
 29. Zhang S, Fan Y, Xia YX, Keyhani NO. 2010. Sulfonylurea resistance as a new selectable marker for the entomopathogenic fungus *Beauveria bassiana*. *Appl. Microbiol. Biotechnol.* 87:1151–1156.