

---

**The isolation and characterization of the *Escherichia coli* DNA adenine methylase (*dam*) gene**

---

Joan E. Brooks, Robert M. Blumenthal<sup>+</sup> and Thomas R. Gingeras

---

P.O. Box 100, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA

---

Received 18 October 1982; Revised and Accepted 6 December 1982

---

**ABSTRACT**

The *E. coli dam* (DNA adenine methylase) enzyme is known to methylate the sequence GATC. A general method for cloning sequence-specific DNA methylase genes was used to isolate the *dam* gene on a 1.14 kb fragment, inserted in the plasmid vector pBR322. Subsequent restriction mapping and subcloning experiments established a set of approximate boundaries of the gene. The nucleotide sequence of the *dam* gene was determined, and analysis of that sequence revealed a unique open reading frame which corresponded in length to that necessary to code for a protein the size of *dam*. Amino acid composition derived from this sequence corresponds closely to the amino acid composition of the purified *dam* protein. Enzymatic and DNA:DNA hybridization methods were used to investigate the possible presence of *dam* genes in a variety of prokaryotic organisms.

**INTRODUCTION**

The *E. coli dam* (DNA adenine methylase) gene codes for an enzyme which methylates within the sequence GATC (1-3). When DNA has been modified by the *dam* methylase, it is no longer susceptible to cleavage by the restriction endonuclease *Mbo*I (2,4).

The *dam* methylase is not part of a restriction modification system, but rather has been thought to act in post-replication mismatch repair. There are several lines of evidence for its involvement in mismatch repair. First, in heteroduplex lambda phage DNA having only one methylated strand, the repair system will usually correct the unmethylated strand to match the methylated strand; fully methylated mismatched heteroduplexes are not corrected (5,6). Second, *E. coli* strains in which the *dam* methylase is either not produced (*dam*<sup>-</sup>) or overproduced (*dam*<sup>S</sup>) are hypermutable (7-10). Third, the combination of the *dam*<sup>3</sup> mutant allele, which has no detectable *dam* activity, with mutants in DNA repair functions such as *polA*, *lexA*, *recA*, *recB* or *recC* is lethal (11).

Furthermore, the *E. coli dam* protein may play a role in DNA replication: the sequence methylated by the *dam* enzyme, GATC, occurs at a very high

---

frequency (11 times within 245 base pairs) at the *E. coli* origin of replication (12,13). It also tends to occur with high frequency near or at the ends of Okazaki fragments (14,15).

In addition to functional considerations, the *E. coli* dam enzyme is noteworthy because it shares sequence specificity with a number of Type II restriction endonucleases and methylases (16). In particular, we were interested in determining whether the dam gene has any sequence homology to genes encoding Type II restriction methylases with the same specificity. Therefore, we undertook the isolation and characterization of the dam methylase gene from *E. coli*.

#### MATERIALS AND METHODS

##### (a) Bacterial strains.

The *E. coli* strains used were GM119 (dam<sup>3</sup>; Marinus, unpublished observation); SK1036 (dam<sup>4</sup>) (10) and HB101, (dam<sup>+</sup>), (17). The origin of the bacterial species found in Table 3 is given in Roberts (16), except for the following species: *E. aerogenes*, *E. carotovora* and *R. meliloti* came from J. Zyskind; *S. typhimurium* came from S. Schlagman.

##### (b) DNA preparation.

The *E. coli* plasmids used in cloning and sequencing were isolated by the cleared lysate method (18), followed by banding on CsCl gradients containing ethidium bromide. "Mini" preparations of plasmid DNA were prepared by the procedure of Klein et al. (19). Chromosomal DNA preparations from the various bacteria were made by the method of Marmur (20).

##### (c) Restriction enzymes and DNA end-labeling.

The enzymes Sau3AI, BclI, RsaI, and TaqI were prepared in this laboratory by P.A. Myers. DdeI was a gift from R. Meagher. Digests with these enzymes were carried out in buffer containing 6 mM Tris-HCl pH 7.9, 6 mM MgCl<sub>2</sub>, and 6 mM SHCH<sub>2</sub>CH<sub>2</sub>OH. The enzymes BamHI, PvuI, PvuII, MboI, HindIII, and SalI were purchased from New England Biolabs; HpaI was purchased from Bethesda Research Labs. For these enzymes, we used the buffers recommended by their respective manufacturers.

Calf alkaline phosphatase was obtained from Boehringer-Mannheim, and purified further by passage over a DEAE-cellulose column. It was used as described previously (21). T4 polynucleotide kinase (Boehringer-Mannheim), was used as described by Chaconas and van de Sande (22). The 5' [ $\gamma$ <sup>32</sup>P] ATP (3000 Ci/mmol) was purchased from Amersham, Inc.

(d) Determination and analysis of base sequence.

The chemical method of Maxam and Gilbert (23) was used to determine the base sequence of the cloned insert in pdam118. The specific reactions used were dimethylsulfate (G), formic acid (G + A), hydrazine (T + C), and hydrazine plus NaCl (C). Reaction products were resolved on 0.2 mm thick 6, 8, or 12% polyacrylamide gels containing 8 M urea and a Tris-borate buffer (24). Gels were autoradiographed using Kodak XR-1 X-Ray Film, sometimes with the aid of DuPont Cronex 'Lightning Plus' intensifying screens.

The resulting data were assembled and analysed using computer programs which are described elsewhere (25-27). Further analysis involved the use of several programs in the MOLGEN-SEQ collection on the SUMEX computer system at Stanford University (28).

(e) Cloning dam in E. coli.

pGG503, containing the dam function on a 23 kb insert into pBR322 (10,29) was our starting material. 10 µg of plasmid was subjected to partial cleavage with Sau3A to give fragments averaging 2-5 kb in length. 1.0 µg of the digested plasmid was ligated to phosphatase-treated pBR322 (0.1 µg) using T4 DNA ligase (N.E. Biolabs) under conditions suggested by the manufacturer. The ligation reaction was used to transform E. coli SK1036 (dam<sup>-</sup>) cells by the CaCl<sub>2</sub>-heat shock method (30). After 1 hour, 5 ml LB broth was added to the transformation mix and the cells grown for 2 hours in the presence of ampicillin (100 µg/ml). At that time the cells were harvested, resuspended in 5 ml LB with ampicillin and grown overnight. Plasmids were isolated from the culture by the method of Klein et al. (19). 2 µg of the recombinant plasmids were extensively cleaved by incubation with 10 units of MboI for 3 hours at 37°C. The reaction was terminated by heating to 68°C for 5 minutes and then used to transform E. coli SK1036 cells. Transformants were selected on LB plates containing 100 µg/ml ampicillin, and screened by replica plating for sensitivity to 2-aminopurine in LB plates (400 µg/ml; Vega Biochemicals) (2-aminopurine is an adenine analog that is lethal to dam<sup>-</sup> cells at high concentrations (8)). Colonies that could grow in the presence of 2-aminopurine were picked from control plates, their plasmids isolated and challenged by MboI. Those plasmids not cleavable by MboI were further analyzed to determine the size of the insert.

(f) Presence of dam in other organisms.

To test for methylation of the GATC sequences in other organisms, 2 µg of bacterial DNA were digested with 3 units of MboI enzyme and the products resolved by electrophoresis on a 1% agarose gel (31). To test for sequence

homology, 2 µg of each DNA were digested with 3 units of HindIII enzyme and the digests run out on 1% agarose gels. These DNAs were then transferred to nitrocellulose filters (Schleicher and Schuell, BA85) by the Southern method (32). Hybridization probes (pBR322, Adenovirus-2 DNA, and the isolated BamI-PvuII fragment from pdam118) were labeled by nick-translation as described (33). Hybridization and filter washes were performed at 42°C as described by Bukhari et al. (34) with the modifications of Chaconas et al. (35). Hybridizations were performed under less stringent conditions in which the filters were first incubated at 15°C and washed extensively at 4°C.

## RESULTS


### (a) Localization of the dam gene.

Details of the cloning procedure are given in the Methods section and diagrammed in Fig. 1. Prior to the second transformation step, only 1 in 10<sup>5</sup> plasmids escapes MboI restriction; of the secondary transformants, 80% possess a Dam<sup>+</sup> phenotype (i.e., their isolated plasmids are resistant to MboI cleavage). The smallest of the Dam<sup>+</sup> plasmids, designated pdam117, contains a 1.6 kb insert. pdam117 was further analyzed to determine the position of the dam gene within the 1.6 kb insert. This was first done by a series of subcloning experiments based upon extensive restriction site analysis of the plasmid.


The results of these subcloning experiments are illustrated in Figure 2. Cleavage of pdam117 with PvuII followed by religation led to a clone (pdam118) which was phenotypically Dam<sup>+</sup>, but carried an insert which was approximately 400 base pairs shorter than that of pdam117 (i.e., 1.14 kb). Two other subclones were constructed by splitting the pdam117 insert at its internal BamHI site. The 500 base pair 'Bam-Bam' fragment was inserted into the BamHI site of pBR322, while the remainder of pdam117 (minus the Bam-Bam piece) was recircularized by ligation. Both plasmids, when used to transform GM119 cells were found to be Dam<sup>-</sup>. These clones were designated as pxdam1 and pxdam2, respectively. Finally, the plasmid pdam118 was cleaved with PvuII and HpaI, the resulting large fragment recircularized, and transformed into GM119 cells. This construct, designated pxdam3, also resulted in cells which were Dam<sup>-</sup>. These experiments suggest that the dam gene is situated with one end between the two BamHI sites and the other between the HpaI and PvuII sites within the pdam118 insert (Figure 2).

### (b) Nucleotide sequence of the dam gene.

The strategy used in sequencing pdam118 by the Maxam-Gilbert method is



**Figure 1:** Cloning strategy for the *E. coli* *dam* methylase gene. A clone bank of the *E. coli* genome was constructed by Clarke and Carbon (50). This bank was screened by Modrich and Herman (29) for clones containing the *trpS* gene, since *trpS* had previously been genetically mapped proximal to the *dam* locus (51,52). A 23 kb *Pst* fragment, containing both the *dam* and *trpS* functions was inserted into pBR322 and the construct designated pGG503 (29).




**Figure 2:** Restriction map of the insert from *pdam117* and derivative plasmids. The position and orientation of the *dam* methylase gene is indicated.

shown in Figure 3. The completed sequence (Figure 4) was analyzed by computer for the presence of an open reading frame which could accommodate a product the size of the dam methylase. The longest available reading frame is 834 nucleotides in length, beginning with an AUG codon at position 195 of the insert, and ending with a UAA terminator at nucleotide 1029. The dam methylase has been purified to homogeneity (29) and the amino acid composition of the dam protein determined (P. Modrich, pers. comm.). We have compared the actual amino acid composition to the predicted composition of the putative protein from the 834 base open reading frame (Table 1): the comparison shows an excellent agreement in both size and composition between the actual and putative proteins.

(c) The presence of dam genes in other organisms.

A variety of prokaryotic organisms were tested for both the presence of a 'dam-like' function and sequence homology to the cloned dam gene. To evaluate whether an organism has a dam methylase, we digested its DNA with MboI; if methylated in vivo, the GATC sequences are completely resistant to MboI cleavage. To determine whether an organism has DNA sequences homologous to the dam gene, we used Southern blot hybridization analysis. The results of both types of experiments are summarized in Table 2. It is clear that there are DNA sequences homologous to dam in all the Enterobacteriaceae and Haemophilus species tested; the DNA of all these bacteria are also protected against MboI cleavage. It is therefore probable that these organisms all



**Figure 3:** Strategy employed in sequencing of Eco dam region. Directionality and extent of sequence determination (see Methods section) from restriction termini are indicated by the small arrows above the restriction map. Open circles correspond to the start of 5'-end labeled DNA fragments. Sequencing from the BclI site (370) used DNA fragments isolated from pxdam1, whereas sequencing from all other sites used pdam117 DNA (Fig. 2). Location and orientation of the major open reading frame containing the dam gene is indicated by the boldfaced arrow. Both strands have been sequenced extensively and covered in many areas by duplication, although the duplications have not been depicted on this map.

5' GAGTCCCTTTTTCTGCGGTATCTGCTGCAACAAAACACCCAGCCCTACAGCGGTGGTTGTGTCGGGATCTGAAGTAACTCAAGGTTATCTCCCGCATGCTGCGAGGTTTCCACAGCCGGGAAGGTTAATAGTTAGTACAG

150

100

50

200

100

50

250

300

350

400

450

500

550

600

650

700

750

800

850

900

950

1000

1050

1100

TTTTCGCCCTGAGTGAAGTACCCGAAAGCCCTGCGAGCTG<sup>3'</sup>

MET LYS ASN ARG ALA PHE LEU LYS TRP ALA GLY LYS TYR PRO LEU LEU ASP ILE LYS ARG HIS LEU PRO LYS GLY GLU LEU VAL GLU VAL PHE VAL GLY SER VAL PHE LEU ASN THR ASP PHE SER ARG  
 ATG ARG AAA MAT CGG GCT TTT TTG AMG TGG GCA GGG GGC MAG TAT CCC CTG CTT GAT GAT ATT AAA CGG CAT TTG CCC MAG GGC GAA TGT CTG GTT GAG CCT TTA GTA GGT GCC GGG TCG GTG TTT CTC AAC ACC GAC TTT TCT CGT  
 TYR TLE LEU ALA ASP TLE ASN SER ASP ILE SER LEU TYR ASN TLE VAL LYS MET ARG THR ASP GLU TYR VAL GLN ALA ALA ARG GLU LEU PHE VAL PRO GLU THR ASN CYS ALA GLU VAL TYR TYR GLN PHE ARG GLU GLU PHE  
 TAC ATT CTT GCC GAT ATC AAT AGC GAG CTG ATC AGT CTC TAT AAC ATT GTG AMG ATG CGT ACT GAT GAG TAC GTA CAG GCC GCA CGC GAG CTG TTT GTT CCC GAA ACA AAT TGC GCC GAG GTT TAC TAT CAG TTC CGC GAA GAG TTC  
 ASN LYS SER GLN ASP PRO PHE ARG ARG ALA VAL LEU LEU TYR LEU ASN ARG TYR GLY TYR ASN GLY LEU CYS ARG TYR ASN LEU ARG GLY GLU PHE ASN VAL PRO PHE GLY ARG TYR LYS LYS PRO TYR PHE PRO GLU ALA GLU  
 AAC AAA AGC CAG GAT CCG TTC CGT CGG GCG GTA CTG TTT TTA TAT TTG AAC CGC TAC GGT TAC AAC GGC CTG TGT GGT TAC PAT CTG CGC GGT GAG TTT AAC GTG CCG TTC GGC GCG TAC AAA CCC TAT TTC CCG GAA GCA GAG  
 LEU TYR HIS PHE ALA GLU LYS ALA GLN ASN ALA PHE PHE TYR CYS GLU SER TYR ALA ASP ALA ARG ALA SER VAL VAL TYR CYS ASP PRO TYR TYR ARG GLU TRP TYR GLN ARG ALA THR ALA ASN PHE THR ALA TYR  
 TTG TAT CAG TTC GCT GAA AAA GCG CAG AAT GCC TTT TTC TAT TGT GAG TCT TAC GCC GAT AGC ATG GCG CGC GCA GAT GAT GCA TCC GTC TAT TGC GAT CCG CCT TAT GCA CGC CTG TCT GCG ACC GCC AAC TTT ACG GCG TAT  
 HIS THR ASN SER PHE THR LEU GLU GLN ALA HIS LEU ALA ILE ALA GLU LYS LEU VAL LEU VAL LEU ILE SER ASN HIS ASP THR MET LEU THR ARG GLU TRP TYR GLN ARG ALA LYS LEU HIS VAL VAL  
 CAC ACA AAC AGT TTT ACG CTT GAA CAA GAG CAT CTG GCG GAG ATC GCC GAA GGT CTG GTT GAG CGC CAT ATT CCA GTG CTG ATC TCC AAT CAC GAT ACG ATG TTA ACG CGT GAG TGG TAT CAG CGC GCA AAA TTG CAT GTC GTC  
 LYS VAL ARG ARG SER ILE SER SER ASN GLY TYR ARG LYS LYS VAL ASP GLU LEU LEU ALA LEU TYR LYS PRO GLY VAL VAL SER PRO ALA LYS LYS \*\*\*  
 AAA GTT CAG CGC AGT ATA AGC ARG AGC GGC ACA GGT AAA AAG GTG GAC GAA CTG CTG GCT TTG TAC AAA CCA GGA GTT TCA CCC GCG AAA AAA TAA TTCTCAAGGAGAAGGCGATCAAGAGTATTGATTCGCCCTCAATCTGTGCGCTGA

**Figure 4:** Nucleotide sequence of the *pdam118* insert. The *dam* methylase gene is predicted to begin at nucleotide 195 and end at nucleotide 1029, giving rise to a protein of 31,000 MW. Also represented, above the nucleotide sequence, is the predicted amino acid sequence of the *dam* protein.

TABLE 1  
Comparison of the Predicted Vs. the Empirically-derived  
Amino Acid Sequence of the dam Protein

<u>Amino Acid</u>	<u>Prediction from</u>		<u>Empirical</u>
	<u>Nucleotide Sequence (%)</u>		<u>Determination (%)<sup>a</sup></u>
Alanine	9.3		9.4
Arginine	6.8		7.2
Asparagine	5.4	10.1	10.5 (ASX)
Aspartic Acid	4.7		
Cysteine	1.8		ND <sup>b</sup>
Glutamine	2.5	9.3	9.5 (GLX)
Glutamic Acid	6.8		
Glycine	4.7		5.3
Histidine	2.5		2.9
Isoleucine	3.2		3.0
Leucine	9.3		9.5
Lysine	6.5		7.7
Methionine	1.4		2.5
Phenylalanine	6.1		6.0
Proline	5.0		5.1
Serine	5.4		4.5
Threonine	3.6		3.4
Tryptophan	0.7		ND <sup>b</sup>
Tyrosine	7.2		6.8
Valine	6.8		6.4

<sup>a</sup> Modrich and Herman, personal communicaton.  
<sup>b</sup> ND, not determined.

contain homologous DNA adenine methylase genes. There were no cases found in which a bacterium contains DNA sequences homologous to the pdam118 insert but has no active dam methylase. There are two cases (Moraxella bovis and Anabaena variabilis) in which the bacterium's DNA was resistant to MboI cleavage but showed no sequence homology to the pdam118 insert, even under nonstringent hybridization conditions.

**DISCUSSION**

(a) Identification of the dam gene.

As stated previously, the nucleotide sequence of pdam118 in conjunction



TABLE 2  
Occurrence of dam Genes in Other Organisms

<u>Organism</u>	<u>Hybridization to</u> <u>pdam 118</u> <u>(dam<sup>+</sup> clone)</u>	<u>Cleavage by</u> <u>MboI enzyme</u>
Agrobacterium tumefaciens	-	+
Anabaena variabilis	-	-
Bacillus caldolyticus	-	+
Bacillus globigii	-	+
Enterobacter aerogenes	+	-
Enterobacter cloacae	+	-
Erwinia carotovora	+	-
Haemophilus gallinarum	+	-
Haemophilus parahaemolyticus	+	-
Haemophilus parainfluenzae	+	-
Klebsiella pneumoniae	+	-
Moraxella bovis	-	- <sup>a</sup>
Proteus vulgaris	+	-
Providencia stuartii	+	-
Pseudomonas aeruginosa	-	+
Rhizobium meliloti	-	+
Salmonella typhimurium	+	-
Serratia marcesens	+	-
Staphylococcus aureus 3A	-	+
Xanthomonas holcicola	-	+
Xanthomonas malvacearum	-	+
Xanthomonas oryzae	-	+

<sup>a</sup> Moraxella bovis differs from the other dam<sup>+</sup> organisms listed above by having a corresponding Type II restriction endonuclease activity.

with the phenotypes of the various subclones has allowed us to set a unique set of boundaries for the dam gene. The agreement between the predicted amino acid composition of the putative protein and the amino acid composition of the purified protein also lends further support to our having correctly identified and sequenced the structural gene for the Eco dam methylase. Nevertheless, there still existed a possibility that the function cloned and sequenced was an E. coli regulatory protein that activated the dam gene rather than the methylase gene itself.

Recent experiments exclude this possibility. The HindIII-PvuII fragment of pdam118, containing the putative dam gene, was ligated to a yeast-E. coli shuttle vector, YEP228 (J. Hicks, unpublished observation) and the construct used to transform competent Saccharomyces cerevisiae cells. Chromosomal and plasmid DNA from these transformants were examined for the presence of dam modification. We have determined that all the MboI sites on the expression plasmid and as much as 30% of the chromosomal DNA is protected from MboI

TABLE 3  
E. coli Codon Usage (in frequency per 1000)

		<u>dam</u>	<u>RIM</u> <sup>a</sup>	<u>RIR</u> <sup>a</sup>	<u>MIX</u> <sup>b</sup>
Arg	CGU	22	0	4	30
	CGC	36	3	0	21
	CGA	4	3	7	3
	CGG	7	0	0	4
	AGA	0	31	22	5
	AGG	0	3	18	2
Leu	CUU	11	16	22	18
	CUC	7	3	11	5
	CUA	0	16	14	7
	CUG	47	9	0	47
	UAA	7	28	40	21
	UUG	22	16	11	6
Ser	UCU	11	25	29	17
	UCC	7	0	4	13
	UCA	4	9	25	8
	UCG	4	12	7	9
	AGU	11	12	7	7
	AGC	18	12	14	9
Thr	ACU	4	9	18	21
	ACC	7	0	0	22
	ACA	11	16	14	5
	ACG	14	3	4	10
Pro	CCU	7	19	18	5
	CCC	18	0	0	4
	CCA	7	16	4	7
	CCG	18	0	0	19
Ala	GCU	11	12	32	37
	GCC	29	3	4	20
	GCA	25	9	14	31
	GCG	29	6	4	23
Gly	GGU	14	19	32	33
	GGC	22	9	4	29
	GGA	4	9	18	4
	GGG	7	16	22	6
Val	GUU	22	40	29	28
	GUC	18	3	7	9
	GUA	11	9	18	21
	GUG	18	6	4	17
Lys	AAA	43	68	61	46
	AAC	22	40	18	18
Asn	AAC	32	28	11	25
	AAU	22	59	61	10

Gln	CAA	7	12	25	13
	CAU	14	16	14	16
Glu	GAA	29	31	36	37
	GAG	40	31	25	18
Asq	GAC	11	6	11	27
	GAU	36	65	54	25
Tyr	UAC	32	16	7	12
	UAU	40	50	22	14
Cys	UGC	7	12	0	6
	UGU	11	9	4	5
Phe	UUC	25	22	4	18
	UUU	36	50	36	18
Ile	AUA	4	22	43	5
	AUC	18	6	14	32
	AUU	11	43	25	24
Met	AUG	14	6	25	22
Trp	UGG	7	6	7	12

<sup>a</sup> See references: 50 and 51.

<sup>b</sup> See references: 38. This is a collection of the codon usage for a total of 25 *E. coli* genes.

cleavage and susceptible to *DpnI* cleavage (R. Kostriken, et.al., unpublished observations.) (*DpnI* cuts DNA at the sequence GATC but only when the adenines within this site are methylated (36). Since yeast DNA is known to be devoid of N<sup>6</sup>-methyladenine in its native state (D. Swinton and S. Hattman, personal communication), the *dam* methylase activity must come from the *pdam118* insert. The phenotypic effects of the adenine methylation in the yeast transformants are now being investigated.

(b) Location of *dam* on the *E. coli* chromosome.

Clones containing the *dam* gene were originally isolated on the basis of their proximity to the *trpS* gene (37). The *trpS* gene, coding for the tryptophanyl tRNA synthetase enzyme has also been isolated and its sequence determined (37,38). A comparison of restriction maps made of clones containing *trpS* and *dam* genes respectively, show the genes are proximal. They are separated by approximately 1 kb of DNA and are both transcribed counterclockwise on the *E. coli* chromosome (38).

(c) Presence of the *dam* gene in other bacteria.

As shown in Table 2, all members of the family *Enterobacteriaceae* and

also the unrelated genus Haemophilus that were tested were found to possess a functional dam methylase, both by resistance of the DNA to MboI cleavage and sequence homology to the cloned dam gene. In no instance did an organism whose DNA had sequence homology to the Eco dam gene contain DNA sensitive to MboI cleavage. There were two cases in which a bacterial DNA was resistant to MboI cleavage but the DNA showed no sequence homology to the pdam118 insert. The first case is that of Moraxella bovis, which is unique among the bacteria surveyed in that it has a Type II restriction modification system recognizing the sequence GATC and is blocked by adenine methylation within that sequence (39,40). The MboI methylase, which is part of a restriction modification system, may differ in origin as well as in function from the Eco dam methylase, and therefore lack any sequence homology. The second case is that of Anabaena variabilis, a cyanobacterium which possesses three known restriction modification systems (41,42). However, no combination of DNA methylation associated with any of these three systems could confer protection to the GATC sequences. Three possible explanations can be given for the presence of a ''dam-like'' activity but the absence of hybridization to Eco dam in Anabaena. First, it is possible that the methylase is part of a restriction modification system which is expressed at a very low level and, like the MboI methylase, is unrelated to the Eco dam. Second, it is possible that the Anabaena methylase represents a new class of adenine methylases recognizing GATC that is functionally and evolutionarily unrelated to either dam or MboI methylases. The third possibility is that the Anabaena methylase is, in fact, evolutionarily related to Eco dam but has undergone sufficient genetic drift so as to no longer hybridize, even under nonstringent conditions.

A third case worthy of mention is that of Staphylococcus aureus 3A. This bacterium also has a restriction system, Sau3A, specific for GATC (43); however, unlike the MboI system, its restriction activity is not blocked by methyladenine within the recognition sequence (16). Staphylococcus aureus 3A DNA is not resistant to MboI cleavage, nor does it contain any sequence homology to pdam118. The hybridization experiments do not indicate any sequence homology exists between the dam gene and modification methylases that belong to restriction systems.

The fact that all the bacteria tested belonging to the family Enterobacteriaceae have a dam methylase is particularly interesting in light of the reports concerning the nucleotide sequences present at their origins of replication (ori). In addition to E. coli, (12,13) origins of replication

for Salmonella typhimurium (44), Enterobacter aerogenes (45), Klebsiella pneumoniae (45), Erwinia carotovora (45), and Vibrio harveyi (Zyskind et al., submitted for publication) have been cloned and sequenced. In all six bacteria, within the approximately 250 bases necessary for oriC function, there are between 11 and 14 occurrences of the dam site, GATC. By random occurrence this sequence would be expected only once in 256 bases. Zyskind and Smith (44) proposed that the high concentration of dam sites in the ori region may be necessary for its sequence conservation. Since the post replication mismatch repair system may act by discriminating between methylated and unmethylated DNA strands, frequent occurrence of the GATC sequence could result in localization of repair enzymes within this region. Therefore, errors introduced during replication would have a greater chance of being repaired.

There is also evidence that dam methylation plays a role in the expression of the mom gene in bacteriophage Mu (46). Unlike the role as repressor of gene activity that methylation is proposed to play in eukaryotes (47,48), dam methylation seems to activate the mom function. One hypothesis currently being investigated is that dam methylation is required for transcription of the mom gene (49, R. Kahmann, pers. comm.). Having the dam gene cloned will undoubtedly assist in understanding its various roles repair, repair and transcription.

#### ACKNOWLEDGMENTS

We wish to thank Drs. P. Modrich and G. Herman for providing the pGG503 clone, purified dam protein and data on the amino acid composition of the dam methylase. We also are indebted to Dr. J. Zyskind for her gift of several bacterial DNAs as well as unpublished nucleotide sequence information about their origins of replication. Our thanks to Drs. G.M. Marinus and S. Schlagman for providing us with the aforementioned bacterial strains. We thank P. Myers for preparing the majority of restriction enzymes used in these experiments. We also thank Dr. R.J. Roberts for his helpful discussions and critical reading of this manuscript. Finally, we wish to acknowledge D. O'Loane for her technical assistance; N. D'Anna, C. Carpenter and M. Ockler for their help in preparation of the manuscript. This work was supported by National Institutes of Health Grant # 5-R01-CA27275-02 (to T.R.G.), National Science Foundation Grant # PCM-7919-882 (to Richard J. Roberts) and National Institutes of Health Postdoctoral Fellowship # CA 09311-02 (to R.M.B.)

† Current address: Department of Microbiology, Medical College of Ohio, Toledo, OH 43699, USA

### REFERENCES

1. Marinus, M.G. and Morris, N.R. (1973) *J. Bacteriol.* 114, 1143-1150
2. Lacks, S. and Greenberg, B. (1977) *J. Mol. Biol.* 114, 153-168
3. Hattman, S., Brooks, J.E. and Masurekar, M. (1978) *J. Mol. Biol.* 126, 367-380
4. Dreiseikelmann, B., Eichenlaub, R. and Wackernagel, W. (1979) *Biochim. Biophys. Acta*, 562, 418-428
5. Wagner, Jr., R. and Meselson, M. (1976) *Proc. Natl. Acad. Sci. USA* 73, 4135-4139
6. Meselson, M., Pukkila, P., Rykowski, M., Peterson, J., Radman, M., Wagner, R., Herman, G. and Modrich, P. (1980) *J. Supramolec. Struct. Suppl.* 4, 311
7. Marinus, M.G. and Morris, N.R. (1975) *Mutat. Res.* 28, 15-26.
8. Glickman, B.W., van den Elsen, P. and Radman, M. (1978) *Molec. Gen. Genet.* 163, 307-312
9. Glickman, B. and Radman, M. (1980) *Proc. Natl. Acad. Sci. USA* 77, 1063-1067
10. Herman, G.E. and Modrich, P. (1981) *J. Bacteriol.* 145, 644-646
11. Marinus, M.G. and Morris, N.R. (1974) *J. Mol. Biol.* 85, 309-322
12. Sugimoto, K., Oka, A., Sugisaki, H., Takanami, M., Nishimura, A., Yasuda, S. and Hirota, Y. (1979) *Proc. Natl. Acad. Sci. USA* 76, 575-579
13. Meijer, M., Beck, E., Hansen, F.G., Bergmans, H.E.N., Messer, W., von Meyenburg, K. and Schaller, H. (1979) *Proc. Natl. Acad. Sci. USA* 76, 580-584
14. Gomez-Eichelmann, M.C. and Lark, K.G. (1977) *J. Mol. Biol.* 117, 621-635
15. Marinus, M.G. (1976) *J. Bacteriol.* 128, 853-854
16. Roberts, R.J. (1981) *Nucl. Acids Res.* 9, r75-96
17. Boyer, H.W. and Roulland-Dussoix, D. (1969) *J. Mol. Biol.* 41, 459-472
18. Clewell, D.B. and Helinski, D.R. (1972) *J. Bacteriol.* 110, 1135-1146
19. Klein, R.D., Selsing, E. and Wells, R.D. (1980) *Plasmid* 3, 88-91
20. Marmur, J. (1961) *J. Mol. Biol.* 3, 208-218
21. Torok, I. and Karch, F. (1980) *Nucleic Acids Res.* 8, 3105-3123
22. Chaconas, G. and van de Sande, H.H. (1980) in *Methods in Enzymology*, Grossman, L. and Moldave, K. Eds., Vol 65, pp. 75-85, Academic Press, New York
23. Maxam, A.M. and Gilbert, W. (1980) *Methods Enzymol.* 65, 499-560
24. Sanger, F. and Coulson, A.R. (1978) *FEBS Letters* 87, 107-110
25. Gingeras, T.R., Milazzo, J.P., Sciaky, D. and Roberts, R.J. (1979) *Nucleic Acids Res.* 7, 529-545
26. Gingeras, T.R., Rice, P., Gelinas, R. and Roberts, R.J. (1982) *Nucleic Acids Res.* 10, 103-113
27. Blumenthal, R., Rice, P. and Roberts, R.J. (1982) *Nucleic Acids Res.* 10, 91-102
28. Brutlag, D.L., Clayton, J., Friedland, P. and Kedes, L.H. (1982) *Nucleic Acids Res.* 10, 279-294
29. Herman, G.E. and Modrich, P. (1982) *J. Biol. Chem.* 257, 2605-2612
30. Cohen, S.N., Chang, A.C.Y. and Hsu, L. (1972) *Proc. Natl. Acad. Sci. USA* 69, 2110-2114
31. Sharp, P.A., Sugden, B. and Sambrook, J. (1973) *Biochemistry* 12, 3055-3063
32. Southern, E.M. (1975) *J. Mol. Biol.* 98, 503-517
33. Kelly, R.B., Cozzarelli, N.R., Deutscher, M.P., Lehman, I.R. and Kornberg, A. (1970) *J. Biol. Chem.* 245, 39-45

- 
34. Bukhari, A.I., Froshauer, S. and Botchan, M. (1976) *Nature (London)* 264, 580-583
  35. Chaconas, G., Harshey, R.M. and Bukhari, A.I. (1980) *Proc. Natl. Acad. Sci. USA* 77, 1778-1782
  36. Lacks, S. and Greenberg, B. (1975) *J. Biol. Chem.* 250, 4060-4072
  37. Hall, C.V. and Yanofsky, C. (1981) *J. Bacteriol.* 148, 941-949
  38. Hall, C.V., van Cleemput, M., Muench, K.H. and Yanofsky, C. (1982) *J. Biol. Chem.* 257, 6132-6136
  39. Gelinas, R.E., Myers, P.A. and Roberts, R.J. (1977) *J. Mol. Biol.* 114, 169-179
  40. Brooks, J.E. and Roberts, R.J. (1982) *Nucl. Acids Res.* 10, 913-934
  41. Hughes, S.G. and Murray, K. (1980) *Biochem. J.* 185, 65-75
  42. Roizes, G., Nardeux, P.-C. and Monier, R. (1979) *FEBS Letters*, 104, 39-44
  43. Sussenbach, J.S., Monfoort, C.H., Schiphof, R. and Stobberingh, E.E. (1976) *Nucl. Acids Res.* 3, 3193-3202
  44. Zyskind, J.W. and Smith, D.W. (1980) *Proc. Natl. Acad. Sci. USA* 77, 2460-2464
  45. Cleary, J.M., Smith, D.W., Harding, N.E. and Zyskind, J.W. (1982) *J. Bacteriol.* 150, 1467-1471
  46. Toussaint, A. (1977) *J. Virol.* 23, 825-826
  47. Razin, A. and Riggs, A.D. (1980) *Science* 210, 604-610
  48. Ehrlich, M. and Wang, R. Y.-H. (1981) *Science* 212, 1350-1357
  49. Hattman, S. (1982) *Proc. Natl. Acad. Sci.* 79, 5518-5521
  50. Clarke, L. and Carbon, J. (1976) *Cell* 9, 91-99
  51. Marinus, M.G. (1973) *Molec. Gen. Genet.* 127, 47-55
  52. Marinus, M.G. and Konrad, E.B. (1976) *Molec. Gen. Genet.* 149, 273-277