

ORIGINAL

Open Access

SimReg1 is a master switch for biosynthesis and export of simocyclinone D8 and its precursors

Liliya Horbal¹, Yuriy Rebets¹, Mariya Rabyk¹, Roman Makitrynskyy¹, Andriy Luzhetskyy², Victor Fedorenko¹ and Andreas Bechthold^{3*}

Abstract

Analysis of the simocyclinone biosynthesis (*sim*) gene cluster of *Streptomyces antibioticus* Tü6040 led to the identification of a putative pathway specific regulatory gene *simReg1*. *In silico* analysis places the SimReg1 protein in the OmpR-PhoB subfamily of response regulators. Gene replacement of *simReg1* from the *S. antibioticus* chromosome completely abolishes simocyclinone production indicating that SimReg1 is a key regulator of simocyclinone biosynthesis. Results of the DNA-shift assays and reporter gene expression analysis are consistent with the idea that SimReg1 activates transcription of simocyclinone biosynthesis, transporter genes, regulatory gene *simReg3* and his own transcription. The presence of extracts (simocyclinone) from *S. antibioticus* Tü6040 × pSSimR1-1 could dissociate SimReg1 from promoter regions. A preliminary model for regulation of simocyclinone biosynthesis and export is discussed.

Keywords: Simocyclinone, angucycline, regulation, transport

Introduction

The actinomycetes, including in particular members of the genus *Streptomyces*, are the industrial source for a large number of bioactive compounds employed as antibiotics and other drugs Horinouchi 2007; Bibb and Hesketh 2009. Actinomycetes produce these molecules as part of their "secondary" or nonessential metabolism van Wezel et al. 2009. Many *Streptomyces* species are capable of producing more than one secondary metabolite Ohnishi et al. 2008; van Wezel et al. 2009. The timing of the production of secondary metabolites and the amount of the accumulated compounds correlates with the environmental conditions and morphological differentiation van Wezel et al. 2009; Bibb et al. 2009; van Wezel et al. 2011. Furthermore, it has also been associated with the accumulation of small signaling molecules, such as ppGpp, microbial hormones, and late intermediates or end-products of the secondary metabolite biosynthetic pathways Ruiz et al. 2008; O'Rourke et al. 2009; Hsiao et al. 2009; Wang et al. 2009. The influence of all aforementioned factors in most cases is reflected to

the activity of the pathway-specific regulatory genes, which are believed to be final checkpoints in the onset of antibiotic production Arias et al. 1999; Nuria et al. 2007; van Wezel et al. 2009; Pulsawat et al. 2007; Wang et al. 2009. Because most antibiotics are potentially lethal to the producing organism, the onset of antibiotic production should be under tight control and mechanisms of self-resistance of producing bacteria must exist. All this requires a precise regulatory network coordinating both, biosynthesis and resistance genes expression Le et al. 2009. That is why very often resistance genes are linked to antibiotic biosynthesis genes Tahlan et al. 2007; Ostash et al. 2008. As our understanding of secondary metabolism advances, it is becoming clear that the relationship between antibiotic production and resistance is more complicated than expected. For example, in *S. coelicolor*, along with the mature antibiotic(s), intermediates of the biosynthetic pathway might activate expression of the export genes, thereby coupling resistance to biosynthesis Hopwood 2007. In *S. cyanogenus* intermediates are able, not only to release repression of the export machinery, but also to de-repress expression of the late biosynthetic enzymes that attach the final sugars to yield mature landomycin A Ostash et al. 2008. However, despite the identification and characterization of numerous genes, which

* Correspondence: andreas.bechthold@pharmazie.uni-freiburg.de

³Institut für Pharmazeutische Wissenschaften, Lehrstuhl für Pharmazeutische Biologie und Biotechnologie, Albert-Ludwigs-Universität Freiburg, Stefan-Meier-Strasse 19, 79104 Freiburg, Germany

Full list of author information is available at the end of the article

affect antibiotic production and resistance, our understanding of the regulatory networks that govern these processes is far from complete.

A biosynthetic gene cluster usually contains at least one regulatory gene Sheldon et al. 2002; Rebets et al. 2003; Rebets et al. 2008; Chen et al. 2008. This is also the case for the gene cluster of the aminocoumarin antibiotic simocyclinone D8 (Figure 1), produced by *S. antibioticus* Tü6040, that has distinct cytostatic and antibiotic activities Trefzer et al. 2002; Galm et al. 2002; Oppedard et al. 2009; Sadig et al. 2010; Edwards et al. 2009. The simocyclinone biosynthetic gene cluster includes three putative regulatory genes: *simReg1*, *simReg2* (hereafter *simR*) and *simReg3* (Figure 2). Recently, the function of SimR was investigated *in vitro* and it was shown to repress the transcription of *simX* gene that encodes simocyclinone efflux pump Le et al. 2009; Le et al. 2011, whereas the function of the two other regulators is still unknown. SimReg1 is the first example of an OmpR-PhoB subfamily regulator, identified in an aminocoumarin biosynthetic gene cluster. Herein, we describe the generation and analysis of the mutant strain deficient in the *simReg1* gene, mobility shift DNA-binding assays of His-SimReg1 to putative promoter regions and propose a putative model for regulation of the biosynthesis and export of simocyclinones.

Materials and methods

Bacterial strains, plasmids, and culture conditions

All strains and plasmids are listed in Table 1. *E. coli* DH5 α (Life Technologies) was used for routine subcloning. *E. coli* ET12567 harboring the conjugative plasmid pUB307 (provided by C. P. Smith, UMIST, Manchester, UK) was used to perform intergeneric conjugation from *E. coli* to *Streptomyces* species Flett et al. 1997; Luzhetskyy et al. 2006. For plasmid and total DNA isolation, *E. coli* and *S. antibioticus* strains were grown as described by Sambrook and Russell (2001), and Kieser et al. (2000). For simocyclinone production, *S. antibioticus* strains were grown in liquid NL5 medium (NaCl 1 g l⁻¹, KH₂PO₄ 1 g l⁻¹, MgSO₄ × 7H₂O 0.5 g l⁻¹, glycerol 25 g l⁻¹, L-glutamin - 5.84 g l⁻¹, trace elements - 2.0 ml, pH 7.3 prior to sterilization) at 30°C. For conjugation, spores of

S. antibioticus strains were harvested from a sporulated lawn grown on soya-mannitol or oatmeal medium Kieser et al. 2000; Luzhetskyy et al. 2006. When it was necessary, bacterial strains were grown in the presence of respective antibiotics. X-gal and IPTG were used for blue-white colony selection in the case of the pBluescript, pSET152, pKC1139, pKC1218E vectors as described elsewhere Kieser et al., 2000; Sambrook et al., 2001.

DNA manipulations

Isolation of genomic DNA from streptomycetes and plasmid DNA from *E. coli* were carried out using standard protocols Kieser et al. 2000. Restriction enzymes and molecular biology reagents were used according to the recommendation of suppliers (NEB, MBI Fermentas, Promega). DIG DNA labeling and Southern hybridization analyses were performed according to the DIG DNA labeling and detection kit (Roche Applied Science).

Construction of the plasmid for *simReg1* inactivation

A 4.3 kb BamHI fragment carrying the entire *simReg1* gene and its flanking regions was cloned from 5JH10 (Table 1) into pUC19 to yield pUCsimR1 with a unique BsaAI site within the coding region of the *simReg1* gene. The plasmid pUCsimR1 was digested with BsaAI and ligated to the hygromycin resistance cassette *hyg*, retrieved as an EcoRV fragment from pHYG1 (Table 1). The resulting plasmid pUCsimR1-*hyg* was digested with BamHI and the fragment containing the *simReg1::hyg* mutant allele was cloned into the shuttle vector pKC1139 to yield pKCsirR1-*hyg*.

Generation of the chromosomal mutant *S. antibioticus* Δ *simReg1*

The gene disruption plasmid pKCsirR1-*hyg* was conjugally transferred from *E. coli* into *S. antibioticus* Tü6040. Exconjugants were selected for resistance to apramycin (10 μ g ml⁻¹). To generate *S. antibioticus* Δ *simReg1* strain, single-crossover mutants were obtained by cultivation of the respective exconjugants at 39°C for 3 days with a further screen for the loss of apramycin resistance as a consequence of a secondary crossover.





Figure 1 Structure of simocyclinone D8.



Complementation of the *simReg1* mutant

The *simReg1* gene with flanking regions was retrieved from the plasmid pKCEsimR1 Rebets et al. 2008 as a 2.3 kb BamHI fragment and cloned into the BamHI sites of pSET152 to yield pSsimR1. A 1.4 kb SmaI fragment harboring only *simReg1* with its promoter region was retrieved from pSsimR1 and cloned into EcoRV linearized pSET152 to yield pSsimR1-1.

Construction of the plasmids for *gusA* reporter fusion system

A 0.5 kb DNA fragment, containing promoter of the *simD4* gene (P_{D4}) was amplified from the chromosome *S. antibioticus* Tü6040 using primers *simD4_for_script* and *simD4_rev_script* (Table 2). The PCR product was digested with XbaI/KpnI and cloned into the respective sites of pGUS Myronovskyi et al. 2011, giving pSimD4-script. In this plasmid transcription of the *gusA* gene is under the control P_{SD4} promoter.

A 0.8 kb fragment, carrying the *simReg1* gene, was amplified from the *S. antibioticus* Tü6040 chromosome using the primers *simReg1_for* and *simReg1_rev* (Table 2). The amplified DNA fragment was cleaved with HindIII/BamHI and cloned into the respective sites of pUWL-oriT (Table 1), yielding pUWLsimReg1. In this plasmid the *simReg1* gene is under the control of P_{ermE} .

Spectrophotometric measurement of glucuronidase activity in cell lysates

For measurement of GusA activity, mycelium of the *S. lividans* strain harboring both pSimD4script and

pUWLsimReg1 plasmids, the control strains *S. lividans* 1326 × pSimD4script, *S. lividans* 1326 × pGUS, and *S. lividans* 1326 × pGUS/pUWLsimReg1 were grown in liquid TSB medium (100 ml) for 2 days at 30°C in a rotary shaker (180 rpm). 1 ml of the pre-culture was inoculated into liquid TSB medium (100 ml) and grown for 5 days at 30°C in a rotary shaker. Mycelium was harvested, washed with distilled water, then resuspended in lysis buffer (50 mM phosphate buffer [pH 7.0], 0.1% Triton X-100, 5 mM DTT, 4 mg ml⁻¹ lysozyme) and incubated for 30 min at 37°C. Lysates were centrifuged for 10 min at 5000 rpm. Then, 0.5 ml of lysate was mixed with 0.5 ml of dilution buffer (50 mM phosphate buffer [pH 7.0], 5 mM DTT, 0.1% Triton X-100) supplemented with 5 μl 0.2 M *p*-nitrophenyl-β-D-glucuronide and used for measuring optical density at λ = 415 nm every minute during 20 min of incubation at 37°C. As a reference, a 1:1 mixture of lysate and dilution buffer was used.

Analysis of secondary metabolites production

Streptomyces strains were grown in liquid TSB medium (50 ml) for 2 days at 30°C in a rotary shaker (180 rpm). Five ml of the pre-cultures were inoculated into liquid NL5 medium (100 ml) and the cultures were grown for 5 days at 30°C in a rotary shaker. The culture broths were extracted three times with 100 ml of ethyl acetate. The extracts were dried in vacuum and dissolved in methanol (200-400 μl). The metabolites were analyzed by high-pressure liquid chromatography-mass spectrometry (HPLC-MS) Schimana et al. 2001. 10 ml of each culture

Table 1 Strains and plasmids

Bacterial strains and plasmids	Description	Source or reference
<i>E. coli</i> DH5 α	<i>supE44</i> Δ <i>lacU169</i> (ϕ 80 <i>lacZ</i> Δ M15)	Hanahan 1985
<i>E. coli</i> BL21 (DE3) pLysS	Host for the heterologous expression of His ₆ -tagged <i>simReg1</i>	Novagen
<i>E. coli</i> ET12567/pUB307	<i>hsdR17 recA1endA1gyrA96 thi-1 relA1 dam-13::Tn9</i> (Cmr) <i>dcm-6 hsdM1</i> ; harbors conjugative plasmid pUB307; Cm ^r , Km ^r	Flett et al. 1997
<i>S. antibioticus</i> Tü6040	Simocyclinone D8 producing strain	Trefzer et al. 2002
<i>S. antibioticus</i> Δ simReg1	Derivative of <i>S. antibioticus</i> Tü6040 with disrupted <i>simReg1</i> gene	This work
<i>S. antibioticus</i> Δ simReg1 \times pSSimR1-1	Δ simReg1 strain carrying plasmid with the intact <i>simReg1</i> gene under its own promoter, used for complementation studies	This work
<i>S. lividans</i> 1326	Wild type	Hopwood et al. 1985
<i>S. lividans</i> \times pSimD4script	Derivative of <i>S. lividans</i> 1326 carrying plasmid with <i>gusA</i> gene under the control of the putative promoter of the <i>simD4</i> gene	This work
<i>S. lividans</i> \times pSimD4script/pUWLSimReg1	Derivative of <i>S. lividans</i> 1326 carrying plasmid with <i>gusA</i> gene under the control of putative promoter of the <i>simD4</i> gene and second plasmid with <i>simReg1</i> gene under the control of P _{ermE}	This work
<i>S. lividans</i> xpGUS	Derivative of <i>S. lividans</i> 1326 carrying plasmid with promoterless reporter gene <i>gusA</i>	This work
<i>S. lividans</i> \times pGUS/pUWLSimReg1	Derivative of <i>S. lividans</i> 1326 carrying plasmid with promoterless reporter gene <i>gusA</i> and plasmid with <i>simReg1</i> gene under the control of the P _{ermE} promoter	This work
pBluescriptIIKS ⁺	General purpose cloning vector; Ap ^r	MBI Fermentas
pUC19	General purpose cloning vector; Ap ^r	MBI Fermentas
pSET152	<i>E. coli</i> / <i>Streptomyces</i> shuttle vector with ϕ C31 integration system for streptomycetes; Am ^r	Bierman et al. 1992
pKC1218E	pKC1218 derivative expression vector with P _{ermE} promoter and SCP2* replicon; Am ^r	Ostash et al. 2004
pHYG1	pLitmus38 containing hygromycin resistance cassette <i>hyg</i>	C. Olano Univ. de Oviedo, Spain
pKC1139	<i>E. coli</i> / <i>Streptomyces</i> shuttle vector with temperature sensitive replicon pSG5, Am ^r	Muth et al. 1989
pUWL-oriT	pUWL-KS derivative harboring <i>oriT</i> from pSET152	Zelyas et al. 2009
pET21d	Vector for His-tagged protein expression	Novagen
5JH10	pUC plus <i>simB3-D4</i> segment	Trefzer et al. 2002
pUCsimR1	pUC19 derivative containing <i>simReg1</i> gene	This work
pUCsimR1-hyg	pUCsimR1 derivative with <i>hyg</i> cassette cloned into the <i>simReg1</i> coding region	This work
pKCSimR1-hyg	pKC1139 derivative with cloned <i>simReg1::hyg</i> construction used for <i>simReg1</i> gene inactivation	This work
pKCEsimR1	pKCE1218 derivative containing <i>simReg1</i> gene under the control of P _{ermE}	Rebets et al. 2008
pSSimR1	pSET152 plus 2.3 kb <i>simD4-X1</i> segment	This work
pSSimR1-1	pSET152 derivative containing <i>simReg1</i> gene under the control of its own promoter	This work
pMA-simR1	plasmid containing synthetic codon-optimized <i>simReg1</i> gene	Mr. Gene, Heidelberg
pETSR1c-15	pET21d derivative containing synthetic codon-optimized <i>simReg1</i> gene	This work
pGUS	pSET152 derivative containing promoterless reporter gene <i>gusA</i>	Myronovskyi et al. 2011
pSimD4script	derivative of pGUS harboring <i>gusA</i> reporter gene under the promoter of the <i>simD4</i> gene	This work
pUWLSimReg1	derivative of pUWL containing gene <i>simReg1</i> under the control of P _{ermE}	This work

were taken and lyophilized. The dry weight of each sample was measured. In all cases amounts of antibiotic were referred back to equal amounts of biomass (dry weight) and are mean values from at least three independent experiments.

Overexpression of SimReg1

The codon-optimized copy of the *simReg1* gene, named *simReg1s*, was synthesized by Mr. GENE Company (Heidelberg, Germany) and was provided on the plasmid pMA-simR1. Gene *simReg1s* was amplified from pMA-simR1 using primers SSR1F and SSR1R (Table 2). The

PCR product was cloned into the pET21d NcoI-EcoRI sites, giving pETSR1c-15.

E. coli BL21(DE3) (pLysS) harboring the pETSR1c-15 plasmid was grown overnight at 37°C. LB (400 mL) containing 50 μ g/mL of ampicillin was inoculated with 2 mL of the overnight culture and incubated at 21°C until the OD_{600 nm} reached 0.7. SimReg1 expression was induced with 1 mM IPTG. After incubation for an additional 16 h, the cells were harvested by centrifugation and washed with ice-cold column buffer (20 mM Tris-HCl [pH 8.0], 50 mM NaCl). Cell lysis and purification of SimReg1 with His-tag-binding resins were performed according to

Table 2 Primers used in this study

Primer	Nucleotide sequence (5'-3')	Purpose	Gene name
SSR1F	ATACCATGGCCCGTGAACGT	SimReg1	<i>simReg1</i>
SSR1R	TTTGAATTCATTAATGGTGATGGT	purification	
SR1D4F	TAGAATTCGTGAGCAGATCATGT	DNA-shift assay	P_{D4}
SR1D4R	TAGAATTCATTGTGAACCATC		
SD2R1F	TAGAATTCATCGCCACGACCATG	DNA-shift assay	P_{R1}
SD2R1R	TAGAATTCGCGGTTTCGGCAGA		
simX5D3F	TAGAATTCGTACAAGGCCTGGT	DNA-shift assay	P_{D3}
simX5D3R	TAGAATTCGCGACAGGAGCCATA		
simEXX4F	TAGAATTCGACGCCTTCCAGTC	DNA-shift assay	P_{X4}
simEXX4R	TAGAATTCAGAACATCGTCC		
SR2ExXF	AAATCTAGATCAAGCCAGTGCTG	DNA-shift assay	P_{R2Ex}
SR2ExXR	TTTGAATTCCTTGACCACCACTTC		
PSR2F	TCGACGAGGTCCTCTTTG	DNA-shift assay	P_{SR2}
PSR2R	TCGTATTCATACACCGTAC		
PEX1F	CCAATTGCGCTACGCTCCT	DNA-shift assay	P_{SEx1}
PEX1R	CCATGTAGGCGGTGACGA		
simA7F	TAAAGCTTCAAACGGGGTGAAC	DNA-shift assay	P_{A7}
simA7R	ATAAGCTTGTGCGATACCGATCTTC		
PEX2F	ACTTCCCAGAAGTA	DNA-shift assay	P_{Ex2}
PEX2R	AGAGGGCAGTAGAC		
PR3F	TTTCTAGATGCACCCGATCCTC	DNA-shift assay	P_{SR3}
PR3R	GAACAGGATTCGCATGAGTACT		
D4For	TATTGGTCGCGCAGTCGTCC	DNA-shift assay	part of the <i>simD4</i> gene
D4Rev	TTGTCCTGCATCATGACGAG		
simD4_for_script	AAATCTAGAGGCGACCGACCCCG GCCGAG	<i>simD4</i> promoter cloning	P_{D4}
simD4_rev_script	AAAGGTACCCGATCCGGCTGGCA TTACTG		
simReg1_for	AAAAAGCTTTACCTGTATCCAGGGC GGGCACTT	<i>simReg1</i> cloning	<i>simReg1</i>
simReg1_rev	AAAGGATCCGCACAAGCGGCAGC AATCG		

Novagen instructions. SimReg1 was eluted with column buffer containing 200 mM imidazole. The purest fractions (as determined by SDS-PAGE and Coomassie blue staining) were pooled, washed with storage buffer (50 mM potassium phosphate [pH 8.0], 300 mM NaCl, 10% glycerol), concentrated using Amicon Ultra (Millipore). Aliquots of SimReg1 fusion protein in storage buffer were stored at - 80°C, or used immediately in DNA-binding assays.

Electrophoretic mobility shift DNA-binding assays (EMSA)

DNA fragments containing putative promoters of *simD4* (P_{D4} , 513 bp), *simReg1* (P_{R1} , 490 bp), *simD3* (P_{D3} , 300 bp), *simX4* (P_{X4} , 350 bp), *simA7* (P_{A7} , 300 bp), *simEx2* (P_{Ex2} , 550 bp), *simB7* (P_{SR3} , 319 bp), *simX* (P_{SEx1} , 280 bp), *simR* (P_{SR2} , 300 bp), and the putative promoter region between *simX* and *simR* genes (P_{R2Ex} , 780 bp) (Figure 2) were used in EMSA. Indicated promoter regions were amplified from

the chromosomal DNA of *S. antibioticus* using primer pairs listed in Table 2. Each EMSA contained 50 ng of a target DNA and 0.9 µg, 1.8 µg, 2.7 µg, 3.6 µg, 4.5 µg of the His-SimReg1 protein in a total volume of 20 µL in a binding buffer (20 mM Tris HCl [pH 8.0], 1 mM EDTA, 1 mM DTT, 100 mM KCl, 1 mM MgCl₂, 10% glycerol). After incubation for 25 min at room temperature, protein-bound and free DNA were separated by electrophoresis at 4°C on a 4.5% nondenaturing polyacrylamide gel in 0.5 × TBE buffer. The gel was stained with ethidium bromide and analyzed using a UV-imaging system (Fluorochem 5330). A negative control assay was carried out in the presence of the part of the *simD4* coding region, amplified with the use of primers D4For and D4Rev (Table 2). Extracts from the strain *S. antibioticus* Tü6040 × pSsimR1-1, containing more than 95% of simocyclinones (Additional file 1), dissolved in methanol (5% and 10% - final volume in a reaction mixture) were tested as SimReg1 ligands.

Results

In silico analysis of the *simReg1* gene product

The putative product of the *simReg1* gene is a 251 aa protein with a molecular mass 27.94 kDa. As evident from BLAST and CDD search results, putative amino acid sequence of the protein has significant similarity to response regulators in two component control systems. The closest homologues of SimReg1 are proteins that act as positive regulators for angucycline-like biosynthesis, including JadR1 from *S. venezuelae* (60% similarity) Wang et al. 2009, LanI from *S. cyanogenus* (58% similarity) Rebets et al. 2008 and LndI from *S. globisporus* (58% similarity) Rebets et al. 2003; Rebets et al. 2005. Analysis of the SimReg1 amino acid sequence using ExPASy Proteomics Server <http://expasy.org> revealed a putative signal receiver domain (the REC domain, aa 15-123) located at the N-terminal part of the protein and a DNA-binding domain at the C-terminus (aa 167-239). The latter is predicted to interact with short conserved regions of the target DNA and with the RNA polymerase. The secondary structure of the C-terminal DNA-binding domain of SimReg1 was similar to that of OmpR (*E. coli*) and PhoB (*E. coli*), which adopt a winged helix-turn-helix (HTH) moiety. In the REC domain of the regulatory protein PhoB, six conserved amino acid residues are believed to be vital for phosphorylation and consequence response Sola-Landa et al. 2003; Wang et al. 2009; Dyer and Dahlquist 2006, but only three of them are present in SimReg1 (Figure 3). Also, no protein kinase encoding genes have been found within the *sim* cluster. Thus, we suppose that SimReg1 belongs to “atypical” response regulators (ARR), like its close homolog JadR1 Wang et al. 2009.

S. antibioticus Δ simReg1 mutant is deficient in simocyclinone production

In order to investigate the function of *simReg1*, the chromosomal copy of the gene was replaced by the mutant allele containing a hygromycin resistance cassette (*hyg*) (Figure 4a). Inactivation of the *simReg1* gene was proven by Southern hybridization. BamHI digested chromosomal DNA of the wild type and *S. antibioticus* Δ simReg1 strains were probed with the DIG-labeled fragment containing *simReg1*, obtained as a KpnI fragment from the plasmid pKCEsimR1 Rebets et al. 2008. A single hybridization signal of the expected size (4.3 kb) was detected in the case of the wild type strain and a 6.3 kb fragment was detected in the Δ simReg1 mutant (Figure 4b). The *S. antibioticus* Δ simReg1 mutant had growth and morphological characteristics identical to those of the wild type. HPLC and TLC analysis (Figure 5a) of the extracts from the mutant strain Δ simReg1 revealed no simocyclinone and its precursors, indicating that this gene is essential for antibiotic production.

To exclude any possibility of polar effects and to confirm that the cessation of simocyclinone production was caused by the inactivation of the *simReg1*, complementation experiment was carried. For this purpose, we used the pSSimR1-1 plasmid (Table 1), which contains the *simReg1* gene under its own promoter cloned in the integrative vector pSET152. This plasmid was transferred into *S. antibioticus* wild type strain by means of conjugation. The recombinant strain *S. antibioticus* Δ simReg1 \times pSSimR1-1 was found to accumulate simocyclinone at a level comparable to those of the wild type (Figure 5b).

It is known that very often overexpression of the positive pathway-specific regulators lead to overproduction of antibiotics (Bibb 2005; Novakova et al., 2011). To analyze the effect of additional copies of *simReg1* gene on simocyclinone biosynthesis, we introduced the plasmid pSSimR1-1 that contains *simReg1* gene under its own promoter, into the wild type strain. Recombinant strain *S. antibioticus* Tü6040 \times pSSimR1-1 produced in average 2.5 times more simocyclinone than the wild type.

SimReg1 binds to the putative promoter regions of structural, transporter genes and its own gene

In order to prove the DNA binding activity of SimReg1, gel mobility-shift assays were carried out. His-SimReg1 was purified (Additional file 2) and an *in vitro* binding assay was performed using His-SimReg1 and DNA fragments containing putative promoters of the regulator gene *simReg1* (P_{R1}), the 3-keto-acyl-reductase gene *simD4* (P_{D4}), the oxygenase gene *simA7* (P_{A7}), the transporter gene *simEx2* (P_{Ex2}), the 3-keto-acyl-reductase gene *simD3* (P_{D3}), the putative gene *simX4* (P_{X4}), the putative oliviosyl-transferase gene *simB7* (P_{SR3}), and the intergenic region between *simR* and the transporter gene *simEx1* (hereafter *simX*) (P_{R2Ex}) (Figure 2). Shifted bands were detected using the promoter regions of the enzyme encoding genes (Figure 6a, c, d, f), the transporter gene *simEx2* (Figure 6g) and the regulatory gene *simReg3*, which is likely co-transcribed with the genes *simB7*, *simB5*, *simB4*, *simX5* and *simX7* (Figure 6h). Furthermore, DNA retardation occurred (Figure 6b) when the promoter of the *simReg1* gene was used in the binding assay, indicating that SimReg1 is an autoregulatory protein. We carried out a set of control assays to demonstrate the specificity of the SimReg1 binding. For instance, none of the compounds in the crude extract of *E. coli* BL21(DE3) binds to any of the putative promoters (data not shown). We also showed that randomly chosen DNA did not interact with SimReg1 (Additional file 3).

SimReg1 was found to bind to the DNA fragment containing the *simR/simX* intergenic region (Figure 6e). However, it was not known whether SimReg1 interacts with the promoters of both genes. A 67 bp fragment




Figure 3 Amino acid sequence comparison of the SimReg1 and PhoB (*E. coli*) proteins. The conserved amino acids which are important for phosphorylation and consequence response are shaded in grey (aa that differ in proteins) and dark grey (aa that are identical in both sequences).

located in front of the start codon of *simR* (P_{SR2}) and a 69 bp fragment located in front of *simX* (P_{SEx1}) (Figure 7a) were used for additional EMSA analysis. No binding was identified with the P_{SR2} promoter, whereas DNA retardation occurred when the P_{SEx1} promoter was used in the assay (Figure 7b). These results indicate that SimReg1 is capable of binding to the promoter region of *simX*.

Effect of culture extracts from *S. antibioticus* Tü6040 × pSSimR1-1 on the activity of SimReg1

Since DNA binding ability of JadR1, which also belongs to ARR and is very similar to SimReg1 (60% similarity), is regulated by jadomycin B Wang et al. 2009, we studied the effects of simocyclinone extracts from the *S. antibioticus* Tü6040 × pSSimR1-1 on the DNA binding activity of SimReg1. For this purpose the culture broth of *S. antibioticus*







Figure 4 Inactivation of the *simReg1* gene. (a) Schematic representation of the *simReg1* gene disruption. (b) Results of the Southern hybridization of KpnI-digested plasmid pKcesimR1 (1), BamHI digested total DNA samples from *S. antibioticus* Δ simReg1 (2, 3) and Tü6040 (4) with 1.4 kb SmaI fragment harboring *simReg1* gene.



Tü6040 \times pSSimR1 strain grown for 72 hours was extracted with an ethyl acetate, dried and dissolved in methanol. In overall the percentage of different types of simocyclinone in such an extract was more than 95% (Additional file 1). Presence of these extracts could

dissociate His-SimReg1 from the promoter regions P_{RI} and P_{A7} , as a result no shifted bands occurred (Figure 8). This effect was not due to methanol, the simocyclinone D8 solvent, as equivalent amounts of methanol had no effect on His-SimReg1-DNA complex formation (Figure 8).






SimReg1 activates expression of a *gusA* reporter gene from P_{D4} promoter

On the basis of the gene inactivation, overexpression and EMSA results we suppose that SimReg1 is a positive regulator of simocyclinone production. To investigate whether SimReg1 can activate the expression of the structural genes under heterologous conditions, a reporter system on the basis of *gusA* was used. For these purpose, we constructed two plasmids pSimD4script and pUWLsimReg1 (Table 1). In the first plasmid the

promoter region of the putative ketoreductase gene *simD4* (P_{D4}) was fused with the coding sequence of the *gusA* gene. As a result expression of the reporter *gusA* is controlled by P_{D4} . In the plasmid pUWLsimReg1 intact gene *simReg1* was cloned under the control of erythromycin resistance gene promoter to make the expression of the regulatory gene constitutive. As it is evident from the EMSA analysis SimReg1 binds to the promoter of the gene *simD4* (Figure 6a) this means that SimReg1 should influence expression from this promoter. To



verify this assumption, both plasmids were transferred into heterologous host *S. lividans* 1326 to avoid influence of two other regulatory proteins SimR and SimReg3 Trefzer et al., 2002. We obtained two strains: *S. lividans* harboring only pSimD4script and *S. lividans* harboring both plasmids pSimD4script and pUWLsimReg1. As a negative control we used strains: *S. lividans* 1326 × pGUS to show that there is no GusA activity when *gusA* gene contains no promoter and *S. lividans* 1326 harboring both plasmids pGUS (Table 1) and pUWLsimReg1 to demonstrate that SimReg1 specifically binds only to *simD4* promoter region and that SimReg1 can't influence *gusA* expression in the absence of this promoter. Aforementioned four strains were grown in liquid TSB medium for 5 days and samples of the strains were used for GusA activity measurement as described in Materials and Methods. In the control strains the activity of GusA was approximately 0.25 ± 0.06 (Figure 9). In the case of the *S. lividans* strain that contains *gusA* gene under P_{D4} activity was in average 3.3 ± 0.24 (Figure 9). In the strain containing both *gusA* gene under P_{D4} promoter and the SimReg1 protein the activity was 6.25 ± 0.43 (Figure 9). It is in overall two times higher than without the protein. On the basis of these results, we may conclude that SimReg1 binds to the *simD4* promoter region.

Discussion

Simocyclinone is a potent antibacterial drug that inhibits DNA gyrase supercoiling Oppegard et al. 2009; Sadig et al. 2010; Edwards et al. 2009; Sissi et al. 2009. The gene cluster responsible for simocyclinone production was cloned and biosynthetic, and regulatory genes were detected Trefzer et al. 2002; Galm et al. 2002. Here, we report on the

function of the gene *simReg1* involved in the regulation of simocyclinone production and export.

SimReg1, to our knowledge, is the first OmpR-PhoB subfamily regulator identified within aminocoumarin biosynthetic gene clusters. It appears to be a key regulator of simocyclinone production since inactivation of *simReg1* completely abolished antibiotic biosynthesis and its overexpression in the wild type strain *S. antibioticus* Tü6040 led to almost 2.5 times increase in simocyclinone production. *In silico* analysis and DNA shift assays showed that SimReg1 is a DNA-binding autoregulatory protein that interacts directly with putative promoter regions of the structural *sim* genes, both transporter genes *simX* and *simEx2*, and the putative regulatory gene *simReg3*. Our results indicate that SimReg1 is an activator of the structural and transporter genes transcription, as expression of the reporter gene *gusA* under P_{D4} in the presence of SimReg1 was at least two times higher, than without it. DNA-binding activity of SimReg1 is abolished in the presence of extracts from *S. antibioticus* Tü6040 × pSSimR1-1. As extracts used in the experiment were enriched with simocyclinones, these might indicate the existence of autoregulation by binding most likely simocyclinone or its intermediates. However to establish this assumption additional experiments are required. Similar autoregulation by binding of the end product was described for JadR1 Wang et al. 2009, the close homolog of SimReg1. An interesting finding is that SimReg1 binds to the promoter region of the exporter gene *simX*. SimR is known to repress expression of *simX* and its own gene by binding to two distinct operators within the *simR/simX* intergenic region Le et al. 2009. SimR was shown to dissociate from the *simX* promoter in the presence of simocyclinone D8 Le et al. 2009; Le et al. 2011a; Le et al.




Figure 9 Glucuronidase activity in cell lysates of *S. lividans* strains: 1 - *S. lividans* × pSimD4script; 2 - *S. lividans* × pGUS; 3 - *S. lividans* × pSimD4script/pUWLsimReg1; 4 - *S. lividans* × pGUS/pUWLsimReg1.

2011b. At the same time SimReg1 is interacting with the 69 bp DNA region upstream to the start codon of *simX*. This means that the operator of SimReg2 partially overlaps with the DNA-binding region of SimReg1. Therefore, it is very likely that in the presence of simocyclinone dissociation of SimReg2 from the promoter region of *simX* is necessary for SimReg1 binding indicating that SimReg1 and SimReg2 compete for the binding to the *simX* promoter.

The presence of distinct regulatory proteins indicates the importance for the cell to strongly control simocyclinone production and transport. The structure of simocyclinone is assembled from products of three distinct biosynthetic routes. To produce such a complex molecule the biosynthetic pathway and the transport have to be precisely tuned and controlled.

Based on our data and the data described by Buttner and coworkers Le et al. 2009; Le et al. 2011a; Le et al. 2011b, we proposed the following preliminary model for the regulation of simocyclinone biosynthesis and export. When the concentration of simocyclinone and/or its intermediates is low the transcription of the exporter gene *simX* is repressed by SimR. At the same time, SimReg1, being the key regulator of simocyclinone biosynthesis, activates expression of the structural *sim*-genes and simocyclinone production. When the cellular concentration of simocyclinone exceeds a certain level, SimR is released from P_{SE_X} that allows SimReg1 to bind to the promoter. This activates *simX* expression, followed by the transport of simocyclinones out of the cell. This mechanism couples the biosynthesis of simocyclinone to its export. In such a way, an additional mechanism of exact tuning of biosynthesis level is exerted ensuring the protection of the producing bacteria from the toxicity of its secondary metabolism product.

The present study portrays a strong link between antibiotic production and export and describes for the first time the function of the atypical response regulator in the control of the biosynthesis of simocyclinone. Furthermore, our data suggest a useful biotechnological approach for optimization of simocyclinone production, as overexpression the gene encoding positive regulator SimReg1 leads to antibiotic overproduction.

Additional material

Additional file 1: HPLC analysis of secondary metabolites produced by *S. antibioticus* Tü6040 × pSSimR1-1. On axis y relative absorption units (AU) are plotted. On axis x retention time of compounds is plotted (in min). Under conditions stated SD8 has R_t of 24.7 min. The overall content of simocyclinones in the extract was around of 95%.

Additional file 2: Purification of the His-tagged SimReg1 protein from *E. coli* BL21 (DE3). Lane 1: molecular mass marker (Pierce Protein Research Products); lane 2: flow through; lane 3: purified SimReg1.

Additional file 3: Results of EMSA performed to detect interactions of SimReg1 to part of the *simD4* gene. Lane 1: *simD4*; lane 2: *simD4* + His-SimReg1; lane 3: *simD4* + His-SimReg1; lane 4: *simD4* + His-SimReg1.

Acknowledgements

This work was supported by the DAAD, grant to L.H. (PKZ A/07/99406) and by the BMBF (grant to A.B.)

Author details

¹Department of Genetics and Biotechnology of Ivan Franko National University of L'viv, Grushevskogo st.4, L'viv 79005, Ukraine ²Helmholtz-Institute for Pharmaceutical Research Saarland (HIPS), Helmholtz Center for Infectious Research (HZI), Department Microbial Natural Products Actinobacteria, Metabolic Engineering Group, Saarland University, Campus C2 3 66123 Saarbrücken, Germany ³Institut für Pharmazeutische Wissenschaften, Lehrstuhl für Pharmazeutische Biologie und Biotechnologie, Albert-Ludwigs-Universität Freiburg, Stefan-Meier-Strasse 19, 79104 Freiburg, Germany

Competing interests

The authors declare that they have no competing interests.

Received: 21 November 2011 Accepted: 3 January 2012

Published: 3 January 2012

References

- Arias P, Miguel A, Fernández-Moreno A, Malpartida F (1999) Characterization of the Pathway-Specific Positive Transcriptional Regulator for Actinorhodin Biosynthesis in *Streptomyces coelicolor* A3(2) as a DNA-Binding Protein. *J Bacteriol* 181:6958–6968
- Bibb M, Hesketh A (2009) Analyzing the regulation of antibiotic production in *Streptomyces*. *Method Enzymol* 458:93–116
- Bierman M, Logan R, O'Brein K, Seno E, Rao R, Schoner B (1992) Plasmid cloning vectors for the conjugal transfer of DNA from *Escherichia coli* to *Streptomyces* spp. *Gene* 116:43–49. doi:10.1016/0378-1119(92)90627-2.
- Blondelet-Rouault M-H, Weiser Y, Lebrühi A, Branny P, Pernodet JL (1997) Antibiotic resistance gene cassettes derived from the U' interposon for use in *Escherichia coli* and *Streptomyces*. *Gene* 190:315–317. doi:10.1016/S0378-1119(97)00014-0.
- Chen L, Lu Y, Chen J, Zhang W, Shu D, Qin Z, Yang S, Jiang W (2008) Characterization of a negative regulator Avel for avermectin biosynthesis in *Streptomyces avermitilis* NRRL8165. *Appl Microbiol Biotechnol* 80:277–286. doi:10.1007/s00253-008-1545-8.
- Dyer CM, Dahlquist FW (2006) Switched or not: the structure of unphosphorylated CheY bound to the N terminus of FliM. *J Bacteriol* 188:7354–7363. doi:10.1128/JB.00637-06.
- Edwards MJ, Flatman RH, Mitchenall LA, Stevenson CE, Le TB, Clarke TA, McKay AR, Fiedler H-P, Buttner MJ, Lawson DM, Maxwell A (2009) A crystal structure of the bifunctional antibiotic simocyclinone D8, bound to DNA gyrase. *Science* 326:1415–1458. doi:10.1126/science.1179123.
- Flett F, Mersinias V, Smith C (1997) High efficiency intergeneric conjugal transfer of plasmid DNA from *Escherichia coli* to methyl DNA-restricting *Streptomyces*. *FEMS Microbiol Lett* 155:223–229. doi:10.1111/j.1574-6968.1997.tb13882.x.
- Galm U, Schimana J, Fiedler H-P, Schmidt J, Li SM, Heide L (2002) Cloning and analysis of the simocyclinone biosynthetic gene cluster of *Streptomyces antibioticus* Tü6040. *Arch Microbiol* 178:102–114. doi:10.1007/s00203-002-0429-z.
- Hanahan D (1985) Techniques for transformation of *Escherichia coli*. In: Glover DM (ed) *DNA Cloning*. Oxford, UK: IRL Press pp 109–135
- Hopwood DA, Bibb MJ, Chater KF, Kieser T, Bruton CJ, Kieser HM, Lydiate DJ, Smith CP, Ward JM, Schrempf H (1985) Genetic manipulation of *streptomyces*: A laboratory manual. Norwich: The John Innes Foundation
- Hopwood DA (2007) How do antibiotic-producing bacteria ensure their self-resistance before antibiotic biosynthesis incapacitates them? *Mol Microbiol* 63:937–940. doi:10.1111/j.1365-2958.2006.05584.x.
- Horinouchi S (2007) Mining and polishing of the treasure trove in the bacterial genus *Streptomyces*. *Biosci Biotechnol Biochem* 71:283–299. doi:10.1271/bbb.60627.

- Hsiao N-H, Gottelt M, Takano E (2009) Regulation of antibiotic production by bacterial hormones. *Meth Enzymol* 458:143–157
- Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA (2000) *Practical Streptomyces genetics*. The John Innes Foundation, Norwich
- Le TB, Fiedler H-P, den Hengst CD, Ahn SK, Maxwell A, Buttner MJ (2009) Coupling of the biosynthesis and export of the DNA gyrase inhibitor simocyclinone in *Streptomyces antibioticus*. *Mol Microbiol* 72:1462–1474. doi:10.1111/j.1365-2958.2009.06735.x
- Le TB, Stevenson CE, Buttner MJ, Lawson DM (2011) Crystallization and preliminary X-ray analysis of the TetR-like efflux pump regulator SimR. *Acta Crystallogr Sect Struct Biol Cryst Commun* 67:307–309. doi:10.1107/S1744309110053078
- Le TB, Stevensoc CE, Fiedler H-P, Maxwell A, Lawson DM, Buttner MJ (2011) Structures of the TetR-like Simocyclinone Efflux Pump Repressor, SimR, and the Mechanism of Ligand-Mediated Derepression. *J Mol Biol* 408:40–56. doi:10.1016/j.jmb.2011.02.035
- Luzhetskyy A, Fedoryshyn M, Gromyko O, Ostash B, Rebets Y, Bechthold A, Fedorenko V (2006) IncP plasmids are most effective in mediating conjugation between *Escherichia coli* and *Streptomyces*. *Genetica* 42:595–601
- Muth G, Nussbaumer B, Wohlleben W, Pühler A (1989) A vector system with temperature-sensitive replication for gene disruption and mutational cloning in *streptomycetes*. *Mol Gen Genet* 6:1–8
- Myronovskiy M, Welle E, Fedorenko V, Luzhetskyy A (2011) β -Glucuronidase as a sensitive and versatile reporter in *actinomycetes*. *Appl Environ Microbiol* 77:5370–5383. doi:10.1128/AEM.00434-11
- Novakova R, Rehakova A, Feckova L, Kutas P, Knischova R, Kormanec J (2011) Genetic manipulation of pathway regulation for overproduction of angucycline-like antibiotic auricin in *Streptomyces aureofaciens* CCM 3239. *Folia Microbiol* 56:276–282. doi:10.1007/s12223-011-0033-1
- Nuria A, Santos-Aberturas J, Mendes M, Guerra S, Martin J, Aparacio J (2007) PimM, a PAS domain positive regulator of pimaricin biosynthesis in *Streptomyces natalensis*. *J Bacteriol* 153:3174–3183
- Ohnishi Y, Ishikawa J, Hara H, Suzuki H, Ikenoya M, Ikeda H, Yamashita A, Hattori M, Horinouchi S (2008) Genome sequence of the streptomycin-producing microorganism *Streptomyces griseus* IFO 13350. *J Bacteriol* 190:4050–4060
- Oppegard LM, Hamann BL, Streck KR, Ellis KC, Khodursky AB, Hiasa H (2009) In vivo and in vitro patterns of the activity of simocyclinone D8, an angucyclinone antibiotic from *Streptomyces antibioticus*. *Antimicrob Agents Chemother* 53:2110–2119. doi:10.1128/AAC.01440-08
- O'Rourke S, Wietzorrek A, Fowler K, Corre C, Challis G, Chater K (2009) Extracellular signaling, translational control two repressors and an activator all contribute to the regulation of methylenomycin production in *Streptomyces coelicolor*. *Mol Microbiol* 71:763–778. doi:10.1111/j.1365-2958.2008.06560.x
- Ostash I, Ostash B, Luzhetskyy A, Bechthold A, Walker S, Fedorenko V (2008) Coordination of export and glycosylation of landomycins in *Streptomyces cyanogenus* S136. *FEMS Microbiol Lett* 285:195–202. doi:10.1111/j.1574-6968.2008.01225.x
- Ostash B, Rix U, Rix LL, Liu T, Lombo F, Luzhetskyy A, Gromyko O, Wang C, Braña AF, Méndez C, Salas JA, Fedorenko V, Rohr J (2004) Generation of new landomycins by combinatorial biosynthetic manipulation of the *IndGT4* gene of the landomycin E cluster in *S. globisporus*. *Chem Biol* 11:547–555. doi:10.1016/j.chembiol.2004.03.011
- Pulsawat N, Kitani S, Nihira T (2007) Characterization of the biosynthetic gene cluster for the production of virginiamycin M, a streptogramin type A antibiotic, in *Streptomyces virginiae*. *Gene* 393:31–42. doi:10.1016/j.gene.2006.12.035
- Rebets Y, Ostash B, Luzhetskyy A, Hoffmeister D, Brana A, Mendez C, Salas JA, Bechthold A, Fedorenko V (2003) Production of landomycins in strains *Streptomyces globisporus* 1912 and *S. cyanogenus* S136 is regulated by genes encoding putative transcriptional activators. *FEMS Microbiol Lett* 222:149–153. doi:10.1016/S0378-1097(03)00258-1
- Rebets Y, Ostash B, Luzhetskyy A, Kushnir S, Fukuhara M, Bechthold A, Nashimoto M, Nakamura T, Fedorenko V (2005) DNA-binding activity of LndI protein and temporal expression of the gene that upregulates landomycin E production in *Streptomyces globisporus* 1912. *Microbiol* 151:281–290. doi:10.1099/mic.0.27244-0
- Rebets Y, Dutko L, Ostash B, Luzhetskyy A, Kulachkovskyy O, Yamaguchi T, Nakamura T, Bechthold A, Fedorenko V (2008) Function of *lanI* gene in regulation of landomycin A biosynthesis in *Streptomyces cyanogenus* S136 and cross-complementation studies of streptomycetes antibiotic regulatory proteins encoding genes. *Archives of Microbiol* 189:111–120. doi:10.1007/s00203-007-0299-5
- Ruiz D, Salinas P, Lopez-Redondo ML, Cayuela M, Marina A, Contreras A (2008) Phosphorylation-independent activation of the atypical response regulator NblR. *Microbiol* 154:3002–3001. doi:10.1099/mic.0.2008/020677-0
- Sadig A, Patel M, Jacobson B, Escobedo M, Ellis K, Oppegard L, Hiasa H, Kratzke R (2010) Anti-proliferative effects of simocyclinone D8 (SD8), a novel catalytic inhibitor of topoisomerase II. *Invest New Drugs* 28:20–25. doi:10.1007/s10637-008-9209-1
- Sambrook J, Russell D (2001) *Molecular cloning, a laboratory manual*. Cold Spring Harbor Laboratory, 3rd edn. Cold Spring Harbor, NY
- Schimana J, Walker M, Zeeck A, Fiedler H-P (2001) Simocyclinones: diversity of metabolites is dependent on fermentation conditions. *J Ind Microbiol Biotechnol* 27:144–148. doi:10.1038/sjijm.7000024
- Sheldon P, Busarow S, Hutchinson R (2002) Mapping the DNA-binding domain and target sequences of the *Streptomyces peuceetius* daunorubicin biosynthesis regulatory protein, DnrI. *Mol Microbiol* 44:449–460. doi:10.1046/j.1365-2958.2002.02886.x
- Sissi C, Vazquez E, Chemello A, Mitchenall LA, Maxwell A, Palumbo M (2009) Mapping simocyclinone D8 interaction with DNA gyrase: evidence for a new binding site on GyrB. *Antimicrob Agents Chemother* 54:213–220
- Sola-Landa A, Moura RS, Martin J (2003) The two-component PhoR-PhoP system controls both primary metabolism and secondary metabolite biosynthesis in *Streptomyces lividans*. *PNAS* 100:6133–6138. doi:10.1073/pnas.0931429100
- Tahlan K, Ahn S, Sing A, Bodnaruk T, Willems A, Davidson A, Nodwell J (2007) Initiation of actinorhodin export in *Streptomyces coelicolor*. *Mol Microbiol* 63:951–961. doi:10.1111/j.1365-2958.2006.05559.x
- Trefzer A, Pelzer S, Schimana J, Stockert S, Bihlmaier C, Fiedler H-P, Welzer K, Vente A, Bechthold A (2002) Biosynthetic gene cluster of simocyclinone, a natural multihybrid antibiotic. *Antimicrobial Agents and Chemother* 46:1174–1182. doi:10.1128/AAC.46.5.1174-1182.2002
- van Wezel GP, McKenzie N, Nodwell J (2009) Applying the genetics of secondary metabolism in model *actinomycetes* to the discovery of novel antibiotics. *Methods Enzym* 458:118–134
- van Wezel PGilles, McDowall JKenneth (2011) The regulation of the secondary metabolism of *Streptomyces*: new links and experimental advances. *Nat Prod Rep* 28:1311–1333. doi:10.1039/c1np00003a
- Wang L, Tian X, Wang J, Yang H, Fan K, Xu G, Yang K, Tan H (2009) Autoregulation of antibiotic biosynthesis by binding of the end product to an atypical response regulator. *PNAS* 106:8617–8622. doi:10.1073/pnas.0900592106
- Zelyas N, Tahlan K, Jensen S (2009) Use of the native *flp* gene to generate in-frame unmarked mutations in *Streptomyces spp.* *Gene* 443:48–54. doi:10.1016/j.gene.2009.03.022

doi:10.1186/2191-0855-2-1

Cite this article as: Horbal et al.: SimReg1 is a master switch for biosynthesis and export of simocyclinone D8 and its precursors. *AMB Express* 2012 2:1.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com