

# Enhanced resistance to bacterial and fungal pathogens by overexpression of a human cathelicidin antimicrobial peptide (hCAP18/LL-37) in Chinese cabbage

Yu-Jin Jung · Soon-Youl Lee · Yong-Sun Moon ·  
Kwon-Kyoo Kang

Received: 6 July 2011 / Accepted: 2 August 2011 / Published online: 11 October 2011  
© The Author(s) 2011. This article is published with open access at Springerlink.com

**Abstract** The human cathelicidin antimicrobial protein hCAP18, which includes the C-terminal peptide LL-37, is a multifunctional protein. As a possible approach to enhancing the resistance to plant disease, a DNA fragment coding for hCAP18/LL-37 was fused at the C-terminal end of the leader sequence of endopolygalacturonase-inhibiting protein under the control of the cauliflower mosaic virus 35S promoter region. The construct was then introduced into *Brassica rapa*. LL-37 expression was confirmed in transgenic plants by reverse transcription-polymerase chain reaction and western blot analysis. Transgenic plants exhibited varying levels of resistance to bacterial and fungal pathogens. The average size of disease lesions in the transgenic plants was reduced to less than half of that in wild-type plants. Our results suggest that the antimicrobial LL-37 peptide is involved in wide-spectrum resistance to bacterial and fungal pathogen infection.

**Keywords** Antimicrobial peptide · Cathelicidin · Disease resistance · Transgenic Chinese cabbage

## Introduction

Bacterial and fungal plant pathogens severely affect crop productivity. For example, *Xanthomonas campestris* pv. *campestris* and *Pectobacterium carotovorum* subsp. *carotovorum*, which cause black rot and soft rot, respectively, are present worldwide and severely damage plants and reduce their yields, especially in cruciferous plants (Boman 2003). Therefore, the development of cruciferous plants that are resistant to black and soft rot diseases has been a major goal of researchers for several decades. Strategies based on transgenic approaches to enhance plant disease resistance involve the use of genes associated with plant defense pathways (Makandar et al. 2006; Zhang et al. 2007) and genes encoding plant or fungal hydrolytic enzymes (Bieri et al. 2003), defense-related transcription factors (Chen and Chen 2002; Sohn et al. 2006) and antimicrobial peptides (Alan et al. 2004).

A large number of antimicrobial peptides from different organisms have been characterized (Simmaco et al. 1998). The human cathelicidin antimicrobial protein hCAP18 is the only member of the mammalian cathelicidin family of proteins that is present in humans (Gudmundsson et al. 1996). The holoprotein consists of a conserved prodomain, a cathelin domain, and the non-conserved C-terminal peptide LL-37, which is enzymatically cleaved after secretion (Sorensen et al. 2001; Yamasaki et al. 2006). Its precursor molecule, an 18 kDa human cationic antimicrobial protein (hCAP-18), is secreted by activated neutrophil granulocytes. After release, the helical C-terminal end of this precursor comprising 37 amino acids is cleaved off, thereby forming the functional antimicrobial peptide LL-37 (Sorensen et al. 2001). Since LL-37 is the only human antimicrobial peptide that is active at physiological or elevated salt concentration conditions, there is a significant

---

Y.-J. Jung · K.-K. Kang (✉)  
Department of Horticulture, Hankyong National University,  
67 Sukjong-dong, Ansung, Kyonggi-do 456-749, South Korea  
e-mail: kykang@hknu.ac.kr

S.-Y. Lee  
Korea School of Applied Life Sciences Genetic Informatics  
Center, GRRC Hankyong National University,  
67 Sukjong-dong, Ansung, Kyonggi-do 456-749,  
South Korea

Y.-S. Moon  
Department of Horticulture, Yeungnam University,  
214-1 Daedong, Gyeongsan, Gyeongbuk 712-749,  
South Korea

interest in using this peptide for pharmaceutical applications (De Smet and Contreras 2005; Reddy et al. 2004; Travis et al. 2000).

In the present study, we report the transgenic expression of human cathelicidin antimicrobial peptide carrying the substitution Met37Leu in Chinese cabbage. The expression of this peptide in cabbage plants significantly inhibited the growth of *Pectobacterium carotovorum* subsp. *carotovorum* on the plant leaves, and it conferred resistance to several fungal pathogens. These results further support the assignment of a defense role to LL-37 and highlight its plant biotechnological potential.

## Materials and methods

### Expression vector construction

The leader sequence of the gene encoding *Phaseolus vulgaris* endopolygalacturonase-inhibiting protein (PGIP) (GenBank Accession No. X64769) was fused upstream of an LL-37-coding DNA fragment to cause the extracellular localization of the mature protein. The PGIP signal peptide (87 bp) was amplified by PCR with primers A linked with *Bam*HI site (5'-CCGGATCCATGACTCAATTCAATATC CCA-3') and B (5'-AGAGAGTGCAGTTCTCAA-3'). The coding region of LL-37 (111 bp) was substituted with Met-LL37-Leu and amplified by PCR from *pFALL37* DNA using primers C (5'-ATGCTGCTGGGTGATTTCTTC-3') and D with *Sac*I site (5'-CGAGAGCTCCTAGGACTCTG TCC TGGG-3'). The two products were ligated into pBlueScript-SK (Stratagene, La Jolla, CA, USA) at the *Bam*HI and *Sac*I restriction sites. The generated LL-37 fragment was further amplified using primers A and D. For *Agrobacterium* transformation, the PCR product was sub-cloned into pBI121 binary vector driven by cauliflower mosaic virus 35S (CaMV35S) promoter (Gelvin 1998). The Ti plasmid vector construct pBI-LL37 was confirmed by DNA sequencing (ABI 377 DNA sequencer; Perkin-Elmer, Cypress, CA, USA).

### Plant transformation and regeneration

The prepared construct was transformed into Chinese cabbage using the protocol described in Min et al. (2007). A total of 168 hypocotyls from in vitro grown seedlings of Chinese cabbage (*Brassica rapa* cv. Osome) were inoculated with *Agrobacterium tumefaciens* strain LBA4404 carrying either pBI-LL37 or pBI121. Green shoots that developed in the selective medium were transferred to a rooting medium containing 100 mg L<sup>-1</sup> kanamycin and 500 mg L<sup>-1</sup> carbenicillin. Rooted shoots were screened by PCR for the presence of the transgene before transfer to plastic pots.

### Estimation of transformants and generation of homozygous lines

Self-pollinated seeds obtained from T<sub>0</sub> plants were sown into plastic pots in the greenhouse. Two weeks after germination, seedlings were sprayed with 400 mg L<sup>-1</sup> kanamycin in water; they were sprayed again 2 days later. Three days after the second spray, the ratios of green seedlings to bleached seedlings were determined, and the results were analyzed by a Chi-square test for goodness of fit to the ratios 3:1, 15:1, or 63:1. In order to obtain transformants homozygous for the LL-37 gene, kanamycin-resistant T<sub>1</sub> progenies were grown to produce selfed T<sub>2</sub> seeds. T<sub>2</sub> lines that had no bleached segregants after kanamycin sprays were assumed to be homozygous for the LL-37 and *nptII* genes.

### Molecular analysis of transformants

PCR analyses were conducted to detect the presence of LL-37- or *nptII*-specific fragments. Primers 35SF (5'-TC CACTGACGTAAGGGATGA-3') and LL-37R (5'-CGAG AGCTCCTAGGACTCTGTCCTGGG-3'), which amplified a fragment of size approximately 750 bp, including sequences from the 3' end of the 35S promoter, signal peptide (SP), and part of the LL-37 gene, were used to screen for putative LL-37 transformants. Putative transformants were screened using primers *nptII*F (5'-TCGGC TATGACTGGGCACAACAGC-3') and *nptII*R (5'-AAGA AGGCGATAGAAGGCGATGCG-3'), which amplified a 722-bp *nptII*-specific fragment. Genomic DNA was isolated from young leaves of Chinese cabbage plants using a DNeasy Plant kit (Qiagen, Germantown, MD, USA.) as per the manufacturer's instructions. PCR was performed and the reaction conditions were followed (1 cycle of 94°C for 1 min; 30 cycles of 94°C for 30 s, 55°C for 30 s, and 72°C for 1 min; and 1 cycle of 72°C for 10 min). The reaction products were electrophoresed on a 2% (w/v) agarose gel with 1× TAE buffer and visualized by staining with ethidium bromide.

In order to analyze gene expression in transgenic plants by RT-PCR, total RNA from wild-type and transgenic plants was reverse transcribed using AMV reverse transcriptase (Roche, USA) with oligo (dT) primers for 1 h at 42°C. The expression level of actin mRNA was used as a quantitative control.

Western blot analysis was performed by following standard molecular techniques (Sambrook et al. 1998). Briefly, for western blotting, 200 ng of purified protein or synthetic peptide was electroblotted onto a polyvinylidene difluoride (PVDF) membrane. Blots were probed with 1:1,000 dilution of polyclonal Hbt Ab antiserum raised in rabbit and then with 1:10,000 dilution of goat anti-rabbit

immunoglobulin-peroxidase conjugate (Vector, Burlingame, CA, USA). The blots were then developed using the enhanced chemiluminescence (ECL) developing system (GE Health Care, USA). The Low Range BioRad 161-0304 markers (BioRad, Hercules, CA, USA) were used as the molecular size markers.

### Pathogen inoculation

*P. carotovorum* subsp. *carotovorum* KACC 10057 obtained from the Korean Agricultural Culture Collection (<http://kacc.rda.go.kr>) was grown in Luria–Bertani (LB) medium until the absorbance at 600 nm ( $A_{600}$ ) was 0.2, which corresponds to a concentration of approximately  $2 \times 10^8$  CFU/mL. Three different concentrations of the culture ( $10^4$ ,  $10^6$ , and  $10^8$  CFU/mL) with 10 mM  $MgCl_2$  were inoculated by syringe infiltration. The inoculated plants were transferred to a growth chamber and incubated at 28°C under continuous light. They were examined for 12–96 h after inoculation. Lesion length (cm) and disease index (DI) were recorded for each individual plant; the disease index ranged from 0 to 6 on the basis of the development of the disease lesions: 0, no lesion; 1, lesion size 0.1–0.5 cm; 2, 0.5–1.5 cm; 3, 1.5–3.5 cm; 4, 3.5–5.5 cm; 5, 5.5–8.5 cm; and 6, over 8.5 cm or plant dead.

The fungi *Fusarium oxysporum* f. sp. *Lycopersici* (KACC 40032), *Colletotrichum higginsianum* (KACC 40807), and *Rhizoctonia solani* (KACC 40107) were inoculated on plant leaves by placing 10  $\mu$ l of an aqueous suspension containing  $10^6$  spores/mL on the leaves. Plants

were maintained in highly humidified conditions (100% RH) at 25°C with 16 h of light in a growth chamber.

### Evaluation of in vitro inhibition assays

Total and extracellular fluids were extracted by the methods described in Alan et al. (2004). Protein concentrations in the leaflet fluids were determined by the Bradford assay (Bradford 1976). The in vitro inhibition assays were evaluated by the protocol from Alan et al. (2004). Briefly, the experiments tested whether total fluid (TF) and extracellular fluid (EF) from three homozygous lines possessed antimicrobial activity. Assays were also performed using TF and EF from wild-type plant and LB medium as controls. A volume of 248  $\mu$ l of TF, EF, and LB was mixed with 2.5  $\mu$ l of  $10^8$  CFU/mL *P. carotovorum* subsp. *carotovorum* in Eppendorf tubes and incubated on a shaker for 4 h. The samples remaining in the tubes were mixed in the ratio 1:9 with LB, returned to the shaker, and incubated overnight at 37°C. The bacterial growth in these tubes was determined by a spectrophotometer at 600 nm.


## Results


### Generation and characterization of LL-37 transgenic plants

The sequence of mature peptide of Met-LL37-Leu (GenBank accession No. NM-004345) is shown in Fig. 1a and the construct of binary pBI121 in Fig. 1b. Morphological

**Fig. 1** Construction of a binary plant expression vector pBI121 for the transformation of Chinese cabbage plants.

**a** Sequence of the peptide (GenBank Accession No. NM-004345). **b** Schematic diagram of the expression construct pBI-LL37. **c** Morphological characteristics of non-transformed Chinese cabbage (wt) and transgenic plants






**Fig. 2** PCR amplification of the foreign genes in the transgenic Chinese cabbage lines. **a** PCR analysis of 14 putative LL-37 Chinese cabbage lines (T0) using the 35SF (5'-ATGGAGTCAAAGATTCAA ATAGAG-3') and LL-37R (5'-CGAGAGCTCCTAGGACTCTGTCC TGGG-3') primers. *Lanes:* M 1 kb ladder, P positive control (pBI LL-37), C non-transformed plant, 1–14 putative transformants. **b** Four T3

(B21, 22, 23, 24) of single transgenic homozygous lines from T2 (B11, 12, 13, 14) were reconfirmed by PCR after genetic segregation events by kanamycin resistance. Amplification products of the 35S promoter, *NptII*, and *LL-37* were separated using a 1.5% agarose gel. *Lanes:* M DNA ladder, C non-transformed plant, P pBI LL-37, B21–B24 independent T3 transgenic homozygous lines

characteristics of non-transformed Chinese cabbage (wt) and transgenic plants are shown in Fig. 1c. Transformation experiments with *Agrobacterium* carrying pBI-LL37 yielded 17 independent kanamycin-resistant putative transformants. Three out of 17 transformants showed abnormal phenotypes and eliminated for further experiment. PCR analysis with 35SF and LL-37R primers revealed that 11 of 14 transformants contained the expected amplified product (Fig. 2a). The PCR-positive plants (T0) were transferred to the greenhouse where they were further observed for their phenotypic characters and were grown for recovery of self-pollinated seeds. Most transformants were phenotypically similar to non-transformed plants. Segregation analysis was indicated that T1 transformants with a single copy showed 3:1 kanamycin resistance. Four of them were randomly selected for selfing, and named lines B11, B12, B13, and B14 (T2). Ten plants per line after selfing were screened on kanamycin to get homozygotes (T3). Additionally, we confirmed the integration of the genes in four homozygous lines (T3; B21, 22, 23, and 24 from T2; B11, 12, 13, and 14) by PCR using the 35S-specific, *nptII*-specific, and *LL-37*-specific primer sets (Fig. 2b). All of these contained the expected amplified product.

#### Expression of *LL-37* in homozygous lines

In order to confirm *LL-37* expression in the transformants, we performed quantitative RT-PCR using total RNA from 4 homozygous lines. The results of RT-PCR indicated that *LL-37* RNA was expressed in the homozygous lines, and



**Fig. 3** Analysis of the *LL-37* gene expression in the transgenic homozygous lines by using reverse transcription polymerase chain reaction. Total RNA was isolated from each plant, and 0.5  $\mu$ g of this RNA was amplified with *LL-37*-specific primers. The Chinese cabbage actin gene was amplified as a loading control. The amplification products of the *LL-37* gene were separated using a 1.5% agarose gel. *Lanes:* M DNA ladder, WT wild-type plant, B21–B24 independent transgenic homozygous lines

the expression levels were similar across all the transformants (Fig. 3). Western blot analysis also confirmed that the four homozygous lines expressed the 4-kDa peptide at varying levels, whereas the control plants did not show the *LL-37*-specific band (Fig. 4). Therefore, we concluded that the antimicrobial *LL-37* gene was stably integrated into the genome of the transformants and was transcribed into mRNA that yielded the *LL-37* peptide.

#### Increased resistance of the transgenic plants to soft rot

The resistance to several rot pathogens was evaluated for the four homozygous lines harboring the human *LL-37* gene by comparing the severity of the disease to that seen

with the nontransgenic plants. Transgenic and control plants were inoculated with conidia, and the sizes of the disease lesions were determined. In order to test the resistance of the transgenic plants, we first cultivated the cabbage soft rot pathogen *P. carotovorum* subsp. *carotovorum* in LB medium for 1–2 days. We tested the resistance of the plants to three different concentrations of the bacterial pathogen:  $10^4$ ,  $10^6$ , and  $10^8$  CFU/mL. We challenged the leaf body and leaf vein with  $10^4$  and  $10^6$  CFU/mL of the bacterial culture, respectively; subsequently, the pathogenesis at the leaf was examined after 24, 48, and 72 h of inoculation. The control cabbage plants were susceptible to the bacterial pathogen and their leaves became softer with visible lesions 12 h after inoculation. After 48 h, the symptom spread all over the leaves, and the plants died owing to softening 72 h after inoculation of the pathogen. On the other hand, the softening symptom was not observed in three of the four transgenic lines, B21, B22, and B24, by 72 h after bacterial inoculation (Fig. 5). When  $10^8$  CFU/mL of the bacterial culture was inoculated, the leaves of the control plants showed the softening symptom 12 h after inoculation; the condition worsened and the leaves became amorphous after 24 h of inoculation. The

transgenic lines only showed mild softening of the leaves around the area of inoculation (Fig. 6). Therefore, it could be concluded that transgenic plants showed significantly increased resistance to the bacterial pathogen *P. carotovorum* subsp. *carotovorum*, which causes soft rot.

#### Increased resistance of the transgenic plants to fungal pathogens

The leaves of the transgenic plants were challenged with several fungal pathogens such as *F. oxysporum*, *C. higginsianum*, and *R. solani* in order to test the resistance of the transgenic plants to fungal pathogens. Three of the four transgenic homozygous lines, B21, B22, and B23, showed higher resistance to *F. oxysporum* than the control plants (Fig. 7). In particular, B21 showed remarkably high resistance to *F. oxysporum*. After being challenged with *C. higginsianum* and *R. solani*, the control plants showed typical symptoms and died after 6 days of inoculation, while the transgenic plants were resistant, as shown in Fig. 7. Therefore, we can conclude that the LL-37 peptide expressed in the transgenic plants acts as an effective antimicrobial peptide.


#### Bacterial inhibition assays in transgenic leaflets


We collected TF and EF from the leaflets of three transgenic homozygous lines and a non-transformed line in order to study their effects on the activity of LL-37. The inhibition of the bacterial (*P. carotovorum* subsp.



**Fig. 4** Western blot analysis of transgenic homozygous lines. Lanes: WT wild-type plant, B21–B24, transgenic homozygous lines containing the LL-37 gene, arrow indicates 4 kDa protein

**Fig. 5** Temporal development of soft rot in Chinese cabbage (“Osome”) plants when the leaves were inoculated with *Pectobacterium carotovorum* subsp. *carotovorum*. B21–B24 Transgenic homozygous lines





**Fig. 6** Disease development in transgenic homozygous lines containing the *LL-37* gene, which were inoculated with  $10^4$ ,  $10^6$ , or  $10^8$  CFU/mL of *P. carotovorum* subsp. *carotovorum*. Disease manifestations were scored 3 days after inoculation. *RG* water, *WT* wild-type plant, *B21*–*B24* *LL-37* transgenic homozygous lines. Disease lesions: 0, no lesion; 1, lesion size 0.1–0.5 cm; 2, 0.5–1.5 cm; 3, 1.5–3.5 cm; 4, 3.5–5.5 cm; 5, 5.5–8.5 cm; and 6, over 8.5 cm or plant dead

*carotovorum*) growth by TF and EF was evaluated using the spectrophotometric method. As expected, bacteria grew normally when they were incubated in LB medium only and in TF or EF obtained from the non-transformed line, while the growth was completely inhibited when EF obtained from the three transgenic lines was added to the LB medium (Fig. 8). Bacteria also grew well in the TF obtained from the three transgenic lines, although their growth was lower than that in LB medium alone.

## Discussion

In response to a microbial attack, plants activate a complex series of responses that lead to the local and systemic induction of a broad spectrum of antimicrobial defenses (Kim and Martin 2004; Kunkel 2002). When induced defense responses are rapidly and coordinately triggered during a given plant–pathogen interaction, plants become broadly resistant to diseases. These defense responses include the strengthening of mechanical barriers, oxidative


burst, and production of antimicrobial compounds (Hammond-Kosack and Parker 2003; Park 2005). Some research has been performed to bolster plant defenses against bacteria and fungi by genetically engineering plants to express antimicrobial peptides (Lee et al. 2008; Prasad et al. 2008).

In this study, we have chosen a variant of the antimicrobial peptide *LL-37* as an interesting candidate for transgenic plant expression; this variant is designed to target the peptide into extracellular spaces. Extracellular targeting was originally intended to prevent possible deleterious effects of the peptide in plant cells. Moreover, secretion into extracellular spaces allows the plant-produced peptide to come into direct contact with pathogens growing and multiplying extracellularly before attacking the cells (Ponti et al. 2003). We developed homozygous Chinese cabbage lines stably expressing *LL-37*, which did not cause adverse effects on the plant phenotypes. These transgenic homozygous lines were tested with four important pathogens of this crop and found to inhibit the growth of the bacterial pathogen causing Chinese cabbage rot and also that of three fungal pathogens.


The results of our pathogenicity assays suggest that the expression of *LL-37* provides a moderate level of resistance against a bacterial pathogen (*P. carotovorum* subsp. *carotovorum*) at the inoculum concentration of  $10^4$  CFU/mL. However, the extent of disease suppression provided by *LL-37* expression was reduced as the inoculum concentrations were increased to  $10^6$  and  $10^8$  CFU/mL. Moreover, we observed reduction in the survival of *P. carotovorum* subsp. *carotovorum* cells incubated with EF from *LL-37*-expressing Chinese cabbage cell lines. However, bacteria grew normally when they were incubated in LB medium only and in TF or EF obtained from a non-transformed cell line. Therefore, leaves appeared to express *LL-37* at sufficient levels in the extracellular spaces to retard bacterial multiplication and hence decrease disease severity. The functionality of this transgene and the presence of antimicrobial activity in the EF indicates that the *LL-37* peptide was properly targeted to the extracellular space even with a foreign plant signal peptide. Furthermore, the mammalian peptide was not subjected to a processing step in the foreign plant cell environment that rendered it inactive.

Until now, studies involving the enhancement of resistance to various bacterial, fungal, and oomycete pathogens by the expression of antimicrobial peptides have been reported for rice, tobacco, poinsettia, banana, and more host species (Chakrabarti et al. 2003; Liang et al. 2002; Smith et al. 1998). However, progress on identifying the defense mechanisms in Chinese cabbage (*B. rapa*), an important vegetable crop in Asia, has been very slow. Previously, we reported an enhancement in the resistance to bacterial soft rot by the expression of the bromelain gene in Chinese cabbage (Jung et al. 2008). Here, we

**Fig. 7** Enhanced disease resistance of transgenic homozygous lines to several fungal pathogens. F.O., *Fusarium oxysporum*; C.h., *Colletotrichum higginsianum*; R.S., *Rhizoctonia solani*. Disease manifestations assessed until 6 days after inoculation



**Fig. 8** Bacterial inhibition assays of total fluid (TF) and extracellular fluid (EF) obtained from the leaflets of transgenic homozygous lines; LB Luria–Bertani medium



demonstrate that expression of the human LL-37 peptide has antimicrobial activity toward both bacterial and fungal pathogens of Chinese cabbage. Since pathogens have the ability to overcome gene-for-gene host defense mechanisms in the field by undergoing mutations in the cognate avirulence genes, any transgenic model based on such resistance-conferring genes may easily be evaded by pathogens. On the other hand, this problem is less likely to occur in transgenic plants overexpressing genes with more general antimicrobial activity. Therefore, the expression of the human LL-37 peptide is expected to confer durable

resistance (i.e., field resistance) to a wide variety of pathogens infecting Chinese cabbage plants.

In order to obtain a high level of resistance as observed in the case of R gene-mediated resistance, the overexpression of multiple antifungal proteins with different functions may be necessary. In a *R. solani* infection assay, tobacco plants coexpressing the barley transgenes (a class II chitinase, a class II  $\beta$ -1,3-glucanase, and a type I ribosome-inactivating protein) were reported to elicit significantly enhanced protection against fungal attack as compared to that of the corresponding isogenic lines expressing a single barley

transgene at a similar level (Brogue et al. 1991). Thus, other antimicrobial genes such as the bromelain gene (Jung et al. 2008) could be stacked with the human *LL-37* gene by crossing different transgenic lines and this may be able to strongly and durably inhibit the growth of pathogens.

**Acknowledgments** We thank to Dr. Philip Becraft for critical reviews and comments. This work was supported by a grant from the Next-Generation BioGreen 21 Program (No. PJ008085), Rural Development Administration, Republic of Korea.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- Alan AR, Blowers A, Earle ED (2004) Expression of a magainin-type antimicrobial peptide gene (MSI-99) in tomato enhances resistance to bacterial speck disease. *Plant Cell Rep* 22:388–396
- Bieri S, Potrykus I, Fütterer J (2003) Effects of combined expression of antifungal barley seed proteins in transgenic wheat on powdery mildew infection. *Mol Breed* 11:37–48
- Boman HG (2003) Antibacterial peptides: basic facts and emerging concepts. *J Intern Med* 254:197–215
- Bradford MM (1976) A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254
- Brogue K, Chet I, Holliday M, Cressman R, Biddle P, Knowlton S, Mauvais CJ, Broglie R (1991) Transgenic plants with enhanced resistance to the fungal pathogen *Rhizoctonia solani*. *Science* 254:1194–1197
- Chakrabarti A, Ganapathi TR, Mukherjee PK, Bapat VA (2003) MSI-99, a magainin analogue, imparts enhanced disease resistance in transgenic tobacco and banana. *Planta* 216:587–596
- Chen C, Chen Z (2002) Potentiation of developmentally regulated plant defense response by AtWRKY18, a pathogen-induced *Arabidopsis* transcription factor. *Plant Physiol* 129:706–716
- De Smet K, Contreras R (2005) Human antimicrobial peptides: defensins, cathelicidins and histatins. *Biotechnol Lett* 27:1337–1347
- Gelvin SB (1998) The introduction and expression of transgenes in plants. *Curr Opin Biotech* 9:227–232
- Gudmundsson GH, Agerberth B, Odeberg J, Bergman T, Olsson B, Salcedo R (1996) The human gene FALL39 and processing of the cathelin precursor to the antibacterial peptide LL-37 in granulocytes. *Eur J Biochem* 238:325–332
- Hammond-Kosack KE, Parker JE (2003) Deciphering plant-pathogen communication: fresh perspectives for molecular resistance breeding. *Curr Opin Biotech* 14:177–193
- Jung YJ, Choi CS, Park JH, Kang HW, Choi JE, Nou IS, Lee SY, Kang KK (2008) Overexpression of the pineapple fruit bromelain gene (BAA) in transgenic Chinese cabbage (*Brassica rapa*) results in enhanced resistance to bacterial soft rot. *Electron J Biotechnol* 11:1–8
- Kim YJ, Martin GB (2004) Molecular mechanisms involved in bacterial speck disease resistance of tomato. *Plant Pathol J* 20:7–12
- Kunkel BN (2002) Brooks cross talk between signaling pathways in pathogen defense. *Curr Opin Plant Biol* 5:325–331
- Lee SC, Hwang IS, Choi HW, Hwang BK (2008) Involvement of the pepper antimicrobial protein *CaAMP1* gene in broad spectrum disease resistance. *Plant Physiol* 143:1004–1020
- Liang H, Catranis CM, Maynard CE, Powell WA (2002) Enhanced resistance to the poplar fungal pathogen, *Septoria musiva*, in hybrid poplar clones transformed with genes encoding antimicrobial peptides. *Biotechnol Lett* 24:383–389
- Makandar R, Essig JS, Schapaugh MA, Trick HN, Shah J (2006) Genetically engineered resistance to *Fusarium* head blight in wheat by expression of *Arabidopsis NPR1*. *Mol Plant Microbe Interact* 19:123–129
- Min BW, Cho YN, Song MJ, Noh TK, Kim BK, Chae WK, Park YS, Choi YD, Harn CH (2007) Successful genetic transformation of Chinese cabbage using phosphomannose isomerase as a selection marker. *Plant Cell Rep* 26:337–344
- Park JM (2005) The hypersensitive response: a cell death during disease resistance. *Plant Pathol J* 21:99–101
- Ponti D, Mangoni ML, Mignogna G, Simmaco M, Barra D (2003) An amphibian antimicrobial peptide variant expressed in *Nicotiana tabacum* confers resistance to phytopathogens. *Biochem J* 370:121–127
- Prasad BD, Jha S, Chattoo BB (2008) Transgenic *indica* rice expressing *Mirabilis jalapa* antimicrobial protein (Mj-AMP2) shows enhanced resistance to the rice blast fungus *Magnaporthe oryzae*. *Plant Sci* 175:364–371
- Reddy KV, Yedery RD, Aranha C (2004) Antimicrobial peptides: premises and promises. *Int J Antimicrob Ag* 24:536–547
- Sambrook J, Fritsch EF, Maniatis T (1998) Molecular cloning: a laboratory manual, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York
- Simmaco M, Mignogna G, Barra D (1998) Antimicrobial peptides from amphibian skin: what do they tell us? *Biopolymers* 47:435–450
- Smith FD, Gadoury DM, Vaneck JM, Blowers A, Sanford JC, Van der Meij J, Eisenreich R (1998) Enhanced resistance to powdery mildew in transgenic poinsettia conferred by antimicrobial peptides. *Phytopathology* 88:S83
- Sohn KH, Lee SC, Jung HW, Hong JK, Hwang BK (2006) Expression and functional roles of the pepper pathogen-induced transcription factor RAV1 in bacterial disease resistance, and drought and salt stress tolerance. *Plant Mol Biol* 61:897–915
- Sorensen OE, Follin P, Johnsen AH, Calafat J, Tjabringa GS, Hiemstra PS, Borregaard N (2001) Human cathelicidin, hCAP-18, is processed to the antimicrobial peptide LL-37 by extracellular cleavage with proteinase 3. *Blood* 97:3951–3959
- Travis SM, Anderson NN, Forsyth WR, Espiritu C, Conway BD, Greenberg EP, McCray PB, Lehrer RI, Welsh MJ, Tack BF (2000) Bactericidal activity of mammalian cathelicidin-derived peptides. *Infect Immun* 68:2748–2755
- Yamasaki K, Schaubert J, Coda A, Lin H, Dorschner RA, Schechter NM, Bonnard C, Descargues P, Hovnanian A, Gallo RL (2006) Kallikrein-mediated proteolysis regulates the antimicrobial effects of cathelicidins in skin. *Fed Am Soc Exp Biol J* 20:2068–2080
- Zhang X, Dai Y, Xiong Y, DeFraia C, Li J, Dong X, Mou Z (2007) Overexpression of *Arabidopsis MAP kinase kinase 7* leads to activation of plant basal and systemic acquired resistance. *Plant J* 52:1066–1079