

Review: Verbal Autopsy for Neurological Diseases

Farrah J. Mateen* and Henry D. Kalter

*Department of International Health, Bloomberg School of Public Health and Department of Neurology,
Johns Hopkins Hospital, The Johns Hopkins University, Baltimore, Maryland*

Abstract. Verbal autopsy is an interview-based technique to determine the cause distribution of death in a population. The use of verbal autopsy for understanding neurological diseases is crucial to burden of disease analyses in many countries, particularly in locations where civil registration systems are non-functioning or absent. We review the purposes, strengths, and weaknesses in the use of verbal autopsy for neurological diseases.

Neurological disease is a major cause of mortality worldwide. The upcoming publication of the Global Burden of Disease analysis depends on accurate and adequate reporting of the cause distribution of deaths internationally. Cerebrovascular disease has traditionally ranked as the third most common cause of death worldwide and accounts for ~5.5 million of the 55 million deaths that occur each year.¹ Other neurological disorders, including epilepsy and dementia, are highly prevalent, but data on the incidence and prevalence of neurological diseases are lacking from many regions. Available information is at best an extrapolation from incomplete civil registration systems. Only one-third of countries globally certify the causes of all deaths. In the Southeast Asia region, less than half of the countries have vital registration. No country in the Africa region has complete data and only six African countries (13%) have any form of vital registration whatsoever.²

The verbal autopsy method. Verbal autopsy is the most commonly used method to determine cause of death in places lacking adequate civil registration, where the majority of deaths occur outside of medical care. The method typically involves asking a respondent knowledgeable of the deceased's fatal illness a series of questions on the signs and symptoms that preceded death (e.g., "During the last 3 months of pregnancy, did she suffer from a convulsion?" or "During the illness that led to death, was he paralyzed on one side?") combined with an open-ended narrative of the illness-related events.³ The respondent must 1) be a close family member or caregiver of the decedent, 2) have been present during the illness leading to the death, and 3) be able to recount and provide details of the decedent's health information. The interviewers usually have a secondary school-level education but no medical qualifications and complete intensive short-term training in the study techniques. Ultimate diagnosis is most commonly made by physicians who review the completed surveys.

Validity, which is the accuracy of the verbal autopsy diagnoses compared with an external reference standard, is the most important measurement of the verbal autopsy instrument. Validation studies may best be accomplished in a hospital setting where objective reference standard diagnoses can be developed based on clinical history, examination, laboratory results, and imaging findings. Reviews of the completed verbal autopsy surveys by an individual or a panel of physicians or, alternatively, diagnoses derived from objective algo-

ri thms (combinations of verbal autopsy signs and symptoms) predetermined by an expert panel are compared with the reference standard diagnoses to determine the accuracy of the verbal autopsy diagnoses. Such validation studies have established various levels of sensitivity and specificity for verbal autopsy diagnoses of major causes of death.⁴ New statistical methods for analyzing verbal autopsy data hold promise for more accurate identification of the cause distribution of deaths and more rapid, computerized processing of high-volume verbal autopsy data.⁵

Existing use of verbal autopsy for neurological disease. The World Health Organization (WHO) recognizes three major goals of verbal autopsy: 1) research and longitudinal analysis measures, 2) assessment of the cross-sectional prevalence or burden of disease, and 3) evaluation of disease-focused treatment programs and interventions.⁶ Although the use of verbal autopsy methods began as early as the 1930s, there has been little research on neurological causes of death, particularly in adulthood. Early studies that included neurological conditions focused on childhood-predominant diseases including neonatal tetanus, cerebral malaria, and meningitis. In recent years, however, increasing attention has been paid to adult causes of neurological death (Table 1). For example, in a study of 48,000 adult deaths in Chennai, India, the death rate from stroke was found to reach 1308.5/100,000 in people over 75 years of age.¹² Compared with death certificates alone, verbal autopsy yielded fewer unspecified causes of death. In China, a validation study identified more stroke deaths by questionnaire analysis than by medical diagnosis (i.e., verbal autopsy led to false positive diagnoses), with sensitivity of 82% and specificity of 94% for the verbal autopsy diagnoses.¹¹ There have been no validation studies of verbal autopsy for deaths caused by epilepsy, traumatic brain and spinal cord injury, or human immunodeficiency virus (HIV) dementia to date.

Estimating sample size for verbal autopsy. Of importance, the numbers needed to understand neurological deaths in large populations may not be exceptionally high. For example, hypothesizing that ischemic stroke is the most common neurological cause of death and accounts for ~10% of all adult deaths, an annual cluster sample of 1,200 deaths of people 18 years of age and above selected with a design effect of 1.4 could estimate the true cause proportion with precision $\pm 2\%$ and $\alpha = 0.05$ ($z = 1.96$). Using a fixed cohort and the assumption that adult mortality is 2% per year, achieving this estimate would require monitoring about 30,000 households. Estimating less common neurological diseases that may lead to death—meningitis, cerebral malaria, or epilepsy, for example—will require different numbers depending on the prevalence

*Address correspondence to Farrah J. Mateen, Departments of Neurology and International Health, The Johns Hopkins University Pathology Building, Room 627, 600 N. Wolfe Street Baltimore, MD 21287. E-mail: fmateen@jhsph.edu

TABLE 1
Large studies that use verbal autopsy in the study of neurological disease

Reference no.	Study location, years	Neurological disease	Study populations size	Main outcomes
7	Inner Mongolia, China in progress	Subarachnoid hemorrhage	1.7 million individuals	Incidence including sex- and age-specific incidence, population-attributable risks for specific exposures, case fatality ratios
8	Rural West China 2005–2009	Convulsive epilepsy	3,568 individuals	Case fatality; proportional mortality ratios by cause of death, standardized mortality ratios for age and cause
9	Tanzania 2003–2006	Stroke	636 strokes	Annual stroke incidence rates
10	Mozambique 2002–2004	Eclampsia, pyogenic meningitis	139 maternal deaths	Sensitivity, specificity, positive and negative predictive values, and diagnostic accuracy compared with necropsy
11	Urban China, 6 cities 2002	Cerebrovascular diseases	3,290 deaths	Validity of verbal autopsy diagnoses, sensitivity, specificity
12	Chennai, India 1995–1997	Cerebrovascular diseases	48,000 deaths	Sex- and age-specific death rates
13	Tanzania, Ethiopia, and Ghana 1993–1995	Rabies, tetanus, meningitis	796 deaths	Cause distribution of deaths, sensitivity, specificity, positive and negative predictive values

of the disease, its case fatality rate, and the age structure of the population.

Costs and benefits of verbal autopsy. Civil registration for the African continent may cost as little as 10 cents per person (total 80 million US\$ annually) but has not achieved sufficient funding to be a quickly realizable goal.¹⁴ The cost of obtaining verbal autopsy data above and beyond that of identifying the deaths was found to be US\$4 per death (8 cents per individual monitored in the verbal autopsy program), including physician review and data analysis.¹⁵ The cost likely would be less using automated techniques without physician involvement.

Verbal autopsy can provide baseline data to assess the prevalence of death caused by neurological and other chronic diseases. Data on preventable and modifiable causes of death can ideally identify new health priorities, lead to more appropriate allocation of sparse resources, and assist in the evaluation of health programs. Such data may be of high value to both policy makers and clinicians. Modules added to the verbal autopsy can determine the prevalence of important risk factors and health-related behaviors in the deceased persons. Expected results could include targeted vaccination and risk factor reduction campaigns and the strengthening of health systems in a locally meaningful way. Reliability of medical diagnoses among physician coders over time has recently been reported to be high.¹⁶

Limitations to verbal autopsy. Verbal autopsy is an interim solution. Because most sites that conduct verbal autopsy cover limited areas within countries, they are most useful for assessing local health needs and contribute less to nationwide or global estimates. Few verbal autopsy instruments have been validated for adult neurological diseases. In general, more work is needed on the comparison between verbal autopsy findings and traditional necropsy, which has been reported only rarely.¹⁰

Verbal autopsy data can be subject to inaccuracies in recall by the interviewed respondent. When verbal autopsy is delayed, as occurs in many reported studies, the recall period of the respondent may stretch into the months or years after the death. The accuracy of the calculations of incidence and prevalence of neurological disease may be compromised by misclassification error and poor sensitivity of the verbal autopsy instrument. For example, assessment of adult meningitis deaths in Tanzania, Ethiopia, and Ghana found a diag-

nostic sensitivity of 40–64%.¹³ If data are inaccurate, findings can misinform public health policy formulation. The underlying cause and subtypes of many neurological diseases may remain unclear (e.g., hemorrhagic versus ischemic stroke in incidence studies of stroke), leading to the need for further and more in-depth studies.

Opportunities. In the coming years, until civil registration and death certification systems are strengthened, verbal autopsy methods can provide information for some of the world's poorest regions where little to nothing is reported on neurological causes of death. Resource-poor regions remain of high importance because of the excess neurological disease and resultant premature deaths that can occur. Embedding verbal autopsy studies in periodic population surveys or existing surveillance systems may be an initial step to strengthen the use of verbal autopsy within existing infrastructure. The ongoing prospective study of one million deaths in India is one example of population-based data that can be used to conceptualize neurological disease rates in the adult population.¹⁷ Although a study of this scale may not be easily set up in other resource poor regions, its feasibility in India suggests that other low-income countries may similarly improve their monitoring of community-level deaths through verbal autopsy in sample registration systems. Despite multiple limitations to the verbal autopsy method, engagement of the growing group of community health workers, researchers, and health professionals can lead to meaningful refinements and valuable epidemiological data based on the verbal autopsy approach.

Received August 18, 2011. Accepted for publication November 18, 2011.

Acknowledgments: We thank Robert E. Black, MD, MPH, Professor and Chair of International Health, Johns Hopkins University, for helpful comments and suggestions, integral to the development of this manuscript.

Authors' addresses: Farrah J. Mateen, Departments of Neurology and International Health, The Johns Hopkins University, Baltimore, MD, E-mail: fmateen@jhsph.edu. Henry D. Kalter, Department of International Health, Bloomberg School of Public Health, The Johns Hopkins University, Baltimore, MD, E-mail: hkalter@jhsph.edu.

REFERENCES

1. World Health Organization, 2004. *Atlas of Heart Disease and Stroke: Deaths from Stroke*. Geneva: WHO Press, 53.

2. Black RE, Cousens S, Johnson HL, Lawn JE, Rudan I, Bassani DG, Jha P, Campbell H, Walker CF, Cibulskis R, Eisele T, Liu L, Mathers C; Child Health Epidemiology Reference Group of WHO and UNICEF, 2010. Global, regional, and national causes of child mortality in 2008: a systematic analysis. *Lancet* 375: 2969–2987.
3. Bang AT, Bang RA, 1992. Diagnosis of causes of childhood deaths in developing countries by verbal autopsy: suggested criteria. The SEARCH Team. *Bull World Health Organ* 70: 499–507.
4. Anker M, Black RE, Coldham C, Kalter HD, Quigley MA, Ross D, Snow RW, 1999. *A Standard Verbal Autopsy Method for Investigating Causes of Death in Infants and Children*. Geneva: WHO.
5. King G, Lu Y, 2008. Verbal autopsy methods with multiple causes of death. *Stat Sci* 23: 78–91.
6. World Health Organization, 2007. *Verbal Autopsy Standards: Ascertaining and Attributing Causes of Death*. Geneva: WHO Press.
7. Shiue I, Zhang JF, Arima H, Wang JG, Liu GR, Li YC, Wang M, Cheng GJ, Anderson CS; CHERISH Collaborative Group, 2010. Design of the China Epidemiology Research in Subarachnoid Hemorrhage (CHERISH) study. *Int J Stroke* 5: 493–498.
8. Mu J, Liu L, Zhang Q, Si Y, Hu J, Fang J, Gao Y, He J, Li S, Wang W, Wu J, Sander JW, Zhou D, 2011. Causes of death among people with convulsive epilepsy in rural west China: a prospective study. *Neurology* 77: 132–137.
9. Walker R, Whiting D, Unwin N, Mugusi F, Swai M, Aris E, Jusabani A, Kabadi G, Gray WK, Lewanga M, Alberti G, 2010. Stroke incidence in rural and urban Tanzania: a prospective, community-based study. *Lancet Neurol* 9: 786–792.
10. Ordi J, Ismail MR, Carrilho C, Romagosa C, Osman N, Machungo F, Bombi JA, Balasch J, Alonso PL, Menéndez C, 2009. Clinico-pathological discrepancies in the diagnosis of causes of maternal death in sub-Saharan Africa: retrospective analysis. *PLoS Med* 6: e10000036.
11. Yang G, Rao C, Ma J, Wang L, Wan X, Dubrovsky G, Lopez AD, 2006. Validation of verbal autopsy procedures for adult deaths in China. *Int J Epidemiol* 35: 741–748.
12. Gajalakshmi V, Peto R, Kanaka S, Balasubramanian S, 2002. Verbal autopsy of 48,000 adult deaths attributable to medical causes in Chennai (formerly Madras), India. *BMC Public Health* 2: 7.
13. Chandramohan D, Maude GH, Rodrigues LC, Hayes RJ, 1998. Verbal autopsies for adult deaths: their development and validation in a multicentre study. *Trop Med Int Health* 3: 436–446.
14. AbouZahr C, Gollogly L, Stevens G, 2010. Better data needed: everyone agrees, but no one wants to pay. *Lancet* 375: 619–620.
15. Marsh D, Husein K, Lobo M, Shah MA, Luby S, 1995. Verbal autopsy in Karachi slums: comparing single and multiple cause of child deaths. *Health Policy Plan* 10: 395–403.
16. Khademi H, Etemadi A, Kamangar F, Nourai M, Shakeri R, Abaie B, Pourshams A, Bagheri M, Hooshyar A, Islami F, Abnet CC, Pharoah P, Brennan P, Boffetta P, Dawsey SM, Malekzadeh R, 2010. Verbal autopsy: reliability and validity estimates for causes of death in the Golestan Cohort Study in Iran. *PLoS ONE* 5: e11183.
17. Jha P, Gajalakshmi V, Gupta PC, Kumar R, Mony P, Dhingra N, Peto R; RGI-CGHR Prospective Study Collaborators, 2006. Prospective study of one million deaths in India: rationale, design, and validation. *PLoS Med* 3: e18.