

Published in final edited form as:

*Psychol Med.* 2010 December ; 40(12): 1967–1978. doi:10.1017/S0033291710000231.

## The varying impact of type, timing and frequency of exposure to childhood adversity on its association with adult psychotic disorder

Helen L. Fisher<sup>1,\*</sup>, Peter B. Jones<sup>2</sup>, Paul Fearon<sup>1</sup>, Thomas K. Craig<sup>1</sup>, Paola Dazzan<sup>1,3</sup>, Kevin Morgan<sup>4</sup>, Gerard Hutchinson<sup>5</sup>, Gillian A. Doody<sup>6</sup>, Peter McGuffin<sup>1</sup>, Julian Leff<sup>1</sup>, Robin M. Murray<sup>1,3</sup>, and Craig Morgan<sup>1,3</sup>

<sup>1</sup>Institute of Psychiatry, King's College London, UK

<sup>2</sup>Department of Psychiatry, University of Cambridge, UK

<sup>3</sup>National Institute of Health Research Biomedical Research Centre for Mental Health, London, UK

<sup>4</sup>Department of Psychology, Westminster University, London, UK

<sup>5</sup>Psychiatry Unit, Faculty of Medical Sciences, University of the West Indies, Trinidad

<sup>6</sup>Division of Psychiatry, University of Nottingham, UK

### Abstract

**Background**—Childhood adversity has been associated with onset of psychosis in adulthood but these studies have used only general definitions of this environmental risk indicator. Therefore, we sought to explore the prevalence of more specific adverse childhood experiences amongst those with and without psychotic disorders using detailed assessments in a large epidemiological case-control sample (ÆSOP).

**Method**—Data were collected on 182 first-presentation psychosis cases and 246 geographically-matched controls in two UK centres. Information relating to the timing and frequency of exposure to different types of childhood adversity (neglect, antipathy, physical and sexual abuse, local authority care, disrupted living arrangements and lack of supportive figure) was obtained using the Childhood Experience of Care and Abuse Questionnaire.

**Results**—Psychosis cases were three times more likely to report severe physical abuse from mother that commenced prior to 12 years of age, even after adjustment for other significant forms of adversity and demographic confounders. A non-significant trend was also evident for greater prevalence of reported severe maternal antipathy amongst those with psychosis. Associations with maternal neglect and childhood sexual abuse disappeared after adjusting for maternal physical abuse and antipathy. Paternal maltreatment and other forms of adversity were not associated with psychosis nor was there evidence of a dose-response effect.

**Conclusions**—These findings suggest that only specific adverse childhood experiences are associated with psychotic disorders and only in a minority of cases. If replicated, this greater precision will ensure that research into the mechanisms underlying the pathway from childhood adversity to psychosis is more fruitful.

---

\*Address for correspondence: PO 80 MRC SGDP Centre, Institute of Psychiatry, De Crespigny Park, London SE5 8AF, United Kingdom. (helen.fisher@iop.kcl.ac.uk) .

## Keywords

child abuse; psychotic disorders; childhood trauma; aetiology; psychosis

---

## Introduction

Despite a growing interest in the role of adverse childhood experiences in the later development of psychotic disorders, the majority of studies have been methodologically limited (Bendall et al., 2008; Morgan & Fisher, 2007). One problem has been the use of composite measures of adversity (e.g., Romero et al., 2009; Weber et al., 2008; Wexler et al., 1997), or a focus on just one or two forms of adversity (e.g., Houston et al., 2008; Kim et al., 2006; Neria et al., 2002), thus preventing assessment of the independent effects of different types of adversity. Even those studies that have included a range of adverse childhood experiences have not considered either the timing or frequency of exposure or both (Heads et al., 1997; Rosenberg et al., 2007; Rubino et al., 2009; Schafer et al., 2006). Moreover, no studies to date have investigated the role of different perpetrators of childhood abuse in relation to psychosis. These over-generalisations may mask true associations and jeopardise future attempts to uncover the mechanisms involved in such aetiological pathways.

Previous research emphasises the importance of taking into account these specific aspects of childhood adversity as they may vary in their impact on subsequent mental wellbeing. For instance, a recent study highlighted that severe maltreatment perpetrated by the main mother figure during childhood has a more central aetiological role in depression than abuse committed by a father figure (Brown et al., 2007). Given the hypothesised key role of mother-child attachment bonds in psychological wellbeing (Bowlby, 1951, 1977), evidence from cohort studies of mothers' perceptions and care of the child being related to adult schizophrenia (Jones et al., 1994; McNeil et al., 2009; Myhrman et al., 1996), and the greater absence of fathers in the childhoods of individuals with psychosis (Agid et al., 1999; Morgan et al., 2007) it seems plausible to hypothesise that abuse perpetrated by mothers will also have a greater association with psychotic disorders. More intrusive forms of adversity such as physical, sexual and emotional abuse have also been postulated to have aetiological significance for psychosis (Harris, 1987). Indeed, higher rates of these types of maltreatment have been found amongst psychosis patients than neglect (Hlastala & McClellan 2005; Schenkel et al., 2005) though the results are mixed (Beattie et al., 2009; Heads et al., 1997; Schafer et al. 2006). Therefore, further exploration is required of the differential associations with clinical disorder controlling for appropriate confounders and other forms of adversity.

Exposure to adversity at a very young age has also been shown to have more detrimental consequences than later exposure for other disorders (Blaauw et al., 2002; Keiley et al., 2001). Furthermore, studies on non-clinical samples have indicated a greater risk of psychotic-like symptoms following occurrence of multiple versus single adverse experiences (Janssen et al., 2004; Shevlin et al., 2008; Spauwen et al., 2006; Whitfield et al., 2005). Consequently, before the research focus can be shifted to the mechanisms underlying the association between childhood adversity and psychosis, it seems imperative to establish more precisely what aspects of adversity are driving this association.

We therefore investigated the prevalence of different forms of childhood adversity amongst cases with first-presentation psychosis and catchment-based healthy controls. We hypothesised that there would be (i) associations between more intrusive forms of childhood adversity and psychotic disorder; and (ii) the associations would be stronger in those exposed to maternally perpetrated, more frequent or earlier experiences of adversity.

## Method

### Participants

The sample was drawn from the Aetiology and Ethnicity in Schizophrenia and Other Psychoses (ÆSOP) study conducted in 1997-2000 (see Morgan et al., 2006 for full details). Briefly, all patients aged 16-65 years who presented to psychiatric services for the first time with a psychotic disorder (codes F20-29 and F30-33 from the International Classification of Diseases [ICD-10]; WHO, 1992) within tightly defined catchment areas in Southeast London and Nottingham were approached. Exclusion criteria included: organic psychosis; IQ under 50; previous contact with services for psychosis; and transient psychotic symptoms resulting from acute intoxication (ICD-10; WHO, 1992). ICD-10 diagnoses were determined on the basis of consensus meetings involving one of ÆSOP's principal investigators (J.L., R.M., P.J.) using data from the Schedules for Clinical Assessment in Neuropsychiatry (SCAN; WHO, 1994). Diagnoses were made masked to ethnicity and abuse history (see Kirkbride et al., 2006).

For the control group a random sample of individuals aged 16-64 years were recruited from the population of the same geographical areas as the cases. The sampling procedure was adapted from that used by the Office of Population and Census Statistics Psychiatric Morbidity Survey (Jenkins & Meltzer, 1995). To ensure that a sufficient number of people of Black Caribbean ethnicity were recruited, we purposely over-sampled this population by continuing recruitment for a longer period. The Psychosis Screening Questionnaire (PSQ; Bebbington & Nayani, 1995) was administered to all potential control group participants; individuals were excluded if they screened positive and were found to have a psychotic disorder.

### Measures

Data on age, gender, ethnicity and parental occupations during the participant's childhood were obtained during face-to-face interviews using the Medical Research Council (MRC) Sociodemographic Schedule (Mallet, 1997). Ethnicity was self-ascribed and standardised using the 16 categories employed by the UK Census in 2001. The most senior occupation that participants' fathers had held was converted into 'highest ever parental social class' using the Office of National Statistics' (2002) Socio-Economic Classification system.

The Childhood Experience of Care Abuse Questionnaire (CECA.Q; Bifulco et al., 2005) was used to retrospectively elicit information from participants concerning childhood experiences before the age of 16 years. These include changes in parental living arrangements (lasting at least a year each), being taken into local authority care, physical abuse, antipathy or neglect by mother and father figures, sexual abuse by any adult or an individual at least 5 years older than the recipient, and lack of a support figure. Both the physical and sexual abuse sections of the CECA.Q begin with screening questions and then positive responses are followed up with more detailed questions. The antipathy and neglect items are rated on a 5-point scale from 'yes, definitely' (1) to 'no, not at all' (5) for the mother and father figure that the respondent lived with for the longest or had the most difficulties with. Antipathy refers to hostility, rejection or coldness as well as 'scapegoating' behaviour and overlaps with the concept of emotional abuse (see Bernstein & Fink, 1998). Neglect comprises a distinct lack of interest in the child's physical care (e.g., food, shelter, clothing, and health), friendships, schoolwork and whereabouts. Full details of the questionnaire are provided in Bifulco et al (2005). The CECA.Q has been shown to have good internal consistency (Smith et al., 2002), satisfactory levels of test-retest reliability over 2 and 3 years in depression samples (Bifulco et al., 2005; Smith et al., 2002) and 7 years in this psychosis sample (Fisher et al., 2009a), and reasonable concurrent validity with

the CECA interview and Parental Bonding Instrument (Bifulco et al., 2005; Fisher et al., 2009a; Smith et al., 2002). This questionnaire was read out to all participants to improve the accuracy of the fixed category responses obtained.

## Analysis

The most conservative cut-off points published by Bifulco et al. (2005) were used to dichotomise responses on the CECA.Q into severe and non-severe categories for each adversity variable. Briefly, physical abuse was rated separately for the main mother and father figure (usually but not necessarily the biological parents) with at least 3 positive responses to the 4 follow-up probes required to warrant categorisation as severe. Sexual abuse was not limited to family members and was considered severe if at least 2 of the 7 follow-up questions were answered positively for either the first or second experience. The ratings for the antipathy and neglect items were totalled for each parent and the following scores were classified as severe:  $\geq 28$  maternal antipathy,  $\geq 30$  paternal antipathy,  $\geq 25$  maternal neglect, and  $\geq 26$  paternal neglect. The total number of parental arrangements were recoded into those with 1 or 2 arrangements as 0 (no/minimal disruption) and those with 3 or more arrangements as 1 (disrupted living arrangements). Individuals who reported being placed in local authority care for any length of time were rated as 1 (taken into care) whilst those who reported no institutional care or being in other forms of institutions were coded as 0 (no care). Lack of a support figure was classed as not being able to identify either an adult or a peer who the respondent could confide in or discuss their problems with prior to 16 years of age.

A 'total adversity' score was created by summing the dichotomous CECA.Q subscale scores (range 0-10) and then recoding the total into 0 (none), 1 (single adverse experience), and 2 (multiple adverse experiences). Frequency variables for physical abuse, antipathy and neglect were created using codes of 0 (neither parent), 1 (maltreated by one parent) and 2 (maltreated by both parents). The following codes were employed to reflect frequency of sexual abuse experience: 0 (no abuse), 1 (abuse from one individual), 2 (abuse from two separate individuals). Additionally, the reported age of first severe sexual abuse and commencement of severe physical abuse from mother and father were grouped into 'childhood' (0-11 years) and 'adolescence' (12-16 years) in accordance with the conventions employed by Thornberry et al. (2001) and Widom et al. (2008).

Logistic regression was used to analyse the relationship between each form of adversity and case-control status. All analyses were weighted to correct for the over-sampling of Black Caribbean controls (see Morgan et al., 2007). In the final logistic regression models, sex (male or female), age (16-35 or 36-64), ethnicity (White British, White Other, Black Caribbean, Black African, Asian [all] or Other), study centre (London or Nottingham) and highest ever parental social class (managerial/professional, intermediate, or routine/manual) were controlled. All analyses were conducted using STATA release 10.1 (StataCorp, 2008).

## Results

### Sample description

A total of 390 psychosis cases and 391 healthy controls were successfully recruited to the case-control arm of the ÆSOP study in the south-east London and Nottingham sites (Morgan et al., 2006). Of these, 182 cases and 246 controls completed the CECA.Q and were included in the analyses. Those who completed the CECA.Q did not differ significantly from those who did not in terms of gender, age, ethnicity, duration of untreated psychosis or length of psychosis at interview (data, not shown, are available from the authors). Moreover, as the CECA.Q appeared near the end of the assessment battery it is

likely that the length of the assessment period rather than these specific demographic factors influenced the likelihood of it being completed. The majority of cases (61.0%) who completed the CECA.Q had a non-affective psychotic diagnosis (narrowly defined schizophrenia: 39.0% of full sample; 7.2% brief reactive psychosis; 5.5% schizoaffective; 5.5% delusional disorder; 2.7% unspecified non-organic psychosis; and 1.1% substance-induced psychosis), with the rest split between depressive (20.3%) and manic (18.7%) psychoses. The basic demographic data by case and control status for those included in the analyses are presented in Table 1.

Compared with controls, and in line with what would be expected, cases were younger, more often men, and more often of non-White ethnicity. There were also more controls from Nottingham than London. Hence, all these variables were controlled for in the adjusted analyses. There was no significant difference between cases and controls in terms of parental social class but this was also included as a confounder as it has previously been correlated with childhood adversity and psychosis (Croudace et al., 2000; Sidebotham et al., 2006).

### Characteristics of childhood adversity

**Type of exposure and perpetrator**—The prevalence of each type of childhood adversity for psychosis cases and healthy controls is provided in Table 2. Psychosis cases were approximately three times more likely than healthy controls to report severe physical abuse from their mother during childhood and this association held after adjustment for potential confounders. Reported maternal antipathy and neglect were each around twice as common amongst psychosis patients as controls. Furthermore, not having a supportive adult or peer was twice as prevalent in the psychosis group but this failed to reach conventional levels of statistical significance after adjustment. Sexual abuse was marginally more common in cases than controls but again was not statistically significant. Restricting this to incidents involving intercourse also failed to demonstrate a significant association with psychotic disorder (OR=1.68, 95% CI 0.75-3.79,  $p=0.209$ ). Moreover, no associations were found between case status and having more than three family living arrangements, being taken into local authority care or reports of physical abuse, antipathy or neglect from the main father figure (Table 2).

Further adjusting for the other significant types of adversity reduced the odds ratios for all three maternally-perpetrated adversities suggesting overlap between experiences. The association between maternal physical abuse and psychotic disorder was slightly attenuated and just failed to reach significance ( $p=0.051$ ) whilst a non-significant trend remained for maternal antipathy. No association remained for maternal neglect, probably due to a significant overlap with maternal antipathy (63.2%;  $X^2=130.17$ ,  $p<0.001$ ) and maternal physical abuse (31.6%;  $X^2=25.78$ ,  $p<0.001$ ).

**Frequency of exposure**—The proportions of psychosis cases and controls reporting one or more experiences from any of the above list (total) along with specific adversities from one or two perpetrators are presented in Table 3. The association with psychotic disorder was similar for participants who reported single and multiple adverse childhood experiences. Visual inspection of the odds ratios indicated that these were not significantly different as the confidence intervals overlapped and included the point estimates (Table 3). For the individual abuse variables, similar odds were found for frequency of exposure to physical abuse and neglect. However, there appeared to be an increasing association with psychosis from one to two severe sexual abuse experiences. A score test for trend did not provide evidence, though, for a linear trend ( $X^2=1.74$ ,  $p=0.187$ ). Similarly, reported severe antipathy



from both parents appeared to be more strongly associated with psychosis than antipathy from just one parent, but no evidence was found for a trend ( $X^2=2.07$ ,  $p=0.150$ ).

**Age of exposure**—Table 4 displays the proportion of cases and healthy controls initially exposed during childhood (0-11 years) or adolescence (12-16 years) to physical or sexual abuse. The size of the unadjusted odds ratio for maternal physical abuse reported as starting during childhood was approximately twice that for adolescent-onset abuse. In other words, the effect of maternal physical abuse appeared stronger when it commenced during childhood than during adolescence. However, the overlapping confidence intervals for these odds ratios, which also included each point estimate, indicated that this was not a significant difference (Table 4). The odds ratio for sexual abuse before the age of 12 was also marginally larger than that for adolescent-onset experiences but again not to a significant degree. No significant association was found between paternal physical abuse and psychosis regardless of timing.

Following adjustment for all confounders, severe physical abuse from the main mother figure that commenced in childhood was the only adversity that remained statistically precise with regards to association with psychotic disorder. A trend was also evident for around a two-fold greater prevalence of reported childhood sexual abuse amongst psychosis cases compared to controls but this failed to reach statistical significance and was further attenuated after adjustment for maternal physical abuse and antipathy (OR=1.60, 95% CI 0.70-3.62,  $p=0.263$ ). However, the association between maternal physical abuse that started before the age of 12 and psychotic disorder remained after controlling for all confounders, childhood sexual abuse, maternal antipathy and neglect (OR=3.01, 95% CI 1.02-8.89,  $p=0.046$ ).

## Discussion

Initially, several adverse experiences related to the main mother figure were found to be significantly associated with psychotic disorder following adjustment for demographic confounders. However, further analysis revealed that reports of severe physical abuse from mother that started prior to 12 years of age had the most robust association with psychotic disorder. Severe antipathy from mother demonstrated a non-significant trend for approximately a two-fold association with psychosis, independent of maternal physical abuse. In absolute terms, though, the vast majority of psychosis cases (86%) and controls (93%) in this study did *not* report maternal physical or emotional abuse. No associations with psychotic disorder were evident for maltreatment by the main father figure during childhood, nor for the other forms of adversity. There was also no clear evidence of a dose-response effect in this sample in terms of total number of adversities experienced or exposure to individual types of childhood adversity by different perpetrators.

### Comparisons with previous research

**Prevalence rates**—Previous studies of individuals diagnosed with psychotic disorders have reported huge variations in prevalence rates of adversity probably reflecting vast differences in the way abuse is defined, length of illness, size and selection of the sample, demographic and diagnostic composition etc. (Bendall et al., 2008; Fisher & Craig, 2008; Morgan & Fisher, 2007). If comparisons are restricted to the largest previous study of first-episode psychosis patients (Neria et al., 2002), then the rates reported in the current study are slightly lower for physical abuse (21.6% vs. 14.0-15.2%) but fairly similar for sexual abuse (15.0% vs. 18.2%). The discrepancy for physical abuse rates could potentially be due to the broader definition used in the Neria et al. (2002) study that may have resulted in

inclusion of less severe abuse than the present study, thus potentially rendering the current estimate more conservative.

It is reassuring to note, though, that the prevalence of physical and sexual abuse reported by healthy controls in this study (5.4-12.3% and 13.1%, respectively) is similar to that found in a recent UK general population survey (May-Chahal & Cawson, 2005: 7-14% physical, 11% sexual) despite the comparatively lower parental social class of our participants. Our rates are also within the range of international abuse estimates (WHO, 2002: 5-20% sexual abuse; Gilbert et al., 2009: 4-16% physical abuse). Therefore, the associations found in the current study do not simply seem to be the result of under-reporting amongst controls and indeed our control sample appears reasonably representative of the general population in terms of abuse prevalence.

**Specific types of adversity associated with psychotic disorder**—Childhood-onset severe maternal physical abuse was the most robust risk indicator of psychosis in this sample. This is in contrast to previous research which has shown associations primarily between childhood sexual abuse and adult psychosis (Briere et al., 1997; Hammersley et al. 2003; Heins et al., 1990). Nevertheless, physical abuse has been found to have stronger associations than sexual abuse with other mental health problems, when the two forms of abuse have been clearly distinguished from each other (Coid et al., 2003; Mulder et al., 1998). Furthermore, Rubino et al. (2009) reported an almost six-fold greater prevalence of physical abuse amongst schizophrenia patients than controls but only a non-significant trend for sexual abuse. Hence, it is possible that previous associations between sexual abuse and psychosis may have been inflated by overlap with physical abuse. Moreover, Shevlin et al. (2007a) reported that physical abuse produced the most robust association with psychosis, as was the case in our clinical sample, with a similar three-fold elevation in risk.

Maternal antipathy was also found to have a reasonably independent association with psychotic disorder. This is consistent with previous studies in which reports of emotional abuse were significantly more prevalent amongst individuals diagnosed with psychosis than general population (Rubino et al., 2009) or medical (Schofield & Balian, 1959) controls. Higher rates of psychotic symptoms have also been found in those exposed to emotional forms of abuse (Colins et al., 2009; Sommer et al., 2008). This 'active' type of maltreatment in combination with maternal physical abuse has been demonstrated to be of central aetiological significance in depression (Brown et al., 2007). Therefore, contrary to Harris' (1987) original proposition about the importance of such intrusive experiences for psychosis rather than depression it may actually be the case that the same types of adversity are key for both disorders. Further exploration of this potential overlap is required, though initial findings indicate that the effect may simply be greater in psychosis (Rubino et al., 2009).

**Relationship to perpetrator**—Early severe abuse from mothers rather than fathers demonstrated the greatest association with psychotic disorder in this sample. This is reasonably consistent with a recent study in depression (Brown et al., 2007) but is contrary to Parker et al. (1982) who found stronger associations between schizophrenia and fathers' parenting style. However, as to our knowledge no research has investigated differential parental effects of severe abuse in psychosis these comparisons are limited. Nevertheless, the importance of the mother in the psychological development of the child has long been advocated as she is usually the main attachment figure in infancy (Bowlby, 1951, 1977). Although fathers often develop significant bonds with their children (Cohen & Campos, 1974; Steele et al., 1996), their greater absence during childhood amongst individuals who later develop psychosis (Agid et al., 1999; Morgan et al., 2007) may mean they have less opportunity to directly inflict abuse upon their offspring. Nonetheless, in the current study maternal abuse is still only present in a minority of participants with clinical disorder

indicating that several other risk factors must be involved. Careful replication of these findings is required along with prospective longitudinal and adoption studies to investigate causality.

**Dose-response effect**—This study did not provide evidence of a dose-response effect for multiple versus single adversity experiences in childhood on development of psychosis nor were there significant effects for number of perpetrators. This is in contrast to previous studies which have demonstrated a cumulative effect of trauma on risk for psychosis in the general population (Janssen et al., 2004; Shevlin et al., 2007a,b, 2008; Spauwen et al., 2006; Whitfield et al., 2005) and one study in patients with schizophrenia (Rubino et al., 2009). However, as formal tests of trend have rarely been conducted it is difficult to draw definitive conclusions from the existing research. Moreover, the current findings are consistent with Schilling et al. (2008) who concluded that the severity of childhood trauma experienced was more important in terms of later mental health outcomes than a simple cumulative adversity score. Similarly, Clausen and Crittenden (1991) argued that single instances of certain types of abuse (e.g., physical or sexual) may be traumatic enough to produce detrimental effects whilst other adverse experiences may require repeated exposure to cause harm to the child.

**Timing of exposure**—The results of the current study tentatively indicated that the timing of some adverse events may be important in terms of later development of psychosis. Childhood-onset adversity has previously been linked with more severe and persistent mental health difficulties (e.g., Blaauw et al., 2002; Keiley et al., 2001) even when affected individuals are subsequently exposed to more beneficial environments (van der Vegt et al., 2009). The evidence is sparse though and therefore further research is essential. It is also plausible that stronger effects in childhood are just a proxy for more prolonged abuse or higher risk of exposure to other adverse experiences. Indeed, Thornberry et al. (2001) determined that only childhood-onset abuse that persists through adolescence had a significant impact on later outcomes. Unfortunately, in the current study no information was available about the length of maltreatment exposure and therefore we were not able to investigate this possibility.

### Methodological considerations

One potential limitation is the reliance on retrospective reporting of abuse which may be particularly problematic in psychosis patients (Howard, 1993; Young et al., 2001). The childhood adversity data obtained on this psychosis sample, though, has been demonstrated to have reasonable levels of test-retest reliability, convergent and concurrent validity, and no measurable impact of current symptoms was found (Fisher et al., 2009a). Nevertheless, the possibility of false positives or negatives cannot be entirely ruled out. Moreover, retrospective dating of the timing of abuse is thought to be fairly unreliable (Hardt & Rutter, 2004). Although, the use of broad age periods (childhood vs. adolescence) should minimise the impact of such memory biases, it seems prudent to only draw tentative conclusions about the relative importance of adversities occurring prior to age 12 years in the later development of psychosis.

Despite reasonable sample sizes for psychosis patients and healthy controls for a clinical study in this field, a large number of analyses were performed on this data giving rise to the possibility of spurious statistical associations. Moreover, there was insufficient power to detect all potential associations with adversity, especially after adjusting for all confounders. Consequently, the results presented here should be interpreted with caution and require replication in an even larger epidemiological sample.



Furthermore, there was insufficient power to investigate associations with specific diagnoses, individual symptoms, or potential mediators or moderators of the association between childhood abuse and psychosis. For instance, substance misuse (Houston et al., 2008; Whitfield et al., 2005), adult victimisation (Briere et al., 1997) and comorbid psychiatric disorders, such as anxiety, depression or PTSD (Garety et al., 2007; Kilcommons & Morrison, 2005; Shevlin et al., 2007a), could have confounded the results of the current study and should be taken into account in future research in this area. It is also possible that an undetected gene-environment correlation (Plomin et al., 1977) may underlie the results, though recent research suggests that controlling for parental mental illness does not attenuate associations between childhood adversity and psychosis (Fisher et al., 2009b; Kelleher et al., 2008).

Additionally, utilisation of the CECA.Q in this study meant that other potentially predictive forms of childhood adversity were not measured, such as bullying, psychological abuse, domestic violence and family discord (see Bebbington et al., 2004; Campbell & Morrison, 2007; Kelleher et al., 2008; Lataster et al., 2006; Rosenberg et al., 2007). Moreover, the reliance on self-reported adversity using a questionnaire format rather than an in-depth interview may have resulted in under-reporting of adversity. Therefore, future research needs to include comprehensive assessments of an even wider range of adverse childhood experiences to fully explore the associations with clinically diagnosed psychosis.

## Summary

In conclusion, reports of severe childhood physical abuse from the main mother figure were found to have the most robust association with psychotic disorders, particularly when the abuse began prior to 12 years of age. Paternal maltreatment was not shown to have any clear aetiological significance for psychotic disorders in adulthood. Therefore, specific characteristics of childhood adversity appear to be important in relation to clinical psychosis. If replicated, these findings have potentially important implications for future research into the aetiological mechanisms operating between childhood adversity and later psychosis. Moreover, the elevated frequency of childhood abuse amongst psychosis patients in this sample supports recent guidance for mental health services in the UK that requires clinicians to routinely assess all patients for early abusive experiences and provide suitable interventions (NHS Confederation, 2008).

## Acknowledgments

We thank all staff members and students past and present who were involved in the *ÆSOP* study. We are also indebted to all individuals who participated in the study and were essential for its successful completion. This work was supported by the UK Medical Research Council (MRC) and the Stanley Medical Research Institute. H.L.F. is jointly funded by the MRC and the Economic and Social Research Council. All authors declare that they have no financial conflicts of interest.

## References

- Agid O, Shapira B, Zislin J, Ritsner M, Hanin B, Murad H, Troudart T, Bloch M, Heresco-Levy U, Lerer B. Environment and vulnerability to major psychiatric illness: a case control study of early parental loss in major depression, bipolar disorder and schizophrenia. *Molecular Psychiatry*. 1999; 4:163–172. [PubMed: 10208448]
- Beattie N, Shannon C, Kavanagh M, Mulholland C. Predictors of PTSD symptoms in response to psychosis and psychiatric admission. *Journal of Nervous and Mental Disease*. 2009; 197(1):56–60. [PubMed: 19155811]
- Bebbington PE, Bhugra D, Brugha T, Singleton N, Farrell M, Jenkins R, Lewis G, Meltzer H. Psychosis, victimization and childhood disadvantage. *British Journal of Psychiatry*. 2004; 185:220–226. [PubMed: 15339826]

- Bebbington P, Nayani T. The Psychosis Screening Questionnaire. *International Journal of Methods in Psychiatric Research*. 1995; 5:11–20.
- Bendall S, Jackson HJ, Hulbert CA, McGorry PD. Childhood trauma and psychotic disorders: a systematic, critical review of the evidence. *Schizophrenia Bulletin*. 2008; 34(3):568–579. [PubMed: 18003630]
- Bernstein, DP.; Fink, L. *Childhood Trauma Questionnaire: A Retrospective Self-report Manual*. The Psychological Corporation; New York, NY: 1998.
- Bifulco A, Bernazzani O, Moran PM, Jacobs C. The Childhood Experiences of Care and Abuse Questionnaire (CECA.Q) – validation in a community series. *British Journal of Clinical Psychology*. 2005; 44:563–581. [PubMed: 16368034]
- Blaauw E, Arensman E, Kraaij V, Winkel FW, Bout R. Traumatic life events and suicide risk among jail inmates: the influence of types of events, time period and significant others. *Journal of Traumatic Stress*. 2002; 15(1):9–16. [PubMed: 11936726]
- Bowlby, J. *Maternal Care and Mental Health*. Schocken; New York, NY: 1951.
- Bowlby J. The making and breaking of affectional bonds. I. Aetiology and psychopathology in the light of attachment theory. An expanded version of the Fiftieth Maudsley Lecture, delivered before the Royal College of Psychiatrists, 19 November 1976. *British Journal of Psychiatry*. 1977; 130:201–210. [PubMed: 843768]
- Briere J, Woo R, McRae B, Foltz J, Sitzman R. Lifetime victimization history, demographics and clinical status in female psychiatric emergency room patients. *Journal of Nervous and Mental Disease*. 1997; 185(2):95–101. [PubMed: 9048701]
- Brown GW, Craig TK, Harris TO, Handley RV, Harvey AL. Development of a retrospective interview measure of parental maltreatment using the Childhood Experience of Care and Abuse (CECA) instrument -- A life-course study of adult chronic depression - I. *Journal of Affective Disorders*. 2007; 103(1-3):205–215. [PubMed: 17651811]
- Campbell ML, Morrison AP. The relationship between bullying, psychotic-like experiences and appraisals in 14–16-year olds. *Behaviour, Research and Therapy*. 2007; 45:1579–1591. [PubMed: 17229400]
- Clausen AH, Crittenden PM. Physical and psychological maltreatment: relations among types of treatment. *Child Abuse and Neglect*. 1991; 15:5–18.
- Cohen LJ, Campos JJ. Father, mother, and stranger as elicitors of attachment behaviors in infancy. *Developmental Psychology*. 1974; 10:146–154.
- Coid J, Petruckevitch A, Chung W-S, Richardson J, Moorey S, Feder G. Abusive experiences and psychiatric morbidity in women primary care attenders. *British Journal of Psychiatry*. 2003; 183:332–339. [PubMed: 14519611]
- Colins O, Vermeiren R, Vreugdenhil C, Schuyten G, Broekaert E, Krabbendam A. Are psychotic experiences among detained juvenile offenders explained by trauma and substance use? *Drug and Alcohol Dependence*. 2009; 100(1-2):39–46. [PubMed: 19010608]
- Croudace TJ, Kayne R, Jones PB, Harrison GL. Non-linear relationship between an index of social deprivation, psychiatric admission prevalence and the incidence of psychosis. *Psychological Medicine*. 2000; 30(1):177–185. [PubMed: 10722188]
- Fisher, H.; Craig, T. Childhood adversity and psychosis. In: Morgan, C.; McKenzie, K.; Fearon, P., editors. *Society and Psychosis*. Cambridge University Press; Cambridge: 2008.
- Fisher HL, Craig TK, Fearon P, Morgan K, Dazzan P, Lappin J, Hutchinson G, Doody GA, Jones PB, McGuffin P, Murray RM, Leff J, Morgan C. Reliability and comparability of psychosis patients' retrospective reports of childhood abuse. *Schizophrenia Bulletin*. 2009a Published online: 7th October 2009. doi: 10.1093/schbul/sbp103.
- Fisher H, Morgan C, Dazzan P, Craig TK, Morgan K, Hutchinson G, Jones PB, Doody GA, Pariante C, McGuffin P, Murray RM, Leff J, Fearon P. Gender differences in the association between childhood abuse and psychosis. *British Journal of Psychiatry*. 2009b; 194:319–325. [PubMed: 19336782]
- Garety PA, Bebbington P, Fowler D, Freeman D, Kuipers E. Implications for neurobiological research of cognitive models of psychosis: a theoretical paper. *Psychological Medicine*. 2007; 37(10): 1377–1391. [PubMed: 17335638]

- Gilbert R, Widom CS, Browne K, Fergusson D, Webb E, Janson S. Burden and consequences of child maltreatment in high-income countries. *Lancet*. 2009; 373(9657):68–81. [PubMed: 19056114]
- Hammersley P, Dias A, Todd G, Bowen-Jones K, Reilly B, Bentall RP. Childhood trauma and hallucinations in bipolar affective disorder: preliminary investigation. *British Journal of Psychiatry*. 2003; 182:543–547. [PubMed: 12777347]
- Hardt J, Rutter M. Validity of adult retrospective reports of adverse childhood experiences: a review of the evidence. *Journal of Child Psychology and Psychiatry*. 2004; 45:260–273. [PubMed: 14982240]
- Harris, T. Recent developments in the study of life events in relation to psychiatric and physical disorders. In: Cooper, B., editor. *Psychiatric Epidemiology: Progress and Prospects*. Croom Helm; London: 1987.
- Heads TC, Taylor PJ, Leese M. Childhood experiences of patients with schizophrenia and a history of violence: a special hospital sample. *Criminal Behaviour and Mental Health*. 1997; 7:117–130.
- Heins T, Gray A, Tennant M. Persisting hallucinations following childhood sexual abuse. *Australian and New Zealand Journal of Psychiatry*. 1990; 24:561–565. [PubMed: 2073233]
- Hlastala SA, McClellan J. Phenomenology and diagnostic stability of youths with atypical psychotic symptoms. *Journal of Child and Adolescent Psychopharmacology*. 2005; 15:497–509. [PubMed: 16092913]
- Houston JE, Murphy J, Adamson G, Stringer M, Shevlin M. Childhood sexual abuse, early cannabis use, and psychosis: testing an interaction model based on the National Comorbidity Survey. *Schizophrenia Bulletin*. 2008; 34(3):580–585. [PubMed: 18024467]
- Howard LM. Allegations of abuse in psychotic patients. *American Journal of Psychiatry*. 1993; 150:839–840. [PubMed: 8480841]
- Janssen I, Krabbendam L, Bak M, Hanssen M, Vollebergh W, de Graaf R, van Os J. Childhood abuse as a risk factor for psychotic experiences. *Acta Psychiatrica Scandinavica*. 2004; 109:38–45. [PubMed: 14674957]
- Jenkins R, Meltzer H. The national survey of psychiatric morbidity in Great Britain. *Social Psychiatry and Psychiatric Epidemiology*. 1995; 30:1–4. [PubMed: 7892612]
- Jones P, Rodgers B, Murray R, Marmot M. Child development risk factors for adult schizophrenia in the British 1946 birth cohort. *Lancet*. 1994; 344(8934):1398–1402. [PubMed: 7968076]
- Keiley MK, Howe TR, Dodge KA, Bates JE, Petti GS. The timing of child physical maltreatment: a cross-domain growth analysis of impact on adolescent externalizing and internalizing problems. *Development and Psychopathology*. 2001; 13(4):891–912. [PubMed: 11771913]
- Kelleher I, Harley M, Lynch F, Arseneault L, Fitzpatrick C, Cannon M. Associations between childhood trauma, bullying and psychotic symptoms among a school-based adolescent sample. *British Journal of Psychiatry*. 2008; 193(5):378–382. [PubMed: 18978317]
- Kilcommons AM, Morrison AP. Relationships between trauma and psychosis: an exploration of cognitive and dissociative factors. *Acta Psychiatrica Scandinavica*. 2005; 112(5):351–359. [PubMed: 16223422]
- Kim D, Kaspar V, Noh S, Nam JH. Sexual and physical abuse among Korean female inpatients with schizophrenia. *Journal of Traumatic Stress*. 2006; 19:279–289. [PubMed: 16612823]
- Kirkbride J, Fearon P, Morgan C, Dazzan P, Morgan K, Tarrant J, Fung WL, Holloway J, Mallett R, Harrison G, Leff J, Jones PB, Murray RM, AESOP Study Group. Heterogeneity in incidence rates of schizophrenia and other psychotic syndromes: findings from the 3-center AeSOP study. *Archives of General Psychiatry*. 2006; 63:250–258. [PubMed: 16520429]
- Lataster T, van Os J, Drukker M, Henquet C, Feron F, Gunther N, Myin-Germeys I. Childhood victimisation and developmental expression of nonclinical delusional ideation and hallucinatory experiences: victimisation and nonclinical psychotic experiences. *Social Psychiatry and Psychiatric Epidemiology*. 2006; 41:423–428. [PubMed: 16572272]
- Mallett, R. *Sociodemographic Schedule*. Section of Social Psychiatry, Institute of Psychiatry; London: 1997.
- May-Chahal C, Cawson P. Measuring child maltreatment in the United Kingdom: a study of the prevalence of child abuse and neglect. *Child Abuse and Neglect*. 2005; 29:969–984. [PubMed: 16165212]

- McNeil TF, Schubert EW, Cantor-Graae E, Brossner M, Schubert P, Henriksson KM. Unwanted pregnancy as a risk factor for offspring schizophrenia-spectrum and affective disorders in adulthood: a prospective high-risk study. *Psychological Medicine*. 2009; 39(6):957–965. [PubMed: 18945377]
- Morgan C, Dazzan P, Morgan K, Jones P, Harrison G, Leff J, Murray R, Fearon P, AESOP study group. First episode psychosis and ethnicity: initial findings from the AESOP study. *World Psychiatry*. 2006; 5(1):40–46. [PubMed: 16757995]
- Morgan C, Fisher H. Environment and schizophrenia: environmental factors in schizophrenia: childhood trauma--a critical review. *Schizophrenia Bulletin*. 2007; 33(1):3–10. [PubMed: 17105965]
- Morgan C, Kirkbride J, Leff J, Craig T, Hutchinson G, McKenzie K, Morgan K, Dazzan P, Doody GA, Jones P, Murray R, Fearon P. Parental separation, loss and psychosis in different ethnic groups: a case-control study. *Psychological Medicine*. 2007; 37(4):495–503. [PubMed: 17094816]
- Mulder RT, Beautrais AL, Joyce PR, Fergusson DM. Relationship between dissociation, childhood sexual abuse, childhood physical abuse, and mental illness in a general population sample. *American Journal of Psychiatry*. 1998; 155:806–811. [PubMed: 9619154]
- Myhrman A, Rantakallio P, Isohanni M, Jones P, Partanen U. Unwantedness of a pregnancy and schizophrenia in the child. *British Journal of Psychiatry*. 1996; 169(5):637–640. [PubMed: 8932895]
- Neria Y, Bromet EJ, Sievers S, Lavelle J, Fochtmann LJ. Trauma exposure and posttraumatic stress disorder in psychosis: findings from a first-admission cohort. *Journal of Consulting and Clinical Psychology*. 2002; 70(1):246–251. [PubMed: 11860051]
- NHS Confederation. Implementing National Policy on Violence and Abuse. The NHS Confederation; London: 2008.
- Office of National Statistics. The National Statistics Socio-economic Classification User Manual. Office for National Statistics; London: 2002.
- Parker G, Fairley M, Greenwood J, Jurd S, Silove D. Parental representations of schizophrenics and their association with onset and course of schizophrenia. *British Journal of Psychiatry*. 1982; 141:573–581. [PubMed: 7159804]
- Plomin R, DeFries JC, Loehlin JC. Genotype-environment interaction and correlation in the analysis of human behaviour. *Psychological Bulletin*. 1977; 84:309–322. [PubMed: 5572111]
- Romero S, Birmaher B, Axelson D, Goldstein T, Goldstein BI, Gill MK, Iosif AM, Strober MA, Hunt J, Esposito-Smythers C, Ryan ND, Leonard H, Keller M. Prevalence and correlates of physical and sexual abuse in children and adolescents with bipolar disorder. *Journal of Affective Disorders*. 2009; 112(1-3):144–150. [PubMed: 18538857]
- Rosenberg SD, Lu W, Mueser KT, Jankowski MK, Cournos F. Correlates of adverse childhood events among adults with schizophrenia spectrum disorders. *Psychiatric Services*. 2007; 58(2):245–253. [PubMed: 17287383]
- Rubino IA, Nanni RC, Pozzi DM, Siracusano A. Early adverse experiences in schizophrenia and unipolar depression. *Journal of Nervous and Mental Disease*. 2009; 197(1):65–68. [PubMed: 19155813]
- Schäfer I, Harfst T, Aderhold V, Briken P, Lehmann M, Moritz S, Read J, Naber D. Childhood trauma and dissociation in female patients with schizophrenia spectrum disorders: an exploratory study. *Journal of Nervous and Mental Disease*. 2006; 194(2):135–138. [PubMed: 16477194]
- Schenkel L, Spaulding W, DiLillo D, Silverstein S. Histories of childhood maltreatment in schizophrenia: relationships with premorbid functioning, symptomatology, and cognitive deficits. *Schizophrenia Research*. 2005; 76(2):273–286. [PubMed: 15949659]
- Schilling EA, Aseltine RH Jr, Gore S. The impact of cumulative childhood adversity on young adult mental health: Measures, models, and interpretations. *Social Science and Medicine*. 2008; 66:1140–1151. [PubMed: 18177989]
- Schofield W, Balian L. A comparative study of the personal histories of schizophrenic and nonpsychiatric patients. *Journal of Abnormal Social Psychology*. 1959; 59:216–225.
- Shevlin M, Dorahy MJ, Adamson G. Trauma and psychosis: an analysis of the National Comorbidity Survey. *American Journal of Psychiatry*. 2007a; 164(1):166–169. [PubMed: 17202562]

- Shevlin M, Dorahy MJ, Adamson G. Childhood traumas and hallucinations: an analysis of the National Comorbidity Survey. *Journal of Psychiatric Research*. 2007b; 41:222–228. [PubMed: 16643948]
- Shevlin M, Houston JE, Dorahy MJ, Adamson G. Cumulative traumas and psychosis: an analysis of the National Comorbidity Survey and the British Psychiatric Morbidity Survey. *Schizophrenia Bulletin*. 2008; 34(1):193–199. [PubMed: 17586579]
- Sidebotham P, Heron J, ALSPAC Study Team. Child maltreatment in the “children of the nineties”: a cohort study of risk factors. *Child Abuse and Neglect*. 2006; 30(5):497–522. [PubMed: 16701895]
- Smith N, Lam D, Bifulco A, Checkley S. Childhood Experience of Care and Abuse Questionnaire (CECA.Q): validation of a screening instrument for childhood adversity in clinical populations. *Social Psychiatry and Psychiatric Epidemiology*. 2002; 37:572–579. [PubMed: 12545234]
- Sommer IE, Daalman K, Rietkerk T, Diederer KM, Bakker S, Wijkstra J, Boks MP. Healthy individuals with auditory verbal hallucinations; who are they? Psychiatric assessments of a selected sample of 103 subjects. *Schizophrenia Bulletin*. 2008 Published online: 9 October 2008. doi:10.1093/schbul/sbn130.
- Spauwen J, Krabbendam L, Lieb R, Wittchen HU, van Os J. Impact of psychological trauma on the development of psychotic symptoms: relationship with psychosis proneness. *British Journal of Psychiatry*. 2006; 188:527–533. [PubMed: 16738342]
- StataCorp. *Stata Statistical Software: Release 10.1*. Stata Corporation; College Station, TX: 2008.
- Steele H, Steele M, Fonagy P. Associations among attachment classifications of mothers, fathers, and their infants. *Child Development*. 1996; 67:541–555. [PubMed: 8625727]
- Thornberry TP, Ireland TO, Smith CA. The importance of timing: the varying impact of childhood and adolescent maltreatment on multiple problem outcomes. *Development and Psychopathology*. 2001; 13:957–979. [PubMed: 11771916]
- van der Vegt EJ, van der Ende J, Ferdinand RF, Verhulst FC, Tiemeier H. Early childhood adversities and trajectories of psychiatric problems in adoptees: evidence for long lasting effects. *Journal of Abnormal Child Psychology*. 2009; 37(2):239–249. [PubMed: 18807165]
- Weber K, Rockstroh B, Borgelt J, Awiszus B, Popov T, Hoffmann K, Schonauer K, Watzl H, Pröpster K. Stress load during childhood affects psychopathology in psychiatric patients. *BMC Psychiatry*. 2008; 8:63. [PubMed: 18651952]
- Wexler BE, Lyons L, Lyons H, Mazure CM. Physical and sexual abuse during childhood and development of psychiatric illnesses during adulthood. *Journal of Nervous and Mental Disease*. 1997; 185(8):522–524. [PubMed: 9284868]
- Whitfield C, Dube S, Felitti V, Anda R. Adverse childhood experiences and hallucinations. *Child Abuse and Neglect*. 2005; 29:797–810. [PubMed: 16051353]
- Widom CS, Czaja SJ, Dutton MA. Childhood victimization and lifetime revictimization. *Child Abuse and Neglect*. 2008; 32(8):785–796. [PubMed: 18760474]
- World Health Organisation. *The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines*. World Health Organisation; Geneva: 1992.
- World Health Organisation. *Schedules for the Clinical Assessment of Neuropsychiatry*. World Health Organisation; Geneva: 1994.
- World Health Organisation. *The World Health Report 2002: Reducing Risks, Promoting Healthy Life*. World Health Organisation; Geneva: 2002.
- Young M, Read J, Barker-Collo S, Harrison R. Evaluating and overcoming barriers to taking abuse histories. *Professional Psychology, Research and Practice*. 2001; 32:407–414.



Table 1

Basic demographic characteristics of psychosis cases and controls.

Demographic variable	Cases (n=182) n (%)	Controls (n=246) n (%)	$\chi^2$	df	P
Sex			6.024	1	0.015*
Male	98 (53.8)	103 (41.9)			
Female	84 (46.2)	143 (58.1)			
Ethnicity			31.700	5	<0.001**
White British	102 (56.1)	182 (74.1)			
Black Caribbean	36 (19.8)	35 (14.2)			
Black African	14 (7.7)	3 (1.2)			
White Other	9 (4.9)	19 (7.7)			
Asian (all)	12 (6.6)	3 (1.2)			
Other	9 (4.9)	4 (1.6)			
Study Centre			6.525	1	0.012*
London	82 (45.1)	81 (32.9)			
Nottingham	100 (54.9)	165 (67.1)			
Parental social class <sup>a</sup>			0.891	2	0.641
Managerial, professional	45 (27.8)	62 (25.9)			
Intermediate	44 (27.2)	58 (24.3)			
Routine, manual	73 (45.0)	119 (49.8)			
Mean Age in years (S.D.)	31 (11.3)	39 (12.7)	6.498	426	<0.001**

<sup>a</sup>Missing values: cases = 20 (11.0%), controls = 7 (2.8%).  $\chi^2$ , chi-square statistic; df, degrees of freedom; P, p value indicating level of significance; S.D., standard deviation; t, t-test statistic.

\*  $P < 0.05$

\*\*  $P < 0.01$ .

Table 2

Association between type of adversity and psychotic disorder.

Type of Adversity	Cases n/N (%)	Controls n/N (%)	Unadjusted OR <sup>a</sup>	95% CI	P	Adjusted OR <sup>a,b</sup>	95% CI	P	Adjusted OR <sup>a,b,c</sup>	95% CI	P
Physical Abuse – mother	25/178 (14.0)	13/242 (5.4)	3.28	1.59 - 6.77	0.001**	2.91	1.25 - 6.79	0.013*	2.35	1.00-5.52	0.051
Physical Abuse – father	27/178 (15.2)	30/243 (12.3)	1.32	0.75 - 2.34	0.340	1.22	0.66 - 2.25	0.521	-	-	-
Sexual Abuse	32/176 (18.2)	31/237 (13.1)	1.40	0.81 - 2.40	0.229	1.60	0.87 - 2.95	0.133	-	-	-
Antipathy – mother	25/182 (13.7)	18/246 (7.3)	2.15	1.12 - 4.13	0.022*	2.72	1.25 - 5.94	0.012*	2.07	0.82-5.18	0.122
Antipathy – father	22/182 (12.1)	29/246 (11.8)	1.04	0.57 - 1.89	0.900	0.92	0.46 - 1.82	0.804	-	-	-
Neglect – mother	20/182 (11.0)	18/246 (7.3)	1.68	0.84 - 3.33	0.141	2.23	1.03 - 4.83	0.041*	1.24	0.50-3.10	0.644
Neglect – father	23/182 (12.6)	38/246 (15.4)	0.84	0.48 - 1.49	0.555	0.77	0.39 - 1.51	0.441	-	-	-
Taken into care	8/181 (4.4)	15/244 (6.1)	0.73	0.30 - 1.79	0.496	0.54	0.19 - 1.52	0.245	-	-	-
Disrupted living arrangements	40/182 (22.0)	40/246 (16.3)	1.50	0.91 - 2.47	0.108	1.28	0.75 - 2.18	0.374	-	-	-
Lack of support figure	25/182 (13.7)	17/246 (6.9)	2.07	1.07 - 3.99	0.030*	2.01	0.90 - 4.46	0.087	-	-	-

<sup>a</sup> Odds ratios (OR) calculated using weighted data.<sup>b</sup> OR adjusted for gender, age, ethnicity, study centre and highest parental social class.<sup>c</sup> OR adjusted for other significant adversities. CI, confidence interval; P, p value indicating level of significance.\*  $p < 0.05$ \*\*  $p < 0.01$ .

Table 3

Association between frequency of exposure to adversity and psychotic disorder.

Type and level of adversity	Cases n/N (%)	Controls n/N (%)	Unadjusted OR <sup>a</sup>	95% CI	P	Adjusted OR <sup>a,b</sup>	95% CI	P
Total								
One type of adversity	41/175 (23.4)	49/235 (20.9)	1.46	0.88 – 2.43	0.146	1.39	0.78 – 2.48	0.259
Multiple adversities	59/175 (33.7)	59/235 (25.1)	1.80	1.13 – 2.88	0.014*	1.75	1.03 – 2.99	0.040*
Physical Abuse								
One parent	32/178 (18.0)	25/242 (10.3)	2.05	1.15 – 3.67	0.015*	1.79	0.92 – 3.47	0.087
Both parents	10/178 (5.6)	9/242 (3.7)	1.88	0.73 – 4.85	0.192	1.63	0.59 – 4.50	0.344
Sexual Abuse								
One incident	21/173 (12.1)	24/236 (10.2)	1.17	0.63 – 2.20	0.621	1.51	0.74 – 3.09	0.257
Multiple incidents	8/173 (4.6)	6/236 (2.5)	1.89	0.63 – 5.64	0.255	1.49	0.46 – 4.91	0.508
Antipathy								
One parent	33/182 (18.1)	37/246 (15.0)	1.33	0.78 – 2.25	0.286	1.37	0.73 – 2.54	0.327
Both parents	7/182 (3.9)	5/246 (5.0)	2.05	0.63 – 6.72	0.236	2.07	0.62 – 6.88	0.236
Neglect								
One parent	29/182 (15.9)	34/246 (13.8)	1.28	0.74 – 2.22	0.381	1.20	0.62 – 2.29	0.588
Both parents	7/182 (3.9)	11/246 (4.5)	0.92	0.34 – 2.47	0.866	1.19	0.39 – 3.68	0.760

<sup>a</sup>Odds ratios (OR) calculated using weighted data.<sup>b</sup>OR adjusted for gender, age, ethnicity, study centre and highest parental social class. CI, confidence interval; P, p value indicating level of significance.

\* P&lt;0.05.

Table 4

Association between timing of adversity and psychotic disorder.

Type of Adversity	Cases n/N (%)	Controls n/N (%)	Unadjusted OR <sup>a</sup>	95% CI	P	Adjusted OR <sup>a,b</sup>	95% CI	P
Physical Abuse – mother								
Childhood	22/178 (12.4)	9/240 (3.8)	4.34	1.89-10.00	0.001**	3.60	1.36 - 9.55	0.010*
Adolescence	3/178 (1.7)	2/240 (0.8)	2.05	0.34-12.44	0.436	3.02	0.42 - 21.54	0.270
Physical Abuse – father								
Childhood	21/176 (11.9)	22/239 (9.2)	1.39	0.73 - 2.64	0.322	1.29	0.64 - 2.59	0.449
Adolescence	5/176 (2.8)	4/239 (1.7)	1.86	0.48 - 7.22	0.371	1.80	0.39 - 8.30	0.449
Sexual Abuse								
Childhood	20/175 (11.4)	15/235 (6.4)	1.83	0.90 - 3.74	0.096	2.09	0.95 - 4.56	0.065
Adolescence	11/175 (6.3)	14/235 (6.0)	1.06	0.47 - 2.41	0.889	1.26	0.49 - 3.28	0.631

<sup>a</sup> Odds ratios (OR) calculated using weighted data.<sup>b</sup> OR adjusted for gender, age, ethnicity, study centre and highest parental social class. CI, confidence interval; P, p value indicating level of significance.\*  $P < 0.05$ \*\*  $P < 0.01$ .