

---

**The sequence of the 5S ribosomal RNA of the crustacean *Artemia salina***

---

---

**Ludo Diels, Raymond De Baere, Antoon Vandenberghe and Rupert De Wachter**

---

---

**Departement Celbiologie, Universiteit Antwerpen, Universiteitsplein 1, B-2610 Wilrijk, Belgium**

---

---

**Received 30 July 1981**

---

**ABSTRACT**

The primary structure of the 5 S rRNA isolated from the cryptobiotic cysts of the brine shrimp *Artemia salina* is pACCAACGGCCAUACCACGUUGAAAGU ACCCAGUCUCGUCAGAUCCUGGAAGUCACACAACGUCGGGCCGGUCAGUACUUGGAUGGGUG ACCGCCUGGGAACACCGGGUGCUGUUGGCAU<sub>OH</sub>.

**INTRODUCTION**

The availability of gel sequencing techniques applicable to small RNA molecules<sup>1-5</sup> is resulting in a rapid growth of the collection of known 5 S rRNA primary structures<sup>6</sup>. These data can be used in phylogenetic studies<sup>7,8</sup>, and in attempts to derive general secondary structure models for 5 S RNA<sup>7,9-12</sup> by a search for common sequence complementarities. We report the primary structure of 5 S rRNA isolated from the cysts of the brine shrimp *Artemia salina* (Phylum arthropoda, class crustacea).

**MATERIALS AND METHODS**

Dry cysts of *Artemia salina* were obtained from San Francisco Bay Brand (Newark, California). The ribosomes were isolated according to Zasloff and Ochoa<sup>13</sup> and ribosomal RNA was obtained from these by phenol extraction<sup>14</sup>. 5 S RNA was isolated by gel filtration on Sephadex G100 as described by Monier<sup>15</sup>, except that the column was not recycled. It was further purified by electrophoresis on slab gels<sup>16</sup> containing 10% polyacrylamide, 7 M urea, 0.05 M Tris-borate buffer pH 8.3 and 0.001 M EDTA.

Three different gel-sequencing methods were used to derive the primary structure: the methods of Diels and De Wachter<sup>5</sup> and of Tanaka, Dyer and Brownlee<sup>4</sup>, both of which are two-dimensional adaptations of Stanley and Vassilenko's procedure<sup>2</sup>, and Peattie's<sup>3</sup> chemical degradation method. RNA labeled at the 3'-terminus by [<sup>32</sup>P] pCp ligation was also used for a confirmation of the terminal sequence by a moving spot analysis<sup>17</sup>. The 5'-terminal

base was determined by alkaline hydrolysis of 20  $\mu\text{g}$  unlabeled RNA, followed by HPLC identification of the resulting nucleoside bisphosphate. The hydrolysate was fractionated on a 4.6 x 250 mm Partisil SAX (Reeve Angel, Clifton, N.J.) column eluted with 120 ml of a concentration gradient of  $\text{H}_3\text{PO}_4$  adjusted to pH 2.2 with  $\text{NH}_3$ , rising from 0.01 M to 1 M in 100 min. Nucleotides are eluted in the order Cp, Ap, Up, Gp, pCp, pAp, pUp, pGp.

### RESULTS

Different gel sequencing methods were used to establish the complete sequence, as summarized in Table 1. Each part of the sequence was confirmed by at least three successful experiments. In the area of positions 62 to 70, which is rich in G's and C's probably involved in base pairing, the chemical degradation method<sup>3</sup> was preferable to the methods based on aspecific hydrolysis<sup>4,5</sup> which gave poor resolution. The 3'-terminal sequence was confirmed by a moving spot analysis of a partial hydrolysate of 5 S RNA labeled at the 3'-end with [<sup>32</sup>P] pCp. We have found it difficult to dephosphorylate the 5 S RNA without partially degrading it, hence we could not label the 5'-terminus satisfactorily. However, the 5'-terminal residue was identified unequivocally as pAp by complete alkaline hydrolysis and HPLC-analysis of unlabeled material. In Fig. 1, the sequence of *Artemia salina* 5 S RNA is compared with that of the two other arthropods thus far investigated, *Drosophila melanogaster*<sup>18</sup> and *Bombyx mori*<sup>19</sup>.

### DISCUSSION

The primary structure of *Drosophila melanogaster* 5 S rRNA was originally reported<sup>20</sup> as GGGCGC<sub>69</sub>, but later corrected<sup>21</sup> to GCGGGC<sub>69</sub>. The recent determination of the 5 S RNA gene sequence<sup>18</sup> and the comparison with other arthropods (Fig. 1) now make it evident that the first version was the correct one. The primary structure of *Artemia* 5 S rRNA differs with that of

Table 1. Methods used to establish the *Artemia* 5 S rRNA sequence

Method	Positions identified
Gel sequencing : Diels & De Wachter <sup>5</sup>	4-25,27-65,70-117
Tanaka et al. <sup>4</sup>	2-61, 70-111
Peattie <sup>3</sup>	4-120
Other : moving spot <sup>17</sup>	102-120
alkaline hydrolysis, HPLC	1




Fig. 1. Conservation of 5 S RNA primary and secondary structure within the phylum arthropods (A.s., *Artemia salina*; B.m., *Bombyx mori*; D.m., *Drosophila melanogaster*). The five double-stranded areas of our secondary structure model are indicated by the boxes A-A', B-B' etc.. Nested boxes delimit bulges and small interior loops within these areas.

*Drosophila* in 28 positions and with that of the silkworm in 21 positions. The two insect sequences show 16 differences. Among the vertebrates, the mean divergence between two organisms belonging to different classes (fish, amphibia, reptiles, birds, and mammals have been investigated) is limited to 9 positions.

A common base pairing scheme can be applied to the three arthropod sequences, as indicated by the boxes in Fig. 1. The resulting secondary structure model is illustrated with the *Artemia* 5 S RNA in Fig. 2. There now seems to be a consensus among the majority of investigators that there is a




Fig. 2. Secondary structure model for *Artemia* 5 S RNA.

## Nucleic Acids Research

---

basic difference between the secondary structures of bacterial and eukaryotic 5 S RNAs, the former having four helical regions, the latter five<sup>9-12</sup>. We do not share this opinion and we present evidence elsewhere (De Wachter, Chen and Vandenberghe, manuscript submitted) that the model of Fig. 2 applies, with minor variations, to all 5 S RNAs hitherto sequenced.

### ACKNOWLEDGMENTS

We are grateful to Drs. G.G. Brownlee and P. Arcari for instructing one of us (A.V.) in the gel sequencing method of Tanaka et al.<sup>4</sup>. This work was supported by the belgian foundation for basic research FKFD.

### REFERENCES

1. Simoncsits, A., Brownlee, G.G., Brown, R.S., Rubin, J.R., Guilley, H. (1977) *Nature* 269, 833-836.
2. Stanley, J., Vassilenko, S. (1978) *Nature* 274, 87-89.
3. Peattie, D.A. (1979) *Proc. Nat. Acad. Sci. U.S.* 76, 1760-1764.
4. Tanaka, Y., Dyer, T.A., Brownlee, G.G. (1980) *Nucl. Acids Res.* 8, 1259-1272.
5. Diels, L., De Wachter, R. (1980) *Arch. Internat. Physiol. Biochim.* 88, B26-B27.
6. Erdmann, V.A. (1981) *Nucl. Acids Res.* 9, r25-r42.
7. Hori, H., Osawa, S. (1979) *Proc. Nat. Acad. Sci. U.S.* 76, 381-385.
8. Küntzel, H., Heidrich, M., Piechulla, B. (1981) *Nucl. Acids Res.* 9, 1451-1461.
9. Fox, G.E., Woese, C.R. (1975) *Nature* 256, 505-507.
10. Hori, H., Osawa, S., Iwabuchi, M. (1980) *Nucl. Acids Res.* 8, 5535-5539.
11. Luehrsen, K.R., Fox, G.E. (1981) *Proc. Nat. Acad. Sci. U.S.* 78, 2150-2154.
12. Garrett, R.A., Douthwaite, S., Noller, H.F. (1981) *Trends Biochem. Sci.* 6, 137-139.
13. Zasloff, M., Ochoa, S. (1971) *Proc. Nat. Acad. Sci. U.S.* 68, 3059-3063.
14. Vandenberghe, A., Nelles, L., De Wachter, R. (1980) *Anal. Biochem.* 107, 369-376.
15. Monier, R. (1971) in *Procedures in Nucleic Acid Research*, Cantoni, G.L. and Davies, D.R. Eds., Vol. 2, pp. 618-622, Harper & Row, New York.
16. De Wachter, R., Fiers, W. (1971) *Methods Enzymol.* 21, 167-178.
17. Pinck, L., Pinck, M. (1979) *FEBS Lett.* 107, 61-65.
18. Tschudi, C., Pirrotta, V. (1980) *Nucl. Acids Res.* 8, 441-451.
19. Komiya, H., Kawakami, M., Shimizu, N., Takemura, S. (1980) *Nucl. Acids Res.* 8, s119-s122.
20. Benhamou, J., Jordan, B.R. (1976) *FEBS Lett.* 62, 146-149.
21. Benhamou, J., Jourdan, R., Jordan, B.R. (1977) *J. Mol. Evol.* 9, 279-298.