

The correspondence section is a public forum and, as such, is not peer-reviewed. EHP is not responsible for the accuracy, currency, or reliability of personal opinion expressed herein; it is the sole responsibility of the authors. EHP neither endorses nor disputes their published commentary.

## FDA Risk Assessment of Seafood Contamination after the BP Oil Spill

<http://dx.doi.org/10.1289/ehp.1104539>

In response to the BP oil spill of 2010, Rotkin-Ellman et al. (2012) provide a thoughtful assessment of the Food and Drug Administration (FDA) risk criteria to protect vulnerable populations from exposure to polycyclic aromatic hydrocarbons (PAHs) through seafood consumption (FDA 2010a). The FDA and the interagency partners involved in developing the seafood safety risk assessment for the BP oil spill shared the authors' goal of establishing criteria that would be protective of all affected populations. Although Rotkin-Ellman et al. present a well-intentioned case for determining PAH levels for at-risk populations, their interpretation of several factors involved in the derivation of levels of concern (LOC) differs substantially from those used in the BP oil spill risk assessment.

Human health risk assessments for environmental chemical contaminants are undertaken to develop exposure levels that are believed to be safe or associated with negligible risk. Uncertainty is inherent in any risk assessment process due to interspecies, intraspecies, and/or high-to-low dose extrapolations required for risk estimation. As a result, chemical risk assessments tend to be conservative with intentional bias on the side of safety. Unnecessarily conservative risk criteria, however, can have unintended negative consequences for human health and society. The role of risk management is to balance what is known and unknown about a particular risk with the interests of public health protection and societal values.

Development of the seafood safety risk assessment after the BP oil spill was a collaborative, highly iterative, and multiagency process including departments of health from the five states bordering the Gulf of Mexico. The criteria agreed upon for PAHs with cancer end points provide conservative estimates of contamination levels and consumption rates that, if sustained for a period of 5 years, may result in an upper bound consumer lifetime cancer risk of  $1 \times 10^{-5}$ . The agreed upon criteria for PAHs with noncancer end points are U.S. Environmental Protection Agency (EPA) reference doses for daily exposure of general and sensitive populations expected to have no significant risk of adverse effect during a lifetime of exposure (U.S. EPA 2000).

A lifetime cancer risk level of  $1 \times 10^{-5}$  implies a risk management decision to accept no more than a conservative estimate

of one additional cancer case attributable to PAH-contaminated seafood in a population of 100,000 people. This risk level is within the acceptable range of values (i.e.,  $1 \times 10^{-4}$  to  $1 \times 10^{-6}$ ) observed by public health risk managers (U.S. EPA 1998a, 2000). Bias toward safety is also inherent in the derivation and selection of a benzo[*a*]pyrene cancer potency factor (7.3 mg/kg/day) based on the 95% upper confidence limit of the dose-response curve, rather than a maximum likelihood estimate. The actual cancer risk is believed to be somewhere below this upper confidence limit, and could be as low as zero (U.S. EPA 1994). The U.S. EPA value is more conservative than recent derivations of benzo[*a*]pyrene cancer potency factors, which incorporate adjustments for exposures during infancy (Office of Environmental Health Hazard Assessment 2010).

In regard to specific issues raised about selected underlying assumptions used in the risk assessment process, the FDA respectfully disagrees with the arguments of Rotkin-Ellman et al. (2012). Various assumptions must be made in risk assessments in order to extrapolate data from animal or human studies using models to estimate population risks. Such models are designed to be overly protective to account for uncertainty and variability (e.g., upper 95% confidence bound on cancer risk values, and inter- and intraspecies uncertainty factors on reference doses for noncarcinogens). Within this context, some numerical assumptions are based on average or mean values (e.g., average adult body weight and averaging time), and other assumptions are based on upper percentile values such as annualized food consumption rates.

For purposes of risk assessment, average adult body weight may be viewed as an estimate of average lifetime body weight—and averaging time as average lifespan—and people comprising a population. Because these factors are all condensed into a single number, there is often a range of values that it may be reasonable to use. Risk calculations include these factors in the derivation of contaminant thresholds for lifetime cancer risk and risk of adverse effect during a lifetime of exposure for noncancer end points. The resulting threshold values reflect risk across the average lifespan of a population, including men, women (including pregnant women), and children.

Seafood consumption data are collected for different purposes using a variety of survey instruments. It is important to note that

surveys most often reflect short-term intake and do not necessarily address seasonality or otherwise directly capture annualized seafood consumption. The BP risk assessment (FDA 2010a) used National Health and Nutrition Examination Survey (NHANES) 90th percentile consumption data for seafood eaters-only adjusted for consumption frequency (Centers for Disease Control and Prevention 2007). Meal portion and frequency (16.4 seafood meals/month) were converted to annualized daily equivalents.

The selection of 5 years for projected exposure duration following the BP oil spill was considered appropriate and conservative in consideration of the nature of the spilled oil (i.e., light crude), physical conditions (e.g., 29.5°C water temperature), offshore location of the spill (50 miles), and metabolic capacities of seafood species potentially impacted. Exposure duration values selected in previous oil spill assessments range from 2 years (e.g., *New Carissa*, Oregon, 1999) to 10 years (*Exxon Valdez*, Alaska, 1989). Corresponding fishery closures range from weeks to > 6 years for select species that were subjected to prolonged exposures (e.g., farmed salmon, burrowing lobster). In one of the areas most heavily contaminated after the *Exxon Valdez* spill (Windy Bay), benzo[*a*]pyrene equivalents decreased to nondetectable levels by 2.2 years after the contamination event (Bolger et al. 1996). Fisheries closures due to the BP oil spill range from 2 weeks for areas experiencing little to no impact to > 15 months for heavily impacted areas. The reopening of Gulf fisheries based on PAH chemical surveillance results confirmed that selection of a 5-year exposure duration was indeed appropriate and conservative (FDA 2010b; National Oceanic and Atmospheric Administration 2010).

The classification of naphthalene as a noncancer risk in the BP oil spill seafood safety risk assessment was based on current information and concurrence from the U.S. EPA (FDA 2010a). The U.S. EPA (1998b) classified naphthalene in "Group C, a possible human carcinogen. This is based on inadequate data of carcinogenicity in humans exposed to naphthalene via the oral or inhalation routes, and the limited evidence of carcinogenicity in animals via the inhalation route." No oral slope factor or inhalation unit risk estimate were derived for naphthalene by the U.S. EPA because of the weakness of the evidence that naphthalene may be carcinogenic to humans (U.S. EPA 1998b). More recent evaluations have maintained the noncancer risk classification of naphthalene, as evidenced by the 64th meeting of the Food and Agriculture Organization of the United Nations and the World Health Organization (FAO/WHO) Joint Expert Committee on

Food Additives (JECFA), which considered but did not include naphthalene among genotoxic and carcinogenic PAHs evaluated (FAO/WHO 2006).

Although well intentioned, the LOC values suggested by Rotkin-Ellman et al. (2012) do not appear to take into account the natural background occurrence of PAHs in foods in many categories. Assessments from the body of scientific literature is perhaps best represented by the deliberations of JECFA (which included experts from the FDA), which found benzo(*a*)pyrene from dietary intake alone to range from 0.16 to 3.3 µg/person/day (Benford et al. 2010; FAO/WHO 1991, 2006). The LOC values proposed by Rotkin-Ellman et al. (2012) would unnecessarily exclude many food groups from consumers, where nutritional benefits far outweighs negligible risk from PAHs.

Public health authorities are responsible for protecting consumers from contaminated commercial and recreational seafood sources, and to that end advisories may be issued to protect consumers. The federal and state interagency risk assessment for seafood safety following the BP oil spill of 2010 was designed and agreed on by all participants to provide conservative criteria that protect the public. The alternative interpretation provided by Rotkin-Ellman et al. (2012) carries a risk of doing more harm than good.

*The author declares that he has no actual or potential competing financial interest.*

**Robert W. Dickey**

FDA Gulf Coast Seafood Laboratory  
Dauphin Island, Alabama  
E-mail: Robert.Dickey@fda.hhs.gov

## REFERENCES

- Benford D, DiNovi M, Setzer RW. 2010. Application of the margin-of-exposure (MOE) approach to substances in food that are genotoxic and carcinogenic e.g.: benzo[*a*]pyrene and polycyclic aromatic hydrocarbons. *Food Chem Toxicol* 48(suppl 1):S42–S48.
- Bolger M, Henry S, Carrington C. 1996. Hazard and risk assessment of crude oil contaminants in subsistence seafood samples from Prince William Sound. In: *Proceedings of the Exxon Valdez Oil Spill Symposium* (Rice SD, Spiess RB, Wolfe DA, Wright BA, eds). Am Fish Soc Symposium 18:837–843.
- Centers for Disease Control and Prevention. 2007. NHANES 2005–2006. Available: [http://www.cdc.gov/nchs/nhanes/nhanes2005-2006/nhanes05\\_06.htm](http://www.cdc.gov/nchs/nhanes/nhanes2005-2006/nhanes05_06.htm) [accessed 10 October 2011].
- FAO/WHO (Food and Agriculture Organization, World Health Organization) 1991. Polycyclic aromatic hydrocarbons. In: *Toxicological Evaluation of Certain Food Additives and Contaminants*. WHO Food Additive Report Series No. 28. Geneva:World Health Organization, International Programme on Chemical Safety. Available: <http://www.inchem.org/documents/jecfa/jecmono/v28j1e18.htm> [accessed 7 October 2011].
- FAO/WHO (Food and Agriculture Organization of the United Nations, World Health Organization) 2006. Evaluation of Certain Food Contaminants. WHO Technical Report Series 930. Geneva:World Health Organization, International Programme on Chemical Safety. Available: [http://whqlibdoc.who.int/trs/WHO\\_TRS\\_930\\_eng.pdf](http://whqlibdoc.who.int/trs/WHO_TRS_930_eng.pdf) [accessed 7 October 2011].
- FDA (Food and Drug Administration). 2010a. Protocol for Interpretation and Use of Sensory Testing and Analytical Chemistry Results for Re-Opening Oil-Impacted Areas Closed to Seafood Harvesting due to the Deepwater Horizon Oil Spill. Available: <http://www.fda.gov/Food/ucm217601.htm> [accessed 10 October 2011].
- FDA (U.S. Food and Drug Administration). 2010b. Gulf of Mexico Oil Spill Update. Available: <http://www.fda.gov/Food/FoodSafety/Product-SpecificInformation/Seafood/ucm210970> [accessed 10 October 2011].
- National Oceanic and Atmospheric Administration. 2010. Deepwater Horizon/BP Oil Spill Information. Available: [http://sero.nmfs.noaa.gov/deepwater\\_horizon\\_oil\\_spill.htm](http://sero.nmfs.noaa.gov/deepwater_horizon_oil_spill.htm) [accessed 10 October 2011].
- Office of Environmental Health Hazard Assessment. 2010. Benzo[*a*]pyrene. Public Health Goals for Chemicals in Drinking Water. Sacramento, CA:Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. Available: <http://oehha.ca.gov/water/phg/pdf/091610Benzopyrene.pdf> [accessed 26 September 2011].
- Rotkin-Ellman M, Wong KK, Solomon GM. 2012. Seafood contamination after the BP Gulf oil spill and risks to vulnerable populations: a critique of the FDA risk assessment. *Environ Health Perspect* 120:157–161.
- U.S. EPA (U.S. Environmental Protection Agency). 1994. Benzo[*a*]pyrene (BaP) (CASRN50-32-8). Available: <http://www.epa.gov/iris/subst/0136.htm> [accessed 10 June 2010].
- U.S. EPA (U.S. Environmental Protection Agency). 1998a. Draft Water Quality Criteria Methodology Revisions: Human Health, Notice. Fed Reg 63(157):43755–43828. Available: <http://www.epa.gov/fedrgstr/EPA-WATER/1998/August/Day-14/w21517.htm> [accessed 10 October 2011].
- U.S. EPA (U.S. Environmental Protection Agency). 1998b. Naphthalene (CAS No. 91-20-3). Available: <http://www.epa.gov/iris/subst/0436.htm> [accessed 10 June 2010].
- U.S. EPA (U.S. Environmental Protection Agency). 2000. National Guidance: Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories. Volume 2. Risk Assessment and Fish Consumption Limits—Third Edition. EPA 823-B-00-007. Washington, DC:U.S. EPA, Office of Science and Technology. Available: [http://water.epa.gov/scitech/swguidance/fishshellfish/techguidance/risk/volume2\\_index.cfm](http://water.epa.gov/scitech/swguidance/fishshellfish/techguidance/risk/volume2_index.cfm) [accessed 10 October 2011].
- “safety” and is concerned that “unnecessarily conservative risk criteria” could harm human health and society as a whole. This viewpoint, which is arguably rooted more in politics than in science, ignores the long history of chemical assessments where new data and approaches have repeatedly demonstrated significantly greater risks than initially believed (Castorina and Woodruff 2003; Grandjean et al. 2010; Hernberg 2000). Furthermore, from a health-cost perspective, there is considerable benefit to assuring that the population is protected from harmful exposures to carcinogens and other toxicants (Landrigan et al. 2002; Trasande et al. 2006).
- A National Research Council (NRC) committee reviewed the status of environmental regulatory risk assessment and concluded that the new science documenting interindividual variability and the vulnerability of the developing fetus and child to chemical contaminants warrants specific changes to risk assessment practices (NRC 2009). These changes were not reflected in the FDA assessment (FDA 2010). The justifications for the FDA’s risk criteria (FDA 2010) that Dickey provides in his letter do not reflect the most current scientific understanding of the health risks from polycyclic aromatic hydrocarbons (PAHs)—or the risk assessment process—and therefore cannot be characterized as biased “on the side of safety.”
- The NRC, the U.S. Environmental Protection Agency (EPA), and the broader scientific community have recognized that children are not just small adults and that calculation of life-stage-specific doses are the most health protective method to ensure public health protection (American Academy of Pediatrics 2011; NRC 1993, 2009; U.S. EPA 2005). This necessitates use of age-specific body weights and intake and specifically refutes the claim that an adult body weight and dose can represent risk across a lifespan.
- The World Health Organization (WHO) and the U.S. EPA have recognized the extremely skewed nature of food consumption curves and the resulting increased health risk to high-end consumers. These agencies recommended that risk assessments be based on either local surveys (if available) or the 95–97th percentile of national surveys (U.S. EPA 2000; WHO 2008). Previous studies that evaluated the utility of dietary data from the National Health and Nutrition Examination Survey (NHANES) against population-specific surveys have concluded that there is a risk of significantly underestimating exposure among children if NHANES data are the sole source of dietary estimates (Riederer et al. 2010). Furthermore, alternative statistical techniques have been shown to allow better characterization of the

## FDA Risk Assessment of Seafood Contamination after the BP Oil Spill: Rotkin-Ellman and Solomon Respond

<http://dx.doi.org/10.1289/ehp.1104539R>

We thank Dickey for the opportunity to discuss differing approaches to evaluating health risks from chemical contaminants in food, such as those in Gulf seafood after the BP oil spill disaster. As we demonstrate in our commentary, “Seafood Contamination after the BP Gulf Oil Spill and Risks to Vulnerable Populations: A Critique of the FDA Risk Assessment” (Rotkin-Ellman et al. 2012), the choice of parameters and methods can significantly alter the conclusions of a risk assessment, thereby having major impacts on resulting policy decisions. In the example we analyzed, a risk assessment using parameters and methods specifically aimed at protecting vulnerable populations and incorporating the latest risk science differs from the Food and Drug Administration’s (FDA) approach (FDA 2010) by up to four orders of magnitude.

Dickey characterizes chemical risk assessments as inherently biased “on the side of