



## First record of *Fusarium verticillioides* as an entomopathogenic fungus of grasshoppers

SA Pelizza<sup>1,2a\*</sup>, SA Stenglein<sup>3b</sup>, MN Cabello<sup>1c</sup>, MI Dinolfo<sup>3d</sup>, CE Lange<sup>2e</sup>

<sup>1</sup>Instituto de Botánica Carlos Spegazzini, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata

<sup>2</sup>Centro de Estudios Parasitológicos y de Vectores (CEPAVE), CCT-La Plata-CONICET-UNLP, Calle 2 # 584, La Plata (1900), Argentina

<sup>3</sup>Laboratorio de Biología Funcional y Biotecnología (BIOLAB)-CEBB-CONICET, Facultad de Agronomía de Azul, UNCPBA, Republica de Italia # 780, Azul (7300), Argentina

### Abstract

*Fusarium verticillioides* (Saccardo) Nirenberg (Ascomycota: Hypocreales) is the most common fungus reported on infected corn kernels and vegetative tissues, but has not yet been documented as being entomopathogenic for grasshoppers. Grasshoppers and locusts represent a large group of insects that cause economic damage to forage and crops. *Tropidacris collaris* (Stoll) (Orthoptera: Acridoidea: Romaleidae) is a large and voracious grasshopper that in recent years has become an increasingly recurrent and widespread pest in progressively more greatly extended areas of some of in Argentina's northern provinces, with chemical insecticides being currently the only means of control. During February and March of 2008-09, nymphs and adults of *T. collaris* were collected with sweep nets in dense woodland vegetation at a site near Tres Estacas in western Chaco Province, Argentina, and kept in screened cages. *F. verticillioides* was isolated from insects that died within 10 days and was cultured in PGA medium. Pathogenicity tests were conducted and positive results recorded. Using traditional and molecular-biological methods, an isolate of *F. verticillioides* was obtained from *T. collaris*, and its pathogenicity in the laboratory was shown against another harmful grasshopper, *Ronderosia bergi* (Stål) (Acridoidea: Acrididae: Melanoplinae). The mortality caused by *F. verticillioides* on *R. bergi* reached  $58 \pm 6.53\%$  by 10 days after inoculation. This is the first record of natural infection caused by *F. verticillioides* in grasshoppers.

**Keywords:** Acrididae, Orthoptera, *Ronderosia bergi*, *Tropidacris collaris*, Chaco Province, Argentina

**Correspondence:** <sup>a\*</sup> [pelizza@cepave.edu.ar](mailto:pelizza@cepave.edu.ar), <sup>b</sup> [stenglein@faa.unicen.edu.ar](mailto:stenglein@faa.unicen.edu.ar), <sup>c</sup> [mcabello@netverk.com.ar](mailto:mcabello@netverk.com.ar),

<sup>d</sup> [ines\\_dinolfo@hotmail.com](mailto:ines_dinolfo@hotmail.com), <sup>e</sup> [carlosl@cepave.edu.ar](mailto:carlosl@cepave.edu.ar), \*Corresponding author

**Received:** 15 April 2010, **Accepted:** 6 October 2010

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 11, Number 70

#### Cite this paper as:

Pelizza SA, Stenglein SA, Cabello MN, Dinolfo MI, Lang CE. 2011. First record of *Fusarium verticillioides* as an entomopathogenic fungus of grasshoppers. *Journal of Insect Science* 11:70 available online: [insectscience.org/11.70](http://insectscience.org/11.70)

## Introduction

The genus *Fusarium* comprises a large group of species of filamentous fungi widely distributed in soil usually in association with plants. Most species are saprotrophic and relatively abundant members of the soil microbiota (Leslie and Summerell 2006). Many *Fusarium* species are well-known as pathogens of plants, insects, and humans (Majumbar et al. 2008), although there are *Fusarium* species that are insect pathogens but are not pathogenic to the plant (Kuruville and Jacob 1979a, 1979b, 1980). More than 13 *Fusarium* species are pathogenic to insects, and the genus has a host range that includes Coleoptera, Diptera, Hemiptera, Hymenoptera and Lepidoptera (Teotor-Barsch and Roberts 1983; Humber 1992). *Fusarium* includes various species/strains that are able to produce potent secondary metabolites, such as trichothecenes (Kilpatrick 1961; Kuno and Ferrer 1973), fumonisins (Kuruville and Jacob 1979a) and beauvericin (Kuruville and Jacob 1979b, 1980; Gupta et al. 1991). The latter is a widespread metabolite among entomopathogenic fungi, such as *Beauveria bassiana* and *Paecilomyces fumosoroseus* (Leath and Newton 1969; Loke et al. 1970). *Fusarium verticillioides* (Saccardo) Nirenberg (Ascomycota: Hypocreales) is often the most common fungus reported from infected corn kernels and vegetative tissues (Foley 1962; Kommedahl and Windels 1981; Nelson et al. 1993; Kedera et al. 1999; Desjardins et al. 2000). In Argentina, its presence has been registered in corn (Peiretti-Uzal et al. 2007) but has not been recorded as an entomopathogen. In France *F. verticillioides* (as *Fusarium moniliforme*) was isolated from *Ostrinia nubilalis* (Vago 1958) and *Bombyx mori* (Vago and Nicot 1954). *Fusarium acridiorum* (Thabut) (= *Trichothecium acridiorum*) is a cuticular parasite of the

desert locust *Schistocerca gregaria* (Akbar et al. 1958), which constitutes the only previous record of a species of *Fusarium* isolated from the Acridoidea.

As in other regions of the world, grasshoppers and locusts are important agricultural pests in different parts of Argentina. Damage to forage and a variety of crops have been reported from at least 15 of the 201 species known for the country (Lange et al. 2005; Carbonell et al. 2006). Eight of the harmful species are melanoplines (Acrididae, Melanoplineae) and three romaleids (Romaleidae: Romaleinae). The romaleid *Tropidacris collaris* (Stoll) (Orthoptera: Acridoidea: Romaleidae) has become in recent years an increasingly recurrent and extended pest in some of the northern provinces, particularly in parts of Córdoba, Santiago del Estero, and Chaco. *Tropidacris collaris*, one of the largest grasshoppers known ( $\sigma^7 = 73\text{-}101$  mm,  $\text{f} = 92\text{-}126$  mm), is strongly gregarious during juvenile development and it is voracious. Although adults tend to prefer hard-leaf trees and bushes, *T. collaris* is actually a polyphagous species (Barrera and Paganini 1975; Carbonell 1986). The bands of nymphs consume virtually all available plant material. Currently, chemical insecticides are the only mean of control. Pathogens have not been reported, and *Paranosema locustae*, a microsporidium developed in the USA as a biocontrol agent of grasshoppers that was introduced and became established in some other areas of the country (Lange and Azzaro 2008), did not produce encouraging results when tested against *T. collaris* under laboratory conditions (Lange et al. 2008).

Here we report obtaining, by traditional and molecular methods, an isolate of *F. verticillioides* in *T. collaris*, and its pathogenicity in the laboratory (under

controlled conditions) against another harmful grasshopper, the melanopline *Ronderosia bergi* (Stål).

## Materials and Methods

During February and March of 2008-09, nymphs and adults of *T. collaris* were collected with sweep nets in dense woodland vegetation at a site (27° 8' 21.9" S; 61° 34' 23.8" W) near Tres Estacas in western Chaco Province, Argentina. The area is within the Chaqueña biogeographic province (Cabrera and Willink 1973), where the annual rainfall is 600 mm and the average temperatures range between 19 and 21 °C with summer maxima reaching 42-46 °C. The samples were immediately taken to the laboratory where the grasshoppers were kept in groups in wire-screened cages in a rearing room under controlled conditions (30 °C, 14:10 light-dark photoperiod, 60% relative humidity). This setting usually favors the expression of entomopathogenic fungi present in field-collected, infected insects (Shah et al. 1997).

The grasshoppers that died within 10 days after collection were superficially sterilized by placing the specimens in 70% ethanol for a few seconds, then washed in sterile distilled water, followed by 0.5% sodium hypochlorite for 1 min, and rinsed again in sterile distilled water according to Lacey and Brooks (1997). They were then placed in a sterile culture chamber consisting of a Petri dish (150 mm diameter) with a filter-paper disk that was periodically moistened with sterile distilled water and incubated at 25° C in the dark. Daily checks were performed during the first 5 days post-mortem. The filamentous fungi emerging from the dead individuals (Figure 1) were transferred to Petri dishes containing potato-dextrose agar (PDA) + antibiotics, and incubated (26° C). The fungal species isolated



**Figure 1.** Adult *Tropidacris collaris* 48 h after its death caused by *Fusarium verticillioides*. Scale bar: 14 mm. High quality figures are available online

from *T. collaris* were identified on the basis of the macromorphological appearance of the colonies—such as color, diameter, mycelial texture—and their micromorphological characteristics were observed under phase-contrast microscopy. Additionally, a specific PCR was performed to confirm the fungal species. For DNA extraction the isolate was grown on PDA medium at  $25 \pm 2$  °C under a 12 h light-dark photoperiod and 6 day-old cultures were quantitatively scraped from the surfaces of three Petri dishes with a scalpel, frozen in liquid nitrogen, and then ground into a fine powder in a mortar. Genomic DNA was extracted by the so-called CTAB method described by Stenglein and Balatti (2006). A *F. verticillioides*-specific PCR was performed with primers 5'-GTCAGAATCCATGCCA GAACG-3'-forward and 5'- CACCCGCAGC AATCCATCAG-3'- reverse (Patiño et al. 2004). The amplification was carried out in a 25- $\mu$ l final volume containing 12-15 ng of genomic DNA, 10X reaction buffer (2 mM Tris-HCl pH 8.0, 10 mM KCl, 0.01 mM EDTA, 1mM DTT, 50% [v/v] glycerol, 0.5% [v/v] Tween 20, 0.5% [v/v] Nonidet P40), 0.5  $\mu$ M of each primer, 200  $\mu$ M of each dNTP (Genbiotech S.R.L.), 2.5 mM MgCl<sub>2</sub>, and 1.25 units of Taq DNA polymerase (Genbiotech, [www.genbiotech.com](http://www.genbiotech.com)). DNA amplification was performed in an XP thermal cycler (Bioer Technology Co., [www.bioer.com.cn](http://www.bioer.com.cn)) with an initial denaturing step at 95° C for 2 min; followed by 29 cycles at 95° C for 30 s, 54° C for 35 s, and 72° C for

45 s; and a final extension cycle at 72° C for 2 min. The specific product of 800 bp was examined by electrophoresis in 1.5% (w/v) agarose gels containing GelRed™ (Genbiotech) at 80 V in 5X Trisborate-EDTA buffer for 3-4 h at room temperature. The fragment was visualized under ultraviolet light. The size of the DNA fragment was estimated by comparing the DNA band with a 100-bp DNA ladder (Genbiotech).

The fungal isolate was deposited at the Fungal Culture Collection of the Spegazzini Institute of Botany as LPS 1057. The viability of the conidial fungi was determined after 24 hours by means of the techniques described by Lane et al. (1988). The germination test was repeated for each stock suspension to increase the accuracy of the viability assessments.

Our efforts to develop a breeding colony of *T. collaris* were unfortunately not successful (Lange et al. 2008). Thus, the insects used for testing pathogenicity in this study were *R. bergi*, bred in a grasshopper colony at CEPAVE.

Three replicates (on different dates) of 50 third-instar nymphs each of healthy *R. bergi* were sprayed in groups of 10 with 1,000 µl of a suspension of  $2.8 \times 10^6$  conidia/ml (in 0.01% [v/v] Tween 20) according to Majumbar et al. (2008). The conidia had been harvested after cultivation on PGA for 10 days at 25 °C in the dark. Three additional replicates of 20 grasshoppers each (in two groups of 10) were sprayed with 1,000 µl of 0.01% [v/v] Tween 20 for use as controls. The grasshoppers were maintained in groups of 10 in acetate tubes (50 x 9 cm) after (Henry 1985) and fed with lettuce (*Lactuca sativa* L.) leaves. Treated and control insects were maintained at 30 °C, 60% relative humidity, and a 14:10-h light-dark photoperiod. Cumulative mortality was

recorded daily for 10 days. Dead grasshoppers were removed and immediately deposited in high-humidity chambers (sterile Petri dishes with filter paper dampened with sterile distilled water). Mycosis was confirmed by microscopical examination of the dead grasshoppers.


## Results and Discussion

The isolated fungus was morphologically identified as *F. verticillioides* on carnation leaf-piece agar and on PDA according to Nelson et al. (1983) and Leslie and Summerell (2006). Their characteristics on the former agar were: macroconidia, formed in pale-orange sporodochia, slightly falcate or else straight with thin walls, and the basal cell foot-shaped, 3-5 septate, 25-55 x 2-4 µm (Figure 2a); microconidia, formed in chains on monophialides, oval to club-shaped with a flattened base, usually 0 septate, 6-10 x 1.5-2.5 µm (Figure, 2b, c). Chlamydo-spores were not found. Characters on PDA were: white aerial mycelium with rapid growth, violet pigmentation with age; lower surface with violet to dark-violet pigmentation.

The isolate determined to be *F. verticillioides* on the basis of morphological characteristics produced a PCR-amplified fragment of 800-bp identical to that observed by Patiño et al. (2004).

The average viability of *F. verticillioides* conidia was 95%. The mortality caused by *F. verticillioides* on *R. bergi* reached  $58 \pm 6.53\%$  by 10 days after inoculation. Phialides and conidia were observed growing out of the dead hosts (Figure 3). No mortality occurred among the controls.

Majumdar et al. (2008) obtained higher mortality rates in laboratory tests reaching



**Figure 2.** (a) Macro- and microconidia of *Fusarium verticillioides* (b and c) Phialides and microconidia in chain and grouped in heads. Scale bar: (a) 10  $\mu\text{m}$ ; (b and c) 30  $\mu\text{m}$ . High quality figures are available online.

80% mortality with the pupae of *Tetanops myopaeformis* 10 days after applying the same concentration ( $2.8 \times 10^6$ ) of *Fusarium solani* conidia. An even greater mortality was observed by Golpalakrishnan and Narayan (1989), who reported 100% in the guava-shield scale, *Pulvinaria* (formerly known as *Chloropulvinaria*) *psidii* Maskel, (Hemiptera: Coccidae) at 5 days after treatment with  $4.8 \times 10^8$  conidia/ml of *Fusarium oxysporum*. Moreover, a 100% mortality was obtained within three days in a field-cage test with *F. oxysporum* against *Nilaparvata lugens* (Kuruville and Jacob 1979a, 1980).

The means by which the infection of *T. collaris* and *R. bergi* by *F. verticillioides* might have been effected is not clear, but Kilpatrick (1961) stated that the entry of the fungus into the insect could occur via the oral route, oviposition tubes or wounds.



**Figure 3.** *Fusarium verticillioides* on third-instar nymphs of *Ronderosia bergi* 48 h after death. Scale bar: 14 mm. High quality figures are available online.

Relatively low temperatures and high moisture levels have been shown to be suitable for the development of *Fusarium* epizootics in other insects (Villacaros and Robin 1989; Venugopal et al. 1989; Pandit and Tarannum 2002; Majumbar et al. 2008). Such conditions do not normally occur in the area sampled by us where the temperatures are much higher (42-46° C) and the rainfall lower (600 mm annually).

At present we do not know if *T. collaris* would be affected by the fungus in the field. However, such possibility should not be ruled out. Although low humidity and elevated temperature are usually unfavorable for most fungi, this particular isolate was recovered living under the conditions affecting the host. Perhaps the fungus is one of the factors preventing further spread of the host into wetter, cooler regions.

In conclusion, the present study provides the first report of *F. verticillioides* as an entomopathogenic fungus in grasshoppers and extends the knowledge of the pathogenic capacity of this *Fusarium* species to that

orthopteran. Future research efforts should investigate the possible existence of mycotoxin production against humans and the potential effects of the *F. verticillioides* isolate LPS 1057 on nontarget fauna. Moreover, future pathogenic studies may clarify whether or not this isolate is capable of infecting plants and ascertain by molecular phylogenetic analyses if this isolate has an identical, similar, or somewhat altered germ-line-genetic lineage to that of the *F. verticillioides* plant-pathogens.

### Acknowledgements

This study was partially supported by Agencia Nacional de Promoción Científica y Técnica (PICT N° 914, PRH32/PICT110), CONICET, CICPBA, and Universidad Nacional de La Plata (11/N 527). M.Cabello and C. Lange are members of Carrera del Investigador CICPBA. The authors wish to thank Dr. Donald F. Haggerty, retired, for editing the final version of the manuscript.

### References

Akbar K, Haque H, Abbas HM. 1958. *Fusarium acridiorum*, a parasite of desert locust. *FAO Plant Protection Bulletin* 6:59.

Barrera M, Paganini HI. 1975. Acridios de Tucumán: notas bioecológicas. *Acta Zoológica Lilloana* 31: 107-123.

Cabrera AL, Willink A. 1973. *Biogeografía de America Latina*. Monografía 13, Serie de Biología Washington, DC: OEA.

Carbonell CS. 1986. Revision of the Neotropical genus *Tropidacris* (Orthoptera, Acridoidea, Romaleidae, Romaleinae). *Proceedings of the Academy of Natural Sciences of Philadelphia* 138: 366-402.

Carbonell C, Cigliano MM, Lange CE. 2006. *Acridomorph (Orthoptera) species of Argentina and Uruguay*. CD-ROM. Publications on Orthopteran Diversity, The Orthopterists Society at the "Museo de La Plata", Argentina.

Desjardins AE, Plattner RD, Gordon TR. 2000. *Gibberella fujikuroi* mating population A and *Fusarium subglutinans* from teosinte species and maize from Mexico and Central America. *Mycological Research* 104: 865-872.

Foley DC. 1962. Systemic infection of corn by *Fusarium moniliforme*. *Phytopathology* 52: 870-872.

Gopalakrishnan C, Narayanan K. 1989. Occurrence of *Fusarium oxysporum* Schlecht and its pathogenicity on guava scale *Chloropulvinaria psidii* Maskell (Hemiptera: Coccidae). *Current Science* 58: 92-93.

Gupta S, Krasnoff SB, Underwood NL, Renwick JAA, Roberts DW. 1991. Isolation of beauvericin as an insect toxin from *Fusarium semitectum* and *Fusarium moniliforme* var. *subglutinans*. *Mycopathologia* 115: 185-189.

Henry JE. 1985. *Melanoplus* spp. In: Singh P, Moore RF, Editors. *Handbook of Insect Rearing*, vol. 1, pp. 451-464. Elsevier.

Humber RA. 1992. *Collection of entomopathogenic fungal cultures: Catalog of strains*, U.S. Department of Agriculture. Agricultural Research Service. Bulletin ARS-110.

Kedera CJ, Plattner RD, Desjardins AE. 1999. Incidence of *Fusarium* sp. and levels of

fumonisin B1 in maize in western Kenya. *Applied and Environmental Microbiology* 64: 41-44.

Kilpatrick RA. 1961. Fungi associated with larvae of *Sitona spp.* *Phytopathology* 51: 640-641.

Kommedahl T, Windels CE. 1981. Root, stalk and ear infecting *Fusarium* species on corn in the USA. In: Nelson PE, Toussoun TA, Cook RJ, Editors. *Fusarium: diseases, biology and taxonomy*, pp. 94-103. Pennsylvania State University Press.

Kuno G, Ferrer MAC. 1973. Pathogenicity of two *Fusarium* fungi to an armoured scale insect *Selenaspidus articulatus*. *Journal of Invertebrate Pathology* 22: 473-474.

Kuruvilla S, Jacob A. 1979a. Comparative susceptibility of nymphs and adults of *Nilaparvata lugens* to *Fusarium oxysporum* and its use in microbial control. *Agricultural Research Journal Kerala* 17: 287-288.

Kuruvilla S, Jacob A. 1979b. Host range of the entomogenous fungus *Fusarium oxysporum* Schlecht and its safety to three crop plants. *Current Science* 48: 603.

Kuruvilla S, Jacob A. 1980. Studies on *Fusarium oxysporum* Schlecht infecting rice brown plant hopper, *Nilaparvata lugens* Stal. *Agricultural Research Journal Kerala* 18: 51-54

Lacey L, Brooks WM. 1997. Initial handling and diagnosis of diseased insects. In: Lacey LA, Editor. *Manual of techniques in insect pathology*. Academic Press.

Lane BS, Humphreys BSAM, Thompson K, Trinci APJ. 1988. ATP content of stored

spores of *Paecilomyces farinosus* and the use of ATP as criterion of spore viability. *Transactions of the British Mycological Society* 90: 109-111.

Lange CE, Cigliano MM, De Wysiecki ML. 2005. Los acridoideos (Orthoptera: Acridoidea) de importancia económica en la Argentina. In: Barrientos Lozano L, Almaguer Sierra P, Editors. *Manejo integrado de la langosta centroamericana (*Schistocerca piceifrons piceifrons*, Walker) y acridoideos plaga en América Latina*. pp. 93-135. Instituto Tecnológico de Ciudad Victoria.

Lange CE, Azzaro FG. 2008. New case of long-term persistence of *Paranosema locustae* (Microsporidia) in melanopline grasshoppers (Orthoptera: Acrididae) of Argentina. *Journal of Invertebrate Pathology* 99: 357-359.

Lange CE, Bardi C, Plischuk S. 2008. Inefectividad del agente de biocontrol *Paranosema locustae* (Microsporidia) para la "tucura quebrachera", *Tropidacris collaris* (Orthoptera: Romaleidae), en condiciones de laboratorio. *Revista de la Sociedad Entomologica Argentina* 67: 151-155.

Leath TK, Newton RC. 1969. Interaction of a fungus gnat *Bradysia* sp. (Sciaridae) with *Fusarium* spp. on alfalfa and red clover. *Phytopathology* 59: 257-258.

Leslie JF, Summerell BA. 2006. *Fusarium verticillioides* (Saccardo) Nirenberg. In: Leslie JF, Summerell BA, Editors. *The Fusarium Laboratory Manual*, pp. 274-279. Blackwell Publishing.

Loke TK, Norris DM, Chu HM. 1970. Sterol metabolism as a basis for a mutualistic symbiosis. *Nature* 225: 661-662.

- Majumbar A, Boetel MA, Jaronski TS. 2008. Discovery of *Fusarium solani* as a naturally occurring pathogen of sugarbeet root maggot (Diptera: Ulidiidae) pupae: Prevalence and baseline susceptibility. *Journal of Invertebrate Pathology* 97: 1-8.
- Nelson PE, Toussoun TA, Marasas WFO. 1983. *An Illustrated Manual for Identification. Fusarium Species*. Pennsylvania State University Press University Park.
- Nelson PE, Desjardins AE, Plattner RD. 1993. Fumonisin, mycotoxins produced by *Fusarium* species: Biology, chemistry, and significance. *Annual Review of Phytopathology* 31: 252.
- Pandit NC, Tarannum Z. 2002. Record of fungi associated with diseased insects of jute and mesta crop. *Journal of Entomological Research* 26: 175–178.
- Patiño B, Mirete S, Gonzalez-Jaen T, Mule G, Vasquez C. 2004. PCR detection assay for fumonisin producing *Fusarium verticillioides* strains. *Journal of Food Protection* 67: 1278-1283.
- Peiretti-Uzal DA, Nazar-Lovera MC, Biasutti-Valenzano CA, Giorda-Lerda LM. 2007. Susceptibilidad a *Fusarium verticillioides* (Sacc.) Nirenberg en la población de maíz MPB-FCA 856. *Agronomía Mesoamericana* 18: 171-176.
- Shah PA, Kooyman C, Paraiso A. 1997. Surveys for fungal pathogens of locust and grasshoppers in Africa and near east. In: Goettel M, Johnson D, Editors. *Microbial control of grasshoppers and locusts*, Memoirs of the Entomological Society of Canada. 171: 27-35.
- Stenglein S, Balatti P. 2006. Genetic diversity of *Phaeoisariopsis griseola* in Argentina as revealed by pathogenic and molecular markers. *Physiological and Molecular Plant Pathology* 68: 158-67.
- Teetor-Barsch GH, Roberts W.D. 1983. *Fusarium Species Pathogens of insects*. Review. *Mycopathologia* 84: 3-16.
- Vago C, Nicot J. 1954. Contribution à l'écopathologie compare des mycoses a propos d' une nouvelle affection du Lépidoptère *Bombyx mori* L. *Revue Mycologie Paris* 19: 255-261.
- Vago C. 1958. Virulence cryptogamique simultanee vis-avis d' un végétal et d' un insecte. *Comptes rendus L' académie des sciences* 247: 1651-1653.
- Venugopal R, Reddy AS, Rao KT. 1989. Natural enemies of cotton whitefly, *Bemisia tabaci* Gennadius, in relation to host populations and weather factors. *Journal Biological Control* 3: 10–12.
- Villacaros LT, Robin RP. 1989. Entomogenous fungi infecting *Heteropsylla cubana* Crawford (Homoptera: Psyllidae) in Leyte, Philippines. *Tropical Pest Management* 35: 120-122.